

Chapter 5: Bayes' Theorem

Meike Niederhausen and Nicky Wakim

2023-10-04

Table of contents

- Learning Objectives
- Introduction
- Bayes' Rule for two events
- Calculating probability with Higher Order Multiplication Rule
- Calculating probability with Law of Total Probability
- General Law of Total Probability
- Calculating probability with generalized Law of Total Probability
- Let's revisit the color-blind example
- Calculate probability with both rules
- Bayes' Rule

Learning Objectives

1. Calculate conditional probability of an event using Bayes' Theorem
2. Utilize additional probability rules in probability calculations, specifically the Higher Order Multiplication Rule and the Law of Total Probabilities

Introduction

- So we learned about conditional probabilities
 - We learned how the occurrence of event A affects event B (B conditional on A)
- Can we figure out information on how the occurrence of event B affects event A?
- We can use the conditional probability ($\mathbb{P}(A|B)$) to get information on the flipped conditional probability ($\mathbb{P}(B|A)$)

Bayes' Rule for two events

Theorem: Bayes' Rule (for two events)

For any two events A and B with nonzero probabilities,

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A) \cdot \mathbb{P}(B|A)}{\mathbb{P}(B)}$$

Calculating probability with Higher Order Multiplication Rule

Example 1

Suppose we draw 5 cards from a standard shuffled deck of 52 cards. What is the probability of a flush, that is all the cards are of the same suit (including straight flushes)?

Higher Order Multiplication Rule

$$\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2|A_1) \cdot \mathbb{P}(A_3|A_1 A_2) \dots \cdot \mathbb{P}(A_n|A_1 A_2 \dots A_{n-1})$$

Calculating probability with Law of Total Probability

Example 2

Suppose 1% of people assigned female at birth (AFAB) and 5% of people assigned male at birth (AMAB) are color-blind. Assume person born is equally likely AFAB or AMAB (not including intersex). What is the probability that a person chosen at random is color-blind?

Law of Total Probability for 2 Events

For events A and B ,

$$\begin{aligned}\mathbb{P}(B) &= \mathbb{P}(B \cap A) + \mathbb{P}(B \cap A^C) \\ &= \mathbb{P}(B|A) \cdot \mathbb{P}(A) + \mathbb{P}(B|A^C) \cdot \mathbb{P}(A^C)\end{aligned}$$

General Law of Total Probability

Law of Total Probability (general)

If $\{A_i\}_{i=1}^n = \{A_1, A_2, \dots, A_n\}$ form a partition of the sample space, then for event B ,

$$\begin{aligned}\mathbb{P}(B) &= \sum_{i=1}^n \mathbb{P}(B \cap A_i) \\ &= \sum_{i=1}^n \mathbb{P}(B|A_i) \cdot \mathbb{P}(A_i)\end{aligned}$$

Calculating probability with generalized Law of Total Probability

Example 3

Individuals are diagnosed with a particular type of cancer that can take on three different disease forms,* D_1 , D_2 , and D_3 . It is known that amongst people diagnosed with this particular type of cancer,

- 20% of people will eventually be diagnosed with form D_1 ,
- 30% with form D_2 , and
- 50% with form D_3 .

The probability of requiring chemotherapy (C) differs among the three forms of disease:

- 80% with D_1 ,
- 30% with D_2 , and
- 10% with D_3 .

Based solely on the preliminary test of being diagnosed with the cancer, what is the probability of requiring chemotherapy (the event C)?

Let's revisit the color-blind example

Example 4

Recall the color-blind example (Example 2), where

- a person is AMAB with probability 0.5,
- AMAB people are color-blind with probability 0.05, and
- all people are color-blind with probability 0.03.

Assuming people are AMAB or AFAB, find the probability that a color-blind person is AMAB.

Calculate probability with both rules

Example 5

Suppose

- 1% of women aged 40-50 years have breast cancer,
- a woman with breast cancer has a 90% chance of a positive test from a mammogram, and
- a woman has a 10% chance of a false-positive result from a mammogram.

What is the probability that a woman has breast cancer given that she just had a positive test?

Bayes' Rule

Theorem: Bayes' Rule

If $\{A_i\}_{i=1}^n$ form a partition of the sample space S , with $\mathbb{P}(A_i) > 0$ for $i = 1 \dots n$ and $\mathbb{P}(B) > 0$, then

$$\mathbb{P}(A_j|B) = \frac{\mathbb{P}(B|A_j) \cdot \mathbb{P}(A_j)}{\sum_{i=1}^n \mathbb{P}(B|A_i) \cdot \mathbb{P}(A_i)}$$

