

Chapter 25: Joint densities

Meike Niederhausen and Nicky Wakim

2023-11-08

Table of contents

- Learning Objectives
- Double Integrals Mini Lesson (1/3)
- Double Integrals Mini Lesson (2/3)
- Double Integrals Mini Lesson (3/3)
- How to define the joint pdf for continuous RVs?
- Important properties of the joint pdf
- What is the joint cumulative distribution function?
- What are the marginal pdf's?
- Example of joint pdf
- Example of joint pdf
- Example of a *more complicated* joint pdf
- Example of a *more complicated* joint pdf
- Let's complicate this even more!
- Let's complicate this even more!
- Let's complicate this even more!
- Let's complicate this even further!

Learning Objectives

1. Solve double integrals in our mini lesson!
2. Calculate probabilities for a pair of continuous random variables
3. Calculate a *joint and marginal* probability density function (pdf)
4. Calculate a *joint and marginal* cumulative distribution function (CDF) from a pdf

Double Integrals Mini Lesson (1/3)

Mini Lesson Example 1

Solve the following integral:

$$\int_2^3 \int_0^1 xy dy dx$$

Double Integrals Mini Lesson (2/3)

Mini Lesson Example 2

Solve the following integral:

$$\int_2^3 \int_0^1 (x + y) dy dx$$

Double Integrals Mini Lesson (3/3)

Do this problem at home for extra practice. The solution is available in Meike's video!

Mini Lesson Example 3

Solve the following integral:

$$\int_2^3 \int_0^1 e^{x+y} dy dx$$

How to define the joint pdf for continuous RVs?

For a single continuous RV X is a function $f_X(x)$, such that for all real values a, b with $a \leq b$,

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f_X(x) dx$$

For two continuous RVs (X and Y), we can define the **joint pdf**, $f_{X,Y}(x, y)$, such that for all real values a, b, c, d with $a \leq b$ and $c \leq d$,

$$\mathbb{P}(a \leq X \leq b, c \leq Y \leq d) = \int_a^b \int_c^d f_{X,Y}(x, y) dy dx$$

Important properties of the joint pdf

1. Note that $f_{X,Y}(x, y) \neq \mathbb{P}(X = x, Y = y)!!!$
2. In order for $f_{X,Y}(x, y)$ to be a pdf, it needs to satisfy the properties

- $f_{X,Y}(x, y) \geq 0$ for all x, y

- $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \underline{f_{X,Y}(x, y)} dx dy = \underline{1}$

What is the joint cumulative distribution function?

Definition: Joint cumulative distribution function (Joint CDF)

The **joint cumulative distribution function (cdf)** of continuous random variables X and Y , is the function $F_{X,Y}(x, y)$, such that for all real values of x and y ,

$$F_{X,Y}(x, y) = \mathbb{P}(X \leq x, Y \leq y) = \int_{-\infty}^x \int_{-\infty}^y \underline{f_{X,Y}(s, t)} dt ds$$

Remarks:

- The definition above for $F_{X,Y}(x, y)$ is a **function** of x and y .
- The joint cdf at the point (a, b) , is

$$F_{X,Y}(a, b) = \mathbb{P}(X \leq a, Y \leq b) = \int_{-\infty}^a \int_{-\infty}^b f_{X,Y}(s, t) dt ds$$

What are the marginal pdf's?

Definition: Marginal pdf's

Suppose X and Y are continuous r.v.'s, with joint pdf $f_{X,Y}(x, y)$. Then the **marginal probability density functions** are

$$\begin{aligned}\underline{f_X(x)} &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) \underline{dy} \\ f_Y(y) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx\end{aligned}$$

int over all y

Example of joint pdf

Example 1.1




Let $f_{X,Y}(x,y) = \frac{3}{2}y^2$, for

$\rightarrow 0 \leq x \leq 2, 0 \leq y \leq 1$.

1. Find

$\rightarrow \mathbb{P}(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2})$.

① & ②



$$\mathbb{P}(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2})$$

$$= \int_0^1 \int_0^{\frac{1}{2}} \frac{3}{2} y^2 dy dx$$

$$= \int_0^1 \left[\frac{1}{2} y^3 \right]_{y=0}^{y=\frac{1}{2}} dx$$

$$= \int_0^1 \left[\frac{1}{2} \left(\frac{1}{2} \right)^3 - \frac{1}{2} (0)^3 \right] dx$$

$$= \int_0^1 \frac{1}{16} dx = \left[\frac{1}{16} x \right]_{x=0}^{x=1}$$

$$= \frac{1}{16} (1) - \frac{1}{16} (0) = \frac{1}{16}$$

$$\mathbb{P}(0 \leq X \leq 1, 0 \leq Y \leq \frac{1}{2}) = \frac{1}{16}$$

steps in problem

① set up domain of pdf

② shade in area of probability of interest

③ set up integral:
 $dy dx$ or $dx dy$?

domain of y and domain of x don't depend on each other & pdf is only in terms of y


$$f_{\mathbf{x}}(\mathbf{x}) = \exp \left(\underline{\hspace{2cm}} \right)$$

Example of joint pdf

Example 1.2

Let $f_{X,Y}(x,y) = \frac{3}{2}y^2$, for
 $0 \leq x \leq 2, 0 \leq y \leq 1$.

2. Find $f_X(x)$ and $f_Y(y)$.



$f_X(x)$: int out y $f_Y(y)$: int out x

$$\begin{aligned} f_X(x) &= \int_0^1 \frac{3}{2} y^2 dy = \left. \frac{1}{2} y^3 \right|_{y=0}^{y=1} \\ &= \frac{1}{2}(1)^3 - \frac{1}{2}(0)^3 \\ &= \frac{1}{2} \end{aligned}$$

$$f_X(x) = \frac{1}{2} \quad \text{for } 0 \leq x \leq 2$$

$$\begin{aligned} f_Y(y) &= \int_0^2 \frac{3}{2} y^2 dx = \left. \frac{3}{2} y^2 x \right|_{x=0}^{x=2} \\ &= \frac{3}{2} y^2 (2) - \frac{3}{2} y^2 (0) \end{aligned}$$

$$f_Y(y) = 3y^2 \quad \text{for } 0 \leq y \leq 1$$

CDF
 $P(y_1 \leq Y \leq y_2)$

$$\rightarrow \left[\underline{P(Y \leq y)} = \int_{-\infty}^y 3t^2 dt \right] \rightarrow F_Y(y)$$

Example of a *more complicated* joint pdf

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 2.1

Let $f_{X,Y}(x, y) = 2e^{-(x+y)}$, for
 $0 \leq x \leq y$.

1. Find $f_X(x)$ and $f_Y(y)$.

Example of a *more complicated* joint pdf

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 2.2

Let $f_{X,Y}(x, y) = 2e^{-(x+y)}$, for
 $0 \leq x \leq y$.

2. Find $\mathbb{P}(Y < 3)$.

Let's complicate this even more!

$$f_{X,Y}(x,y) = \frac{1}{16} \text{ for } 0 \leq X \leq 4, 0 \leq Y \leq 4$$

② shade in prob $P(|X-Y| < 2)$

$$-2 < x - y < 2$$

$$\begin{array}{r} -2 < x - y \\ + y \quad + y \\ \hline -2 < x + 2 \end{array}$$

$$y < x + 2$$

$$\begin{array}{r} x - y < 2 \\ -2 < -y \\ + y \quad + y \\ \hline x - 2 < y \end{array}$$

$$x - 2 < y$$

$$4 \times 4 \cdot b = 1$$

Example 3.1

Let X and Y have constant density on the square

$$0 \leq X \leq 4, 0 \leq Y \leq 4.$$

1. Find $P(|X - Y| < 2)$


③ set up integral for probability

$$P(|X - Y| < 2)$$

$$= \int_0^2 \int_0^{x+2} \frac{1}{16} dy dx + \int_2^4 \int_{x-2}^4 \frac{1}{16} dy dx$$

$$= \int_0^2 \left[\frac{1}{16} y \right]_{y=0}^{y=x+2} dx + \int_2^4 \left[\frac{1}{16} y \right]_{y=x-2}^{y=4} dx$$


$$= \int_0^2 \frac{1}{16} (x+2) dx + \int_2^4 \left[\frac{1}{16} 4 - \frac{1}{16} (x-2) \right] dx$$



$$= \int_0^2 \frac{1}{16}(x+2)dx + \int_2^4 \left(\frac{1}{4} - \frac{1}{16}(x-2) \right) dx$$


$$= \left[\frac{1}{32}x^2 + \frac{1}{8}x \right]_{x=0}^{x=2} + \left[\frac{1}{4}x - \frac{1}{32}x^2 - \frac{1}{8}x \right]_{x=2}^{x=4}$$

solve this @ home




$$\int_0^2 \int_0^{y+2} \frac{1}{16} dx dy + \int_2^4 \int_{y-2}^4 \frac{1}{16} dx dy$$

ALT ALT ROUTE



$$\frac{1}{2}(2)(2) = 2$$

$$f_{X,Y}(x,y) = \frac{1}{16} \rightarrow (4 \cdot 4) - 2 - 2 = 12$$



$$P(|X-Y| < 2)$$

$$= \frac{12}{16} = \frac{3}{4}$$

Let's complicate this even more!

$$Z = \frac{X+Y}{2} \quad \frac{1+2}{2+1}$$

• M is a transformation of X & Y

• CDF method: start by finding the CDF (which includes probabilities) & then find pdf from


CDF : $F_M(m) = \int f_M(m) dm \rightarrow f_M(m) = \frac{d}{dm} F_M(m)$

$$F_M(m) = P(M \leq m) = P(\max(X, Y) \leq m) \\ = P(X \leq m, Y \leq m)$$

Example 3.1

Let X and Y have constant density on the square $0 \leq X \leq 4, 0 \leq Y \leq 4$.

2. Let $M = \max(X, Y)$. Find the pdf for M , that is $f_M(m)$



Let's complicate this even more!

Do this problem at home for extra practice. The solution is available in Meike's video!

Example 3.3

Let X and Y have constant density on the square
 $0 \leq X \leq 4, 0 \leq Y \leq 4$.

3. Let $Z = \min(X, Y)$. Find the pdf for Z , that is $f_Z(z)$.

Let's complicate this even further!

Example 4

Let X and Y have joint density $f_{X,Y}(x, y) = \frac{8}{5}(x + y)$ in the region $0 < x < 1, \frac{1}{2} < y < 1$. Find the pdf of the r.v. Z , where $Z = XY$.

