

Chapter 2: Probability

Meike Niederhausen and Nicky Wakim

2023-09-27

Class Overview

- Probabilities of equally likely events
- Probability Axioms
- Some probability properties
- Partitions
- Venn Diagram Probabilities

Probabilities of equally likely events

Pick an *equally likely* card, any *equally likely* card

Example 1

Suppose you have a regular well-shuffled deck of cards. What's the probability of drawing:

1. any heart
2. the queen of hearts
3. any queen

Let's break down this probability

If S is a finite sample space, with **equally likely outcomes**, then

$$\mathbb{P}(A) = \frac{|A|}{|S|}.$$

A probability is a function...

$\mathbb{P}(A)$ is a function with

- **Input:** event A from the sample space S , ($A \subseteq S$)
- **Output:** a number between 0 and 1 (inclusive)

$$\mathbb{P}(A) : S \rightarrow [0, 1]$$

A function that follows some specific rules though!

See Probability Axioms on next slide.

Probability Axioms

Probability Axioms

Axiom 1

For every event A , $0 \leq \mathbb{P}(A) \leq 1$.

Axiom 2

For the sample space S , $\mathbb{P}(S) = 1$.

Axiom 3

If A_1, A_2, A_3, \dots , is a collection of **disjoint** events, then

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i).$$

Some probability properties

Some probability properties

Using the Axioms, we can prove all other probability properties!

Proposition 1

For any event A , $\mathbb{P}(A) = 1 - \mathbb{P}(A^C)$

Proposition 4

$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$

Proposition 2

$\mathbb{P}(\emptyset) = 0$

Proposition 5

$\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C) + \mathbb{P}(A \cap B \cap C)$

Proposition 3

If $A \subseteq B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$

Proposition 1 Proof

Proposition 1

For any event A , $\mathbb{P}(A) = 1 - \mathbb{P}(A^C)$

Proposition 2 Proof

Proposition 2

$$\mathbb{P}(\emptyset) = 0$$

Proposition 3 Proof

Proposition 3

If $A \subseteq B$, then $\mathbb{P}(A) \leq \mathbb{P}(B)$

Proposition 4 Visual Proof

Proposition 4

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$$

Proposition 5 Visual Proof

Proposition 5

$$\mathbb{P}(A \cup B \cup C) = \mathbb{P}(A) + \mathbb{P}(B) + \mathbb{P}(C) - \mathbb{P}(A \cap B) - \mathbb{P}(A \cap C) - \mathbb{P}(B \cap C) + \mathbb{P}(A \cap B \cap C)$$

Partitions

Partitions

Definition: Partition

A set of events $\{A_i\}_{i=1}^n$ create a **partition** of A , if

- the A_i 's are disjoint (mutually exclusive) and
- $\bigcup_{i=1}^n A_i = A$

Example 2

- If $A \subset B$, then $\{A, B \cap A^C\}$ is a partition of B .
- If $S = \bigcup_{i=1}^n A_i$, and the A_i 's are disjoint, then the A_i 's are a partition of the sample space.

Creating partitions is sometimes used to help calculate probabilities, since by Axiom 3 we can add the probabilities of disjoint events.

Venn Diagram Probabilities

Weekly medications

Example 3

If a subject has an

- 80% chance of taking their medication *this* week,
- 70% chance of taking their medication *next* week, and
- 10% chance of *not* taking their medication *either* week,

then find the probability of them taking their medication exactly one of the two weeks.

Hint: Draw a Venn diagram labelling each of the parts to find the probability.

