

Lesson 5: Random variables and Binomial distribution

TB sections 3.1-3.2


Nicky Wakim

2025-10-13

Learning Objectives

1. Define random variables and how they map to probability distributions
2. Calculate the expected value and variance of discrete random variables
3. Calculate the expected value and variance of linear combinations of discrete random variables
4. Calculate probabilities for different events using a Binomial distribution

Where are we?



Learning Objectives

1. Define random variables and how they map to probability distributions
2. Calculate the expected value and variance of discrete random variables
3. Calculate the expected value and variance of linear combinations of discrete random variables
4. Calculate probabilities for different events using a Binomial distribution

Random variables

Random variable (RV or r.v.)

A **random variable (r.v.)** assigns numerical values (probability) to the outcome of a random phenomenon

Notation: A random variable is usually denoted with a capital letter such as X , Y , or Z .

end of alphabet

From Lesson 3: Probability distributions

Probability distribution


A **probability distribution** consists of all disjoint outcomes and their associated probabilities.

- We've already seen one **in our heads and tails example**

Rules for a probability distribution

A probability distribution is a list of all possible outcomes and their associated probabilities that satisfies three rules:

1. The outcomes listed must be disjoint
2. Each probability must be between 0 and 1
3. The probabilities must total to 1



$\left\{ \begin{array}{l} Y = \# \text{ tails in 6} \\ \text{coin tosses} \end{array} \right.$

$X = \text{tossed heads}$

In the coin toss example...

- We can start to define the probability distribution
- Let's define the coin flip with the random variable X
 - Where $X = 1$ if we get a heads and $X = 0$ if we get a tails

- We can create a table for the random variable and probabilities of each outcome:

Coin flip (x)	$x = 1$	$x = 0$
Probability ($P(X = x)$)	0.5	0.5

- **Note:** I use X to refer to the random variable and x to refer to the realized value it takes
- Then we write $P(X = x)$ to discuss the probability for each realized value (x) of the random variable (X)

- Also note that the sum of the probabilities equal 1: $\sum_{x=0}^1 P(X = x) = 1$

$$\begin{aligned} & \hookrightarrow P(X = 1) + P(X = 0) = 0.5 + 0.5 \\ & \quad = 1 \end{aligned}$$

RV $\uparrow$ realized value $\uparrow$

$$P(X = x)$$

$\downarrow$

$x = 1$

$$P(X = 1)$$

Poll Everywhere Question 1

09:41 Tue Jan 9

100%



Join by Web PollEv.com/nickywakim275



Which of the following best describes a random variable?

A specific outcome of an experiment or random process




29%

[A function that assigns numerical values to the outcomes of a random process ✓




53%

A fixed number determined by a random event




6%

[*probability distribution*
A graph showing the distribution of possible outcomes of an experiment



12%

Let's extend this to rolling a die



Example 1: Rolling a die

Suppose you roll a fair die. Let the random variable (r.v.) X be the outcome of the roll, i.e. the value of the face showing on the die.

1. What is the probability distribution of the r.v. X ?



x	1	2	3	4	5	6
$P(X=x)$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$

$$P(X=x) = \frac{1}{6} \quad \text{for } x=1, 2, 3, 4, 5, 6$$

Learning Objectives

1. Define random variables and how they map to probability distributions
2. Calculate the expected value and variance of discrete random variables
3. Calculate the expected value and variance of linear combinations of discrete random variables
4. Calculate probabilities for different events using a Binomial distribution

Discrete vs. continuous random variables

- Probability distributions are usually either **discrete** or **continuous**, depending on whether the random variable (RV) is discrete or continuous.

Discrete random variable (RV)

A **discrete** RV X takes on a finite number of values or countably infinite number of possible values.

Think:

- Number of heads in a set of coin tosses
- Number of people who have had chicken pox

finite/inf &
countable

$$P(X=x) = \text{real prob value}$$

Continuous random variable (RV)

A **continuous** RV X can take on any real value in an interval of values or unions of intervals.

Think:

- Height
- Blood pressure

infinite &
NOT countable

$$P(X=\underline{x}) = 0$$

8 hrs

7.9 hrs

7.99 hrs

7.999

7.9999

Expectation of random variables discrete RVs

- We call the mean of a random variable its **expected value**
- The expected value is calculated as a **weighted average** weighted by probability

Expected value of a discrete random variable

If X takes on outcomes $x_1, \dots, x_k$ with probabilities $P(X = x_1), \dots, P(X = x_k)$, the expected value of X is the sum of each outcome multiplied by its corresponding probability:

$$\begin{aligned}\mu = E[X] &= x_1 P(X = x_1) + x_2 P(X = x_2) + \dots + x_k P(X = x_k) \\ &= \sum_{i=1}^k x_i P(X = x_i)\end{aligned}$$

$$\begin{aligned}x_1 &= 1 \quad H \\ x_2 &= 0 \quad T\end{aligned}$$

Back to rolling a die

Example 1: Rolling a die

Let's go back to our fair die with RV X as the value of the face showing on the die.

- ② What is the expected outcome of the RV X ?
3. Now suppose the 6-sided die is not fair. How would we calculate the expected outcome?

x	$\mathbb{P}(X = x)$
1	0.10
2	0.20
3	0.05
4	0.05
5	0.25
6	0.35

x	1	2	3	4	5	6
$P(X=x)$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

②

$$\begin{aligned} E(X) &= 1 \cdot \left(\frac{1}{6}\right) + 2 \cdot \left(\frac{1}{6}\right) + 3 \cdot \left(\frac{1}{6}\right) \\ &\quad + 4 \cdot \left(\frac{1}{6}\right) + 5 \cdot \left(\frac{1}{6}\right) + 6 \cdot \left(\frac{1}{6}\right) \\ &= \frac{1 + 2 + 3 + 4 + 5 + 6}{6} \end{aligned}$$

$$= 3.5$$

exp val does NOT need to be a possible outcome of X

③

$$\begin{aligned} E(X) &= 1 \cdot (0.1) + 2 \cdot (0.2) + 3 \cdot (0.05) \\ &\quad + 4 \cdot (0.05) + 5 \cdot (0.25) + 6 \cdot (0.35) \\ &= 4.2 \end{aligned}$$

Variability of random variables

Just like with data, the variability of a r.v. is described with its variance or standard deviation

Variance of a discrete random variable

If X takes on outcomes $x_1, \dots, x_k$ with probabilities $P(X = x_1), \dots, P(X = x_k)$ and expected value $\mu = E(X)$, then the variance of X , denoted by $\text{Var}(X)$ or σ^2 , is

$$\begin{aligned}\text{Var}(X) &= (x_1 - \mu)^2 P(X = x_1) + \dots + (x_k - \mu)^2 P(X = x_k) \\ &= \sum_{i=1}^k (x_i - \mu)^2 P(X = x_i)\end{aligned}$$

Standard deviation of a discrete random variable

The standard deviation of X , labeled $SD(X)$ or σ , is

$$\sigma = SD(X) = \sqrt{\text{Var}(X)}$$


Back to rolling a die

Example 1: Rolling a die

Suppose you roll a fair 6-sided die. Let the random variable (r.v.)* X be the outcome of the roll, i.e. the value of the face showing on the die.

4. Find the variance and standard deviation of X .

x	$\mathbb{P}(X = x)$
1	$\frac{1}{6}$
2	$\frac{1}{6}$
3	$\frac{1}{6}$
4	$\frac{1}{6}$
5	$\frac{1}{6}$
6	$\frac{1}{6}$



$$\begin{aligned}\text{Var}(X) &= \sum_{i=1}^6 (\underline{x_i} - \underline{\mu})^2 P(X = x_i) \\ &= \underline{(1 - 3.5)^2} \underline{\frac{1}{6}} + \underline{(2 - 3.5)^2} \underline{\frac{1}{6}} + \\ &\quad (3 - 3.5)^2 \left(\frac{1}{6}\right) + (4 - 3.5)^2 \left(\frac{1}{6}\right) \\ &\quad + (5 - 3.5)^2 \left(\frac{1}{6}\right) + (6 - 3.5)^2 \left(\frac{1}{6}\right)\end{aligned}$$

$$\sigma^2 = \underline{2.917}$$

$$\begin{aligned}\sigma &= \text{SD}(X) = \sqrt{\text{Var}(X)} = \sqrt{2.917} \\ &= 1.71\end{aligned}$$

Learning Objectives

1. Define random variables and how they map to probability distributions
2. Calculate the expected value and variance of discrete random variables
3. Calculate the expected value and variance of linear combinations of discrete random variables
4. Calculate probabilities for different events using a Binomial distribution

Linear combinations of random variables

Linear combinations of random variables

If X and Y are random variables and a and b are constants, then

$$aX + bY$$

is a linear combination of the random variables.

X + Y
fair die unfair

Theorem: Expected value of a linear combination of random variables

If X and Y are random variables and a and b are constants, then

$$E(aX + bY) = aE(X) + bE(Y)$$

$$E(a) = a$$

$$E(b) = b$$

and

expected val of

sum is sum of
exp values

$$E(aX + b) = aE(X) + b$$

$$X + Y : E(X + Y) = E(X) + E(Y)$$

Poll Everywhere Question2

09:41 Tue Jan 9


100%



Join by Web PollEv.com/nickywakim275



Let X and Y be two random variables with $E(X) = 3$ and $E(Y) = 5$, and constants $a = 2$ and $b = -4$. What is the expected value of the linear combination $aX + bY$?



Powered by Poll Everywhere

$$\begin{aligned} E(aX + bY) &= aE(X) + bE(Y) \\ &= 2(3) + (-4) \cdot 5 \\ &= 6 - 20 \\ &= -14 \end{aligned}$$

Keep rolling dice!

DO NOT } might be
DO THIS } tempted to
do

$$1 \cdot X_1 + 2 X_2 = \underline{3 X_1}$$

↓ realized
↑ can be diff than

Example: Expected money for rolling 3 dice

Let the random variables X_1, X_2, X_3 be the values shown on rolls for 2 fair 6-sided dice and 1 unfair die (as described in our previous example). Suppose you are given in dollars the amount of the first roll, plus twice the value of the second roll, plus 4 times the value of the unfair die roll. How much money do you expect to get?

X_1, X_2, X_3 let M be linear combo
↳ unfair

linear combo: $M = 1 X_1 + 2 X_2 + 4 X_3$

$$E(M) = E[X_1 + 2 X_2 + 4 X_3]$$

$$= E(X_1) + 2 E(X_2) + 4 E(X_3)$$

$$= 3.5 + 2 \cdot 3.5 + 4 \cdot 4.2$$

$$= \$27.30$$

$$E(X_1) = 3.5$$

$$E(X_2) = 3.5$$

$$E(X_3) = 4.2$$

Variance of a linear combination

Theorem: Variance of a linear combination of random variables

If X and Y are **independent** random variables and a and b are constants, then

$$\text{Var}(aX + bY) = a^2 \text{Var}(X) + b^2 \text{Var}(Y)$$

$$aX + b$$

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

$$\text{Var}(b) = 0$$

$$\text{Var}(aX) = a^2 \text{Var}(X)$$

BONUS

$$\text{Var}(X) = E(X^2) - [E(X)]^2$$

$$\text{Var}(aX) = E[(aX)^2] - [E(aX)]^2 = a^2 E(X^2) - \dots \text{more to it}$$

Keep keep rolling dice!

one roll will not affect others $\Rightarrow$ independent

Example: Expected money for rolling 3 dice

Let the random variables X_1, X_2, X_3 be the values shown on rolls for 2 fair 6-sided dice and 1 unfair die (as described in our previous example). Suppose you are given in dollars the amount of the first roll, plus twice the value of the second roll, plus 4 times the value of the unfair die roll. What are the variance and standard deviation of the amount you get from the 3 rolls?

$$M = X_1 + 2X_2 + 4X_3 \rightarrow$$

$$\text{Var}(M) = \text{Var}(X_1 + 2X_2 + 4X_3)$$

$$= \text{Var}(X_1) + 2^2 \text{Var}(X_2) + 4^2 \text{Var}(X_3)$$

$$= \text{Var}(X_1) + 4 \text{Var}(X_2) + 16 \text{Var}(X_3)$$

$$5 \text{Var}(X_1)$$

$$\text{if } \text{Var}(X_1) = \text{Var}(X_2)$$

WRONG WAY TO DO!

$$M = 3X_1 + 4X_3$$

$$\text{Var}(3X_1 + 4X_3)$$

$$= 9 \text{Var}(X_1) + 16 \text{Var}(X_3)$$

Learning Objectives

1. Define random variables and how they map to probability distributions
2. Calculate the expected value and variance of discrete random variables
3. Calculate the expected value and variance of linear combinations of discrete random variables
4. Calculate probabilities for different events using a Binomial distribution

Binomial random variable

- One specific type of discrete random variable is a binomial random variable

Binomial random variable

- X is a binomial random variable if it represents the number of successes in n independent replications (or trials) of an experiment where
 - Each replicate has two possible outcomes: either success or failure
 - The probability of success is p
 - The probability of failure is $q = 1 - p$

- A binomial random variable takes on values $0, 1, 2, \dots, n$.
- If a r.v. X is modeled by a Binomial distribution, then we write in shorthand $X \sim \text{Binom}(n, p)$
- Quick example: The number of heads in 3 tosses of a fair coin is a binomial random variable with parameters $n = 3$ and $p = 0.5$.

↳ let $X = \# \text{ heads in 3 tosses}$
 $x = 0, 1, 2, 3$

"simulated as" or
"distributed as"

$$X \sim \text{binom}(3, 0.5)$$

Poll Everywhere Question 3

13:34 Wed Oct 15

...

65%



Join by Web PollEv.com/nickywakim275



Which of the following scenarios can be modeled by a Binomial distribution?

Normal measuring numeric charact. (no succ/fail)
Measuring the height of students in a class 12%

* Poisson
Counting the number of cats in a household 0%

Flipping a coin 5 times and counting the number of heads ☒ 88%
set reps "successes"

Recording the temperature throughout the day 0%

Powered by Poll Everywhere

Bernoulli distribution

- When $n = 1$, aka we have a single trial, we give a different name to the random variable: Bernoulli

reps = 1

Bernoulli random variable

Bernoulli random variable. If X is a random variable that takes value 1 with probability of success p and 0 with probability $1 - p$ (or q), then X is a Bernoulli random variable.

- We call the probability of success p the **parameter** of the Bernoulli distribution.
- If a r.v. X is modeled by a Bernoulli distribution, then we write in shorthand $X \sim \text{Bernoulli}(p)$ or $X \sim \text{Bern}(p)$

Mean and SD of a Bernoulli r.v.

If* X is a Bernoulli r.v. with probability of success p , then $E(X) = p$ and $\text{Var}(X) = p(1 - p)$

Relationship between Bernoulli and Binomial¹

- The **Bernoulli distribution** is a special case of the Binomial distribution where $n = 1$
 - Specifically:

$$\text{Binomial}(1, p) = \text{Bernoulli}(p)$$

- To get a **Binomial distribution**, we simply extend the scenario from a **single trial** to **multiple independent trials**.
 - If we conduct n independent Bernoulli trials with the same success probability p , the total number of successes across these n trials will follow a Binomial distribution
- Quick example:
 - **Bernoulli**: If you flip a coin once, with probability $p = 0.5$ of landing heads, that is a Bernoulli trial.
 - **Binomial**: If you flip the coin 5 times, and you want to know how many times it will land heads, the number of heads will follow a Binomial distribution with parameters $n = 5$ and $p = 0.5$

$$\begin{array}{ccccccc} X & = & \sum_{i=1}^5 & Y_i & = & Y_1 + Y_2 + Y_3 + Y_4 + Y_5 \\ \text{binomial RV} & & & \downarrow & & & \\ & & & \text{bernoulli RV} & & & \end{array}$$

Binomial distribution

Distribution of a Binomial random variable

Let X be the total number of successes in n independent trials, each with probability p of a success. Then probability of observing exactly k successes in n independent trials is

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, 2, \dots, n$$

5 trials, want to know
coin flip

$$P(X=3) = \binom{5}{3} 0.5^3 (1-0.5)^{5-3}$$

- The parameters of a binomial distribution are p and n .
- If a r.v. X is modeled by a binomial distribution, then we write in shorthand

Mean and variance of a Binomial r.v

If X is a binomial r.v. with probability of success p , then $E(X) = np$ and $\text{Var}(X) = np(1-p)$

$$E\left(\sum_{i=1}^n Y_i\right) = E(Y_1) + E(Y_2) + E(Y_3) + E(Y_4) + E(Y_5)$$

Binomial distribution: R commands

R commands with their **input** and **output**:

R code

What does it return?

`rbinom()`

returns sample of random variables with specified binomial distribution

`dbinom()`


returns probability of getting certain number of successes

`pbinom()`

returns cumulative probability of getting certain number or less successes

`qbinom()`

returns number of successes corresponding to desired quantile



↳ 90th
75th

↳ always bottom
up, so
 $0 \rightarrow 75 \rightarrow \text{bottom}$
 $100 \rightarrow 75 \rightarrow \text{top 25}$

Binomial distribution example (1/5)

$$n=10 \quad p=0.25$$

Vaccinated people testing positive for Covid-19

About 25% of people that test positive for Covid-19 are vaccinated for Covid-19. Suppose 10 people have tested positive for Covid-19 (independently of each other). Let X denote the number of people that are vaccinated among the 10 that tested positive.

1. What is the expected value of X ?
2. What is the SD of X ?
3. What is the probability that exactly 4 of the 10 people that tested positive are vaccinated?
4. What is the probability that at most 3 of the 10 people that tested positive are vaccinated?
5. What is the probability that at least 5 of the 10 people that tested positive are vaccinated?

$X = \# \text{ vaccinated of } 10 \text{ ppl}$

$$X \sim \text{binom}(n=10, p=0.25)$$

Binomial distribution example (2/5)

Vaccinated people testing positive for Covid-19

About 25% of people that test positive for Covid-19 are vaccinated for Covid-19. Suppose 10 people have tested positive for Covid-19 (independently of each other). Let X denote the number of people that are vaccinated among the 10 that tested positive.

1. What is the expected value of X ?
2. What is the SD of X ?

$$\textcircled{1} \quad E(X) = np = 10 \cdot 0.25 = 2.5 \text{ ppl vaccinated}$$

$$\textcircled{2} \quad \text{Var}(X) = n(p)(1-p) = 10 \cdot 0.25 \cdot (1-0.25) = 1.875$$

$$\text{SD}(X) = \sqrt{\text{Var}(X)} = \sqrt{1.875} = 1.369$$

Binomial distribution example (3/5)

Vaccinated people testing positive for Covid-19

About 25% of people that test positive for Covid-19 are vaccinated for Covid-19. Suppose 10 people have tested positive for Covid-19 (independently of each other). Let X denote the number of people that are vaccinated among the 10 that tested positive.

3. What is the probability that exactly 4 of the 10 people that tested positive are vaccinated?

$$P(X = 4) = \binom{10}{4} 0.25^4 (1 - 0.25)^{10-4} = \underline{0.146}$$

Handwritten notes: "10 choose 4" above the binomial coefficient; "y" next to the 0.25 term.

```
1 dbinom(x = 4, size = 10, prob = 0.25) # d for distribution
```

```
[1] 0.145998
```

• In general, for $P(X = k)$ we code: `dbinom(x = k, size = n, prob = p)`

Handwritten: # trials

Handwritten: prob of success

Binomial distribution example (4/5)

Vaccinated people testing positive for Covid-19

About 25% of people that test positive for Covid-19 are vaccinated for Covid-19. Suppose 10 people have tested positive for Covid-19 (independently of each other). Let X denote the number of people that are vaccinated among the 10 that tested positive.

4. What is the probability that *at most* 3 of the 10 people that tested positive are vaccinated?

$$\begin{aligned} P(X \leq 3) &= P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3) \\ &= \binom{10}{0} 0.25^0 (0.75)^{10} + \binom{10}{1} 0.25^1 (0.75)^9 + \binom{10}{2} 0.25^2 (0.75)^8 + \binom{10}{3} 0.25^3 (0.75)^7 \\ &= 0.7758 \end{aligned}$$

```
1 pbinom(q = 3, size = 10, prob = 0.25, lower.tail = T)
```

```
[1] 0.7758751
```

- In general, for $P(X \leq k)$ we code: `pbinom(q = k, size = n, prob = p)` with `lower.tail = T` as a default option

> lower.tail
= F

<
=

Binomial distribution example (5/5)

Vaccinated people testing positive for Covid-19

About 25% of people that test positive for Covid-19 are vaccinated for Covid-19. Suppose 10 people have tested positive for Covid-19 (independently of each other). Let X denote the number of people that are vaccinated among the 10 that tested positive.

5. What is the probability that *at least* 5 of the 10 people that tested positive are vaccinated?

$$\begin{aligned} P(X \geq 5) &= P(X = 5) + P(X = 6) + P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10) \\ &= \binom{10}{5} 0.25^5 (0.75)^5 + \binom{10}{6} 0.25^6 (0.75)^4 + \dots + \binom{10}{10} 0.25^{10} (0.75)^0 \\ &= \cancel{0.7758} \rightarrow 0.078 \end{aligned}$$

```
1 pbinom(q = 4, size = 10, prob = 0.25, lower.tail = F) # for greater than!  
[1] 0.07812691
```

```
1 1 - pbinom(q = 4, size = 10, prob = 0.25, lower.tail = T)  
[1] 0.07812691
```

$$P(X \leq 4) + P(X \geq 5) = 1$$

complementary

$$P(X \geq 5) = P(X > 4)$$

Resources!

- Seeing theory: random variables