

Quantum Theory Angular Momentum
Electronic version 2026

originally by:
D. A. Varshalovich
A. N. Moskalev
V. K. Khersonskii
This version:
Niels R. Walet

Contents

1	Elements of Vector and Tensor Theory	7
1.1	Coordinate Systems. Basis Vectors	7
1.1.1	Cartesian Coordinate System	7
1.1.2	Polar Coordinate System	8
1.1.3	Spherical Coordinate System	9
1.1.4	Helicity Basis Vectors	10
1.1.5	Relations Between Different Basis Vectors	10
1.2	Vectors. Tensors	14
1.2.1	Vector Components	14
1.2.2	Scalar Product of Vectors	15
1.2.3	Vector Product of Vectors	16
1.2.4	Products Involving Three or More Vectors	17
1.2.5	Tensors δ_{ik} and ε_{ikl}	18
1.3	Differential Operations	18
1.3.1	Operator ∇	18
1.3.2	Laplace Operator	19
1.3.3	Differential Operations on Scalars and Vectors	20
1.4	Rotations of Coordinate System	22
1.4.1	Description of Rotations in Terms of the Euler Angles	22
1.4.2	Description of Rotations in Terms of Rotation Axis and Rotation Angle	23
1.4.3	Description of Rotations in Terms of Unitary 2×2 Matrices. Cayley-Klein Parameters.	25
1.4.4	Relations Between Different Descriptions of Rotations	26
1.4.5	Rotation Operator	27
1.4.6	Transformation of Cartesian Vectors and Tensors Under Rotations of Coordinate Systems. Rotation Matrix A	28
1.4.7	Addition of Rotations	31
2	Angular momentum operators	35
2.1	Total angular momentum operator	35
2.1.1	Definition	35
2.1.2	Commutation Relations	35
2.1.3	Coordinate Inversion. Time Reversal	36
2.1.4	Total Angular Momentum of a System. Orbital and Spin Angular Momenta	37
2.2	Orbital angular momentum operator	37
2.2.1	Definition	37
2.2.2	Commutation Relations	38
2.2.3	Explicit Form	39
2.3	Spin angular momentum operator	40
2.3.1	Definition	40
2.3.2	Commutation Relations	40
2.3.3	Explicit Form	40
2.3.4	Traces of Products of Spin Matrices	41
2.4	Polarization operators	41
2.4.1	Definition	41
2.4.2	Explicit Form	42
2.4.3	Properties of $\hat{T}_{LM}(S)$ under Transformations of the Coordinate System	42
2.4.4	Expansion Series of Polarization Operators	43
2.4.5	Commutators and Anticommutators	43
2.4.6	Traces of Products of Polarization Operators	44

2.5	Spin matrices for $S = 1/2$	44
2.5.1	Explicit Form	44
2.5.2	Commutators and Anticommutators	44
2.5.3	Products of Spin Matrices	45
2.5.4	Functions of Spin Matrices	46
2.5.5	Rotation Operators	46
2.5.6	Euler angles	46
2.5.7	Result of rotation operator	47
2.5.8	Traces of Products of Spin Matrices ($S = \frac{1}{2}$)	47
2.6	Spin matrices and polarization operators for $S = 1$	48
2.6.1	Spin $S = 1$	48
2.6.2	Explicit Form	48
2.6.3	Products of Spin and Polarization Matrices	51
2.6.4	Functions of Spin Matrices	53
2.6.5	Operators of Coordinate Rotations	53
2.6.6	Traces of Products of Spin Matrices	55
3	Irreducible tensors	57
3.1	Definition and properties of irreducible tensors	57
3.1.1	Definition	57
3.1.2	Covariant and Contravariant Components	58
3.1.3	Transformation of Irreducible Tensors Under a Rotation of the Coordinate System	58
3.1.4	Transformation of Irreducible Tensors Under Inversion of the Coordinate System	58
3.1.5	Double Tensors	58
3.1.6	Examples of Irreducible Tensors	59
3.1.7	Direct and Irreducible Tensor Products. Commutators of Tensor Products	59
3.1.8	Scalar Products of Irreducible Tensors	60
3.2	Relation between the irreducible tensor algebra and vector and tensor theory	60
3.2.1	Vectors and Irreducible Tensors	60
3.2.2	Cartesian Tensors of Second and Third Ranks	62
3.2.3	Differential Operations as Irreducible Tensor Products	63
3.3	Recoupling in irreducible tensor products	64
3.3.1	Relations Valid for Commuting as well as Non-Commuting Tensors	64
3.3.2	Relations for Commuting Tensors	65
3.3.3	Relations for Non-Commuting Tensors	66
4	Wigner D-functions	67
4.1	Definition of $D_{MM'}^J(\alpha, \beta, \gamma)$	67
4.2	Differential equations for $D_{MM'}^J(\alpha, \beta, \gamma)$	69
4.3	Explicit forms of the Wigner D-functions	70
4.3.1	Expressions for $d_{MM'}^J(\beta)$ Involving Trigonometric Functions	70
4.3.2	Differential Representations of $d_{MM'}^J(\beta)$	71
4.3.3	Integral Representations of $d_{MM'}^J(\beta)$	72
4.3.4	Relation Between $d_{MM'}^J(\beta)$ and the Jacobi Polynomials	72
4.3.5	Relations Between $d_{MM'}^J(\beta)$ and Hypergeometric Functions	72
4.4	Symmetries of $d_{MM'}^J(\beta)$ and $D_{MM'}^J(\alpha, \beta, \gamma)$	73
4.5	Rotation matrix U_{MM}^J , in terms of angles ω, Θ, Φ	74
4.5.1	Definition	74
4.5.2	Explicit form	74
4.5.3	Differential Equations	75
4.5.4	Orthogonality and Completeness	75
4.5.5	Principal Properties	76
4.5.6	Special Cases	77
4.6	Sums involving D -functions	77
4.6.1	The Clebsch-Gordan Series	77
4.6.2	Some Applications of the Clebsch-Gordan Expansion	77
4.6.3	Generalization of the Clebsch-Gordan Expansion	78
4.6.4	Determinant of Matrix $D_{MM'}^J$	79
4.7	Addition of rotations	79
4.7.1	The Addition Theorem for $D_{MM'}^J(\alpha, \beta, \gamma)$	79
4.7.2	The Addition Theorem for $d_{MM'}^J(\beta)$	80

4.7.3	Addition of Two Identical Rotations	80
4.7.4	The Multiplication Theorem for $d_{MM'}^J(\beta)$	81
4.7.5	Sums Involving the D-Functions of Different Arguments	81
4.7.6	The Ponzano-Regge sum	81
4.8	Recursion relations for $D_{MM'}^J$	82
4.8.1	Relations between D^J and $D^{J\pm 1}$	82
4.8.2	Relations Between D^J and $D^{J\pm 1/2}$	83
4.8.3	Relations Between $D_{MM'}^J$ and $D_{M\pm 1M'\mp 1}^J$	84
4.9	Differential relations for $D_{MM'}^J(\alpha, \beta, \gamma)$	85
4.10	Orthogonality and completeness of the D -functions	86
4.11	Integrals involving the D -functions	87
4.11.1	Integration of Products of $D_{MM'}^J$	87
4.11.2	Integrals Involving $d_{MM'}^J(\beta)$	87
4.12	Invariant summation of integrals involving $D_{MM'}^J(\alpha, \beta, \gamma)$	88
4.13	Generating functions for $d_{MM}^J(\beta)$	89
4.14	Characters $\chi^J(R)$ of irreducible representations of rotation group	90
4.14.1	Definition	90
4.14.2	Explicit Forms	90
4.14.3	Principal Properties	91
4.14.4	Differential Equation	92
4.14.5	Differential Relations	92
4.14.6	Algebraic Relations	92
4.14.7	Orthogonality and Completeness	92
4.14.8	Integrals Involving $\chi^J(\omega)$	93
4.14.9	Sums Involving $\chi^J(\omega)$	93
4.14.10	$\chi^J(\omega)$ for Particular Values of ω	94
4.14.11	Special Cases of $\chi^J(\omega)$	94
4.15	Generalized characters, $\chi_\lambda^J(R)$, of irreducible representations of the rotation group	95
4.15.1	Definition	95
4.15.2	Explicit Forms	95
4.15.3	Principal Properties	96
4.15.4	Differential Equation	97
4.15.5	Orthogonality and Completeness	97
4.15.6	The Addition Theorem for $\chi_\lambda^J(\omega)$	97
4.15.7	Sums and Infinite Series Involving $\chi_\lambda^J(\omega)$	98
4.15.8	Special Cases of $\chi_\lambda^J(\omega)$ for Particular λ	99
4.15.9	Special Cases of $\chi_\lambda^J(\omega)$ for Particular J	100
4.16	$D_{MM'}^J(\alpha, \beta, \gamma)$ for particular values of the arguments	100
4.17	Special cases of $D_{MM'}^J$ for particular M or M'	101
4.18	Asymptotics of $D_{MM'}^J(\alpha, \beta, \gamma)$	103
4.18.1	Large Angular Momentum	103
4.18.2	Small Variation of Rotation Axis	103
4.18.3	Infinitesimal Rotations	103
4.19	Definitions of $D_{MM'}^J(\alpha, \beta, \gamma)$ by other authors	104
4.20	Special cases of $d_{MM'}^J(\beta)$ for particular J, M and M'	104
4.21	Tables of $d_{MM'}^J(\beta)$ for $\beta = \pi/2$	104
4.22	Special cases of $U_{MM'}^J(\omega; \Theta, \Phi)$	106
5	Spherical harmonics	113
5.1	Definition	113
5.1.1	Commutation Relations	113
5.1.2	Differential Equations	113
5.1.3	Boundary Conditions	114
5.1.4	Normalization	114
5.1.5	Choice of Phase	114
5.1.6	Zonal, Sectorial and Tesseral Harmonics	115
5.1.7	Solutions of Some Differential Equations in Terms of $Y_{lm}(\vartheta, \varphi)$	115
5.2	Explicit forms of the spherical harmonics and their relations to other functions	116
5.2.1	Differential Expressions for $Y_{lm}(\vartheta, \varphi)$	116
5.2.2	Representations of $Y_{lm}(\vartheta, \varphi)$ as a Power Series of Trigonometric Functions of $\vartheta/2$	116
5.2.3	Representations of $Y_{lm}(\vartheta, \varphi)$ as a Power Series of Trigonometric Functions of ϑ	117

5.2.4	$Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Trigonometric Functions of $\vartheta/2$	118
5.2.5	$Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Trigonometric Functions of ϑ	118
5.2.6	$Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Exponential Functions	119
5.2.7	$Y_{lm}(\vartheta, \varphi)$ and Other Special Functions	119
5.2.8	$Y_{lm}(\vartheta, \varphi)$ as an Irreducible Tensor Product	120
5.3	Integral representations of the spherical harmonics [4,22,27]	120
5.3.1	$Y_{lm}(\vartheta, \varphi)$ in the Form of Indefinite Integrals	120
5.3.2	$Y_{lm}(\vartheta, \varphi)$ in the Form of Definite Integrals	120
5.3.3	$Y_{lm}(\vartheta, \varphi)$ in the Form of Improper Integrals	121
5.4	Symmetry properties	121
5.5	Behaviour of $Y_{lm}(\vartheta, \varphi)$ under transformations of coordinate systems	122
5.5.1	Rotation	122
5.5.2	Inversion	123
5.5.3	Parallel Translation	123
5.5.4	Special Cases of Coordinate-System Transformations	123
5.6	Expansions in series of the spherical harmonics	124
5.6.1	General Relations	124
5.6.2	Expansion of Products of the Spherical Harmonics	125
5.7	Recursion relations	126
5.8	Differential relations	127
5.8.1	Action of the Operator of Orbital Angular Momentum on $Y_{lm}(\vartheta, \varphi)$	127
5.8.2	First and Second Order Derivatives of $Y_{lm}(\vartheta, \varphi)$	127
5.8.3	Vector Differentiation Operations	127
5.9	Some integrals involving spherical harmonics	128
5.9.1	Integrals over Total Solid Angle	128
5.9.2	Fourier Transformations for Some Functions Which Contain $Y_{lm}(\vartheta, \varphi)$	129
5.9.3	Integrals with Respect to ϑ	129
5.10	Sums involving spherical harmonics	129
5.10.1	Sums over m (with fixed l)	129
5.10.2	Sums over l (with fixed $m \geq 0$)	130
5.11	Generating functions for $Y_{lm}(\vartheta, \varphi)$	130
5.12	Asymptotic expressions for $Y_{lm}(\vartheta, \varphi)$	131
5.12.1	$Y_{lm}(\vartheta, \varphi)$ for Large l	131
5.12.2	Behaviour of $Y_{lm}(\vartheta, \varphi)$ in Neighbourhoods of $\vartheta = 0, \pi$ and $\pi/2$	131
5.12.3	Asymptotic Expression for Fixed $m, l \rightarrow \infty, \vartheta \rightarrow 0$ and Finite $l\vartheta$	132
5.13	$Y_{lm}(\vartheta, \varphi)$ for special values of l and m	132
5.13.1	$Y_{lm}(\vartheta, \varphi)$ for $l \leq 5$	132
5.13.2	$Y_{lm}(\vartheta, \varphi)$ for $ m = 0, 1, 2, 3, 4$ and Any Integer l	134
5.13.3	$Y_{lm}(\vartheta, \varphi)$ for $ m = l, l-1, l-2, l-3, l-4, l-5$ and Any Integer l	134
5.14	$Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ for special ϑ	135
5.15	Zeros of $Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$	135
5.15.1	Zeros of $Y_{lm}(\vartheta, \varphi)$	135
5.15.2	Zeros of $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$	136
5.16	Bipolar and tripolar spherical harmonics	136
5.16.1	Bipolar Spherical Harmonics	136
5.16.2	Tripolar Spherical Harmonics	137
5.17	Expansions of functions which depend on two vectors	138
5.17.1	Preliminary Remarks	138
5.17.2	Addition Theorem for the Spherical Harmonics	139
5.17.3	Expansions of Some Functions Which Depend on $(\mathbf{r}_1 \cdot \mathbf{r}_2)$	140
5.17.4	Expansions of Some Functions Which Depend on $r = \mathbf{r}_1 - \mathbf{r}_2 $	140
5.17.5	Expansions of $r^n \equiv \mathbf{r}_1 - \mathbf{r}_2 ^n$	141
5.17.6	Expansions of Spherical Waves	142
5.17.7	Expansions of $r^N Y_{LM}(\vartheta, \varphi)$	142

6	Spin functions	145
6.1	Spin functions of particles with arbitrary spin	145
6.1.1	Definition	145
6.1.2	Basis Spin Functions	145
6.1.3	Helicity Basis Functions	147
6.1.4	General Spin Functions	148
6.1.5	Polarization Density Matrix	149
6.1.6	Two Particles with Arbitrary Spins	150
6.2	Spin functions for $S = 1/2$	151
6.2.1	Basis Spin Functions	151
6.2.2	Expansions of Products of Basis Functions	152
6.2.3	Action of Spin Operators on Basis Functions	152
6.2.4	Transformation of the Basis Functions Under Rotations of the Coordinate System	152
6.2.5	Helicity Basis Functions	153
6.2.6	General Spin Functions for $S = \frac{1}{2}$	154
6.2.7	Polarization Density Matrix	156
6.3	Spin functions for $S = 1$	156
6.3.1	Basis Spin Functions	156
6.3.2	Expansions of Products of Spin Functions	158
6.3.3	Action of Spin Operators on Basis Functions	159
6.3.4	Action of Quadrupole Operators on Basis Functions	160
6.3.5	Transformation of Basis Functions Under Rotations of the Coordinate Systems	161
6.3.6	Helicity Basis Functions for $S = 1$	161
6.3.7	General Spin Functions for $S = 1$	164
7	Tensor spherical harmonics	167
7.1	General properties of tensor spherical harmonics	167
7.1.1	Definition	167
7.1.2	Components of Tensor Spherical Harmonics	168
7.1.3	Complex Conjugation	168
7.1.4	Transformations of Coordinate Systems	169
7.1.5	Differential Equations	169
7.1.6	Action of Operators ∇, n and Angular Momentum Operators	169
7.1.7	Sums of Tensor Spherical Harmonics	171
7.1.8	Orthogonality, Normalization and Completeness	171
7.1.9	Expansion in a Series of Tensor Spherical Harmonics	171
7.2	Spinor spherical harmonics	171
7.2.1	Definition	171
7.2.2	Components of Spinor Spherical Harmonics	172
7.2.3	Complex Conjugation. Time Reversal	173
7.2.4	Transformation of Coordinate Systems.	173
7.2.5	Action of ∇ and Angular Momentum Operators	174
7.2.6	Recursion Relations	174
7.2.7	Orthonormality and Completeness. Expansion in a Series of Spinor Spherical Harmonics	175
7.2.8	Clebsch-Gordan Series	175
7.2.9	Addition Theorems	175
7.2.10	Quadratic Forms of Spinor Spherical Harmonics	176
7.3	Vector spherical harmonics	177
7.3.1	Definition	177
7.3.2	Components of Vector Spherical Harmonics	179
7.3.3	Complex Conjugation	182
7.3.4	Transformations of Coordinate Systems	183
7.3.5	Differential Equations	183
7.3.6	Differential Operations	184
7.3.7	Action of Angular Momentum Operators	185
7.3.8	Algebraic Relations	186
7.3.9	Sums of Vector Spherical Harmonics	187
7.3.10	Clebsch-Gordan Series	189
7.3.11	Addition Theorems for Vector Spherical Harmonics	189
7.3.12	Integrals Involving Vector Spherical Harmonics	192
7.3.13	Orthogonality, Normalization and Completeness	193

7.3.14	Expansion in Series of Vector Spherical Harmonics	194
7.3.15	Vector Spherical Harmonics for $\vartheta = 0$ or $\vartheta = \pi$	195
7.3.16	Vector Spherical Harmonics at $J = 0, 1$	195
7.3.17	Quadratic Forms of the Vector Spherical Harmonics	196
7.4	Other notations for tensor spherical harmonics	197
8	Clebsch-Gordan coefficients and $3jm$ symbols	199
8.1	Definition of vector addition coefficients	199
8.1.1	The Clebsch-Gordan Coefficients	199
8.1.2	The Wigner $3jm$ Symbols	200
8.1.3	Regge R-Symbols	200
8.2	EXPLICIT forms of the Clebsch-Gordan coefficients and their relations to other functions	201
8.2.1	Representations of the Clebsch-Gordan Coefficients in the Form of Algebraic Sums	201
8.2.2	Quasi-Binomial Representation of the Clebsch-Gordan Coefficients	202
8.2.3	Clebsch-Gordan Coefficients and Finite Differences	202
8.2.4	Expressions for the Clebsch-Gordan Coefficients in Terms of Binomial Coefficients [116]	203
8.2.5	Representations of the Clebsch-Gordan Coefficients in Terms of the Hypergeometric Functions	203
8.2.6	Representations of the $3jm$ Symbols in the Form of Algebraic Sums	204
8.2.7	Quasi-binomial Representations of the $3jm$ Symbols	205
8.3	Integral representations	205
8.3.1	Integrals Involving Algebraic Functions	205
8.3.2	Integrals Involving the Wigner D-Functions	205
8.3.3	Integral Involving the Spherical Harmonics	205
8.3.4	Integral Representations for Products of the Clebsch-Gordan Coefficients	206
8.4	Symmetry properties	206
8.4.1	Symmetry Properties of the R-Symbols	206
8.4.2	Symmetry Properties of the $3jm$ Symbols	206
8.4.3	Symmetry Properties of the Clebsch-Gordan Coefficients	207
8.4.4	"Mirror" Symmetry	208
8.4.5	Properties of the Vector-Addition Coefficients under Transformations of the Coordinate System and Time Reversal	208
8.5	Explicit forms of the Clebsch-Gordan coefficients for special values of the arguments	209
8.5.1	Special Values of Momenta a, b, c	209
8.5.2	Special Values of Momentum Projections	212
8.6	Recursion relations for the Clebsch-Gordan coefficients	213
8.6.1	General Recursion Relations	213
8.6.2	Arguments α, β, γ Change by 1	213
8.6.3	Arguments Change by $1/2$	214
8.6.4	The Case When $\alpha = \beta = \gamma = 0$	215
8.6.5	Arguments a, b, c Change by 1	215
8.6.6	Arguments a, b, α, β Change by 1	216
8.6.7	Arguments c, b, γ, β Change by 1	216
8.6.8	Recursion Relations for the Regge Symbols	217
8.7	Sums of products of Clebsch-Gordan coefficients	218
8.7.1	Sums Involving One Clebsch-Gordan Coefficient	218
8.7.2	Sums Involving Products of Two Clebsch-Gordan Coefficients	219
8.7.3	Sums Involving Products of Three Clebsch-Gordan Coefficients	219
8.7.4	Sums Involving Products of Four Clebsch-Gordan Coefficients	220
8.7.5	Sums Involving Products of the Clebsch-Gordan Coefficients and One $6j$ Symbol	221
8.7.6	Sums Involving Products of the Clebsch-Gordan Coefficients and One $9j$ Symbol	221
8.7.7	Some Additional Sums of Products of Two Clebsch-Gordan Coefficients	221
8.8	Generating functions of the Clebsch-Gordan coefficients	222
8.8.1	Regge Determinant to Power J	222
8.8.2	Products of Binomials	222
8.8.3	Exponential Function	222
8.8.4	Hypergeometric Function	223
8.8.5	Wigner D-Function	223
8.8.6	Generating Function for the Coefficients C_{a0b0}^{c0}	223
8.9	Classical limit and asymptotic expressions for the Clebsch-Gordan coefficients	223
8.9.1	Asymptotic Expressions for $a, c \gg b$	223

8.9.2	Asymptotic Expressions for $a, b, c, \gamma \gg a + b - \gamma$	224
8.9.3	Semiclassical Formulas for $a, b, c \gg 1$	224
8.9.4	Squares of the Clebsch-Gordan Coefficients in the Classical Limit	225
8.10	Zeros of the vector-addition coefficients	226
8.11	Connection of the Clebsch-Gordan coefficients and the $3jm$ symbols with analogous functions of other authors	227
8.12	Algebraic tables of the Clebsch-Gordan coefficients	227
8.13	Numerical tables of the clebsch-gordan coefficients	227
9	6j Symbols and the Racah Coefficients	231
9.1	Definition	231
9.1.1	6j Symbols	231
9.1.2	Racah Coefficients	232
9.1.3	R Symbols	232
9.2	General Expressions for the $6j$ Symbols. Relations Between the $6j$ Symbols and Other Functions	233
9.2.1	Expressions for the $6j$ Symbols in Terms of Finite Sums	233
9.2.2	Bargmann Formula [61]	234
9.2.3	Relations Between the $6j$ Symbols and the Generalized Hypergeometric Functions	235
9.2.4	Relations Between the $6j$ Symbols and the $3jm$ Symbols	235
9.2.5	Quasi-Binomial Representation of the $6j$ Symbols	235
9.3	Integral Representations of the $6j$ Symbols	236
9.4	Symmetries of the $6j$ Symbols and the Racah Coefficients	237
9.4.1	R-Symbols	237
9.4.2	6j Symbols	237
9.4.3	Racah Coefficients	238
9.4.4	"Mirror" Symmetry	238
9.5	Explicit Forms of the $6j$ Symbols for Certain Arguments	239
9.5.1	One of Arguments is Equal to Zero	239
9.5.2	One of Arguments is Equal to the Sum of Two Others	239
9.5.3	One of Arguments is Smaller by Unity than the Sum of Two Others	240
9.5.4	Arguments a, b, d, e are Equal in Pairs	241
9.6	Recursion Relations	243
9.6.1	Relations in Which Arguments are Changed by $1/2$	243
9.6.2	Relations in Which Arguments are Changed by 1	243
9.7	Generating Function	244
9.8	Sums Involving the $6j$ Symbols	244
9.9	Asymptotics of the $6j$ Symbols for Large Angular Momenta	245
9.9.1	Asymptotic Relations Between the $6j$ Symbols and the Clebsch-Gordan Coefficients	245
9.9.2	Asymptotic Expressions for the $6j$ Symbols	246
9.10	Relations Between the Wigner $6j$ Symbols and Analogous Functions of Other Authors	249
9.11	Tables of Algebraic Expressions for the $6j$ Symbols	249
10	9j and 12j Symbols	251
10.1	Definition of the $9j$ Symbols	251
10.1.1	9j Symbols as Recoupling Coefficients	251
10.1.2	9j Symbol and r Symbol	252
10.2	Explicit Forms of the $9j$ Symbols and Their Relations to Other Functions	253
10.2.1	Expressions for the $9j$ Symbols in the Forms of Algebraic Sums	253
10.2.2	Wu Formulas [77]	254
10.2.3	9j Symbols as Sums of Products of the Clebsch-Gordan Coefficients or the $3jm$ Symbols [54]	255
10.3	Integral Representations of the $9j$ Symbols	256
10.3.1	Representation Involving the Generalised Hypergeometric Function	256
10.3.2	Representation Involving the Wigner D Functions	256
10.3.3	Integrals Involving Characters of Irreducible Representations of Rotation Group	257
10.4	Symmetry Properties of the $9j$ Symbols	257
10.4.1	Permutation Symmetry	257
10.4.2	Symmetries of the r Symbol	259
10.4.3	"Mirror" Symmetry	260
10.5	Recursion Relations for the $9j$ Symbols	261
10.5.1	General Form of the Recursion Relation	261

10.5.2	Relations Involving Four $9j$ Symbols	261
10.5.3	Relations Involving Five $9j$ Symbols [20]	262
10.5.4	Relation Involving Six $9j$ Symbols	263
10.5.5	Recursion Relations for Some Special Cases	264
10.6	Generating Function of the $9j$ Symbols	265
10.7	Asymptotic Expression for a $9j$ Symbol	265
10.8	Explicit Forms of the $9j$ Symbols at Some Relations Between Arguments	266
10.8.1	Two Rows (or Columns) are Identical	266
10.8.2	One Degenerate Triad	266
10.8.3	Two Degenerate Triads	267
10.8.4	Three Degenerate Triads	269
10.8.5	Four Degenerate Triads	269
10.8.6	Five or Six Degenerate Triads	270
10.9	Explicit Forms of the $9j$ Symbols for Special Values of the Arguments	271
10.9.1	One of Arguments Equals Zero	271
10.9.2	One of Arguments Equals Unity [11]	272
10.9.3	One of Triads Equals ($1/2, 1/2, 1$)	272
10.10	Relations Between the Wigner $9j$ Symbols and Analogous Functions of Other Authors	273
10.11	Tables of Algebraic Formulas of the $9j$ Symbols	273
10.12	Tables of Numerical Values of the $9j$ Symbols	273
10.13	$12j$ Symbols	274
10.13.1	Some Remarks on the $3nj$ Symbols	274
10.13.2	$12j$ Symbols of the First Kind ($12j(\text{I})$ -Symbols)	275
10.13.3	$12j$ Symbols of the Second Kind ($12j(\text{II})$ Symbols)	279
11	The graphical method in angular momentum theory	285
11.1	Graphical representation of functions	285
11.1.1	Basic Elements of Diagrams	285
11.1.2	Diagrams of the Basic Functions of the Theory	286
11.2	Graphical representation of the main operations of the theory	287
11.2.1	Multiplication	287
11.2.2	Invariant Summation over Projections	287
11.2.3	Summation over Angular Momentum	289
11.2.4	Invariant Integration over Directions	293
11.2.5	Integration over Rotation Parameters	293
11.3	Rules of the graphical technique	295
11.3.1	Deformation of Diagrams	295
11.3.2	Change of Node Sign	295
11.3.3	Change of Direction of External Lines	298
11.3.4	Change of Direction of Integral Lines	298
11.3.5	Linking Subdiagrams	299
11.3.6	Cutting Diagram into Subdiagrams	300
11.3.7	Graphical Method of Summation	301
11.3.8	Replacing An Internal Ω -Line by a Thick j -Line	302
11.3.9	Elimination of $j = 0$ Line	302
11.3.10	Elimination of the $R = 1$ Line	302
11.4	Summary of the graphical technique	303
11.4.1	General Properties of Diagrams	303
11.4.2	Generalized Wigner-Eckart Theorem in Diagrammatic Form	303
11.4.3	Scheme for the Application of the Graphical Technique	317
12	Sums involving vector addition and recoupling coefficients	321
12.1	Summation of products of $3jm$ symbols	321
12.1.1	Sum Involving One $3jm$ Symbol	321
12.1.2	Sums Involving Products of Two $3jm$ Symbols	321
12.1.3	Sums Involving Products of Three $3jm$ Symbols	322
12.1.4	Sums Involving Products of Four $3jm$ Symbols	323
12.1.5	Sums Involving Products of Five $3jm$ Symbols	324
12.1.6	Sums Involving Products of Six $3jm$ Symbols	325
12.2	Summation of products of $6j$ and $9j$ symbols	330
12.2.1	Sums Involving One $3nj$ Symbol	330

12.2.2	Sums Involving Products of Two $3nj$ Symbols	331
12.2.3	Sums Involving Products of Three $3nj$ Symbols	333
12.2.4	Sums Involving Products of Four $3nj$ Symbols	337
13	Matrix elements of irreducible tensor operators	343
13.1	The Wigner-Eckart theorem and the evaluation of matrix elements	343
13.1.1	Wigner-Eckart Theorem	343
13.1.2	Sum Rules	344
13.1.3	Matrix Elements of Products of Irreducible Tensor Operators	344
13.1.4	Matrix Elements of Operators Which Depend on Variables of Two Subsystems	346
13.1.5	Matrix Elements of Operators Which Depend on Variables of One of the Subsystems	350
13.2	Matrix elements of basic tensor operators	352
13.2.1	Some Introductory Remarks	352
13.2.2	Matrix Elements of the Unit Operator $\hat{\mathbf{I}}$	353
13.2.3	Matrix Elements of the Unit Vector $\hat{\mathbf{n}}(\vartheta, \varphi) \equiv \hat{\mathbf{n}}_1(\vartheta, \varphi)$	353
13.2.4	Matrix Elements of the Operator $\hat{\mathbf{V}}(r, \vartheta, \varphi) \equiv \hat{\mathbf{V}}_1(r, \vartheta, \varphi)$	355
13.2.5	Matrix Elements of the Total Angular Momentum Operator $\hat{\mathbf{J}} \equiv \hat{\mathbf{J}}_1$	357
13.2.6	Matrix Elements of the Orbital Angular Momentum Operator $\hat{\mathbf{L}} \equiv \hat{\mathbf{L}}_1$	360
13.2.7	Matrix Elements of the Spin Angular Momentum Operator $\hat{\mathbf{S}} \equiv \hat{\mathbf{S}}_1$	363
13.2.8	Matrix Elements of the Spherical Harmonic Operator $\hat{Y}_{L\nu} \equiv Y_{L\nu}(\vartheta, \varphi)$	364
13.2.9	Matrix Elements of Some Scalar and Vector Products	370
14	Computer generation of numerical and algebraic expressions	375
14.1	Scope	375
14.2	Algebraic generation	375
14.2.1	The Clebsch-Gordan coefficients	375
14.2.2	The $6j$ symbols	376
14.2.3	The $9j$ symbols	376
14.3	Numerical evaluation	376
14.4	Verification	377
14.5	The programs	377
14.6	Use of the notebooks	377
14.6.1	Algebraic expressions	377
14.6.2	Numerical values	377

Originally published by

World Scientific Publishing Co. Pte. Ltd.

5 Toh Tuck Link, Singapore 596224

USA office: 27 Warren Street, Suite 401-402, Hackensack, NJ 07601

UK office: 57 Shelton Street, Covent Garden, London WC2H 9HE

Open Access

This project is open access and is licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

It is based on the original work published by World Scientific Publishing Co. Pte. Ltd., Singapore, 1988, ISBN 978-9971-5-0107-5 (hardcover), ISBN 978-9971-5-0996-5 (paperback), ISBN 978-981-4415-49-1 (ebook for institutions), ISBN 978-981-4578-28-8 (ebook for individuals).

Open Access funded by SCOAP (Sponsoring Consortium for Open Access Publishing in Particle Physics)

The original ebook was converted to Open Access in 2021 through the sponsorship of SCOAP , licensed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made. Original copyright © 1988 by World Scientific Publishing Co. Pte. Ltd.

Preface

In the years since its publication, this book has become the standard reference on the quantum theory of angular momentum. The book provides a comprehensive treatment of the subject, covering both the theoretical foundations and practical applications of angular momentum in quantum mechanics. As an open access publication, it is freely available to researchers and students worldwide, ensuring that the knowledge contained within its pages can be accessed by anyone interested in the field. The book has been widely cited and used in research and teaching, and it continues to be a valuable resource for those studying quantum mechanics and related areas of physics.

However, there are some drawbacks to the form in which it was provided. The scanned version of the book is not searchable, and the quality of the scanned images is not always optimal. This can make it difficult to find specific information or to read the text clearly. Additionally, the scanned version may not be compatible with all devices or software, which can limit its accessibility. Also, there is no verification of the accuracy of the scanned text, which can lead to errors or misinterpretations of the content. Finally, the scanned version may not be easily editable or adaptable for different purposes, such as creating new editions or translations.

Thus this attempt to provide a more accessible and user-friendly version of the book, with improved searchability, image quality, and compatibility with different devices and software. The text has been carefully reviewed and verified for accuracy, and it is provided in a format that can be easily edited or adapted for different purposes. We hope that this version of the book will be a valuable resource for researchers, students, and anyone interested in the quantum theory of angular momentum. The author would like to thank the original publisher, World Scientific Publishing Co. Pte. Ltd., for their support in making this open access version available to the public.

Clearly in the era of AI, I have made substantial use of AI tools to assist in the preparation of this version of the book. I have used AI tools to help with tasks such as text recognition, formatting, and editing. These tools have helped to improve the accuracy and quality of the text, as well as to streamline the process generating code to verify equations. However, I have also carefully reviewed and verified the output of these tools to ensure that the final version of the book is as accurate and reliable. Overall, I believe that the use of AI tools has been a valuable aid in the preparation of this new open access version of the book.

Of course, the author would like to acknowledge the staggering effort in authoring the original work by D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, and that this new version is very closely based on their work. The author has made every effort to ensure that the content of this new version is faithful to the original, while also improving its accessibility and usability for modern readers.

I do not see this as a static resource—there are undoubtedly errors in the text, and I would be grateful for any corrections or suggestions for improvement. Please post these as an “[issue](#)” at the GitHub repository for this project at <https://github.com/nwalet/Varshalovich>. The intention is to improve the accessibility and usability of this important work, and I welcome any feedback, and contributions, that can help achieve that goal.

Manchester, UK, September 2026

acknowledgements

The following colleagues have provided corrections: Wes Campbell, UCLA;

Original Preface

This book deals with one of the basic topics of quantum mechanics: the theory of angular momentum and irreducible tensors. Being rather versatile, the mathematical apparatus of this theory is widely used in atomic and molecular physics, in nuclear physics and elementary particle theory. It enables one to calculate atomic, molecular and nuclear structures, energies of ground and excited states, fine and hyperfine splittings, etc. The apparatus is also very handy for evaluating the probabilities of radiative transitions, cross sections of various processes such as elastic and nonelastic scattering, different decays and reactions (both chemical and nuclear) and for studying angular distributions and polarizations of particles.

Today this apparatus is finding ever increasing use in solving practical problems relating to quantum chemistry, kinetics, plasma physics, quantum optics, radiophysics and astrophysics.

The basic ideas of the theory of angular momentum were first put forward by M. Born, P. Dirac, W. Heisenberg and W. Pauli. However, the modern version of its mathematical apparatus was developed mainly in the works of E. Wigner, J. Racah, L. Biedenharn and others who applied group theoretical methods to problems in quantum mechanics.

At present a number of good books on the theory of angular momentum have been already published. The general principles and results of the theory may be found in the books by M. Rose [31], A. Edmonds [16], U. Fano

and G. Racah [18], A. P. Yutsis, I. B. Levinson and V. V. Vanagas [44], A. P. Yutsis and A. A. Bandzaitis [45], D. Brink and G. Satcher [9]. Nevertheless, many formulas and relationships essential for practical calculations have escaped these books and are either scattered in various editions, or included as appendices in papers discussing somewhat disparate topics, making them generally inaccessible. Even greater difficulties arise when one tries to use the results, as each author employs his own phase conventions, initial definitions and symbols.

The authors of this book aimed at collecting and compiling ample material on the quantum theory of angular momentum within the framework of a single system of phases and definitions. This is why, in addition to the basic theoretical results, the book also includes a great number of formulas and relationships essential for practical applications.

This edition is the translated version of our book published in the USSR in 1975. In the course of its preparation we have tried to comply with a number of suggestions from our readers. For instance, each chapter opens with a comprehensive listing of its contents to ease the search for information needed. We also included some new results relating to different aspects of angular momentum theory which have recently appeared in journals. Unfortunately the limited volume of the present book prevented us from covering all the aforementioned results. We offer sincere apologies to the authors whose results we failed to include.

The monograph is a kind of handbook. Consequently the material is presented in concise form. Most of the formulas and relationships are given without proof. Their full derivation may be found in the literature listed at the end of the text. Some results which have become generally known are given without references, and for this we also apologize.

The sequence adopted is as follows: chapter, section and subsection. Many chapters are self-contained and can be read independently of the others. Sections have double numbering: the first figure denotes the number of the chapter, the second, the number of the section. Equations are numbered within the confines of the section they are included in. When referring to an equation from the same section only the number of the equation is given, e.g., (3), (27); when reference is made to an equation from another section the numbers of the chapter, section and equation are given, e.g., Eq. 4.2.(17). A similar system is adopted when referring to individual subsections, e.g., Sec. 1.2.5. For convenience the book also contains a glossary of all symbols used in the text with references to the pages where their corresponding definitions are given. The list of references is divided into parts: the first part lists books and reviews; the second, papers on different subjects; the third, tables; the fourth, references added during translation.

The authors hope that many specialists will find in the book some fresh and interesting information. The material is prepared and arranged so as to make it useful to those less familiar with theory and for students of physics. These readers can effectively use the monograph as a supplementary text to their main courses.

For those who wish to thoroughly familiarize themselves with the fundamentals of angular momentum theory we recommend the excellent new book by L. Biedenharn and J. Louck [132] *Angular Momentum in Quantum Physics. Theory and Applications*.

The authors wish to express their deep appreciation to D. G. Yakovlev who took the trouble of reading the English translation of the book and gave some valuable suggestions on its preparation.

Leningrad

INTRODUCTION: BASIC CONCEPTS

The evaluation of many physical quantities, such as expectation values of energy, electric and magnetic multipole moments, transition probabilities, etc., is considerably simplified by making use of the transformation properties of these quantities under coordinate rotation and inversion in three-dimensional space.

The transformation properties of physical quantities with respect to rotations reveal themselves either through rotations of the given physical system relative to some fixed reference frame or through rotations of the coordinate axes relative to the physical system.

Inversion characterizes the behavior of a physical quantity under transformation from a right-handed coordinate system to a left-handed one, or vice versa.

Many physical quantities, by their nature, are invariants under coordinate rotations. In particular, the properties of any closed physical system should be independent of rotations, as follows from the isotropy of space. As a consequence of this fundamental property of space, the total angular momentum of such a system is an integral of motion.

A similar situation occurs with regard to coordinate inversion. Excluding phenomena connected with the weak interaction, a wealth of atomic, molecular and nuclear processes look alike in right-handed and left-handed coordinate systems. As a result of this "mirror" symmetry, quantum states of atoms, molecules, nuclei and elementary particles may be characterized by definite parity.

Strictly speaking, a quantum mechanical wave function $\Psi(r)$ of any closed physical system may be characterized by four quantum numbers (ε, π, j, m) which are the eigenvalues of four commuting operators: the Hamiltonian $\hat{H}$, the parity operator $\hat{P}_r$, the operator $\hat{\mathbf{J}}^2$ of the square of the angular momentum and the operator $\hat{J}_z$ of the projection of this momentum onto a quantization axis. Thus $\Psi(r) \equiv \Psi_{\varepsilon\pi\alpha jm}(r)$ obeys the equations

$$\begin{aligned}\hat{H}\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}) &= \varepsilon\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}), \\ \hat{P}_r\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}) &= \pi\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}), \\ \hat{\mathbf{J}}^2\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}) &= j(j+1)\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}), \\ \hat{J}_z\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}) &= m\Psi_{\varepsilon\pi\alpha jm}(\mathbf{r}),\end{aligned}$$

where α denotes all other quantum numbers (if available) and r represents a variety of arguments.

ε, π, j and m are the integrals of motion not only for a closed system, but also for any system exposed to the action of an external spherically-symmetric field. Moreover, the $(\varepsilon\pi jm)$ -representation appears to be very convenient for practical calculation even if the external field is not spherically-symmetric.

For a given j , there exist $2j+1$ wave functions which correspond to different m . These functions describe the quantum states of the system which differ only by the orientation of the angular momentum in space.

Under coordinate rotations these functions undergo linear mutual transformations which do not involve the functions of other quantum numbers $(\varepsilon, \pi, \alpha, j)$ and may be written as

$$\Psi_{c\pi\alpha jm'}(\mathbf{r}') = \sum_m \Psi_{c\pi\alpha jm}(\mathbf{r}) D_{mm'}^j(\alpha, \beta, \gamma),$$

where the transformations coefficients $D_{mm'}^j(\alpha, \beta, \gamma)$ are called Wigner D -functions. They are the elements of the finite rotation matrix in the j -representation and depend on the Euler angles α, β, γ which determine rotation.

Let a quantum-mechanical system which possesses some fixed angular momentum j and its projection m consist of two subsystems, with certain angular momenta j_1 and j_2 , respectively. In this case the wave function $\Psi_{j_1 j_2 jm}(\mathbf{r}_1, \mathbf{r}_2)$ may be constructed from the wave functions of subsystems according to the relation

$$\Psi_{j_1 j_2 jm}(\mathbf{r}_1, \mathbf{r}_2) = \sum_{m_1 m_2} C_{j_1 m_1 j_2 m_2}^{j m} \Psi_{j_1 m_1}(\mathbf{r}_1) \Psi_{j_2 m_2}(\mathbf{r}_2).$$

The quantities $C_{j_1 m_1 j_2 m_2}^{j m}$ are called the Clebsch-Gordan coefficients. They are very important in quantum mechanics because they allow one to construct wave functions for various complex systems (nuclei, atoms, molecules, etc.). The addition of many angular momenta into some resultant total angular momentum may be performed in different ways, or, in other words, in accordance with different coupling schemes of the momenta in question. Each scheme may be associated with a certain representation of these angular momenta. The unitary transformations which relate various representations and describe the recoupling of angular momenta are realized by matrices expressed in terms of the $6j$ - $9j$ -symbols and others $3nj$ -symbols of higher order.

One of the principal concepts of the quantum theory of angular momentum is that of an irreducible tensor. By definition, an irreducible tensor of rank λ has $2\lambda + 1$ components which transform under coordinate rotation as

$$\mathfrak{M}_{\lambda\mu} = \sum_{\mu'} \mathfrak{M}_{\lambda\mu'} D_{\mu'\mu}^{\lambda}(\alpha, \beta, \gamma).$$

In particular, the wave functions Ψ_{jm} with fixed j but different m constitute an irreducible tensor of rank j . Moreover, irreducible tensors may be constructed from various physical quantities. For instance, energy is a tensor of zero rank (scalar), spin and magnetic moments are tensors of first rank (vectors), the quadrupole moment is a tensor of second rank, etc. Generally, any physical quantity or operator which corresponds to these quantities may be represented as a linear combination of irreducible tensors.

The introduction of irreducible tensors into the analysis of physical quantities or corresponding operators substantially simplifies the evaluation of matrix elements of irreducible tensor operators,

$$\langle \varepsilon' \pi' \alpha' j' m' | \widehat{\mathfrak{M}}_{\lambda\mu} | \varepsilon \pi \alpha j m \rangle = \int \Psi_{\varepsilon' \pi' \alpha' j' m'}^*(\mathbf{r}) \widehat{\mathfrak{M}}_{\lambda\mu} \Psi_{\varepsilon \pi \alpha j m}(\mathbf{r}) d\mathbf{r},$$

where $\int \dots d\mathbf{r}$ denotes integration over continuous variables and summation over discrete ones. These matrix elements determine expectation values of physical quantities in definite quantum states, the probabilities of quantum transitions between various states, etc.

According to the Wigner-Eckart theorem, all the dependence of matrix elements on the orientation of coordinate axes, i.e., on quantum numbers m, m', μ which determine the projections of angular momenta, is entirely contained in the Clebsch-Gordan coefficient

$$\langle \varepsilon' \pi' \alpha' j' m' | \widehat{\mathfrak{M}}_{\lambda\mu} | \varepsilon \pi \alpha j m \rangle = C_{j m \lambda \mu}^{j' m'} \langle \varepsilon' \pi' \alpha' j' | | \widehat{\mathfrak{M}}_{\lambda} | | \varepsilon \pi \alpha j \rangle.$$

In this case $\langle \varepsilon' \pi' \alpha' j' | | \widehat{\mathfrak{M}}_{\lambda} | | \varepsilon \pi \alpha j \rangle$ is an invariant factor called a reduced matrix element. The Wigner-Eckart theorem allows one to reduce evaluation of transition probabilities, angular distributions, polarizations, etc., to the calculation of standard sums of the vector addition coefficients and recoupling coefficients.

All the facts mentioned above reveal that the quantum theory of angular momentum provides the formalism which is universal and extremely convenient for various practical calculations.

Chapter 1

Elements of Vector and Tensor Theory

The theory of angular momenta and irreducible tensors represents, in principle, a development of the classical theory of vectors and tensors. In this chapter only the basic definitions and relations of the vector and tensor theory are represented which will be used throughout. For more detailed analysis see corresponding monographs (e.g., Refs. [1, 2, 3]).

1.1 Coordinate Systems. Basis Vectors

In the quantum theory of angular momentum cartesian, polar and spherical coordinate systems are widely used.

1.1.1 Cartesian Coordinate System

In a rectangular cartesian coordinate system the position of a point is specified by three real numbers x, y, z which represent the distances between the point and coordinate planes (Fig. 1.1). The position vector (radius vector) of a point $\mathbf{r}$ may be written as

$$\mathbf{r} = x\mathbf{e}_x + y\mathbf{e}_y + z\mathbf{e}_z. \quad (1.1.1)$$




Figure 1.1. Cartesian coordinate system.

The covariant cartesian basis (base) vectors $\mathbf{e}_x, \mathbf{e}_y, \mathbf{e}_z$ form a real orthonormal basis

$$\mathbf{e}_i \mathbf{e}_k = \delta_{ik}, (i, k = x, y, z), \quad (1.1.2)$$

$$\mathbf{e}_i^* = \mathbf{e}_i, (i = x, y, z). \quad (1.1.3)$$

The contravariant cartesian basis (base) vectors $\mathbf{e}^i (i = x, y, z)$ coincide with the covariant ones

$$\mathbf{e}^i = \mathbf{e}_i. \quad (1.1.4)$$

Throughout this book the right-handed coordinate system will be used. In this system

$$[\mathbf{e}_i \times \mathbf{e}_k] = \varepsilon_{ikl} \mathbf{e}_l, \quad (i, k, l = x, y, z), \quad (1.1.5)$$

$$\varepsilon_{ikl} = [\mathbf{e}_i \times \mathbf{e}_k] \cdot \mathbf{e}_l. \quad (1.1.6)$$

A detailed form of (1.1.5) is

$$[\mathbf{e}_x \times \mathbf{e}_y] = \mathbf{e}_z, \quad [\mathbf{e}_y \times \mathbf{e}_z] = \mathbf{e}_x, \quad [\mathbf{e}_z \times \mathbf{e}_x] = \mathbf{e}_y. \quad (1.1.7)$$

1.1.2 Polar Coordinate System

In a polar coordinate system ¹ the position of a point is determined by r, ϑ, φ , where r is the position vector length, ϑ is the colatitude, and φ is the longitude (Fig. 1.2). The angles ϑ and φ are called the polar angles of vector $\mathbf{r}$. The relations between cartesian and polar coordinates are




Figure 1.2. Polar coordinate system.

$$\begin{aligned} x &= r \sin \vartheta \cos \varphi, & r &= \sqrt{x^2 + y^2 + z^2}, & 0 &\leq r < \infty, \\ y &= r \sin \vartheta \sin \varphi, & \vartheta &= \arccos \frac{z}{\sqrt{x^2 + y^2 + z^2}}, & 0 &\leq \vartheta \leq \pi, \\ z &= r \cos \vartheta, & \varphi &= \arccos \frac{x}{\sqrt{x^2 + y^2}}, \left(\tan \varphi = \frac{y}{x} \right), & 0 &\leq \varphi < 2\pi. \end{aligned} \quad (1.1.8)$$

The position vector r may be written as

$$\mathbf{r} = r \mathbf{e}_r. \quad (1.1.9)$$

The covariant polar basis vectors $\mathbf{e}_r, \mathbf{e}_\vartheta, \mathbf{e}_\varphi$ are shown in Fig. 1.2. They form a real orthonormal basis

$$\mathbf{e}_\alpha \mathbf{e}_\beta = \delta_{\alpha\beta}, \quad (\alpha, \beta = r, \vartheta, \varphi), \quad (1.1.10)$$

$$\mathbf{e}_\alpha^* = \mathbf{e}_\alpha, \quad (\alpha = r, \vartheta, \varphi). \quad (1.1.11)$$

The contravariant polar basis vectors $\mathbf{e}^r, \mathbf{e}^\vartheta, \mathbf{e}^\varphi$ coincide with the covariant ones

$$\mathbf{e}^\alpha = \mathbf{e}_\alpha \quad (\alpha = r, \vartheta, \varphi). \quad (1.1.12)$$

The unit vectors $\mathbf{e}_r, \mathbf{e}_\vartheta, \mathbf{e}_\varphi$ form the right-handed basis

$$[\mathbf{e}_r \times \mathbf{e}_\vartheta] = \mathbf{e}_\varphi, \quad [\mathbf{e}_\vartheta \times \mathbf{e}_\varphi] = \mathbf{e}_r, \quad [\mathbf{e}_\varphi \times \mathbf{e}_r] = \mathbf{e}_\vartheta. \quad (1.1.13)$$

¹ Note that this coordinate system is often called “spherical”. To avoid misunderstanding we prefer to call it the “polar” system reserving the name “spherical” only for the coordinate system considered in the next section.

The polar basis vectors $\mathbf{e}_r, \mathbf{e}_\vartheta, \mathbf{e}_\varphi$, contrary to the cartesian ones, depend on the angles ϑ, φ . This should be taken into account when evaluating the derivatives

$$\begin{aligned} \frac{\partial}{\partial r} \mathbf{e}_r &= 0, & \frac{\partial}{\partial r} \mathbf{e}_\vartheta &= 0, & \frac{\partial}{\partial r} \mathbf{e}_\varphi &= 0, \\ \frac{\partial}{\partial \vartheta} \mathbf{e}_r &= \mathbf{e}_\vartheta, & \frac{\partial}{\partial \vartheta} \mathbf{e}_\vartheta &= -\mathbf{e}_r, & \frac{\partial}{\partial \vartheta} \mathbf{e}_\varphi &= 0, \\ \frac{\partial}{\partial \varphi} \mathbf{e}_r &= \mathbf{e}_\varphi \sin \vartheta, & \frac{\partial}{\partial \varphi} \mathbf{e}_\vartheta &= \mathbf{e}_\varphi \cos \vartheta, & \frac{\partial}{\partial \varphi} \mathbf{e}_\varphi &= -\mathbf{e}_r \sin \vartheta - \mathbf{e}_\vartheta \cos \vartheta. \end{aligned} \quad (1.1.14)$$

The results of applying the ∇ operator (see Sec. 1.3) to the polar basis vectors are presented in the form

$$(\nabla \cdot \mathbf{e}_r) = \frac{2}{r}, \quad (\nabla \cdot \mathbf{e}_\vartheta) = \frac{1}{r} \cot \vartheta, \quad (\nabla \cdot \mathbf{e}_\varphi) = 0, \quad (1.1.15)$$

$$[\nabla \times \mathbf{e}_r] = 0, \quad [\nabla \times \mathbf{e}_\vartheta] = \frac{1}{r} \mathbf{e}_\varphi, \quad [\nabla \times \mathbf{e}_\varphi] = \frac{1}{r} \cot \vartheta \mathbf{e}_r - \frac{1}{r} \mathbf{e}_\vartheta. \quad (1.1.16)$$

1.1.3 Spherical Coordinate System

Spherical coordinates are widely used in the angular momentum theory.

The covariant spherical coordinates x_μ (with $\mu = \pm 1, 0$) are defined by the relations

$$\begin{aligned} x_{+1} &= -\frac{1}{\sqrt{2}}(x + iy) = -\frac{1}{\sqrt{2}}r \sin \vartheta e^{i\varphi}, \\ x_0 &= z = r \cos \vartheta, \\ x_{-1} &= \frac{1}{\sqrt{2}}(x - iy) = \frac{1}{\sqrt{2}}r \sin \vartheta e^{-i\varphi}. \end{aligned} \quad (1.1.17)$$

The contravariant spherical coordinates x^μ (where $\mu = \pm 1, 0$) are given by

$$\begin{aligned} x^{+1} &= -\frac{1}{\sqrt{2}}(x - iy) = -\frac{1}{\sqrt{2}}r \sin \vartheta e^{-i\varphi}, \\ x^0 &= z = r \cos \vartheta, \\ x^{-1} &= \frac{1}{\sqrt{2}}(x + iy) = \frac{1}{\sqrt{2}}r \sin \vartheta e^{i\varphi}. \end{aligned} \quad (1.1.18)$$

The relations between covariant and contravariant spherical coordinates are as follows

$$\begin{aligned} x^\mu &= (-1)^\mu x_{-\mu}, & x_\mu &= (-1)^\mu x^{-\mu}, \\ x^\mu &= x_\mu^*, & x_\mu &= x^{\mu*}, \end{aligned} \quad (\mu = \pm 1, 0). \quad (1.1.19)$$

The covariant spherical basis vectors $\mathbf{e}_\mu$ ($\mu = \pm 1, 0$) are defined by

$$\begin{aligned} \mathbf{e}_{+1} &= -\frac{1}{\sqrt{2}}(\mathbf{e}_x + i\mathbf{e}_y), \\ \mathbf{e}_0 &= \mathbf{e}_z, \\ \mathbf{e}_{-1} &= \frac{1}{\sqrt{2}}(\mathbf{e}_x - i\mathbf{e}_y). \end{aligned} \quad (1.1.20)$$

The contravariant spherical basis vectors $\mathbf{e}^\mu$ ($\mu = \pm 1, 0$) are given by

$$\begin{aligned} \mathbf{e}^{+1} &= -\frac{1}{\sqrt{2}}(\mathbf{e}_x - i\mathbf{e}_y), \\ \mathbf{e}^0 &= \mathbf{e}_z, \\ \mathbf{e}^{-1} &= \frac{1}{\sqrt{2}}(\mathbf{e}_x + i\mathbf{e}_y). \end{aligned} \quad (1.1.21)$$

Relations between the covariant and contravariant spherical basis vectors read

$$\begin{aligned} \mathbf{e}^\mu &= (-1)^\mu \mathbf{e}_{-\mu}, & \mathbf{e}_\mu &= (-1)^\mu \mathbf{e}^{-\mu}, \\ \mathbf{e}^\mu &= \mathbf{e}_\mu^*, & \mathbf{e}_\mu &= \mathbf{e}^{\mu*}, \end{aligned} \quad (\mu = \pm 1, 0). \quad (1.1.22)$$

The spherical basis vectors form a complex orthonormal basis

$$\mathbf{e}_\mu \mathbf{e}^\nu = \mathbf{e}_\mu \mathbf{e}_\nu^* = \delta_{\mu\nu}, \quad (\mu, \nu = \pm 1, 0). \quad (1.1.23)$$

Vector products of spherical basis vectors may be written with the use of the Clebsch-Gordan coefficients (see Chap. 8) in the form

$$\begin{aligned} \mathbf{e}_\mu \times \mathbf{e}_\nu &= i\sqrt{2}C_{1\mu 1\nu}^{1\lambda} \mathbf{e}_\lambda, \\ \mathbf{e}^\mu \times \mathbf{e}^\nu &= -i\sqrt{2}C_{1\mu 1\nu}^{1\lambda} \mathbf{e}^\lambda, \end{aligned} \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (1.1.24)$$

One may also rewrite these formulas in a form similar to (1.1.5)

$$\begin{aligned} \mathbf{e}_\mu \times \mathbf{e}_\nu &= -i\varepsilon_{\mu\nu\lambda} \mathbf{e}^\lambda, \quad (\mu, \nu, \lambda = \pm 1, 0) \\ \mathbf{e}^\mu \times \mathbf{e}^\nu &= i\varepsilon_{\mu\nu\lambda} \mathbf{e}_\lambda, \end{aligned} \quad (\mu, \nu, \lambda = \pm 1, 0), \quad (1.1.25)$$

where $\varepsilon_{\mu\nu\lambda} = +1$ if the combination of indices μ, ν, λ is obtained by an even permutation of $+1, 0, -1$, $\varepsilon_{\mu\nu\lambda} = -1$ for an odd permutation, and $\varepsilon_{\mu\nu\lambda} = 0$ if at least two indices among μ, ν, λ are equal.

A detailed form of (1.1.25) is as follows:

$$\begin{aligned} \mathbf{e}_{+1} \times \mathbf{e}_0 &= i\mathbf{e}_{+1}, & \mathbf{e}_0 \times \mathbf{e}_{-1} &= i\mathbf{e}_{-1}, \\ \mathbf{e}^0 \times \mathbf{e}_{+1}^+ &= i\mathbf{e}_{-1}^+ = i\mathbf{e}_0^+, & \mathbf{e}^{-1} \times \mathbf{e}^0 &= i\mathbf{e}^{-1}, \\ \mathbf{e}^{-1} \times \mathbf{e}^{+1} &= i\mathbf{e}^0. \end{aligned} \quad (1.1.26)$$

Covariant and contravariant spherical components (see Sec. 1.2) of the basis vectors $\mathbf{e}_\mu$ and $\mathbf{e}^\mu$ are given by

$$\begin{aligned} [\mathbf{e}_\mu]_\sigma &= (-1)^\sigma \delta_{\sigma-\mu}, & [\mathbf{e}_\mu]^\sigma &= \delta_{\sigma\mu} \\ [\mathbf{e}^\mu]_\sigma &= \delta_{\sigma\mu}, & [\mathbf{e}^\mu]^\sigma &= (-1)^\sigma \delta_{\sigma-\mu}. \end{aligned} \quad (1.1.27)$$

1.1.4 Helicity Basis Vectors

By analogy with Eqs. (1.1.20), (1.1.21) one may construct the following combinations of the polar basis vectors $\mathbf{e}_r, \mathbf{e}_\vartheta, \mathbf{e}_\varphi$

$$\begin{aligned} \mathbf{e}'_{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta + i\mathbf{e}_\varphi), & \mathbf{e}'^{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta - i\mathbf{e}_\varphi), \\ \mathbf{e}'_0 &= \mathbf{e}_r, & \mathbf{e}'^0 &= \mathbf{e}_r, \\ \mathbf{e}'_{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta - i\mathbf{e}_\varphi), & \mathbf{e}'^{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta + i\mathbf{e}_\varphi). \end{aligned} \quad (1.1.28)$$

The vectors $\mathbf{e}'_\mu (\mu = \pm 1, 0)$ are called the covariant helicity basis vectors and $\mathbf{e}'^\mu (\mu = \pm 1, 0)$ are called the contravariant helicity basis vectors (the explanation of the term “helicity” is given below in Sec. 6.3.6).

The helicity basis vectors $\mathbf{e}'_\mu$ and $\mathbf{e}'^\mu$ satisfy the same relations (1.1.22)-(1.1.26) as the spherical basis vectors $\mathbf{e}_\mu$ and $\mathbf{e}^\mu$.

1.1.5 Relations Between Different Basis Vectors

(a) Cartesian vectors and other basis vectors

Cartesian and Polar Basis Vectors

$$\begin{aligned} \mathbf{e}_x &= \mathbf{e}_r \sin \vartheta \cos \varphi + \mathbf{e}_\vartheta \cos \vartheta \cos \varphi - \mathbf{e}_\varphi \sin \varphi, \\ \mathbf{e}_y &= \mathbf{e}_r \sin \vartheta \sin \varphi + \mathbf{e}_\vartheta \cos \vartheta \sin \varphi + \mathbf{e}_\varphi \cos \varphi, \\ \mathbf{e}_z &= \mathbf{e}_r \cos \vartheta - \mathbf{e}_\vartheta \sin \vartheta. \end{aligned} \quad (1.1.29)$$

Cartesian and Spherical Basis Vectors

$$\begin{aligned} \mathbf{e}_x &= \frac{1}{\sqrt{2}} (\mathbf{e}_{-1} - \mathbf{e}_{+1}) = \frac{1}{\sqrt{2}} (\mathbf{e}^{-1} - \mathbf{e}^{+1}), \\ \mathbf{e}_y &= \frac{i}{\sqrt{2}} (\mathbf{e}_{-1} + \mathbf{e}_{+1}) = -\frac{i}{\sqrt{2}} (\mathbf{e}^{-1} + \mathbf{e}^{+1}), \\ \mathbf{e}_z &= \mathbf{e}_0 = \mathbf{e}^0. \end{aligned} \quad (1.1.30)$$

Cartesian and Helicity Covariant Basis Vectors

$$\begin{aligned} \mathbf{e}_x &= -\mathbf{e}'_{+1} \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi + i \sin \varphi) + \mathbf{e}'_0 \sin \vartheta \cos \varphi + \mathbf{e}'_{-1} \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi - i \sin \varphi), \\ \mathbf{e}_y &= -\mathbf{e}'_{+1} \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi - i \cos \varphi) + \mathbf{e}'_0 \sin \vartheta \sin \varphi + \mathbf{e}'_{-1} \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi + i \cos \varphi), \\ \mathbf{e}_z &= \mathbf{e}'_{+1} \frac{1}{\sqrt{2}} \sin \vartheta + \mathbf{e}'_0 \cos \vartheta - \mathbf{e}'_{-1} \frac{1}{\sqrt{2}} \sin \vartheta. \end{aligned} \quad (1.1.31)$$

Cartesian and Helicity Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_x &= -\mathbf{e}'^{+1} \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi - i \sin \varphi) + \mathbf{e}'^0 \sin \vartheta \cos \varphi + \mathbf{e}'^{-1} \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi + i \sin \varphi), \\
\mathbf{e}_y &= -\mathbf{e}'^{+1} \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi + i \cos \varphi) + \mathbf{e}'^0 \sin \vartheta \sin \varphi + \mathbf{e}'^{-1} \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi - i \cos \varphi), \\
\mathbf{e}_z &= \mathbf{e}'^{+1} \frac{1}{\sqrt{2}} \sin \vartheta + \mathbf{e}'^0 \cos \vartheta - \mathbf{e}'^{-1} \frac{1}{\sqrt{2}} \sin \vartheta.
\end{aligned} \tag{1.1.32}$$

(b) Polar Vectors and other basis vectors

Polar and Cartesian Basis Vectors

$$\begin{aligned}
\mathbf{e}_r &= \mathbf{e}_x \sin \vartheta \cos \varphi + \mathbf{e}_y \sin \vartheta \sin \varphi + \mathbf{e}_z \cos \vartheta, \\
\mathbf{e}_\vartheta &= \mathbf{e}_x \cos \vartheta \cos \varphi + \mathbf{e}_y \cos \vartheta \sin \varphi - \mathbf{e}_z \sin \vartheta, \\
\mathbf{e}_\varphi &= -\mathbf{e}_x \sin \varphi + \mathbf{e}_y \cos \varphi.
\end{aligned} \tag{1.1.33}$$

Polar and Spherical Covariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_r &= -\mathbf{e}_{+1} \frac{1}{\sqrt{2}} \sin \vartheta e^{-i\varphi} + \mathbf{e}_0 \cos \vartheta + \mathbf{e}_{-1} \frac{1}{\sqrt{2}} \sin \vartheta e^{i\varphi}, \\
\mathbf{e}_\vartheta &= -\mathbf{e}_{+1} \frac{1}{\sqrt{2}} \cos \vartheta e^{-i\varphi} - \mathbf{e}_0 \sin \vartheta + \mathbf{e}_{-1} \frac{1}{\sqrt{2}} \cos \vartheta e^{i\varphi}; \\
\mathbf{e}_\varphi &= \mathbf{e}_{+1} \frac{i}{\sqrt{2}} e^{-i\varphi} + \mathbf{e}_{-1} \frac{i}{\sqrt{2}} e^{i\varphi}.
\end{aligned} \tag{1.1.34}$$

Polar and Spherical Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_r &= -\mathbf{e}^{+1} \frac{1}{\sqrt{2}} \sin \vartheta e^{i\varphi} + \mathbf{e}^0 \cos \vartheta + \mathbf{e}^{-1} \frac{1}{\sqrt{2}} \sin \vartheta e^{-i\varphi}, \\
\mathbf{e}_\vartheta &= -\mathbf{e}^{+1} \frac{1}{\sqrt{2}} \cos \vartheta e^{i\varphi} - \mathbf{e}^0 \sin \vartheta + \mathbf{e}^{-1} \frac{1}{\sqrt{2}} \cos \vartheta e^{-i\varphi}, \\
\mathbf{e}_\varphi &= -\mathbf{e}^{+1} \frac{i}{\sqrt{2}} e^{i\varphi} - \mathbf{e}^{-1} \frac{i}{\sqrt{2}} e^{-i\varphi}.
\end{aligned} \tag{1.1.35}$$

Polar and Helicity Basis Vectors

$$\begin{aligned}
\mathbf{e}_r &= \mathbf{e}'_0 = \mathbf{e}'^0, \\
\mathbf{e}_\vartheta &= \frac{1}{\sqrt{2}} (\mathbf{e}'_{-1} - \mathbf{e}'_{+1}) = \frac{1}{\sqrt{2}} (\mathbf{e}'^{-1} - \mathbf{e}'^{+1}), \\
\mathbf{e}_\varphi &= \frac{i}{\sqrt{2}} (\mathbf{e}'_{-1} + \mathbf{e}'_{+1}) = -\frac{i}{\sqrt{2}} (\mathbf{e}'^{-1} + \mathbf{e}'^{+1}).
\end{aligned} \tag{1.1.36}$$

(c) Spherical Vectors and other basis vectors

Spherical and Cartesian Basis Vectors

$$\begin{aligned}
\mathbf{e}_{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_x + i\mathbf{e}_y), & \mathbf{e}^{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_x - i\mathbf{e}_y) \\
\mathbf{e}_0 &= \mathbf{e}_z, & \mathbf{e}^0 &= \mathbf{e}_z, \\
\mathbf{e}_{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_x - i\mathbf{e}_y), & \mathbf{e}^{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_x + i\mathbf{e}_y).
\end{aligned} \tag{1.1.37}$$

Spherical Covariant and Polar Basis Vectors

$$\begin{aligned}
\mathbf{e}_{+1} &= -\mathbf{e}_r \frac{1}{\sqrt{2}} \sin \vartheta e^{i\varphi} - \mathbf{e}_\vartheta \frac{1}{\sqrt{2}} \cos \vartheta e^{i\varphi} - \mathbf{e}_\varphi \frac{i}{\sqrt{2}} e^{i\varphi}, \\
\mathbf{e}_0 &= \mathbf{e}_r \cos \vartheta - \mathbf{e}_\vartheta \sin \vartheta, \\
\mathbf{e}_{-1} &= \mathbf{e}_r \frac{1}{\sqrt{2}} \sin \vartheta e^{-i\varphi} + \mathbf{e}_\vartheta \frac{1}{\sqrt{2}} \cos \vartheta e^{-i\varphi} - \mathbf{e}_\varphi \frac{i}{\sqrt{2}} e^{-i\varphi}.
\end{aligned} \tag{1.1.38}$$

Spherical Contravariant and Polar Basis Vectors

$$\begin{aligned}
\mathbf{e}^{+1} &= -\mathbf{e}_r \frac{1}{\sqrt{2}} \sin \vartheta e^{-i\varphi} - \mathbf{e}_\vartheta \frac{1}{\sqrt{2}} \cos \vartheta e^{-i\varphi} + \mathbf{e}_\varphi \frac{i}{\sqrt{2}} e^{-i\varphi}, \\
\mathbf{e}^0 &= \mathbf{e}_r \cos \vartheta - \mathbf{e}_\vartheta \sin \vartheta, \\
\mathbf{e}^{-1} &= \mathbf{e}_r \frac{1}{\sqrt{2}} \sin \vartheta e^{i\varphi} + \mathbf{e}_\vartheta \frac{1}{\sqrt{2}} \cos \vartheta e^{i\varphi} + \mathbf{e}_\varphi \frac{i}{\sqrt{2}} e^{i\varphi}.
\end{aligned} \tag{1.1.39}$$

Spherical Covariant and Spherical Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_{+1} &= -\mathbf{e}^{-1}, & \mathbf{e}^{+1} &= -\mathbf{e}_{-1}, \\
\mathbf{e}_0 &= \mathbf{e}^0, & \mathbf{e}^0 &= \mathbf{e}_0, \\
\mathbf{e}_{-1} &= -\mathbf{e}^{+1}, & \mathbf{e}^{-1} &= -\mathbf{e}_{+1}.
\end{aligned} \tag{1.1.40}$$

Spherical Covariant and Helicity Covariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_{+1} &= \mathbf{e}'_{+1} \frac{1 + \cos \vartheta}{2} e^{i\varphi} - \mathbf{e}'_0 \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} + \mathbf{e}'_{-1} \frac{1 - \cos \vartheta}{2} e^{i\varphi}, \\
\mathbf{e}_0 &= \mathbf{e}'_{+1} \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}'_0 \cos \vartheta - \mathbf{e}'_{-1} \frac{\sin \vartheta}{\sqrt{2}}, \\
\mathbf{e}_{-1} &= \mathbf{e}'_{+1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi} + \mathbf{e}'_0 \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} + \mathbf{e}'_{-1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi}.
\end{aligned} \tag{1.1.41}$$

Spherical Contravariant and Helicity Covariant Basis Vectors

$$\begin{aligned}
\mathbf{e}^{+1} &= -\mathbf{e}'_{+1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi} - \mathbf{e}'_0 \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} - \mathbf{e}'_{-1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi}, \\
\mathbf{e}^0 &= \mathbf{e}'_{+1} \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}'_0 \cos \vartheta - \mathbf{e}'_{-1} \frac{\sin \vartheta}{\sqrt{2}}, \\
\mathbf{e}^{-1} &= -\mathbf{e}'_{+1} \frac{1 + \cos \vartheta}{2} e^{i\varphi} + \mathbf{e}'_0 \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} - \mathbf{e}'_{-1} \frac{1 - \cos \vartheta}{2} e^{i\varphi}.
\end{aligned} \tag{1.1.42}$$

Spherical Covariant and Helicity Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}_{+1} &= -\mathbf{e}'^{+1} \frac{1 - \cos \vartheta}{2} e^{i\varphi} - \mathbf{e}'^0 \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} - \mathbf{e}'^{-1} \frac{1 + \cos \vartheta}{2} e^{i\varphi}, \\
\mathbf{e}_0 &= \mathbf{e}'^{+1} \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}'^0 \cos \vartheta - \mathbf{e}'^{-1} \frac{\sin \vartheta}{\sqrt{2}}, \\
\mathbf{e}_{-1} &= -\mathbf{e}'^{+1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi} + \mathbf{e}'^0 \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} - \mathbf{e}'^{-1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi}.
\end{aligned} \tag{1.1.43}$$

Spherical Contravariant and Helicity Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}^{+1} &= \mathbf{e}'^{+1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi} - \mathbf{e}'^0 \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} + \mathbf{e}'^{-1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi}, \\
\mathbf{e}^0 &= \mathbf{e}'^{+1} \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}'^0 \cos \vartheta - \mathbf{e}'^{-1} \frac{\sin \vartheta}{\sqrt{2}}, \\
\mathbf{e}^{-1} &= \mathbf{e}'^{+1} \frac{1 - \cos \vartheta}{2} e^{i\varphi} + \mathbf{e}'^0 \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} + \mathbf{e}'^{-1} \frac{1 + \cos \vartheta}{2} e^{i\varphi}.
\end{aligned} \tag{1.1.44}$$

Equations (1.1.41)-(1.1.44) may be written in a more compact form using the Wigner D -functions (see Chap. 4).

$$\begin{aligned}
\mathbf{e}_\mu &= \sum_\nu D^1_{-\nu-\mu}(0, \vartheta, \varphi) \mathbf{e}'_\nu = \sum_\nu (-1)^\nu D^1_{\nu-\mu}(0, \vartheta, \varphi) \mathbf{e}''_\nu, \\
\mathbf{e}^\mu &= (-1)^\mu \sum_\nu D^1_{-\nu\mu}(0, \vartheta, \varphi) \mathbf{e}'_\nu = \sum_\nu (-1)^{\mu+\nu} D^1_{\nu\mu}(0, \vartheta, \varphi) \mathbf{e}''_\nu, \\
&(\mu, \nu = \pm 1, 0).
\end{aligned} \tag{1.1.45}$$

(d) Helicity Vectors and other basis vectors

Helicity Covariant and Cartesian Basis Vectors

$$\begin{aligned}
\mathbf{e}'_{+1} &= -\mathbf{e}_x \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi - i \sin \varphi) - \mathbf{e}_y \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi + i \cos \varphi) + \mathbf{e}_z \frac{1}{\sqrt{2}} \sin \vartheta, \\
\mathbf{e}'_0 &= \mathbf{e}_x \sin \vartheta \cos \varphi + \mathbf{e}_y \sin \vartheta \sin \varphi + \mathbf{e}_z \cos \vartheta, \\
\mathbf{e}'_{-1} &= \mathbf{e}_x \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi + i \sin \varphi) + \mathbf{e}_y \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi - i \cos \varphi) - \mathbf{e}_z \frac{1}{\sqrt{2}} \sin \vartheta.
\end{aligned} \tag{1.1.46}$$

Helicity Contravariant and Cartesian Basis Vectors

$$\begin{aligned}
\mathbf{e}'^{+1} &= -\mathbf{e}_x \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi + i \sin \varphi) - \mathbf{e}_y \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi - i \cos \varphi) + \mathbf{e}_z \frac{1}{\sqrt{2}} \sin \vartheta, \\
\mathbf{e}'^0 &= \mathbf{e}_x \sin \vartheta \cos \varphi + \mathbf{e}_y \sin \vartheta \sin \varphi + \mathbf{e}_z \cos \vartheta, \\
\mathbf{e}'^{-1} &= \mathbf{e}_x \frac{1}{\sqrt{2}} (\cos \vartheta \cos \varphi - i \sin \varphi) + \mathbf{e}_y \frac{1}{\sqrt{2}} (\cos \vartheta \sin \varphi + i \cos \varphi) - \mathbf{e}_z \frac{1}{\sqrt{2}} \sin \vartheta.
\end{aligned} \tag{1.1.47}$$

Helicity and Polar Basis Vectors

$$\begin{aligned}
\mathbf{e}'_{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta + i\mathbf{e}_\varphi), & \mathbf{e}'^{+1} &= -\frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta - i\mathbf{e}_\varphi), \\
\mathbf{e}'_0 &= \mathbf{e}_r, & \mathbf{e}'^0 &= \mathbf{e}_r, \\
\mathbf{e}'_{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta - i\mathbf{e}_\varphi), & \mathbf{e}'^{-1} &= \frac{1}{\sqrt{2}} (\mathbf{e}_\vartheta + i\mathbf{e}_\varphi).
\end{aligned} \tag{1.1.48}$$

Helicity Covariant and Spherical Covariant Basis Vectors

$$\begin{aligned}
\mathbf{e}'_{+1} &= \mathbf{e}_{+1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi} + \mathbf{e}_0 \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}_{-1} \frac{1 - \cos \vartheta}{2} e^{i\varphi}, \\
\mathbf{e}'_0 &= -\mathbf{e}_{+1} \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} + \mathbf{e}_0 \cos \vartheta + \mathbf{e}_{-1} \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi}, \\
\mathbf{e}'_{-1} &= \mathbf{e}_{+1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi} - \mathbf{e}_0 \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}_{-1} \frac{1 + \cos \vartheta}{2} e^{i\varphi}.
\end{aligned} \tag{1.1.49}$$

Helicity Contravariant and Spherical Covariant Basis Vectors

$$\begin{aligned}
\mathbf{e}'^{+1} &= -\mathbf{e}_{+1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi} + \mathbf{e}_0 \frac{\sin \vartheta}{\sqrt{2}} - \mathbf{e}_{-1} \frac{1 + \cos \vartheta}{2} e^{i\varphi}, \\
\mathbf{e}'^0 &= -\mathbf{e}_{+1} \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi} + \mathbf{e}_0 \cos \vartheta + \mathbf{e}_{-1} \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi}, \\
\mathbf{e}'^{-1} &= -\mathbf{e}_{+1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi} - \mathbf{e}_0 \frac{\sin \vartheta}{\sqrt{2}} - \mathbf{e}_{-1} \frac{1 - \cos \vartheta}{2} e^{i\varphi}.
\end{aligned} \tag{1.1.50}$$

Helicity Covariant and Spherical Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}'_{+1} &= -\mathbf{e}^{+1} \frac{1 - \cos \vartheta}{2} e^{i\varphi} + \mathbf{e}^0 \frac{\sin \vartheta}{\sqrt{2}} - \mathbf{e}^{-1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi}, \\
\mathbf{e}'_0 &= -\mathbf{e}^{+1} \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} + \mathbf{e}^0 \cos \vartheta + \mathbf{e}^{-1} \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi}, \\
\mathbf{e}'_{-1} &= -\mathbf{e}^{+1} \frac{1 + \cos \vartheta}{2} e^{i\varphi} - \mathbf{e}^0 \frac{\sin \vartheta}{\sqrt{2}} - \mathbf{e}^{-1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi}.
\end{aligned} \tag{1.1.51}$$

Helicity Contravariant and Spherical Contravariant Basis Vectors

$$\begin{aligned}
\mathbf{e}'^{+1} &= \mathbf{e}^{+1} \frac{1 + \cos \vartheta}{2} e^{i\varphi} + \mathbf{e}^0 \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}^{-1} \frac{1 - \cos \vartheta}{2} e^{-i\varphi}, \\
\mathbf{e}'^0 &= -\mathbf{e}^{+1} \frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} + \mathbf{e}^0 \cos \vartheta + \mathbf{e}^{-1} \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi}, \\
\mathbf{e}'^{-1} &= \mathbf{e}^{+1} \frac{1 - \cos \vartheta}{2} e^{i\varphi} - \mathbf{e}^0 \frac{\sin \vartheta}{\sqrt{2}} + \mathbf{e}^{-1} \frac{1 + \cos \vartheta}{2} e^{-i\varphi}.
\end{aligned} \tag{1.1.52}$$

Equations (1.1.49)-(1.1.52) may be written in a more compact form using the Wigner D -functions (see Chap. 4).

$$\begin{aligned} \mathbf{e}'_\mu &= \sum_\nu D_{\nu\mu}^1(\varphi, \vartheta, 0) \mathbf{e}_\nu = \sum_\nu (-1)^\nu D_{-\nu\mu}^1(\varphi, \vartheta, 0) \mathbf{e}^\nu, \\ \mathbf{e}'^\mu &= \sum_\nu (-1)^\mu D_{\nu-\mu}^1(\varphi, \vartheta, 0) \mathbf{e}_\nu = \sum_\nu (-1)^{\mu+\nu} D_{-\nu-\mu}^1(\varphi, \vartheta, 0) \mathbf{e}^\nu, \\ &(\mu, \nu = \pm 1, 0). \end{aligned} \quad (1.1.53)$$

Helicity Covariant and Helicity Contravariant Basis Vectors

$$\begin{aligned} \mathbf{e}'_{+1} &= -\mathbf{e}'^{-1}, & \mathbf{e}'^{+1} &= -\mathbf{e}'_{-1}, \\ \mathbf{e}'_0 &= \mathbf{e}'^0, & \mathbf{e}'^0 &= \mathbf{e}'_0, \\ \mathbf{e}'_{-1} &= -\mathbf{e}'^{+1}, & \mathbf{e}'^{-1} &= -\mathbf{e}'_{-1}. \end{aligned} \quad (1.1.54)$$

1.2 Vectors. Tensors

Vectors and tensors are usually defined by transformation properties of their components under rotations of coordinate systems. The transformation rule for cartesian components of vectors and tensors is given below in Sec. 1.4 (Eqs. (1.4.46)-(1.4.51)). The transformation properties of spherical components of vectors and irreducible tensors are discussed in Chap. 3.

1.2.1 Vector Components

Any vector can be expanded in terms of basis vectors, i.e., written as

$$\mathbf{A} = \sum_\alpha A^\alpha \mathbf{e}_\alpha = \sum_\alpha A_\alpha \mathbf{e}^\alpha. \quad (1.2.1)$$

The expansion coefficients A_α are called the covariant components of the vector, and A^α are the contravariant vector components

$$A_\alpha = \mathbf{A} \cdot \mathbf{e}_\alpha, \quad A^\alpha = \mathbf{A} \cdot \mathbf{e}^\alpha. \quad (1.2.2)$$

In a cartesian coordinate system one has

$$\mathbf{A} = A_x \mathbf{e}_x + A_y \mathbf{e}_y + A_z \mathbf{e}_z = A^x \mathbf{e}_x + A^y \mathbf{e}_y + A^z \mathbf{e}_z. \quad (1.2.3)$$

The covariant cartesian components of a vector coincide with the contravariant ones.

In a polar coordinate system

$$\mathbf{A} = A_r \mathbf{e}_r + A_\vartheta \mathbf{e}_\vartheta + A_\varphi \mathbf{e}_\varphi = A^r \mathbf{e}_r + A^\vartheta \mathbf{e}_\vartheta + A^\varphi \mathbf{e}_\varphi. \quad (1.2.4)$$

The covariant polar components coincide with the contravariant ones.

For a spherical coordinate system

$$\mathbf{A} = A^{+1} \mathbf{e}_{+1} + A^0 \mathbf{e}_0 + A^{-1} \mathbf{e}_{-1} = A_{+1} \mathbf{e}^{+1} + A_0 \mathbf{e}^0 + A_{-1} \mathbf{e}^{-1} \quad (1.2.5)$$

The relations between covariant and contravariant spherical components are given by

$$A_\mu = (-1)^\mu A^{-\mu}, \quad A^\mu = (-1)^\mu A_{-\mu}, \quad (\mu = \pm 1, 0). \quad (1.2.6)$$

If $\mathbf{A}$ is a real vector, i.e., if $\mathbf{A}^* = \mathbf{A}$, then

$$A_\mu^* = A^\mu, \quad A^{\mu*} = A_\mu, \quad (\mu = \pm 1, 0). \quad (1.2.7)$$

If $\mathbf{A}$ is a complex vector, then

$$A_\mu^* = (\mathbf{A}^*)^\mu, \quad A^{\mu*} = (\mathbf{A}^*)_\mu, \quad (\mu = \pm 1, 0). \quad (1.2.8)$$

An expansion of a real vector $\mathbf{A}$ in terms of spherical basis vectors is written as

$$\begin{aligned} \mathbf{A} &= \sum_\mu A_\mu \mathbf{e}^\mu = \sum_\mu A^\mu \mathbf{e}_\mu = \sum_\mu A_\mu^* \mathbf{e}^{\mu*} = \sum_\mu A^{\mu*} \mathbf{e}_\mu^* \\ &= \sum_\mu A_\mu \mathbf{e}_\mu^* = \sum_\mu A_\mu^* \mathbf{e}_\mu = \sum_\mu A^\mu \mathbf{e}^{\mu*} = \sum_\mu A^{\mu*} \mathbf{e}^\mu \\ &= \sum_\mu (-1)^\mu A_{-\mu} \mathbf{e}_\mu = \sum_\mu (-1)^\mu A^{-\mu} \mathbf{e}^\mu. \end{aligned} \quad (1.2.9)$$

An expansion of an arbitrary vector $\mathbf{A}$ in terms of helicity basis vectors is given by

$$\mathbf{A} = A'^{+1}\mathbf{e}'_{+1} + A'^0\mathbf{e}'_0 + A'^{-1}\mathbf{e}'_{-1} = A'_{+1}\mathbf{e}'^{+1} + A'_0\mathbf{e}'^0 + A'_{-1}\mathbf{e}'^{-1}. \quad (1.2.10)$$

The helicity components of a vector satisfy the same relations (1.2.6)-(1.2.9) as the spherical components. The relations between vector components in different bases are the same as the relations between basis vectors. These relations are given by Eqs. (1.1.29)-(1.1.54) in which one should replace $\mathbf{e}_\alpha \rightarrow A_\alpha$ and $\mathbf{e}^\alpha \rightarrow A^\alpha$. In particular,

$$\begin{aligned} A_{+1} &= -A^{-1} = -\frac{1}{\sqrt{2}}(A_x + iA_y), & A_x &= \frac{1}{\sqrt{2}}(A_{-1} - A_{+1}) = \frac{1}{\sqrt{2}}(A^{-1} - A^{+1}), \\ A_0 &= A^0 = A_z, & A_y &= \frac{i}{\sqrt{2}}(A_{-1} + A_{+1}) = \frac{-i}{\sqrt{2}}(A^{-1} + A^{+1}), \\ A_{-1} &= -A^{+1} = \frac{1}{\sqrt{2}}(A_x - iA_y), & A_z &= A_0 = A^0. \end{aligned} \quad (1.2.11)$$

The matrices of transformations between cartesian, contravariant spherical and polar components of vectors are given in Tables 1.1 and 1.2.

Spherical components of a real vector $\mathbf{A}$ which contains no derivatives and is independent of spin variables are

$$\begin{aligned} A_{\pm 1} &= \mp |\mathbf{A}| \frac{\sin \vartheta}{\sqrt{2}} e^{\pm i\varphi}, & A^{\pm 1} &= \mp |\mathbf{A}| \frac{\sin \vartheta}{\sqrt{2}} e^{\mp i\varphi}, \\ A_0 &= |\mathbf{A}| \cos \vartheta, & A^0 &= |\mathbf{A}| \cos \vartheta, \end{aligned} \quad (1.2.12)$$

where ϑ, φ are the polar angles of the vector $\mathbf{A}$.

Equation (1.2.12) may be written in terms of spherical harmonics (see Chap. 5) as

$$\begin{aligned} A_\mu &= \sqrt{\frac{4\pi}{3}} |\mathbf{A}| Y_{1\mu}(\vartheta, \varphi) \\ A^\mu &= \sqrt{\frac{4\pi}{3}} |\mathbf{A}| Y_{1\mu}^*(\vartheta, \varphi), \quad (\mu = \pm 1, 0) \end{aligned} \quad (1.2.13)$$

The expressions for cartesian components of $\mathbf{A}$ in terms of spherical harmonics read

$$\begin{aligned} A_x &= \sqrt{\frac{2\pi}{3}} |\mathbf{A}| \{Y_{1-1}(\vartheta, \varphi) - Y_{1+1}(\vartheta, \varphi)\}, \\ A_y &= i\sqrt{\frac{2\pi}{3}} |\mathbf{A}| \{Y_{1-1}(\vartheta, \varphi) + Y_{1+1}(\vartheta, \varphi)\}, \\ A_z &= \sqrt{\frac{4\pi}{3}} |\mathbf{A}| Y_{10}(\vartheta, \varphi). \end{aligned} \quad (1.2.14)$$

Table 1.1. Matrix form of the transformations for vector components in different bases.

Cartesian coordinates	Spherical coordinates	Polar coordinates
$\mathbf{A} = A_x \mathbf{e}_x + A_y \mathbf{e}_y + A_z \mathbf{e}_z$	$\mathbf{A} = A^{+1} \mathbf{e}_{+1} + A^0 \mathbf{e}_0 + A^{-1} \mathbf{e}_{-1}$	$\mathbf{A} = A_r \mathbf{e}_r + A_\vartheta \mathbf{e}_\vartheta + A_\varphi \mathbf{e}_\varphi$
In terms of spherical components	In terms of cartesian components	In terms of cartesian components
$\begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix} = M(x, y, z \leftarrow +1, 0, -1) \begin{pmatrix} A^{+1} \\ A^0 \\ A^{-1} \end{pmatrix}$	$\begin{pmatrix} A^{+1} \\ A^0 \\ A^{-1} \end{pmatrix} = M(+1, 0, -1 \leftarrow x, y, z) \begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix}$	$\begin{pmatrix} A_r \\ A_\vartheta \\ A_\varphi \end{pmatrix} = M(r, \vartheta, \varphi \leftarrow x, y, z) \begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix}$
In terms of polar components	In terms of polar components	In terms of spherical components
$\begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix} = M(x, y, z \leftarrow r, \vartheta, \varphi) \begin{pmatrix} A_r \\ A_\vartheta \\ A_\varphi \end{pmatrix}$	$\begin{pmatrix} A^{+1} \\ A^0 \\ A^{-1} \end{pmatrix} = M(+1, 0, -1 \leftarrow r, \vartheta, \varphi) \begin{pmatrix} A_r \\ A_\vartheta \\ A_\varphi \end{pmatrix}$	$\begin{pmatrix} A_r \\ A_\vartheta \\ A_\varphi \end{pmatrix} = M(r, \vartheta, \varphi \leftarrow +1, 0, -1) \begin{pmatrix} A^{+1} \\ A^0 \\ A^{-1} \end{pmatrix}$

1.2.2 Scalar Product of Vectors

The scalar product of vectors $\mathbf{A}$ and $\mathbf{B}$ in an arbitrary orthonormal basis is defined by

$$\mathbf{A} \cdot \mathbf{B} = \sum_{\alpha} A_{\alpha} B^{\alpha} = \sum_{\alpha} A^{\alpha} B_{\alpha}. \quad (1.2.15)$$

Table 1.2. Matrices of transformations between cartesian, spherical contravariant and polar components of vectors.

$M(x, y, z \leftarrow +1, 0, -1)$			$M(+1, 0, -1 \leftarrow x, y, z)$		
	+1	0	-1		$x \quad y \quad z$
x	$-\frac{1}{\sqrt{2}}$	0	$\frac{1}{\sqrt{2}}$	-1	$-\frac{1}{\sqrt{2}} \quad -\frac{i}{\sqrt{2}} \quad 0$
y	$-\frac{i}{\sqrt{2}}$	0	$-\frac{i}{\sqrt{2}}$	0	$0 \quad 0 \quad 1$
z	0	1	0	1	$\frac{1}{\sqrt{2}} \quad -\frac{i}{\sqrt{2}} \quad 0$
$M(x, y, z \leftarrow r, \vartheta, \varphi)$			$M(r, \vartheta, \varphi \leftarrow x, y, z)$		
	r	ϑ	φ		$x \quad y \quad z$
x	$\sin \vartheta \cos \varphi$	$\cos \vartheta \cos \varphi$	$-\sin \varphi$	r	$\sin \vartheta \cos \varphi \quad \sin \vartheta \sin \varphi \quad \cos \vartheta$
y	$\sin \vartheta \sin \varphi$	$\cos \vartheta \sin \varphi$	$\cos \varphi$	ϑ	$\cos \vartheta \cos \varphi \quad \cos \vartheta \sin \varphi \quad -\sin \vartheta$
z	$\cos \vartheta$	$-\sin \vartheta$	0	φ	$-\sin \varphi \quad \cos \varphi \quad 0$
$M(+1, 0, -1 \leftarrow r, \vartheta, \varphi)$			$M(r, \vartheta, \varphi \leftarrow +1, 0, -1)$		
	r	ϑ	φ		+1 0 -1
+1	$-\frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi}$	$-\frac{\cos \vartheta}{\sqrt{2}} e^{-i\varphi}$	$\frac{i}{\sqrt{2}} e^{-i\varphi}$	r	$-\frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi} \quad \cos \vartheta \quad \frac{\sin \vartheta}{\sqrt{2}} e^{-i\varphi}$
0	$\cos \vartheta$	$-\sin \vartheta$	0	ϑ	$-\frac{\cos \vartheta}{\sqrt{2}} e^{i\varphi} \quad -\sin \vartheta \quad \frac{\cos \vartheta}{\sqrt{2}} e^{-i\varphi}$
-1	$\frac{\sin \vartheta}{\sqrt{2}} e^{i\varphi}$	$\frac{\cos \vartheta}{\sqrt{2}} e^{i\varphi}$	$\frac{i}{\sqrt{2}} e^{i\varphi}$	φ	$-\frac{i}{\sqrt{2}} e^{i\varphi} \quad 0 \quad -\frac{i}{\sqrt{2}} e^{-i\varphi}$

In a cartesian coordinate system we have

$$\mathbf{A} \cdot \mathbf{B} = A_x B_x + A_y B_y + A_z B_z. \quad (1.2.16)$$

For polar coordinates the scalar product is given by

$$\mathbf{A} \cdot \mathbf{B} = A_r B_r + A_\vartheta B_\vartheta + A_\varphi B_\varphi. \quad (1.2.17)$$

Equation (1.2.17) is valid only if $\mathbf{A}$ and $\mathbf{B}$ do not contain derivatives because the polar basis vectors depend on the polar angles ϑ, φ (see Section 1.1). In spherical coordinates we have

$$\mathbf{A} \cdot \mathbf{B} = \sum_{\mu} A^{\mu} B_{\mu} = \sum_{\mu} A_{\mu} B^{\mu} = \sum_{\mu} (-1)^{\mu} A_{\mu} B_{-\mu} = \sum_{\mu} (-1)^{\mu} A^{\mu} B^{-\mu}, \quad (\mu = \pm 1, 0), \quad (1.2.18)$$

or, in a more detailed form,

$$\mathbf{A} \cdot \mathbf{B} = -A_{+1} B_{-1} + A_0 B_0 - A_{-1} B_{+1}, \quad (1.2.19)$$

The scalar product in terms of helicity components of vectors is similar to (1.2.18)-(1.2.19). The scalar product of vectors is invariant with respect to rotations of the coordinate system.

1.2.3 Vector Product of Vectors

In a cartesian coordinate system the vector product of vectors $\mathbf{A}$ and $\mathbf{B}$ is defined by

$$\mathbf{A} \times \mathbf{B} = \begin{vmatrix} \mathbf{e}_x & \mathbf{e}_y & \mathbf{e}_z \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{vmatrix} = \sum_{i=x,y,z} [\mathbf{A} \times \mathbf{B}]_i \mathbf{e}_i, \quad (1.2.20)$$

where

$$\begin{aligned} [\mathbf{A} \times \mathbf{B}]_x &= A_y B_z - A_z B_y, \\ [\mathbf{A} \times \mathbf{B}]_y &= A_z B_x - A_x B_z, \\ [\mathbf{A} \times \mathbf{B}]_z &= A_x B_y - A_y B_x. \end{aligned} \quad (1.2.21)$$

Equation (1.2.21) may be written in a more compact form as

$$[\mathbf{A} \times \mathbf{B}]_i = \sum_{kl} \varepsilon_{ikl} A_k B_l, \quad (i, k, l = x, y, z). \quad (1.2.22)$$

In the polar coordinate system

$$\mathbf{A} \times \mathbf{B} = \begin{vmatrix} \mathbf{e}_r & \mathbf{e}_\vartheta & \mathbf{e}_\varphi \\ A_r & A_\vartheta & A_\varphi \\ B_r & B_\vartheta & B_\varphi \end{vmatrix} = \sum_{\alpha=r,\vartheta,\varphi} [\mathbf{A} \times \mathbf{B}]_\alpha \mathbf{e}_\alpha, \quad (1.2.23)$$

where

$$\begin{aligned} [\mathbf{A} \times \mathbf{B}]_r &= A_\vartheta B_\varphi - A_\varphi B_\vartheta, \\ [\mathbf{A} \times \mathbf{B}]_\vartheta &= A_\varphi B_r - A_r B_\varphi, \\ [\mathbf{A} \times \mathbf{B}]_\varphi &= A_r B_\vartheta - A_\vartheta B_r. \end{aligned} \quad (1.2.24)$$

Equation (1.2.24) is valid only if $\mathbf{A}$ and $\mathbf{B}$ are not differential operators.

In the spherical coordinate system

$$\mathbf{A} \times \mathbf{B} = i \begin{vmatrix} \mathbf{e}_{+1} & \mathbf{e}_0 & \mathbf{e}_{-1} \\ A_{+1} & A_0 & A_{-1} \\ B_{+1} & B_0 & B_{-1} \end{vmatrix} = -i \begin{vmatrix} \mathbf{e}^{+1} & \mathbf{e}^0 & \mathbf{e}^{-1} \\ A^{+1} & A^0 & A^{-1} \\ B^{+1} & B^0 & B^{-1} \end{vmatrix} = \sum_{\mu=\pm 1,0} [\mathbf{A} \times \mathbf{B}]_\mu \mathbf{e}^\mu = \sum_{\mu=\pm 1,0} [\mathbf{A} \times \mathbf{B}]^\mu \mathbf{e}_\mu, \quad (1.2.25)$$

where

$$\begin{aligned} [\mathbf{A} \times \mathbf{B}]_{+1} &= i (A_0 B_{+1} - A_{+1} B_0) = i (A^{-1} B^0 - A^0 B^{-1}), \\ [\mathbf{A} \times \mathbf{B}]_0 &= i (A_{-1} B_{+1} - A_{+1} B_{-1}) = i (A^{+1} B^{-1} - A^{-1} B^{+1}), \\ [\mathbf{A} \times \mathbf{B}]_{-1} &= i (A_{-1} B_0 - A_0 B_{-1}) = i (A^0 B^{+1} - A^{+1} B^0). \end{aligned} \quad (1.2.26)$$

$$\begin{aligned} [\mathbf{A} \times \mathbf{B}]^{+1} &= i (A_0 B_{-1} - A_{-1} B_0) = i (A^{+1} B^0 - A^0 B^{+1}), \\ [\mathbf{A} \times \mathbf{B}]^0 &= i (A_{-1} B_{+1} - A_{+1} B_{-1}) = i (A^{+1} B^{-1} - A^{-1} B^{+1}), \\ [\mathbf{A} \times \mathbf{B}]^{-1} &= i (A_{+1} B_0 - A_0 B_{+1}) = i (A^0 B^{-1} - A^{-1} B_0). \end{aligned} \quad (1.2.27)$$

Equations (1.2.26)-(1.2.27) may be written in a more compact form using the Clebsch-Gordan coefficients (see Chap. 8)

$$\begin{aligned} [\mathbf{A} \times \mathbf{B}]_\mu &= -i\sqrt{2} \sum_{\nu\lambda} C_{1\nu 1\lambda}^{1\mu} A_\nu B_\lambda, \\ [\mathbf{A} \times \mathbf{B}]^\mu &= i\sqrt{2} \sum_{\nu\lambda} C_{1\nu 1\lambda}^{1\mu} A^\nu B^\lambda, \end{aligned} \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (1.2.28)$$

Helicity components of the vector product are given by equations analogous to (1.2.25)-(1.2.28).

1.2.4 Products Involving Three or More Vectors

$$\mathbf{A} \cdot [\mathbf{B} \times \mathbf{C}] = \mathbf{B} \cdot [\mathbf{C} \times \mathbf{A}] = \mathbf{C} \cdot [\mathbf{A} \times \mathbf{B}] = -\mathbf{A} \cdot [\mathbf{C} \times \mathbf{B}] = -\mathbf{B} \cdot [\mathbf{A} \times \mathbf{C}] = -\mathbf{C} \cdot [\mathbf{B} \times \mathbf{A}], \quad (1.2.29)$$

$$\mathbf{A} \times [\mathbf{B} \times \mathbf{C}] = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}), \quad (1.2.30)$$

$$[\mathbf{A} \times \mathbf{B}] \cdot [\mathbf{C} \times \mathbf{D}] = (\mathbf{A} \cdot \mathbf{C})(\mathbf{B} \cdot \mathbf{D}) - (\mathbf{A} \cdot \mathbf{D})(\mathbf{B} \cdot \mathbf{C}), \quad (1.2.31)$$

$$[\mathbf{A} \times \mathbf{B}] \times [\mathbf{C} \times \mathbf{D}] = \mathbf{B}(\mathbf{A} \cdot [\mathbf{C} \times \mathbf{D}]) - \mathbf{A}(\mathbf{B} \cdot [\mathbf{C} \times \mathbf{D}]) = \mathbf{C}(\mathbf{A} \cdot [\mathbf{B} \times \mathbf{D}]) - \mathbf{D}(\mathbf{A} \cdot [\mathbf{B} \times \mathbf{C}]). \quad (1.2.32)$$

$$\begin{aligned} (\mathbf{A} \cdot [\mathbf{B} \times \mathbf{C}])(\mathbf{a} \cdot [\mathbf{b} \times \mathbf{c}]) &= \begin{vmatrix} \mathbf{A} \cdot \mathbf{a} & \mathbf{A} \cdot \mathbf{b} & \mathbf{A} \cdot \mathbf{c} \\ \mathbf{B} \cdot \mathbf{a} & \mathbf{B} \cdot \mathbf{b} & \mathbf{B} \cdot \mathbf{c} \\ \mathbf{C} \cdot \mathbf{a} & \mathbf{C} \cdot \mathbf{b} & \mathbf{C} \cdot \mathbf{c} \end{vmatrix} \\ &= (\mathbf{A} \cdot \mathbf{a})(\mathbf{B} \cdot \mathbf{b})(\mathbf{C} \cdot \mathbf{c}) - (\mathbf{A} \cdot \mathbf{a})(\mathbf{B} \cdot \mathbf{c})(\mathbf{C} \cdot \mathbf{b}) - (\mathbf{B} \cdot \mathbf{b})(\mathbf{A} \cdot \mathbf{c})(\mathbf{C} \cdot \mathbf{a}) \\ &\quad - (\mathbf{C} \cdot \mathbf{c})(\mathbf{A} \cdot \mathbf{b})(\mathbf{B} \cdot \mathbf{a}) + (\mathbf{A} \cdot \mathbf{b})(\mathbf{B} \cdot \mathbf{c})(\mathbf{C} \cdot \mathbf{a}) + (\mathbf{A} \cdot \mathbf{c})(\mathbf{B} \cdot \mathbf{a})(\mathbf{C} \cdot \mathbf{b}). \end{aligned} \quad (1.2.33)$$

1.2.5 Tensors δ_{ik} and ε_{ikl}

In a cartesian basis two basic tensors δ_{ik} and ε_{ikl} are widely used. The first tensor, δ_{ik} , is the symmetric unit tensor of rank 2. The second tensor, ε_{ikl} , is the totally antisymmetric unit tensor of rank 3.

The tensor δ_{ik} is called the Kronecker δ -symbol and is defined by

$$\delta_{ik} = \begin{cases} 1, & i = k, \\ 0, & i \neq k, \end{cases} \quad (i, k = x, y, z) \quad (1.2.34)$$

The components δ_{ik} are invariant with respect to rotations and inversion of coordinate systems.

The tensor (or, more precisely, the pseudo-tensor) ε_{ikl} is often called the Levi-Civita tensor. It is antisymmetric with respect to permutations of any pair of indices. Thus, $\varepsilon_{ikl} = 0$ if at least two of the indices i, k, l are equal, and $\varepsilon_{ikl} \neq 0$ only if all indices i, k, l are different. The components ε_{ikl} are given by

$$\begin{aligned} \varepsilon_{iii} &= 0, \quad (i = x, y, z) \quad (3 \text{ components}), \\ \varepsilon_{iik} &= \varepsilon_{iki} = \varepsilon_{kii} = 0, \quad (i, k = x, y, z) \quad (18 \text{ components}), \\ \varepsilon_{xyz} &= \varepsilon_{yzx} = \varepsilon_{zxy} = -\varepsilon_{xzy} = -\varepsilon_{yxz} = -\varepsilon_{zyx} = 1 \quad (6 \text{ components}). \end{aligned} \quad (1.2.35)$$

The components ε_{ikl} are invariant with respect to rotations and inversion of coordinate systems.

The tensor ε_{ikl} has the following properties: The product of two tensors ε_{ikl} and ε_{rst} may be written in the form of a determinant

$$\varepsilon_{ikl}\varepsilon_{rst} = \begin{vmatrix} \delta_{ir} & \delta_{is} & \delta_{it} \\ \delta_{kr} & \delta_{ks} & \delta_{kt} \\ \delta_{lr} & \delta_{ls} & \delta_{lt} \end{vmatrix} = \delta_{ir}\delta_{ks}\delta_{lt} + \delta_{is}\delta_{kt}\delta_{lr} + \delta_{it}\delta_{kr}\delta_{ls} - \delta_{ir}\delta_{kt}\delta_{ls} - \delta_{is}\delta_{kr}\delta_{lt} - \delta_{it}\delta_{ks}\delta_{lr}. \quad (1.2.36)$$

By summing over a pair of indices, one obtains

$$\sum_i \varepsilon_{ikl}\varepsilon_{ist} = \begin{vmatrix} \delta_{ks} & \delta_{kt} \\ \delta_{ls} & \delta_{lt} \end{vmatrix} = \delta_{ks}\delta_{lt} - \delta_{kt}\delta_{ls}. \quad (1.2.37)$$

Summation over two pairs of indices yields

$$\sum_{i,k} \varepsilon_{ikl}\varepsilon_{ikt} = 2\delta_{lt}. \quad (1.2.38)$$

Finally, the summation over three pairs of indices gives

$$\sum_{i,k,l} \varepsilon_{ikl}\varepsilon_{ikl} = 6. \quad (1.2.39)$$

For an arbitrary 3×3 matrix $\|A_{ik}\|$ ($i, k = x, y, z$) the following relation holds

$$\sum_{i,k,l} A_{xi}A_{yk}A_{zl}\varepsilon_{ikl} = \det \|A_{ik}\| = \begin{vmatrix} A_{xx} & A_{xy} & A_{xz} \\ A_{yx} & A_{yy} & A_{yz} \\ A_{zx} & A_{zy} & A_{zz} \end{vmatrix}. \quad (1.2.40)$$

1.3 Differential Operations

1.3.1 Operator ∇

The operator ∇ (nabla) is the basic vector differential operator.

Cartesian components of ∇ are given by

$$\nabla_x = \frac{\partial}{\partial x}, \quad \nabla_y = \frac{\partial}{\partial y}, \quad \nabla_z = \frac{\partial}{\partial z}. \quad (1.3.1)$$

These components may be expressed in terms of polar coordinates as

$$\begin{aligned} \nabla_x &= \sin \vartheta \cos \varphi \frac{\partial}{\partial r} + \frac{\cos \vartheta \cos \varphi}{r} \frac{\partial}{\partial \vartheta} - \frac{\sin \varphi}{r \sin \vartheta} \frac{\partial}{\partial \varphi}, \\ \nabla_y &= \sin \vartheta \sin \varphi \frac{\partial}{\partial r} + \frac{\cos \vartheta \sin \varphi}{r} \frac{\partial}{\partial \vartheta} + \frac{\cos \varphi}{r \sin \vartheta} \frac{\partial}{\partial \varphi}, \\ \nabla_z &= \cos \vartheta \frac{\partial}{\partial r} - \frac{\sin \vartheta}{r} \frac{\partial}{\partial \vartheta}. \end{aligned} \quad (1.3.2)$$

An expansion of the operator ∇ in terms of spherical basis vectors reads

$$\nabla = \sum_{\mu} (-1)^{\mu} \mathbf{e}_{\mu} \nabla_{-\mu} = -\mathbf{e}_{+1} \nabla_{-1} + \mathbf{e}_0 \nabla_0 - \mathbf{e}_{-1} \nabla_{+1}, \quad (1.3.3)$$

where spherical components of ∇ are given by

$$\begin{aligned} \nabla_{+1} &= -\frac{1}{\sqrt{2}} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right), \\ \nabla_0 &= \frac{\partial}{\partial z}, \\ \nabla_{-1} &= \frac{1}{\sqrt{2}} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right). \end{aligned} \quad (1.3.4)$$

Spherical components of ∇ in a polar coordinate system have the form

$$\begin{aligned} \nabla_{+1} &= -\frac{e^{i\varphi}}{\sqrt{2}} \left\{ \sin \vartheta \frac{\partial}{\partial r} + \frac{\cos \vartheta}{r} \frac{\partial}{\partial \vartheta} + \frac{i}{r \sin \vartheta} \frac{\partial}{\partial \varphi} \right\}, \\ \nabla_0 &= \cos \vartheta \frac{\partial}{\partial r} - \frac{\sin \vartheta}{r} \frac{\partial}{\partial \vartheta}, \\ \nabla_{-1} &= \frac{e^{-i\varphi}}{\sqrt{2}} \left\{ \sin \vartheta \frac{\partial}{\partial r} + \frac{\cos \vartheta}{r} \frac{\partial}{\partial \vartheta} - \frac{i}{r \sin \vartheta} \frac{\partial}{\partial \varphi} \right\}. \end{aligned} \quad (1.3.5)$$

An expansion of ∇ in terms of polar basis vectors may be written as

$$\nabla = \mathbf{e}_r \nabla_r + \mathbf{e}_{\vartheta} \nabla_{\vartheta} + \mathbf{e}_{\varphi} \nabla_{\varphi}, \quad (1.3.6)$$

where

$$\nabla_r = \frac{\partial}{\partial r}, \quad \nabla_{\vartheta} = \frac{1}{r} \frac{\partial}{\partial \vartheta}, \quad \nabla_{\varphi} = \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \varphi}. \quad (1.3.7)$$

The order of operator components relative to the basis vectors in Eq. (1.3.6) is essential because $\mathbf{e}_r, \mathbf{e}_{\vartheta}, \mathbf{e}_{\varphi}$ depend on ϑ, φ .

The operator ∇ may be written in the form

$$\nabla = \mathbf{n} \frac{\partial}{\partial r} + \frac{1}{r} \nabla_{\Omega}, \quad (1.3.8)$$

where ∇_{Ω} is the angular part of ∇ , and $\mathbf{n} = \mathbf{r}/r$ is the unit vector determined by angles ϑ and φ . The operator ∇_{Ω} acts only on variables ϑ and φ . In the polar coordinate system it has only two components

$$(\nabla_{\Omega})_{\vartheta} = \frac{\partial}{\partial \vartheta}, \quad (\nabla_{\Omega})_{\varphi} = \frac{1}{\sin \vartheta} \frac{\partial}{\partial \varphi}. \quad (1.3.9)$$

The operator ∇_{Ω} may be written as

$$\nabla_{\Omega} = -i \mathbf{n} \times \hat{\mathbf{L}}, \quad (1.3.10)$$

where $\hat{\mathbf{L}}$ is the orbital angular momentum operator (see Sec. 2.2).

1.3.2 Laplace Operator

The Laplace operator (Laplacian) Δ is a scalar differential operator

$$\Delta = \nabla^2. \quad (1.3.11)$$

In the cartesian coordinate system Δ has the form

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}. \quad (1.3.12)$$

In the polar coordinate system it is given by

$$\Delta = \frac{1}{r^2} \frac{\partial}{\partial r} \left\{ r^2 \frac{\partial}{\partial r} \right\} + \frac{1}{r^2 \sin \vartheta} \frac{\partial}{\partial \vartheta} \left\{ \sin \vartheta \frac{\partial}{\partial \vartheta} \right\} + \frac{1}{r^2 \sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2}. \quad (1.3.13)$$

The operator Δ may also be written as

$$\Delta = \frac{1}{r^2} \frac{\partial}{\partial r} \left\{ r^2 \frac{\partial}{\partial r} \right\} + \frac{1}{r^2} \Delta_\Omega, \quad (1.3.14)$$

where Δ_Ω is the angular part of Δ

$$\Delta_\Omega = \nabla_\Omega^2 = \frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} \left\{ \sin \vartheta \frac{\partial}{\partial \vartheta} \right\} + \frac{1}{\sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2} \quad (1.3.15)$$

expressed in terms of the orbital angular momentum operator $\hat{\mathbf{L}}$ as

$$\Delta_\Omega = -\hat{\mathbf{L}}^2 \quad (1.3.16)$$

1.3.3 Differential Operations on Scalars and Vectors

The gradient of a scalar function $\Phi(\mathbf{r})$ is the vector defined in terms of the operator **abla** as

$$\text{grad } \Phi(\mathbf{r}) = \nabla \Phi(\mathbf{r}) \quad (1.3.17)$$

The components of $\text{grad } \Phi$ may be obtained by use of Eqs. (1.3.1)-(1.3.7) for the components of ∇ . If Φ depends only on $r = |\mathbf{r}|$ (spherically-symmetric field), then

$$\nabla \Phi(r) = \mathbf{n} \frac{d\Phi(r)}{dr} \quad (1.3.18)$$

where $\mathbf{n} = \mathbf{r}/r$.

The directional derivative of a scalar function $\Phi(\mathbf{r})$ in the direction specified by a unit vector $\mathbf{u}$ is the scalar defined by

$$\frac{d}{ds} \Phi(\mathbf{r}) = (\mathbf{u} \cdot \nabla) \Phi(\mathbf{r}). \quad (1.3.19)$$

The divergence of a vector field $\mathbf{A}(\mathbf{r})$ is the scalar product of ∇ and $\mathbf{A}$

$$\text{div } \mathbf{A} = \nabla \cdot \mathbf{A}. \quad (1.3.20)$$

The expression for $\text{div } \mathbf{A}$ in a cartesian coordinate system is

$$\text{div } \mathbf{A} = \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} = \sum_{i=x,y,z} \frac{\partial A_i}{\partial x_i}, \quad (1.3.21)$$

and in a spherical coordinate system it has the form

$$\text{div } \mathbf{A} = -\nabla_{+1} A_{-1} + \nabla_0 A_0 - \nabla_{-1} A_{+1} = \sum_{\mu=\pm 1,0} (-1)^\mu \nabla_\mu A_{-\mu}, \quad (1.3.22)$$

where the spherical components ∇_μ are given by Eqs. (1.3.4)-(1.3.5). In the polar coordinate system we have

$$\text{div } \mathbf{A} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 A_r) + \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta A_\vartheta) + \frac{1}{r \sin \vartheta} \frac{\partial A_\varphi}{\partial \varphi}. \quad (1.3.23)$$

The curl of a vector field $\mathbf{A}(\mathbf{r})$ is the vector product of ∇ and $\mathbf{A}$

$$\text{curl } \mathbf{A} = \nabla \times \mathbf{A}. \quad (1.3.24)$$

The cartesian components of $\text{curl } \mathbf{A}$ are given by

$$\begin{aligned} [\text{curl } \mathbf{A}]_x &= \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z}, \\ [\text{curl } \mathbf{A}]_y &= \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x}, \\ [\text{curl } \mathbf{A}]_z &= \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y}, \end{aligned} \quad (1.3.25)$$

or, in a more compact form,

$$[\text{curl } \mathbf{A}]_i = \sum_{kl} \varepsilon_{ikl} \frac{\partial A_l}{\partial x_k}. \quad (1.3.26)$$

Moreover, $\text{curl } \mathbf{A}$ may also be written in the form

$$\text{curl } \mathbf{A} = \begin{vmatrix} \mathbf{e}_x & \mathbf{e}_y & \mathbf{e}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_x & A_y & A_z \end{vmatrix}. \quad (1.3.27)$$

The spherical components of $\text{curl } \mathbf{A}$ are given by

$$\begin{aligned} [\text{curl } \mathbf{A}]_{+1} &= i (\nabla_0 A_{+1} - \nabla_{+1} A_0), \\ [\text{curl } \mathbf{A}]_0 &= i (\nabla_{-1} A_{+1} - \nabla_{+1} A_{-1}), \\ [\text{curl } \mathbf{A}]_{-1} &= i (\nabla_{-1} A_0 - \nabla_0 A_{-1}), \end{aligned} \quad (1.3.28)$$

or in a more compact form involving the Clebsch-Gordan coefficients

$$[\text{curl } \mathbf{A}]_\mu = -i\sqrt{2} \sum_{\nu\lambda} C_{1\nu 1\lambda}^{1\mu} \nabla_\nu A_\lambda \quad (\mu, \nu, \lambda = \pm 1, 0) \quad (1.3.29)$$

The spherical components of $\text{curl } \mathbf{A}$ may also be written as

$$\text{curl } \mathbf{A} = i \begin{vmatrix} \mathbf{e}_{+1} & \mathbf{e}_0 & \mathbf{e}_{-1} \\ \nabla_{+1} & \nabla_0 & \nabla_{-1} \\ A_{+1} & A_0 & A_{-1} \end{vmatrix}. \quad (1.3.30)$$

The polar components of $\text{curl } \mathbf{A}$ read

$$\begin{aligned} [\text{curl } \mathbf{A}]_r &= \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta A_\varphi) - \frac{1}{r \sin \vartheta} \frac{\partial A_\vartheta}{\partial \varphi}, \\ [\text{curl } \mathbf{A}]_\vartheta &= \frac{1}{r \sin \vartheta} \frac{\partial A_r}{\partial \varphi} - \frac{1}{r} \frac{\partial}{\partial r} (r A_\varphi), \\ [\text{curl } \mathbf{A}]_\varphi &= \frac{1}{r} \frac{\partial}{\partial r} (r A_\vartheta) - \frac{1}{r} \frac{\partial A_r}{\partial \vartheta}. \end{aligned} \quad (1.3.31)$$

The above equations are summarized in Table 1.3.

Note also the following differential operations of the second order

$$\text{div grad } \Phi = \nabla \cdot (\nabla \Phi) = \Delta \Phi. \quad (1.3.32)$$

Table 1.3. Differential operations.

	Cartesian coordinates	Spherical coordinates	Polar coordinates
$\mathbf{r}$	$\mathbf{e}_x x + \mathbf{e}_y y + \mathbf{e}_z z$	$-\mathbf{e}_{+1} x_{-1} + \mathbf{e}_0 x_0 - \mathbf{e}_{-1} x_{+1}$	$\mathbf{e}_r r$
$d\mathbf{r}$	$\mathbf{e}_x dx + \mathbf{e}_y dy + \mathbf{e}_z dz$	$-\mathbf{e}_{+1} dx_{-1} + \mathbf{e}_0 dx_0 - \mathbf{e}_{-1} dx_{+1}$	$\mathbf{e}_r dr + \mathbf{e}_\vartheta r d\vartheta + \mathbf{e}_\varphi r \sin \vartheta d\varphi$
ds^2	$dx^2 + dy^2 + dz^2$	$-2dx_{+1} dx_{-1} + dx_0 dx_0$	$dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\varphi^2$
dV	$dx dy dz$	$i dx_{+1} dx_0 dx_{-1}$	$r^2 \sin \vartheta dr d\vartheta d\varphi$
$\nabla \Phi$	$\mathbf{e}_x \frac{\partial \Phi}{\partial x} + \mathbf{e}_y \frac{\partial \Phi}{\partial y} + \mathbf{e}_z \frac{\partial \Phi}{\partial z}$	$-\mathbf{e}_{+1} \nabla_{-1} \Phi + \mathbf{e}_0 \nabla_0 \Phi - \mathbf{e}_{-1} \nabla_{+1} \Phi$	$\mathbf{e}_r \frac{\partial \Phi}{\partial r} + \mathbf{e}_\vartheta \frac{1}{r} \frac{\partial \Phi}{\partial \vartheta} + \mathbf{e}_\varphi \frac{1}{r \sin \vartheta} \frac{\partial \Phi}{\partial \varphi}$
$(\nabla \cdot \mathbf{A})$	$\frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z}$	$-\nabla_{+1} A_{-1} + \nabla_0 A_0 - \nabla_{-1} A_{+1}$	$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 A_r) + \frac{1}{r \sin \vartheta} \frac{\partial}{\partial \vartheta} (A_\vartheta \sin \vartheta) + \frac{1}{r \sin \vartheta} \frac{\partial A_\varphi}{\partial \varphi}$
$[\nabla \times \mathbf{A}]$	$\begin{vmatrix} \mathbf{e}_x & \mathbf{e}_y & \mathbf{e}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_x & A_y & A_z \end{vmatrix}$	$i \begin{vmatrix} e_{+1} & e_0 & e_{-1} \\ \nabla_{+1} & \nabla_0 & \nabla_{-1} \\ A_{+1} & A_0 & A_{-1} \end{vmatrix}$	$\mathbf{e}_r \frac{1}{r \sin \vartheta} \left[\frac{\partial}{\partial \vartheta} (A_\varphi \sin \vartheta) - \frac{\partial A_\vartheta}{\partial \varphi} \right] + \mathbf{e}_\vartheta \frac{1}{r} \left[\frac{1}{\sin \vartheta} \frac{\partial A_r}{\partial \varphi} - \frac{\partial}{\partial r} (r A_\varphi) \right] + \mathbf{e}_\varphi \frac{1}{r} \left[\frac{\partial}{\partial r} (r A_\vartheta) - \frac{\partial A_r}{\partial \vartheta} \right]$
$\Delta \Phi$	$\frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial y^2} + \frac{\partial^2 \Phi}{\partial z^2}$	$-2\nabla_{+1} \nabla_{-1} \Phi + \nabla_0 \nabla_0 \Phi$	$\frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial \Phi}{\partial r}) + \frac{1}{r^2 \sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta \frac{\partial \Phi}{\partial \vartheta}) + \frac{1}{r^2 \sin^2 \vartheta} \frac{\partial^2 \Phi}{\partial \varphi^2}$

$$\text{curl grad } \Phi = \nabla \times (\nabla \Phi) = 0, \quad (1.3.33)$$

$$\text{div curl } \mathbf{A} = \nabla \cdot [\nabla \times \mathbf{A}] = 0, \quad (1.3.34)$$

$$\begin{aligned} \text{curl curl } \mathbf{A} &= \nabla \times [\nabla \times \mathbf{A}] = \nabla(\nabla \cdot \mathbf{A}) - \Delta \mathbf{A} \\ &= \text{grad div } \mathbf{A} - \Delta \mathbf{A}. \end{aligned} \quad (1.3.35)$$

1.4 Rotations of Coordinate System

An arbitrary rotation of a coordinate system about the origin is completely specified by three real parameters. The most useful description of rotation is that in terms of the Euler angles α, β, γ . Note that two other sets of parameters are also widely used to describe rotations:

- direction of the rotation axis $n(\Theta, \Phi)$ (2 parameters) and the rotation angle ω (1 parameter);
- the Cayley-Klein parameters.

1.4.1 Description of Rotations in Terms of the Euler Angles

Any rotation of the coordinate system $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ may be performed by three successive rotations about the coordinate axes (Fig. 1.3)

- A** $\left\{ \begin{array}{l} \text{(a) rotation about the } z\text{-axis through an angle } \alpha (0 \leq \alpha < 2\pi), \\ \text{(b) rotation about the new } y_1\text{-axis through an angle } \beta (0 \leq \beta \leq \pi), \\ \text{(c) rotation about the new axis } z_2 = z' \text{ through an angle } \gamma (0 \leq \gamma < 2\pi). \end{array} \right.$

The same rotation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ may also be performed by another succession of rotations (Fig. 1.4),




Figure 1.3. Succession of rotations of a coordinate system according to scheme A.

- B** $\left\{ \begin{array}{l} \text{(a) rotation about the } z\text{-axis through an angle } \gamma (0 \leq \gamma < 2\pi), \\ \text{(b) rotation about the initial } y\text{-axis through an angle } \beta (0 \leq \beta \leq \pi), \\ \text{(c) rotation about the initial } z\text{-axis through an angle } \alpha (0 \leq \alpha < 2\pi). \end{array} \right.$

Here the angles α, β, γ are the same as those in the first case.

The relative orientations of initial and final coordinate axes $S\{x, y, z\}$ and $S'\{x', y', z'\}$, obtained in both cases, **A** and **B**, are shown in Fig. 1.5.

The angles α, β, γ are called the Euler angles. They completely define the rotation of the coordinate system. The inverse rotation which returns the coordinate system $S'\{x', y', z'\}$ back into $S\{x, y, z\}$ is specified by the Euler angles $-\gamma, -\beta, -\alpha$, or, equivalently, by the angles $\pi - \gamma, \beta, -\pi - \alpha$.

Sometimes the following successive rotations are used to obtain the general rotation of the coordinate system:

- (a) rotation about the z -axis through an angle α' ;
- (b) rotation about the new x_1 -axis through an angle β' ;
- (c) rotation about the new axis $z_2 = z'$ through an angle γ' .





Figure 1.4. Succession of rotations of a coordinate system according to scheme B.

Figure 1.5. The Euler angles α, β, γ .

The angles α', β', γ' describe the same rotation of the coordinate system as the Euler angles α, β, γ if

$$\alpha' = \alpha + \frac{\pi}{2}, \quad \beta' = \beta, \quad \gamma' = \gamma - \frac{\pi}{2}. \quad (1.4.1)$$

The absolute value of a vector is invariant with respect to rotations, but the polar angles ϑ, φ , which determine the vector direction, change. The relations between angles ϑ, φ and ϑ', φ' which specify vector directions in the initial and final coordinate systems, $S\{x, y, z\}$ and $S'\{x', y', z'\}$, are given by

$$\begin{aligned} \cos \vartheta' &= \cos \vartheta \cos \beta + \sin \vartheta \sin \beta \cos(\varphi - \alpha), \\ \cot(\varphi' + \gamma) &= \cot(\varphi - \alpha) \cos \beta - \frac{\cot \vartheta \sin \beta}{\sin(\varphi - \alpha)}. \end{aligned} \quad (1.4.2)$$

The inverse relations are

$$\begin{aligned} \cos \vartheta &= \cos \vartheta' \cos \beta - \sin \vartheta' \sin \beta \cos(\varphi' + \gamma), \\ \cot(\varphi - \alpha) &= \cot(\varphi' + \gamma) \cos \beta + \frac{\cot \vartheta' \sin \beta}{\sin(\varphi' + \gamma)}. \end{aligned} \quad (1.4.3)$$

1.4.2 Description of Rotations in Terms of Rotation Axis and Rotation Angle

Any rotation of a coordinate system $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ may be treated as one rotation through an angle ω ($0 \leq \omega \leq \pi$) about some axis $\mathbf{n}(\Theta, \Phi)$. The direction of this rotation axis $\mathbf{n}$ is defined by the polar angles




Figure 1.6. Rotation of a coordinate system through an angle ω about an axis $\mathbf{n}(\Theta, \Phi)$.

$\Theta, \Phi (0 \leq \Theta \leq \pi, 0 \leq \Phi < 2\pi)$ which are the same in the initial coordinate system $S\{x, y, z\}$ and in the final one $S'\{x', y', z'\}$ (Fig. 1.6).

The angles ω, Θ, Φ completely determine the relative orientation of the initial and final coordinate axes. The rotation defined by the angles $-\omega, \pi - \Theta, \pi + \Phi$ is identical to the rotation defined by the angles ω, Θ, Φ . The inverse rotation $S'\{x', y', z'\} \rightarrow S\{x, y, z\}$ is specified by the angles $-\omega, \Theta, \Phi$ or, equivalently, by the angles $\omega, \pi - \Theta, \pi + \Phi$.

The direction cosines of the unit vector $\mathbf{n}(\Theta, \Phi)$ in the initial coordinate system $S\{x, y, z\}$ coincide with those in the final coordinate system $S'\{x', y', z'\}$.

$$\begin{aligned} \mathbf{n} \cdot \mathbf{e}_x &= \mathbf{n} \cdot \mathbf{e}'_x = \sin \Theta \cos \Phi, \\ \mathbf{n} \cdot \mathbf{e}_y &= \mathbf{n} \cdot \mathbf{e}'_y = \sin \Theta \sin \Phi, \\ \mathbf{n} \cdot \mathbf{e}_z &= \mathbf{n} \cdot \mathbf{e}'_z = \cos \Theta. \end{aligned} \quad (1.4.4)$$

The polar angles ϑ, φ of vectors not parallel to the $\mathbf{n}$ -axis vary under coordinate rotations. The relations between the polar angles ϑ, φ and ϑ', φ' which specify the direction of a vector with respect to $S\{x, y, z\}$ and $S'\{x', y', z'\}$, respectively, are given by

$$\begin{aligned} \cos \vartheta' &= \cos \vartheta (\cos \omega \sin^2 \Theta + \cos^2 \Theta) + \sin \vartheta \sin \Theta [(1 - \cos \omega) \cos \Theta \cos(\varphi - \Phi) - \sin \omega \sin(\varphi - \Phi)], \\ \cot(\varphi' - \Phi) &= \frac{\cos(\varphi - \Phi) [\cos \omega \cos^2 \Theta + \sin^2 \Theta] + \sin(\varphi - \Phi) \sin \omega \cos \Theta - \cot \vartheta (\cos \omega - 1) \sin \Theta \cos \Theta}{-\cos(\varphi - \Phi) \sin \omega \cos \Theta + \sin(\varphi - \Phi) \cos \omega + \cot \vartheta \sin \omega \sin \Theta}. \end{aligned} \quad (1.4.5)$$

The inverse relations are

$$\begin{aligned} \cos \vartheta &= \cos \vartheta' (\cos \omega \sin^2 \Theta + \cos^2 \Theta) + \sin \vartheta' \sin \Theta [(1 - \cos \omega) \cos \Theta \cos(\varphi' - \Phi) + \sin \omega \sin(\varphi' - \Phi)], \\ \cot(\varphi - \Phi) &= \frac{\cos(\varphi' - \Phi) [\cos \omega \cos^2 \Theta + \sin^2 \Theta] - \sin(\varphi' - \Phi) \sin \omega \cos \Theta - \cot \vartheta' (\cos \omega - 1) \sin \Theta \cos \Theta}{\cos(\varphi' - \Phi) \sin \omega \cos \Theta + \sin(\varphi' - \Phi) \cos \omega - \cot \vartheta' \sin \omega \sin \Theta}. \end{aligned} \quad (1.4.6)$$

Introducing parameters $\omega, \mathbf{n}(\theta, \Phi)$ to describe rotations, we are able to write the transformation properties of components of the position vector $\mathbf{r}$ in compact vector form:

$$\begin{aligned} \mathbf{r}' &= \mathbf{r} \cos \omega + \mathbf{n}(\mathbf{n} \cdot \mathbf{r})(1 - \cos \omega) + [\mathbf{n} \times \mathbf{r}] \sin \omega, \\ \mathbf{r} &= \mathbf{r}' \cos \omega + \mathbf{n}(\mathbf{n} \cdot \mathbf{r}') (1 - \cos \omega) - [\mathbf{n} \times \mathbf{r}'] \sin \omega. \end{aligned} \quad (1.4.7)$$

Equations (1.4.5)-(1.4.6) may be derived by projecting Eq. (1.4.7) onto the coordinate axes.

1.4.3 Description of Rotations in Terms of Unitary 2×2 Matrices. Cayley-Klein Parameters.

The position vector of an arbitrary point $\mathbf{r} = x\mathbf{e}_x + y\mathbf{e}_y + z\mathbf{e}_z$ may be represented by the following Hermitian 2×2 matrix X :

$$X = X^\dagger = \begin{pmatrix} z & x + iy \\ x - iy & -z \end{pmatrix} = \sum_{i=x,y,z} x_i \tilde{\sigma}_i, \quad (1.4.8)$$

where $\tilde{\sigma}_i (i = x, y, z)$ are the transposed Pauli matrices (Eq. (2.5.4)). Note that

$$-\det X = \mathbf{r}^2 = x^2 + y^2 + z^2$$

Each rotation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ may be represented by a unitary transformation U of matrix X into X'

$$X' = UXU^{-1}. \quad (1.4.9)$$

Here U is the unitary unimodular 2×2 matrix

$$U^\dagger = U^{-1}, \quad \det U = 1. \quad (1.4.10)$$




Figure 1.7. Stereographic projection of a point on a sphere.

Bearing in mind that $\mathbf{r}$ is real, one can easily prove that U , indeed, is unitary. The relation $\det U = 1$ ensures the invariance of the absolute value of the position vector under coordinate rotations.

Equations (1.4.10) imply the following form of U

$$U = \begin{pmatrix} a & b \\ -b^* & a^* \end{pmatrix} \quad (1.4.11)$$

where a and b are complex numbers which satisfy the condition

$$|a|^2 + |b|^2 = 1 \quad (1.4.12)$$

Thus, the matrix U depends on three real independent parameters. The numbers a and b are called the Cayley-Klein parameters. They uniquely determine rotation of the coordinate system. The inverse statement is not true because the parameters $-a$ and $-b$ describe the same rotation as a and b .

The inverse rotation $S' \{x', y', z'\} \rightarrow S \{x, y, z\}$ corresponds to the matrix

$$U^{-1} = U^+ = \begin{pmatrix} a^* & -b \\ b^* & a \end{pmatrix}. \quad (1.4.13)$$

The Cayley-Klein parameters permit us to propose an alternative interpretation of coordinate rotations. Let us consider a sphere (of unit diameter) about the origin. Each point of the sphere with coordinates x, y, z ($x^2 + y^2 + z^2 = 1/4$) corresponds to the point $\varsigma = \xi + i\eta$ on the complex plane which is called the stereographic projection of the point x, y, z (Fig. 1.7). The complex number ς is related to x, y, z by

$$\varsigma = \frac{x + iy}{\frac{1}{2} - z} = \frac{\frac{1}{2} + z}{x - iy} \quad (1.4.14)$$

The rotation of the coordinate system which transforms the coordinates x, y, z of the point on the sphere into x', y', z' generates the following complex-plane bilinear transformation

$$\varsigma' = \frac{a\varsigma + b}{-b^*\varsigma + a^*}. \quad (1.4.15)$$

The coefficients of this bilinear transformation are just the Cayley-Klein parameters a and b .

1.4.4 Relations Between Different Descriptions of Rotations

(a) *Relations Between Angles ω, Θ, Φ and Euler Angles α, β, γ*

The angles ω, Θ, Φ are expressed in terms of the Euler angles α, β, γ by

$$\begin{aligned} \cos \frac{\omega}{2} &= \cos \frac{\beta}{2} \cos \frac{\alpha + \gamma}{2} \\ \tan \Theta &= \frac{\tan \frac{\beta}{2}}{\sin \frac{\alpha + \gamma}{2}} \\ \Phi &= \frac{\pi}{2} + \frac{\alpha - \gamma}{2} \end{aligned} \quad (1.4.16)$$

The inverse relations are written as

$$\begin{aligned} \sin \frac{\beta}{2} &= \sin \Theta \sin \frac{\omega}{2}, \\ \tan \frac{\alpha + \gamma}{2} &= \cos \Theta \tan \frac{\omega}{2}, \\ \frac{\alpha - \gamma}{2} &= \Phi - \frac{\pi}{2}. \end{aligned} \quad (1.4.17)$$

Note also the following useful relations between the angles α, β, γ and ω, Θ, Φ .

$$\begin{aligned} \frac{\partial \alpha}{\partial \omega} = \frac{\partial \gamma}{\partial \omega} &= \frac{\cos \Theta}{2 \cos^2 \frac{\beta}{2}}, & \frac{\partial \beta}{\partial \omega} &= \frac{\sin^2 \Theta \sin \omega}{\sin \beta}, \\ \frac{\partial \alpha}{\partial \Theta} = \frac{\partial \gamma}{\partial \Theta} &= -\frac{1}{2} \tan \Theta \sin(\alpha + \gamma), & \frac{\partial \beta}{\partial \Theta} &= \frac{2 \sin 2\Theta \sin^2 \frac{\omega}{2}}{\sin \beta}, \\ \frac{\partial \alpha}{\partial \Phi} = -\frac{\partial \gamma}{\partial \Phi} &= 1, & \frac{\partial \beta}{\partial \Phi} &= 0, \end{aligned} \quad (1.4.18)$$

$$\begin{aligned} \frac{\partial \omega}{\partial \alpha} = \frac{\partial \omega}{\partial \gamma} &= \cos \Theta, & \frac{\partial \omega}{\partial \beta} &= \frac{\tan \frac{\beta}{2}}{\tan \frac{\omega}{2}}, \\ \frac{\partial \Theta}{\partial \alpha} = \frac{\partial \Theta}{\partial \gamma} &= -\frac{\sin \Theta}{2 \tan \frac{\omega}{2}}, & \frac{\partial \Theta}{\partial \beta} &= \frac{\cos \Theta \sin \Theta}{\sin \beta}, \\ \frac{\partial \Phi}{\partial \alpha} = -\frac{\partial \Phi}{\partial \gamma} &= \frac{1}{2}, & \frac{\partial \Phi}{\partial \beta} &= 0. \end{aligned} \quad (1.4.19)$$

The Jacobian of the transformation is equal to

$$\left\| \frac{\partial(\alpha, \beta, \gamma)}{\partial(\omega, \Theta, \Phi)} \right\| = \left\| \frac{\partial(\omega, \Theta, \Phi)}{\partial(\alpha, \beta, \gamma)} \right\|^{-1} = \frac{4 \sin \Theta}{\sin \beta} \sin^2 \frac{\omega}{2}. \quad (1.4.20)$$

A volume element of the three-dimensional rotation group is given by

$$dR \equiv \sin \beta d\alpha d\beta d\gamma = 4 \sin^2 \frac{\omega}{2} d\omega \sin \Theta d\Theta d\Phi \quad (1.4.21)$$

The total volume of the three-dimensional rotation group is equal to

$$\int dR = \int_0^{2\pi} d\alpha \int_0^\pi \sin \beta d\beta \int_0^{2\pi} d\gamma = 4 \int_0^\pi \sin^2 \frac{\omega}{2} d\omega \int_0^\pi \sin \Theta d\Theta \int_0^{2\pi} d\Phi = 8\pi^2. \quad (1.4.22)$$

(b) *Relations Between Cayley-Klein Parameters a, b and Euler Angles α, β, γ*

The parameters a and b are expressed in terms of the Euler angles α, β, γ as

$$\begin{aligned} a &= \cos \frac{\beta}{2} e^{-i \frac{\alpha+\gamma}{2}}, \\ b &= \sin \frac{\beta}{2} e^{i \frac{\alpha-\gamma}{2}}. \end{aligned} \quad (1.4.23)$$

while the inverse relations are

$$\begin{aligned} \cos \beta &= |a|^2 - |b|^2, \quad \cos \frac{\beta}{2} = |a|, \quad \sin \frac{\beta}{2} = |b| \\ \cot \frac{\alpha+\gamma}{2} &= -\frac{\operatorname{Re} a}{\operatorname{Im} a}, \quad \cot \frac{\alpha-\gamma}{2} = \frac{\operatorname{Re} b}{\operatorname{Im} b} \end{aligned} \quad (1.4.24)$$

The parameters a and b may be expressed in terms of the Wigner D -functions (see Chap. 4)

$$a = D_{\frac{1}{2}\frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma), \quad b = D_{-\frac{1}{2}\frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma). \quad (1.4.25)$$

The unitary matrix U , according to Eq. (1.4.11), coincides with the transposed rotation matrix $\widehat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ which transforms spin functions of particles of spin $\frac{1}{2}$ (see Eq. (2.5.32)).

(c) *Relations Between Cayley-Klein Parameters a, b and Angles ω, Θ, Φ*

The parameters a, b are expressed in terms of the angles ω, Θ, Φ as

$$\begin{aligned} a &= \cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \Theta, \\ b &= -i \sin \frac{\omega}{2} \sin \Theta e^{i\Phi}. \end{aligned} \quad (1.4.26)$$

and the inverse relations have the form

$$\begin{aligned} \cos \omega &= 2(\operatorname{Re} a)^2 - 1, \quad \cos \frac{\omega}{2} = \operatorname{Re} a \\ \cos \Theta &= -\frac{\operatorname{Im} a}{\sqrt{1 - (\operatorname{Re} a)^2}}, \\ \cot \Phi &= -\frac{\operatorname{Im} b}{\operatorname{Re} b}. \end{aligned} \quad (1.4.27)$$

The unitary matrix U , according to Eq. (1.4.11), coincides with the transposed rotation matrix $\widehat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi)$ which transforms spin functions of spin $\frac{1}{2}$ (see Eq. (2.5.36)).

1.4.5 Rotation Operator

Under rotations of coordinate systems quantum-mechanical quantities are transformed by the rotation operators $\widehat{D}(\alpha, \beta, \gamma)$ or $\widehat{U}(\omega; \Theta, \Phi)$.

Wave functions (state vectors) Ψ' and operators $\widehat{O}'$ in a rotated coordinate system are related to wave functions Ψ and operators $\widehat{O}$ in an initial coordinate system by

$$\Psi' = \widehat{D}(\alpha, \beta, \gamma)\Psi, \widehat{O}' = \widehat{D}(\alpha, \beta, \gamma)\widehat{O}[\widehat{D}(\alpha, \beta, \gamma)]^{-1}, \quad (1.4.28)$$

$$\Psi' = \widehat{U}(\omega; \Theta, \Phi)\Psi, \widehat{O}' = \widehat{U}(\omega; \Theta, \Phi)\widehat{O}[\widehat{U}(\omega; \Theta, \Phi)]^{-1}. \quad (1.4.29)$$

If the Euler angles α, β, γ are chosen to describe the rotation, then the rotation operator $\hat{D}(\alpha, \beta, \gamma)$ may be written as

$$\hat{D}(\alpha, \beta, \gamma) = e^{-i\gamma\hat{J}_z} e^{-i\beta\hat{J}_y} e^{-i\alpha\hat{J}_z} \quad (1.4.30)$$

or, equivalently,

$$\hat{D}(\alpha, \beta, \gamma) = e^{-i\alpha\hat{J}_z} e^{-i\beta\hat{J}_y} e^{-i\gamma\hat{J}_z}. \quad (1.4.31)$$

Here $\hat{J}_i$ is the projection of the total angular momentum operator (see Chap. 2) on an i -axis. The equivalence of Eqs. (1.4.30) and (1.4.31) follows from the fact that, according to (1.4.28),

$$\begin{aligned} e^{-i\beta\hat{J}_y} &= \hat{D}(\alpha, 0, 0) e^{-i\beta\hat{J}_y} [\hat{D}(\alpha, 0, 0)]^{-1} = e^{-i\alpha\hat{J}_z} e^{-i\beta\hat{J}_y} e^{i\alpha\hat{J}_z}, \\ e^{-i\gamma\hat{J}_z} &= \hat{D}(\alpha, \beta, 0) e^{-i\gamma\hat{J}_z} [\hat{D}(\alpha, \beta, 0)]^{-1} = e^{-i\alpha\hat{J}_z} e^{-i\beta\hat{J}_y} e^{-i\gamma\hat{J}_z} e^{i\beta\hat{J}_y} e^{i\alpha\hat{J}_z}. \end{aligned} \quad (1.4.32)$$

If the direction of the rotation axis $n(\Theta, \Phi)$ and the rotation angle ω are chosen to describe the coordinate rotation, the rotation operator $\hat{U}$ may be written in the form

$$\hat{U}(\omega; \Theta, \Phi) = e^{-i\omega n \cdot \hat{\mathbf{J}}}, \quad (1.4.33)$$

where $\hat{\mathbf{J}}$ is the total angular momentum operator (Chap. 2). Note that $\hat{D}(\alpha, \beta, \gamma) \equiv \hat{U}(\omega; \Theta, \Phi)$. The rotation operator written in the forms (1.4.30), (1.4.31) or (1.4.33) is an unitary operator.

$$\begin{aligned} \hat{D}^+(\alpha, \beta, \gamma) &= [\hat{D}(\alpha, \beta, \gamma)]^{-1} = \hat{D}(\pi - \gamma, \beta, -\pi - \alpha) = \hat{D}(-\gamma, -\beta, -\alpha), \\ \hat{U}^+(\omega; \Theta, \Phi) &= [\hat{U}(\omega; \Theta, \Phi)]^{-1} = \hat{U}(\omega; \pi - \Theta, \pi + \Phi) = \hat{U}(-\omega; \Theta, \Phi). \end{aligned} \quad (1.4.34)$$

Matrix elements of $\hat{D}$ between eigenstates of the operators $\hat{J}^2, \hat{J}_z$ are the Wigner D -functions (see Chap. 4)

$$\langle J' M' | \hat{D}(\alpha, \beta, \gamma) | J M \rangle = \delta_{JJ'} D_{M' M}^J(\alpha, \beta, \gamma). \quad (1.4.35)$$

Matrix elements of $\hat{D}$ between states corresponding to the cartesian basis vectors $e_i (i = x, y, z)$ coincide with elements of the rotation matrix a_{ik} (see Sec. 1.4.6)

$$\langle \mathbf{e}_i | \hat{D}(\alpha, \beta, \gamma) | \mathbf{e}_k \rangle = a_{ik}, \quad (i, k = x, y, z) \quad (1.4.36)$$

Effects of the rotation operator on various wave functions and quantum-mechanical operators are considered in Chaps. 3, 5-7.

1.4.6 Transformation of Cartesian Vectors and Tensors Under Rotations of Coordinate Systems. Rotation Matrix $\mathbf{A}$

An arbitrary vector $\mathbf{A}$ may be written as a column

$$\mathbf{A} = \begin{pmatrix} A_x \\ A_y \\ A_z \end{pmatrix}, \quad (1.4.37)$$

where A_x, A_y, A_z are cartesian components of $\mathbf{A}$. In this representation the Cartesian basis vectors $\mathbf{e}_x, \mathbf{e}_y, \mathbf{e}_z$ have the form

$$\mathbf{e}_x = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{e}_y = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \mathbf{e}_z = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (1.4.38)$$

The effect of the rotation operator on the basis vectors written in such a form is equivalent to an action of some 3×3 matrix a which may be regarded as one of the representations of the rotation operator

$$a = \begin{pmatrix} a_{xx} & a_{xy} & a_{xz} \\ a_{yx} & a_{yy} & a_{yz} \\ a_{zx} & a_{zy} & a_{zz} \end{pmatrix}. \quad (1.4.39)$$

The matrix a is real

$$a^* = a, \quad a_{ik}^* = a_{ik}, \quad (i, k = x, y, z). \quad (1.4.40)$$

and unitary

$$a^+ a = a a^+ = 1. \quad (1.4.41)$$

Equations (1.4.40) and (1.4.41) result in the orthogonality condition

$$\tilde{a}a = a\tilde{a} = 1, \quad (1.4.42)$$

where $\tilde{a}$ is the transpose of a . Equation (1.4.42) written in a component form gives six independent relations for the elements a_{ik}

$$\sum_i a_{ik}a_{il} = \delta_{kl}, \quad (i, k, l = x, y, z), \quad (1.4.43)$$

or the equivalent relations

$$\sum_k a_{ik}a_{lk} = \delta_{il}, \quad (i, k, l = x, y, z). \quad (1.4.44)$$

Relations (1.4.43) or (1.4.44) reveal that only three of the nine matrix elements a_{ik} are independent. This result is in agreement with the fact that any rotation of the coordinate system is completely determined by three real parameters.

The matrix a is unimodular, i.e.,

$$\det a = \begin{vmatrix} a_{xx} & a_{xy} & a_{xz} \\ a_{yx} & a_{yy} & a_{yz} \\ a_{zx} & a_{zy} & a_{zz} \end{vmatrix} = 1. \quad (1.4.45)$$

The relations between cartesian basis vectors e'_i in a rotated coordinate system S' and basis vectors e_i in an initial coordinate system S are given by

$$\mathbf{e}'_i = a\mathbf{e}_i = \sum_k a_{ki}\mathbf{e}_k, \quad (i, k = x, y, z). \quad (1.4.46)$$

The transformation properties of cartesian vector components are given by

$$A'_i = \sum_k a_{ki}A_k, \quad (i, k = x, y, z), \quad (1.4.47)$$

where A_k are the components of $\mathbf{A}$ in the initial coordinate system and A'_i are the components of this vector in the rotated coordinate system. Equations (1.4.43) and (1.4.44) ensure that the absolute value of $\mathbf{A}$ is unchanged by the rotation.

The transformation rule for cartesian components of a tensor of rank n (n is integer) has the form

$$A'_{i_1 i_2 \dots i_n} = \sum_{k_1 k_2 \dots k_n} a_{k_1 i_1} a_{k_2 i_2} \dots a_{k_n i_n} A_{k_1 k_2 \dots k_n}. \quad (1.4.48)$$

The inverse transformation which corresponds to the rotation $S' \rightarrow S$ is performed by the transposed matrix $\tilde{a} = a^{-1}$. The inverse relations are

$$\mathbf{e}_k = \sum_i a_{ki}\mathbf{e}'_i, \quad (i, k = x, y, z), \quad (1.4.49)$$

$$A_k = \sum_i a_{ki}A'_i, \quad (i, k = x, y, z), \quad (1.4.50)$$

$$A_{k_1 k_2 \dots k_n} = \sum_{i_1 i_2 \dots i_n} a_{k_1 i_1} a_{k_2 i_2} \dots a_{k_n i_n} A'_{i_1 i_2 \dots i_n}, \quad (i_1, k_1, i_2, k_2, \dots, i_n, k_n = x, y, z). \quad (1.4.51)$$

The elements of the rotation matrix a_{ik} may be evaluated from

$$a_{ik} = \mathbf{e}_i \mathbf{e}'_k, \quad (i, k = x, y, z), \quad (1.4.52)$$

Thus, the elements a_{ik} are cosines of angles between the basis vectors in the initial (S) and rotated (S') coordinate systems. An equivalent definition of a_{ik} in terms of coordinates in S and S' has the form

$$a_{ik} = \frac{\partial x'_i}{\partial x_k} = \frac{\partial x_k}{\partial x'_i}, \quad (x'_i, x_k = x, y, z) \quad (1.4.53)$$

The rotation matrix a is given in terms of the Euler angles by

$$a = \begin{pmatrix} \cos \alpha \cos \beta \cos \gamma - \sin \alpha \sin \gamma & -\cos \alpha \cos \beta \sin \gamma - \sin \alpha \cos \gamma & \cos \alpha \sin \beta \\ \sin \alpha \cos \beta \cos \gamma + \cos \alpha \sin \gamma & -\sin \alpha \cos \beta \sin \gamma + \cos \alpha \cos \gamma & \sin \alpha \sin \beta \\ -\sin \beta \cos \gamma & \sin \beta \sin \gamma & \cos \beta \end{pmatrix}. \quad (1.4.54)$$

The inverse matrix a^{-1} may be obtained from Eq. (1.4.54) by transposing or, equivalently, by replacing the Euler angles $\alpha, \beta, \gamma \rightarrow -\gamma, -\beta, -\alpha$.

The expression for the rotation matrix a in terms of the angles Θ, Φ which describe the direction of the rotation axis, and the rotation angle ω has the form

$$a = \begin{pmatrix} (1 - \cos \omega) \sin^2 \Theta \cos^2 \Phi + \cos \omega & (1 - \cos \omega) \sin^2 \Theta \cos \Phi \sin \Phi + \sin \omega \cos \Theta & (1 - \cos \omega) \sin \Theta \cos \Theta \cos \Phi + \sin \omega \sin \Theta \sin \Phi \\ (1 - \cos \omega) \sin^2 \Theta \cos \Phi \sin \Phi + \sin \omega \cos \Theta & (1 - \cos \omega) \sin^2 \Theta \sin^2 \Phi + \cos \omega & (1 - \cos \omega) \sin \Theta \cos \Theta \sin \Phi - \sin \omega \sin \Theta \cos \Phi \\ (1 - \cos \omega) \sin \Theta \cos \Theta \cos \Phi + \sin \omega \sin \Theta \sin \Phi & (1 - \cos \omega) \sin \Theta \cos \Theta \sin \Phi - \sin \omega \sin \Theta \cos \Phi & (1 - \cos \omega) \cos^2 \Theta + \cos \omega \end{pmatrix}. \quad (1.4.55)$$

Equation (1.4.55) may be expanded into

$$a = \cos \omega \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + (1 - \cos \omega) \begin{pmatrix} n_x^2 & n_x n_y & n_x n_z \\ n_y n_x & n_y^2 & n_y n_z \\ n_z n_x & n_z n_y & n_z^2 \end{pmatrix} + \sin \omega \begin{pmatrix} 0 & -n_z & n_y \\ n_z & 0 & -n_x \\ -n_y & n_x & 0 \end{pmatrix} \quad (1.4.56)$$

where n_x, n_y, n_z are components of the unit vector n which determines direction of the rotation axis. Using (1.4.56) one can easily derive the following expressions for the matrix elements a_{ik}

$$a_{ik} = \cos \omega \delta_{ik} + (1 - \cos \omega) n_i n_k - \sin \omega \varepsilon_{ikl} n_l, \quad (i, k, l = x, y, z). \quad (1.4.57)$$

The inverse matrix a^{-1} may be obtained from (1.4.55) and (1.4.56) by transposing or, equivalently, by replacing $\omega, \Theta, \Phi \rightarrow -\omega, \Theta, \Phi$ or $\omega, \Theta, \Phi \rightarrow \omega, \pi - \Theta, \pi + \Phi$.

The expressions for ω, Θ, Φ in terms of the matrix elements a_{ik} read

$$\begin{aligned} \cos \omega &= \frac{1}{2} [\text{Tr } a - 1] = \frac{1}{2} (a_{xx} + a_{yy} + a_{zz} - 1), \\ n_x \sin \omega &\equiv \sin \omega \sin \Theta \cos \Phi = \frac{1}{2} (a_{zy} - a_{yz}), \\ n_y \sin \omega &\equiv \sin \omega \sin \Theta \sin \Phi = \frac{1}{2} (a_{xz} - a_{zx}), \\ n_z \sin \omega &\equiv \sin \omega \cos \Theta = \frac{1}{2} (a_{yx} - a_{xy}), \\ n_i \sin \omega &= -\frac{1}{2} \sum_{kl} \varepsilon_{ikl} a_{kl}, \quad (i, k, l = x, y, z). \end{aligned} \quad (1.4.58)$$

The rotation matrix a may be rewritten in terms of the Cayley-Klein parameters as

$$a = \begin{pmatrix} \frac{1}{2} (a^2 - b^2 + a^{*2} - b^{*2}) & \frac{i}{2} (-a^2 + b^2 + a^{*2} - b^{*2}) & ab^* + a^*b \\ \frac{i}{2} (a^2 + b^2 - a^{*2} - b^{*2}) & \frac{1}{2} (a^2 + b^2 + a^{*2} + b^{*2}) & i(ab^* - a^*b) \\ -(ab + a^*b^*) & i(ab - a^*b^*) & aa^* - bb^* \end{pmatrix}. \quad (1.4.59)$$

One can see that the parameters a, b and $-a, -b$ correspond to the same rotation matrix.

(a) Particular Forms of Rotation Matrix

(a) Rotation through an angle Ψ about the x -axis:

$$a_x(\Psi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \Psi & -\sin \Psi \\ 0 & \sin \Psi & \cos \Psi \end{pmatrix}. \quad (1.4.60)$$

(b) Rotation through an angle Ψ about the y -axis:

$$a_y(\Psi) = \begin{pmatrix} \cos \Psi & 0 & \sin \Psi \\ 0 & 1 & 0 \\ -\sin \Psi & 0 & \cos \Psi \end{pmatrix}. \quad (1.4.61)$$

(c) Rotation through an angle Ψ about the z -axis:

$$a_z(\Psi) = \begin{pmatrix} \cos \Psi & -\sin \Psi & 0 \\ \sin \Psi & \cos \Psi & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (1.4.62)$$

For an arbitrary rotation determined by the Euler angles α, β, γ the rotation matrix, in accordance with Eq. (1.4.31), may be written in the form

$$a = a_n(\omega) = a_z(\alpha)a_y(\beta)a_z(\gamma). \quad (1.4.63)$$

Equation (1.4.63) represents a particular case of addition of coordinate rotations (see Sec. a).

1.4.7 Addition of Rotations

Let us consider two successive rotations of the coordinate system. Let the first rotation transform the coordinate system $S\{x, y, z\}$ into $S'\{x', y', z'\}$ and the second one transform $S'\{x', y', z'\}$ into $S''\{x'', y'', z''\}$.

Below the parameters describing the resultant rotation $S\{x, y, z\} \rightarrow S''\{x'', y'', z''\}$ will be given in terms of the parameters specifying the rotations $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ and $S'\{x', y', z'\} \rightarrow S''\{x'', y'', z''\}$.

(a) *Description of rotations in terms of Euler angles*

Let both rotations be performed according to the scheme **B(p.22)**. Let the first rotation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ be described by the Euler angles $\alpha_1, \beta_1, \gamma_1$, the second one, $S'\{x', y', z'\} \rightarrow S''\{x'', y'', z''\}$, by the Euler angles $\alpha_2, \beta_2, \gamma_2$ and the resultant rotation, $S\{x, y, z\} \rightarrow S''\{x'', y'', z''\}$ by the Euler angles α, β, γ . The Euler angles $\alpha, \beta, \gamma, \alpha_1, \beta_1, \gamma_1$ and $\alpha_2, \beta_2, \gamma_2$ are supposed to be defined with respect to an initial coordinate system $S\{x, y, z\}$.

The operator of resultant rotation has the form

$$\widehat{D}(\alpha, \beta, \gamma) = \widehat{D}(\alpha_2, \beta_2, \gamma_2) \widehat{D}(\alpha_1, \beta_1, \gamma_1), \quad (1.4.64)$$

or in more detail

$$e^{-i\alpha\hat{J}_z}e^{-i\beta\hat{J}_y}e^{-i\gamma\hat{J}_z} = e^{-i\alpha_2\hat{J}_z}e^{-i\beta_2\hat{J}_y}e^{-i\gamma_2\hat{J}_z}e^{-i\alpha_1\hat{J}_z}e^{-i\beta_1\hat{J}_y}e^{-i\gamma_1\hat{J}_z}. \quad (1.4.65)$$

In Eq. (1.4.65) $\hat{J}_i$ is the projection of the total angular momentum operator on an i -axis of the coordinate system $S\{x, y, z\}$. The angles of the resultant rotation α, β, γ are expressed in terms of $\alpha_1, \beta_1, \gamma_1$ and $\alpha_2, \beta_2, \gamma_2$ as

$$\begin{aligned} \cot(\alpha - \alpha_2) &= \cos \beta_2 \cot(\alpha_1 + \gamma_2) + \cot \beta_1 \frac{\sin \beta_2}{\sin(\alpha_1 + \gamma_2)}, \\ \cos \beta &= \cos \beta_1 \cos \beta_2 - \sin \beta_1 \sin \beta_2 \cos(\alpha_1 + \gamma_2), \\ \cot(\gamma - \gamma_1) &= \cos \beta_1 \cot(\alpha_1 + \gamma_2) + \cot \beta_2 \frac{\sin \beta_1}{\sin(\alpha_1 + \gamma_2)}. \end{aligned} \quad (1.4.66)$$

The following relations are useful for evaluation of α, β, γ .

$$\frac{\sin(\alpha - \alpha_2)}{\sin \beta_1} = \frac{\sin(\gamma - \gamma_1)}{\sin \beta_2} = \frac{\sin(\alpha_1 + \gamma_2)}{\sin \beta}, \quad (1.4.67)$$

$$\begin{aligned} \cos \beta_1 &= \cos \beta \cos \beta_2 + \sin \beta \sin \beta_2 \cos(\alpha - \alpha_2), \\ \cos \beta_2 &= \cos \beta \cos \beta_1 + \sin \beta \sin \beta_1 \cos(\gamma - \gamma_1), \end{aligned} \quad (1.4.68)$$

$$\begin{aligned} \cos \beta &= \cos \beta_1 \cos \beta_2 - \sin \beta_1 \sin \beta_2 \cos(\alpha_1 + \gamma_2), \\ \cos(\gamma - \gamma_1) &= \cos(\alpha_1 + \gamma_2) \cos(\alpha - \alpha_2) + \sin(\alpha_1 + \gamma_2) \sin(\alpha - \alpha_2) \cos \beta_2, \\ \cos(\alpha - \alpha_2) &= \cos(\alpha_1 + \gamma_2) \cos(\gamma - \gamma_1) + \sin(\alpha_1 + \gamma_2) \sin(\gamma - \gamma_1) \cos \beta_1, \\ \cos(\alpha_1 + \gamma_2) &= \cos(\gamma - \gamma_1) \cos(\alpha - \alpha_2) - \sin(\gamma - \gamma_1) \sin(\alpha - \alpha_2) \cos \beta, \end{aligned} \quad (1.4.69)$$

$$\begin{aligned} \frac{\tan \frac{\beta - \beta_1}{2}}{\tan \frac{\beta + \beta_1}{2}} &= \frac{\tan \frac{\alpha_1 + \gamma_2 + \alpha - \alpha_2}{2}}{\tan \frac{\alpha_1 + \gamma_2 - \alpha + \alpha_2}{2}}, \\ \frac{\tan \frac{\beta - \beta_2}{2}}{\tan \frac{\beta + \beta_2}{2}} &= \frac{\tan \frac{\alpha_1 + \gamma_2 + \gamma - \gamma_1}{2}}{\tan \frac{\alpha_1 + \gamma_2 - \gamma + \gamma_1}{2}}, \\ \frac{\tan \frac{\beta_1 - \beta_2}{2}}{\tan \frac{\beta_1 + \beta_2}{2}} &= \frac{\tan \frac{\alpha - \alpha_2 + \gamma - \gamma_1}{2}}{\tan \frac{\alpha - \alpha_2 - \gamma + \gamma_1}{2}}. \end{aligned} \quad (1.4.70)$$

Equations (1.4.67)-(1.4.70) may be easily interpreted in terms of the geometry on a sphere. Each rotation may be completely determined by a point of intersection of the z' -axis with the spherical surface and by a unit vector in the direction of the x' -axis which lies on a plane tangent to the surface at this point. In this case the determination of α, β, γ is reduced to constructing the corresponding spherical triangle, (Fig. 1.8). Equations (1.4.67)-(1.4.70) represent the formulas of sines, cosines and tangents for the spherical triangle.




Figure 1.8. Addition of rotations in terms of the spherical geometry.

Another expression for the angles of the resultant rotation will be obtained if successive rotations are performed according to the scheme **B** (p. 22) but the Euler angles $\alpha_2, \beta_2, \gamma_2$ specifying the second rotation $S' \{x', y', z'\} \rightarrow S'' \{x'', y'', z''\}$ are defined with respect to the intermediate coordinate system $S' \{x', y', z'\}$ rather than the initial system $S \{x, y, z\}$. In this case the operator of the resultant rotation has the form

$$\hat{D}(\alpha, \beta, \gamma) = \hat{D}'(\alpha_2, \beta_2, \gamma_2) \hat{D}(\alpha_1, \beta_1, \gamma_1), \quad (1.4.71)$$

where prime indicates that the operator of the second rotation is taken in the coordinate system $S' \{x', y', z'\}$. According to (1.4.28), the operator $\hat{D}'$ is related to the operator in the initial coordinate system by

$$\hat{D}'(\alpha_2, \beta_2, \gamma_2) = \hat{D}(\alpha_1, \beta_1, \gamma_1) \hat{D}(\alpha_2, \beta_2, \gamma_2) [\hat{D}(\alpha_1, \beta_1, \gamma_1)]^{-1}. \quad (1.4.72)$$

Substitution of this expression into Eq. (1.4.71) yields

$$\hat{D}(\alpha, \beta, \gamma) = \hat{D}(\alpha_1, \beta_1, \gamma_1) \hat{D}(\alpha_2, \beta_2, \gamma_2), \quad (1.4.73)$$

i.e., the operator of the resultant rotation differs from (1.4.64) in the order of operators of the first and second rotations. Thus, for such a description of the successive rotations the Euler angles α, β, γ may be obtained from (1.4.66)-(1.4.70) by interchange of indices $1 \rightleftharpoons 2$.

Finally, if successive rotations are performed according to the scheme **A** (p. 21), i.e., if each rotation is made about the corresponding new axis, the operator of resultant rotation is given by Eq. (1.4.73). In this case the Euler angles α, β, γ of the resultant rotation may also be derived from (1.4.66)-(1.4.70) by the interchange of indices $1 \rightleftharpoons 2$.

(b) *Description of Rotations in Terms of Rotation Axis $\mathbf{n}(\Theta, \Phi)$ and Rotation Angle ω*

Let the first rotation, $S \{x, y, z\} \rightarrow S' \{x', y', z'\}$, be performed about an axis $\mathbf{n}_1$ through an angle ω_1 , and the second one, $S' \{x', y', z'\} \rightarrow S'' \{x'', y'', z''\}$, be about an axis $\mathbf{n}_2$ through an angle ω_2 . The resultant rotation $S \{x, y, z\} \rightarrow S'' \{x'', y'', z''\}$ may be treated as a rotation about an axis $\mathbf{n}$ through an angle ω .

The operator of the resultant rotation has the form

$$e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{J}}} = e^{-i\omega_2 \mathbf{n}_2 \cdot \hat{\mathbf{J}}} e^{-i\omega_1 \mathbf{n}_1 \cdot \hat{\mathbf{J}}}. \quad (1.4.74)$$

The angle of the resultant rotation ω and the axis of this rotation $\mathbf{n}$ are determined by

$$\begin{aligned} \cos \frac{\omega}{2} &= \cos \frac{\omega_1}{2} \cos \frac{\omega_2}{2} - (\mathbf{n}_1 \cdot \mathbf{n}_2) \sin \frac{\omega_1}{2} \sin \frac{\omega_2}{2}, \\ \mathbf{n} \sin \frac{\omega}{2} &= \mathbf{n}_1 \sin \frac{\omega_1}{2} \cos \frac{\omega_2}{2} + \mathbf{n}_2 \sin \frac{\omega_2}{2} \cos \frac{\omega_1}{2} - [\mathbf{n}_1 \times \mathbf{n}_2] \sin \frac{\omega_1}{2} \sin \frac{\omega_2}{2}. \end{aligned} \quad (1.4.75)$$

It follows from Eq. (1.4.75) that the resultant rotation is independent of the order of successive rotations (i.e., the rotation operators commute) if and only if $n_1 \times n_2 = 0$, i.e., the axes of both rotations are parallel or antiparallel. In this case

$$\omega = \omega_1 \pm \omega_2$$

If directions of the rotation axes $\mathbf{n}_1, \mathbf{n}_2, \mathbf{n}$ are specified by the polar angles $\Theta_1, \Phi_1; \Theta_2, \Phi_2$ and Θ, Φ , respectively, and the polar angles are defined with respect to the initial coordinate system $S\{x, y, z\}$, then

$$\begin{aligned} \cos \frac{\omega}{2} &= \cos \frac{\omega_1}{2} \cos \frac{\omega_2}{2} - \sin \frac{\omega_1}{2} \sin \frac{\omega_2}{2} [\cos \Theta_1 \cos \Theta_2 + \sin \Theta_1 \sin \Theta_2 \cos (\Phi_1 - \Phi_2)] \\ \sin \frac{\omega}{2} \cos \Theta &= \sin \frac{\omega_1}{2} \cos \frac{\omega_2}{2} \cos \Theta_1 + \sin \frac{\omega_2}{2} \cos \frac{\omega_1}{2} \cos \Theta_2 \\ &\quad + \sin \frac{\omega_1}{2} \sin \frac{\omega_2}{2} \sin \Theta_1 \sin \Theta_2 \sin (\Phi_1 - \Phi_2) \\ \cot \Phi &= \frac{\sin \Theta_1 (\cot \frac{\omega_2}{2} \cos \Phi_1 - \cos \Theta_2 \sin \Phi_1) + \sin \Theta_2 (\cot \frac{\omega_1}{2} \cos \Phi_2 + \cos \Theta_1 \sin \Phi_2)}{\sin \Theta_1 (\cot \frac{\omega_2}{2} \sin \Phi_1 + \cos \Theta_2 \cos \Phi_1) + \sin \Theta_2 (\cot \frac{\omega_1}{2} \sin \Phi_2 - \cos \Theta_1 \cos \Phi_2)}. \end{aligned} \quad (1.4.76)$$

If directions of the rotation axes n_1, n are defined by the polar angles Θ_1, Φ_1 and Θ, Φ with respect to the initial coordinate system $S\{x, y, z\}$ and the direction of the axis of the second rotation $\mathbf{n}_2$ is defined by the polar angles Θ_2, Φ_2 with respect to the intermediate coordinate system $S'\{x', y', z'\}$ then the angles ω, Θ, Φ of the resultant rotation will be given by Eqs. (1.4.76) with interchanged indices $1 \rightleftharpoons 2$. This situation is similar to the case when rotations are described by Euler angles.

(c) Description of Rotation in Terms of Cayley-Klein Parameters

Let the first rotation, $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ be determined by the Cayley-Klein parameters a_1, b_1 (1.4.3) and the second one, $S'\{x', y', z'\} \rightarrow S''\{x'', y'', z''\}$, by the parameters a_2, b_2 . Then the resultant rotation $S\{x, y, z\} \rightarrow S''\{x'', y'', z''\}$ will be determined by the parameters a, b such as

$$\begin{aligned} a &= a_1 a_2 - b_1^* b_2, \\ b &= a_1^* b_2 + b_1 a_2. \end{aligned} \quad (1.4.77)$$

The matrix U (see Eq. (1.4.11)) which describes the resultant rotation is a product of matrices corresponding to the first and second rotations

$$U(a, b) = U(a_2, b_2) U(a_1, b_1). \quad (1.4.78)$$

In this case all the matrices are supposed to be given in an initial coordinate system.

(d) Addition Theorem for Rotation Matrices a

Let us carry out two successive rotations $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ and $S'\{x', y', z'\} \rightarrow S''\{x'', y'', z''\}$. The matrix which transforms cartesian components of vectors and tensors under the resultant rotation $S\{x, y, z\} \rightarrow S''\{x'', y'', z''\}$ represents a product of the matrices $a(1)$ and $a(2)$ corresponding to the first and second rotations. The order of these matrices in the product depends on the convention used for the rotation angles. If all angles which describe rotations are referred to the initial coordinate system, i.e., the operator of the resultant rotation is given by Eq. (1.4.64), then

$$a = a(2)a(1), \quad (1.4.79)$$

or, in terms of matrix elements,

$$a_{ik} = \sum_l a_{il}(2) a_{lk}(1), \quad (i, k, l = x, y, z). \quad (1.4.80)$$

The rotation matrices in terms of the rotation angles are given by Eqs. (1.4.54) and (1.4.56). The angles which determine the resultant rotation are related to the angles of the first and second rotations via Eqs. (1.4.66) and (1.4.76).

If the angles which determine the first and resultant rotations, $S \rightarrow S'$ and $S \rightarrow S''$, are defined with respect to the initial system $S\{x, y, z\}$ but the angles of the second rotation $S' \rightarrow S''$ are defined with respect to the intermediate system $S'\{x', y', z'\}$ (i.e., the operator of the resultant rotation is given by Eq. (1.4.73)), then

$$a = a(1)a(2), \quad (1.4.81)$$

or, equivalently,

$$a_{ik} = \sum_l a_{il}(1) a_{lk}(2), \quad (i, k, l = x, y, z). \quad (1.4.82)$$

In this case the relationships between the angles of the resultant rotation and the angles of the first and second rotations may be derived from (1.4.66) and (1.4.76) by an interchange of indices $1 \rightleftharpoons 2$.

Chapter 2

Angular momentum operators

2.1 Total angular momentum operator

2.1.1 Definition

In quantum mechanics the total angular momentum operator $\hat{\mathbf{J}}$ is defined as an operator which generates transformations of wave functions (state vectors) and quantum operators under infinitesimal rotations of the coordinate system (see Eqs. (2.1.1) and (2.1.2)).

A transformation of an arbitrary wave function Ψ under rotation of the coordinate system through an infinitesimal angle $\delta\omega$ about an axis n may be written as

$$\Psi \rightarrow \Psi' = (1 - i\delta\omega \mathbf{n} \cdot \hat{\mathbf{J}})\Psi. \quad (2.1.1)$$

where $\hat{\mathbf{J}}$ is the total angular momentum operator.

A transformation of an arbitrary quantum operator $\hat{O}$ under an infinitesimal rotation has the form

$$\hat{O} \rightarrow \hat{O}' = \hat{O} - i\delta\omega \mathbf{n} \cdot [\hat{\mathbf{J}}, \hat{O}]. \quad (2.1.2)$$

The finite rotation operator can also be written in terms of the total angular momentum operator (see Eqs. 1.4.29, 1.4.30, 1.4.32). The total angular momentum operator is Hermitian,

$$\hat{\mathbf{J}}^+ = \hat{\mathbf{J}}. \quad (2.1.3)$$

This property for cartesian and spherical components of the operator $\hat{J}$ can be expressed as

$$\left(\hat{J}_i\right)^+ = \hat{J}_i, \quad (i = x, y, z); \quad \left(\hat{J}_\mu\right)^+ = \hat{J}^\mu = (-1)^\mu \hat{J}_{-\mu}, \quad (\mu = \pm 1, 0). \quad (2.1.4)$$

The eigenfunctions of the operators $\hat{\mathbf{J}}^2$ and $\hat{J}_z$ represent well-known tensor spherical harmonics (see Chap. 7).

2.1.2 Commutation Relations

Using the definition of the total angular momentum operator (2.1.1) and equations for the rotation addition (see Sec. a), one may obtain the commutation rules for the operator $\hat{\mathbf{J}}$. These rules may be written as

$$[(\mathbf{a} \cdot \hat{\mathbf{J}}), (\mathbf{b} \cdot \hat{\mathbf{J}})] = i \cdot [\mathbf{a} \times \mathbf{b}] \cdot \hat{\mathbf{J}}, \quad (2.1.5)$$

$\mathbf{a}$ and $\mathbf{b}$ being arbitrary constant vectors. The commutation relations for the total angular momentum operator may also be written symbolically as

$$[\hat{\mathbf{J}} \times \hat{\mathbf{J}}] = i\hat{\mathbf{J}}. \quad (2.1.6)$$

Cartesian components of $\hat{\mathbf{J}}$ satisfy the commutation relations

$$[\hat{J}_i, \hat{J}_k] = i\varepsilon_{ikl}\hat{J}_l, \quad [\hat{\mathbf{J}}^2, \hat{J}_i] = 0 \quad (i, k, l = x, y, z). \quad (2.1.7)$$

Equations (2.1.7) can be derived from (2.1.5) by substituting $\mathbf{a} = \mathbf{e}_i$, $\mathbf{b} = \mathbf{e}_k$ ($i, k = x, y, z$) where $\mathbf{e}_i$ and $\mathbf{e}_k$ are the cartesian basis vectors. In more detailed form Eq. (2.1.7) reads

$$\begin{aligned} [\hat{J}_x, \hat{J}_x] &= [\hat{J}_y, \hat{J}_y] = [\hat{J}_z, \hat{J}_z] = 0, \\ [\hat{J}_x, \hat{J}_y] &= -[\hat{J}_y, \hat{J}_x] = i\hat{J}_z, \quad [\hat{J}_x, \hat{J}_z] = -[\hat{J}_z, \hat{J}_x] = -i\hat{J}_y, \\ [\hat{J}_y, \hat{J}_z] &= -[\hat{J}_z, \hat{J}_y] = i\hat{J}_x, \\ [\hat{\mathbf{J}}^2, \hat{J}_x] &= [\hat{\mathbf{J}}^2, \hat{J}_y] = [\hat{\mathbf{J}}^2, \hat{J}_z] = 0. \end{aligned} \quad (2.1.8)$$

The square of the total angular momentum operator $\hat{\mathbf{J}}^2$ may be expressed in terms of cartesian components $\hat{J}_i$ ($i = x, y, z$) as

$$\hat{\mathbf{J}}^2 = \sum_i \hat{J}_i^2 = \hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2. \quad (2.1.9)$$

Covariant spherical components of $\hat{\mathbf{J}}$ satisfy the following commutation relations

$$[\hat{J}_\mu, \hat{J}_\nu] = -\sqrt{2}C_{1\mu 1\nu}^1 \hat{J}_\lambda, \quad [\hat{J}^2, \hat{J}_\mu] = 0, \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (2.1.10)$$

These relations may be obtained from (2.1.5) by putting $\mathbf{a} = \mathbf{e}_\mu$, $\mathbf{b} = \mathbf{e}_\nu$ ($\mu, \nu = \pm 1, 0$), $\mathbf{e}_\mu$ and $\mathbf{e}_\nu$ being covariant spherical basis vectors. A more detailed form of (2.1.10) is

$$\begin{aligned} [\hat{J}_{+1}, \hat{J}_{+1}] &= [\hat{J}_0, \hat{J}_0] = [\hat{J}_{-1}, \hat{J}_{-1}] = 0, \\ [\hat{J}_{+1}, \hat{J}_0] &= -[\hat{J}_0, \hat{J}_{+1}] = -\hat{J}_{+1}, \quad [\hat{J}_{+1}, \hat{J}_{-1}] = -[\hat{J}_{-1}, \hat{J}_{+1}] = -\hat{J}_0, \\ [\hat{J}_0, \hat{J}_{-1}] &= -[\hat{J}_{-1}, \hat{J}_0] = -\hat{J}_{-1}, \\ [\hat{J}^2, \hat{J}_{+1}] &= [\hat{J}^2, \hat{J}_0] = [\hat{J}^2, \hat{J}_{-1}] = 0. \end{aligned} \quad (2.1.11)$$

For contravariant spherical components of $\hat{\mathbf{J}}$ the commutation relations may be written as

$$[\hat{J}^\mu, \hat{J}^\nu] = \sqrt{2}C_{1\mu 1\nu}^1 \hat{J}^\lambda, \quad [\hat{\mathbf{J}}^2, \hat{J}^\mu] = 0, \quad (\mu, \nu, \lambda = \pm 1, 0) \quad (2.1.12)$$

or, in a more detailed form

$$\begin{aligned} [\hat{J}^{+1}, \hat{J}^{+1}] &= [\hat{J}^0, \hat{J}^0] = [\hat{J}^{-1}, \hat{J}^{-1}] = 0, \\ [\hat{J}^{+1}, \hat{J}^0] &= -[\hat{J}^0, \hat{J}^{+1}] = \hat{J}^{+1}, \quad [\hat{J}^{+1}, \hat{J}^{-1}] = -[\hat{J}^{-1}, \hat{J}^{+1}] = \hat{J}^0, \\ [\hat{J}^0, \hat{J}^{-1}] &= -[\hat{J}^{-1}, \hat{J}^0] = \hat{J}^{-1}, \\ [\hat{J}^2, \hat{J}^{+1}] &= [\hat{J}^2, \hat{J}^0] = [\hat{J}^2, \hat{J}^{-1}] = 0. \end{aligned} \quad (2.1.13)$$

The operator $\hat{\mathbf{J}}^2$ is expressed in terms of spherical components $\hat{J}_\mu$ ($\mu = \pm 1, 0$) as

$$\hat{\mathbf{J}}^2 = \sum_\mu (-1)^\mu \hat{J}_{-\mu} \hat{J}_\mu = -\hat{J}_{+1} \hat{J}_{-1} + \hat{J}_0 \hat{J}_0 - \hat{J}_{-1} \hat{J}_{+1} = \hat{J}_0^2 - \hat{J}_0 - 2\hat{J}_{+1} \hat{J}_{-1} = \hat{J}_0^2 + \hat{J}_0 - 2\hat{J}_{-1} \hat{J}_{+1}. \quad (2.1.14)$$

2.1.3 Coordinate Inversion. Time Reversal

The total angular momentum operator $\hat{\mathbf{J}}$ is an axial vector, i.e., it is invariant with respect to coordinate inversion ($\mathbf{r} \rightarrow -\mathbf{r}$)

$$\begin{aligned} \hat{P}_r \hat{J}_i \hat{P}_r^{-1} &= \hat{J}_i, \quad (i = x, y, z), \\ \hat{P}_r \hat{J}_\mu \hat{P}_r^{-1} &= \hat{J}_\mu, \quad (\mu = \pm 1, 0). \end{aligned} \quad (2.1.15)$$

Under time reversal ($t \rightarrow -t$) $\hat{\mathbf{J}}$ changes its sign

$$\begin{aligned} \hat{P}_t \hat{J}_i \hat{P}_t^{-1} &= -\hat{J}_i, \quad (i = x, y, z) \\ \hat{P}_t \hat{J}_\mu \hat{P}_t^{-1} &= -\hat{J}_\mu, \quad (\mu = \pm 1, 0) \end{aligned} \quad (2.1.16)$$

In Eqs. (2.1.15) and (2.1.16) $\hat{P}_r$ and $\hat{P}_t$ represent the operators of coordinate inversion and time reversal, respectively.

2.1.4 Total Angular Momentum of a System. Orbital and Spin Angular Momenta

The total angular momentum of some system consisting of N subsystems with angular momenta $\hat{\mathbf{j}}(1), \hat{\mathbf{j}}(2), \dots, \hat{\mathbf{j}}(N)$ is the vector sum

$$\hat{\mathbf{J}} = \sum_{n=1}^N \hat{\mathbf{j}}(n). \quad (2.1.17)$$

For such a system the following commutation relations hold

$$\begin{aligned} [\hat{j}_i(n), \hat{j}_k(n')] &= i\delta_{nn'}\varepsilon_{ikl}\hat{j}_l(n), \quad (i, k, l = x, y, z), \\ [\hat{J}_i, \hat{j}_k(n)] &= i\varepsilon_{ikl}\hat{j}_l(n), \end{aligned} \quad (2.1.18)$$

$$\begin{aligned} [\hat{j}_\mu(n), \hat{j}_\nu(n')] &= -\sqrt{2}\delta_{nn'}C_{1\mu 1\nu}^1\lambda\hat{j}_\lambda(n), \quad (\mu, \nu, \lambda = \pm 1, 0). \\ [\hat{J}_\mu, \hat{j}_\nu(n)] &= -\sqrt{2}C_{1\mu 1\nu}^1\lambda\hat{j}_\lambda(n), \end{aligned} \quad (2.1.19)$$

In particular, Eq. (2.1.17) determines the total angular momentum operator as the sum of the orbital angular momentum operator $\hat{\mathbf{L}}$ and the spin angular momentum operator $\hat{\mathbf{S}}$.

$$\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}. \quad (2.1.20)$$

The properties of the operators $\hat{\mathbf{L}}$ and $\hat{\mathbf{S}}$ are considered below (Secs. 2.2 and 2.3). The commutation relations for $\hat{\mathbf{L}}$ and $\hat{\mathbf{J}}$ are

$$\begin{aligned} [\hat{J}_i, \hat{L}_k] &= i\varepsilon_{ikl}\hat{L}_l, \quad (i, k, l = x, y, z), \\ [\hat{J}_\mu, \hat{L}_\nu] &= -\sqrt{2}C_{1\mu 1\nu}^1\lambda\hat{L}_\lambda, \quad (\mu, \nu, \lambda = \pm 1, 0). \end{aligned} \quad (2.1.21)$$

or, in more detail

$$\begin{aligned} [\hat{J}_x, \hat{L}_x] &= [\hat{J}_y, \hat{L}_y] = [\hat{J}_z, \hat{L}_z] = 0, \\ [\hat{J}_x, \hat{L}_y] &= -[\hat{J}_y, \hat{L}_x] = i\hat{L}_z, \quad [\hat{J}_x, \hat{L}_z] = -[\hat{J}_z, \hat{L}_x] = -i\hat{L}_y, \\ [\hat{J}_y, \hat{L}_z] &= -[\hat{J}_z, \hat{L}_y] = i\hat{L}_x \end{aligned} \quad (2.1.22)$$

$$\begin{aligned} [\hat{J}_{+1}, \hat{L}_{+1}] &= [\hat{J}_0, \hat{L}_0] = [\hat{J}_{-1}, \hat{L}_{-1}] = 0, \\ [\hat{J}_{+1}, \hat{L}_0] &= -[\hat{J}_0, \hat{L}_{+1}] = -\hat{L}_{+1}, \quad [\hat{J}_{+1}, \hat{L}_{-1}] = -[\hat{J}_{-1}, \hat{L}_{+1}] = -\hat{L}_0, \\ [\hat{J}_0, \hat{L}_{-1}] &= -[\hat{J}_{-1}, \hat{L}_0] = -\hat{L}_{-1}. \end{aligned} \quad (2.1.23)$$

Appropriate relations for the spin operator $\hat{\mathbf{S}}$ and the total angular momentum operator $\hat{\mathbf{J}}$ may be obtained from Eqs. (2.1.21)-(2.1.23) by replacing $\hat{\mathbf{L}} \rightarrow \hat{\mathbf{S}}$. The orbital angular momentum operator $\hat{\mathbf{L}}$ commutes with the spin operator $\hat{\mathbf{S}}$

$$\begin{aligned} [\hat{L}_i, \hat{S}_k] &= 0, \quad (i, k = x, y, z) \\ [\hat{L}_\mu, \hat{S}_\nu] &= 0, \quad (\mu, \nu = \pm 1, 0) \end{aligned} \quad (2.1.24)$$

2.2 Orbital angular momentum operator

2.2.1 Definition

In classical mechanics the angular momentum $\mathbf{L}$ of a particle is defined as

$$\mathbf{L} = [\mathbf{r} \times \mathbf{p}], \quad (2.2.1)$$

where $\mathbf{r}$ is the position vector of the particle and $\mathbf{p}$ is its linear momentum.

In the quantum mechanics the orbital angular momentum operator $\hat{\mathbf{L}}$ of a particle is obtained from Eq. (2.2.1) by replacing $\mathbf{r} \rightarrow \hat{\mathbf{r}}$ and $\mathbf{p} \rightarrow \hat{\mathbf{p}}$, where $\hat{\mathbf{r}}$ and $\hat{\mathbf{p}}$ are the position and momentum operators, respectively.

In the coordinate representation we have $\hat{\mathbf{r}} = \mathbf{r}$, $\hat{\mathbf{p}} = -i\nabla$,

$$\hat{\mathbf{L}} = -i[\mathbf{r} \times \nabla]. \quad (2.2.2)$$

In the momentum representation we have $\hat{\mathbf{r}} = i\nabla_{\mathbf{p}}$, $\hat{\mathbf{p}} = \mathbf{p}$,

$$\hat{\mathbf{L}} = -i[\mathbf{p} \times \nabla_{\mathbf{p}}]. \quad (2.2.3)$$

The formulas below are valid for both representations. The explicit form of $\hat{\mathbf{L}}$ (see Sec. 2.2.3) in the momentum representation is obtained by substituting $\mathbf{r} \rightarrow \mathbf{p}$, $\nabla \equiv \partial/\partial \mathbf{r} \rightarrow \nabla_{\mathbf{p}} \equiv \partial/\partial \mathbf{p}$.

The operator $\hat{\mathbf{L}}$ is Hermitian and purely imaginary.

$$\hat{\mathbf{L}}^+ = \hat{\mathbf{L}}, \quad \hat{\mathbf{L}}^* = -\hat{\mathbf{L}}. \quad (2.2.4)$$

For cartesian components of $\hat{\mathbf{L}}$, Eqs. (2.2.4) become

$$\hat{L}_i^+ = \hat{L}_i, \quad \hat{L}_i^* = -\hat{L}_i, \quad (i = x, y, z). \quad (2.2.5)$$

and for spherical components Eqs. (2.2.4) yield

$$\begin{aligned} (\hat{L}_\mu)^+ &= (-1)^\mu (\hat{L}^+)_{-\mu} = (-1)^\mu \hat{L}_{-\mu}, \\ (\hat{L}_\mu)^* &= (-1)^\mu (\hat{L}^*)_{-\mu} = (-1)^{\mu+1} \hat{L}_{-\mu}, \quad (\mu = \pm 1, 0). \end{aligned} \quad (2.2.6)$$

The orbital angular momentum operator $\hat{\mathbf{L}}$ generates transformations of scalar (spinless) wave functions under rotations of the coordinate system. A rotation through an infinitesimal angle $\delta\omega$ about the $\mathbf{n}$ -axis transforms the position vector $\mathbf{r}$ into $\mathbf{r} + \delta\mathbf{r}$, with $\delta\mathbf{r} = -\delta\omega[\mathbf{n} \times \mathbf{r}]$. The corresponding transformation of the scalar wave functions reads

$$\Psi(\mathbf{r}) \rightarrow \Psi(\mathbf{r} + \delta\mathbf{r}) = (1 + \delta\mathbf{r} \cdot \nabla)\Psi(\mathbf{r}) = (1 - i\delta\omega \mathbf{n} \cdot \hat{\mathbf{L}})\Psi(\mathbf{r}). \quad (2.2.7)$$

In the case of a finite rotation through an angle ω about the axis $\mathbf{n}$, scalar functions are transformed by the operator $e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{L}}}$ which is the particular form of the rotation operator 1.4.32. The eigenfunctions of the operators $\hat{\mathbf{L}}^2$ and $\hat{L}_z$ represent spherical harmonics $Y_{lm}(\vartheta, \varphi)$ (Chap. 5) which depend on the polar angles ϑ, φ .

2.2.2 Commutation Relations

The orbital angular momentum operator $\hat{\mathbf{L}}$ satisfies the same commutation relations as the total angular momentum operator $\hat{\mathbf{J}}$. Appropriate expressions for the commutators are readily obtained from Eqs. (2.1.5)-(2.1.14) by substituting $\hat{\mathbf{J}} \rightarrow \hat{\mathbf{L}}$.

The commutation rules for the operator $\hat{\mathbf{L}}$ with the position operator $\hat{\mathbf{r}}$ and the momentum operator $\hat{\mathbf{p}}$ are the following.

For cartesian components

$$\begin{aligned} [\hat{L}_i, \hat{x}_k] &= i\varepsilon_{ikl}x_l, \\ [\hat{L}_i, \hat{p}_k] &= i\varepsilon_{ikl}p_l, \end{aligned} \quad (i, k, l = x, y, z). \quad (2.2.8)$$

Note also that $\hat{\mathbf{r}}$ and $\hat{\mathbf{p}}$ obey the commutation relations

$$[\hat{x}_i, \hat{x}_k] = 0, \quad [\hat{p}_i, \hat{p}_k] = 0, \quad [\hat{x}_i, \hat{p}_k] = i\delta_{ik}, \quad (i, k = x, y, z). \quad (2.2.9)$$

For spherical components

$$\begin{aligned} [\hat{L}_\mu, \hat{x}_\nu] &= -\sqrt{2}C_{1\mu 1\nu}^1 \hat{x}_\lambda, \\ [\hat{L}_\mu, \hat{p}_\nu] &= -\sqrt{2}C_{1\mu 1\nu}^1 \hat{p}_\lambda, \end{aligned} \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (2.2.10)$$

The commutation relations for the spherical components of $\hat{\mathbf{r}}$ and $\hat{\mathbf{p}}$ are given by

$$[\hat{x}_\mu, \hat{x}_\nu] = 0, \quad [\hat{p}_\mu, \hat{p}_\nu] = 0, \quad [\hat{x}_\mu, \hat{p}_\nu] = i\delta_{\mu-\nu}(-1)^\mu, \quad (\mu, \nu = \pm 1, 0). \quad (2.2.11)$$

Equation (2.2.8) and (2.2.10) show that the operators $\hat{\mathbf{r}}$ and $\hat{\mathbf{p}}$ are vectors.

The commutation relations for square of angular momentum operator $\hat{\mathbf{L}}^2$ are

$$\begin{aligned} [\hat{\mathbf{L}}^2, \hat{\mathbf{r}}] &= i[\hat{\mathbf{r}} \times \hat{\mathbf{L}}] - i[\hat{\mathbf{L}} \times \hat{\mathbf{r}}], \\ [\hat{\mathbf{L}}^2, \hat{\mathbf{p}}] &= i[\hat{\mathbf{p}} \times \hat{\mathbf{L}}] - i[\hat{\mathbf{L}} \times \hat{\mathbf{p}}]. \end{aligned} \quad (2.2.12)$$

The operators $\hat{\mathbf{r}}^2$ and $\hat{\mathbf{p}}^2$ commute with $\hat{\mathbf{L}}$:

$$[\hat{\mathbf{L}}, \hat{\mathbf{r}}^2] = 0, \quad [\hat{\mathbf{L}}, \hat{\mathbf{p}}^2] = 0$$

The commutation rules of $\hat{\mathbf{L}}$ with the total angular momentum operator $\hat{\mathbf{J}}$ and the spin operator $\hat{\mathbf{S}}$ have been considered above (Eqs. (2.1.21)-(2.1.24)).

2.2.3 Explicit Form

The cartesian components of the operator $\hat{\mathbf{L}}$ are given by

$$\hat{L}_x = -i \left(y \frac{\partial}{\partial z} - z \frac{\partial}{\partial y} \right), \quad \hat{L}_y = -i \left(z \frac{\partial}{\partial x} - x \frac{\partial}{\partial z} \right), \quad \hat{L}_z = -i \left(x \frac{\partial}{\partial y} - y \frac{\partial}{\partial x} \right). \quad (2.2.13)$$

or in a more compact form

$$\hat{L}_i = -i \sum_{k,l} \varepsilon_{ikl} x_k \frac{\partial}{\partial x_l}, \quad (i, k, l = x, y, z). \quad (2.2.14)$$

The cartesian components of $\hat{\mathbf{L}}$ are expressed in terms of polar angles ϑ, φ as

$$\begin{aligned} \hat{L}_x &= i \left(\sin \varphi \frac{\partial}{\partial \vartheta} + \cot \vartheta \cos \varphi \frac{\partial}{\partial \varphi} \right) \\ \hat{L}_y &= i \left(-\cos \varphi \frac{\partial}{\partial \vartheta} + \cot \vartheta \sin \varphi \frac{\partial}{\partial \varphi} \right) \\ \hat{L}_z &= -i \frac{\partial}{\partial \varphi} \end{aligned} \quad (2.2.15)$$

The spherical components of $\hat{\mathbf{L}}$ are written as

$$\begin{aligned} \hat{L}_{+1} &= x_0 \nabla_{+1} - x_{+1} \nabla_0, \\ \hat{L}_0 &= x_{-1} \nabla_{+1} - x_{+1} \nabla_{-1}, \\ \hat{L}_{-1} &= x_{-1} \nabla_0 - x_0 \nabla_{-1}. \end{aligned} \quad (2.2.16)$$

or, in a more compact form

$$\hat{L}_\mu = -\sqrt{2} \sum_{\nu, \lambda} C_{1\nu 1\lambda}^{1\mu} x_\nu \nabla_\lambda, \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (2.2.17)$$

The spherical components of $\hat{\mathbf{L}}$ may also be expressed in terms of polar angles ϑ, φ as

$$\begin{aligned} \hat{L}_{+1} &= -\frac{1}{\sqrt{2}} e^{i\varphi} \left\{ \frac{\partial}{\partial \vartheta} + i \cot \vartheta \frac{\partial}{\partial \varphi} \right\}, \\ \hat{L}_0 &= -i \frac{\partial}{\partial \varphi}, \\ \hat{L}_{-1} &= -\frac{1}{\sqrt{2}} e^{-i\varphi} \left\{ \frac{\partial}{\partial \vartheta} - i \cot \vartheta \frac{\partial}{\partial \varphi} \right\}. \end{aligned} \quad (2.2.18)$$

For polar components of $\hat{\mathbf{L}}$ we have

$$\hat{L}_r = 0, \quad \hat{L}_\vartheta = \frac{i}{\sin \vartheta} \frac{\partial}{\partial \varphi}, \quad \hat{L}_\varphi = -i \frac{\partial}{\partial \vartheta}. \quad (2.2.19)$$

The square of the orbital angular momentum operator $\hat{\mathbf{L}}^2$ is expressed in terms of its components as

$$\begin{aligned} \hat{\mathbf{L}}^2 &= \sum_i \hat{L}_i^2 = \hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2, \\ \hat{\mathbf{L}}^2 &= \sum_\mu (-1)^\mu \hat{L}_\mu \hat{L}_{-\mu} = -\hat{L}_{+1} \hat{L}_{-1} + \hat{L}_0 \hat{L}_0 - \hat{L}_{-1} \hat{L}_{+1}. \end{aligned} \quad (2.2.20)$$

We may also write $\hat{\mathbf{L}}^2$ in the form

$$\hat{\mathbf{L}}^2 = - \left[\frac{1}{\sin \vartheta} \cdot \frac{\partial}{\partial \vartheta} \left(\sin \vartheta \frac{\partial}{\partial \vartheta} \right) + \frac{1}{\sin^2 \vartheta} \cdot \frac{\partial^2}{\partial \varphi^2} \right]. \quad (2.2.21)$$

This operator differs only in sign from the angular part of the Laplacian operator (Eq. (1.3.15))

$$\hat{\mathbf{L}}^2 = -\Delta_\Omega. \quad (2.2.22)$$

The operator $\hat{\mathbf{L}}^2$ is Hermitian and real

$$(\hat{\mathbf{L}}^2)^+ = (\hat{\mathbf{L}}^2)^* = \hat{\mathbf{L}}^2. \quad (2.2.23)$$

The representation of the orbital angular momentum $\hat{\mathbf{L}}$ in the form of a differential operator is only one of possible representations. Alternatively, $\hat{\mathbf{L}}$ may be represented by a set of three matrices (because it has three components). These matrix elements of $\hat{\mathbf{L}}$ will be given in Chap. 13.

2.3 Spin angular momentum operator

2.3.1 Definition

The spin angular momentum operator or briefly the spin operator $\hat{\mathbf{S}}$ is usually represented by a set of three (since the vector $\hat{\mathbf{S}}$ has three components) square $(2S+1) \times (2S+1)$ matrices, S being the particle spin. These matrices act on the spin functions (see Chap. 6) and satisfy the same commutation relations as the components of the total angular momentum operator given in Sec.2.1. The spin operator $\hat{\mathbf{S}}$ is Hermitian:

$$\hat{\mathbf{S}}^+ = \hat{\mathbf{S}} \quad (2.3.1)$$

For the cartesian components of $\hat{\mathbf{S}}$ the Hermitian property (2.3.1) has the form

$$(\hat{S}_i)^+ = \hat{S}_i, \quad (i = x, y, z) \quad (2.3.2)$$

and for spherical components

$$(\hat{S}_\mu)^+ = (-1)^\mu \hat{S}_{-\mu}, \quad (\mu = \pm 1, 0). \quad (2.3.3)$$

The behaviour of $\hat{\mathbf{S}}$ under complex conjugation depends on the representation of $\hat{\mathbf{S}}$ (see, e.g., Sec. 2.6.2). The eigenfunctions of the operators $\hat{\mathbf{S}}^2$ and $\hat{S}_z$ are spin functions χ_{Sm} (see Chap. 6) which depend on the spin variable σ . These functions have $(2S+1)$ components and describe polarization states of a particle.

2.3.2 Commutation Relations

The commutation relations for the spin operator $\hat{\mathbf{S}}$ are given by Eqs. (2.1.5)-(2.1.14) in which one must replace $\hat{\mathbf{J}}$ by $\hat{\mathbf{S}}$. The commutation rules for $\hat{\mathbf{S}}$ and the orbital angular momentum operator $\hat{\mathbf{L}}$ have already been considered in Sec. 2.1.4. The spin operator $\hat{\mathbf{S}}$ commutes with the position operator $\hat{\mathbf{r}}$ and the momentum operator $\hat{\mathbf{p}}$,

$$[\hat{S}_i, \hat{x}_k] = 0, \quad [\hat{S}_i, \hat{p}_k] = 0, \quad (i, k = x, y, z), \quad (2.3.4)$$

$$[\hat{S}_\mu, \hat{x}_\nu] = 0, \quad [\hat{S}_\mu, \hat{p}_\nu] = 0, \quad (\mu, \nu = \pm 1, 0). \quad (2.3.5)$$

2.3.3 Explicit Form

The spherical components of the spin operator $\hat{S}$ may be expressed in terms of the basis spin functions χ_{Sm} (Chap. 6) as

$$\hat{S}_\mu = \sqrt{S(S+1)} \sum_{m, m'} C_{S m' 1 \mu}^{S m'} \chi_{S m'} \chi_{S m}^+, \quad (\mu = \pm 1, 0). \quad (2.3.6)$$

Cartesian components of $\hat{\mathbf{S}}$ may be obtained from (2.3.6), using the relations

$$\hat{S}_x = \frac{1}{\sqrt{2}} (\hat{S}_{-1} - \hat{S}_{+1}), \quad \hat{S}_y = \frac{i}{\sqrt{2}} (\hat{S}_{-1} + \hat{S}_{+1}), \quad \hat{S}_z = \hat{S}_0. \quad (2.3.7)$$

Equations (2.3.6) and (2.3.7) are independent of representation used for spin functions, while an explicit form of the spin matrices depends on this representation. For an arbitrary spin S the simplest form of the spin matrices is that in the spherical basis representation. The basis spin functions χ_{Sm} in this representation are given by

$$\chi_{Sm}(\sigma) = \delta_{m\sigma}, \quad (m, \sigma = -S, -S+1, \dots, S-1, S), \quad (2.3.8)$$

and the matrix elements of the spin matrices are

$$\left(\hat{S}_\mu\right)_{\sigma'\sigma} = \sqrt{S(S+1)}C_{S\sigma 1\mu}^{S\sigma'}, \quad (\sigma, \sigma' = -S, -S+1, \dots, S-1, S) \quad (2.3.9)$$

The elements are arranged in the matrices as follows

$$A = \begin{pmatrix} A_{SS} & \dots & A_{S-S} \\ \vdots & \ddots & \vdots \\ A_{-SS} & \dots & A_{-S-S} \end{pmatrix}. \quad (2.3.10)$$

Explicit forms of the spin matrices for spin values $\frac{1}{2}$ and 1 are given below in Secs. 2.5 and 2.6.

2.3.4 Traces of Products of Spin Matrices

The traces of products of cartesian components of the spin matrices can be evaluated for an arbitrary spin S by use of the following formulas (where i, k, l , etc. take the values x, y, z)

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_i \right\} &= 0 \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \right\} &= \frac{S(S+1)(2S+1)}{3} \delta_{ik}, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \right\} &= i \frac{S(S+1)(2S+1)}{6} \varepsilon_{ikl}, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \hat{S}_j \right\} &= \frac{S(S+1)(2S+1)}{15} \left\{ \left[S(S+1) + \frac{1}{2} \right] (\delta_{ik} \delta_{lj} + \delta_{ij} \delta_{kl}) + [S(S+1) - 2] \delta_{il} \delta_{kj} \right\}. \end{aligned} \quad (2.3.11)$$

For spherical components of the spin operator $\hat{S}$ the following relations hold (μ, ν, λ , etc. take the values $\pm 1, 0$)

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_\mu \right\} &= 0, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \right\} &= \frac{S(S+1)(2S+1)}{3} (-1)^\mu \delta_{\mu-\nu}, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \right\} &= -\frac{S(S+1)(2S+1)}{\sqrt{6}} \begin{pmatrix} 1 & 1 & 1 \\ \mu & \nu & \lambda \end{pmatrix} = \frac{S(S+1)(2S+1)}{3\sqrt{2}} (-1)^{1+\lambda} C_{1\mu 1\nu}^{1-\lambda}, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \hat{S}_\rho \right\} &= \frac{S(S+1)(2S+1)}{15} \left\{ \left[S(S+1) + \frac{1}{2} \right] (-1)^{\mu+\lambda} (\delta_{\mu-\nu} \delta_{\lambda-\rho} + \delta_{\mu-\rho} \delta_{\nu-\lambda}) \right. \\ &\quad \left. + [S(S+1) - 2] (-1)^{\mu+\nu} \delta_{\mu-\lambda} \delta_{\nu-\rho} \right\}. \end{aligned} \quad (2.3.12)$$

Some analytic expressions for traces of products of many spin matrices in the spherical basis are given in Refs. [131, 132, 133].

2.4 Polarization operators

2.4.1 Definition

To describe a polarization (i.e., spin) state of a particle the so-called polarization operators are widely used. The polarization operators $\hat{T}_{LM}(S)$ ($M = -L, -L+1, \dots, L-1, L$ and $L = 0, 1, \dots, 2S$ with L and M integer) are $(2S+1) \times (2S+1)$ matrices which act on spin functions (see Chap. 6) and transform under rotations of the coordinate system according to the representation D^L . In other words, $\hat{T}_{LM}(S)$ are irreducible tensors of rank L . Such transformation properties of $\hat{T}_{LM}(S)$ with respect to rotations of the coordinate system imply the following commutation relations with spherical components of the spin operator $\hat{S}_\mu$ ($\mu = \pm 1, 0$)

$$\left[\hat{S}_\mu, \hat{T}_{LM}(S) \right] = \sqrt{L(L+1)} C_{L M 1 \mu}^{L M+\mu} \hat{T}_{L M+\mu}(S). \quad (2.4.1)$$

Let us normalize the polarization operators by the condition

$$\text{Tr} \left\{ \hat{T}_{LM}^+(S) \hat{T}_{L'M'}(S) \right\} = \delta_{LL'} \delta_{MM'}, \quad (2.4.2)$$

and choose the phase factors to satisfy the relations

$$\hat{T}_{LM}^+(S) = (-1)^M \hat{T}_{L-M}(S). \quad (2.4.3)$$

The conditions (2.4.1)-(2.4.3) completely determine the polarization operators. The full complement of polarization operators $\hat{T}_{LM}(S)$ with $-L \leq M \leq L, 0 \leq L \leq 2S$ constitutes a complete set of $\sum_{L=0}^{2S} (2L+1) = (2S+1)^2$ linearly independent square $(2S+1) \times (2S+1)$ matrices.

2.4.2 Explicit Form

The polarization operators $\hat{T}_{LM}(S)$ may be constructed from products of the spin matrices as

$$\hat{T}_{LM}(S) = N_L(S) (\hat{\mathbf{S}} \cdot \nabla)^L \{ r^L Y_{LM}(\vartheta, \varphi) \}, \quad (2.4.4)$$

where $\hat{\mathbf{S}}$ is the spin operator, and $N_L(S)$ is the normalization factor given by

$$N_L(S) = \frac{2^L}{L!} \left[\frac{4\pi(2S-L)!}{(2S+L+1)!} \right]^{\frac{1}{2}}. \quad (2.4.5)$$

Note that $\hat{T}_{LM}(S)$ are actually independent of r , although the vector r enters the right-hand side of Eq. (2.4.4). The operators $\hat{T}_{LM}(S)$ may be expressed in terms of the basis spin functions χ_{Sm} (Chap. 6) by

$$\hat{T}_{LM}(S) = \sqrt{\frac{2L+1}{2S+1}} \sum_{m,m'} C_{S m L M}^{S m'} \chi_{Sm'} \chi_{Sm}^+. \quad (2.4.6)$$

The inverse relation is

$$\chi_{Sm'} \chi_{Sm}^+ = \sum_L \sqrt{\frac{2L+1}{2S+1}} C_{S m L M}^{S m'} \hat{T}_{LM}(S). \quad (2.4.7)$$

An explicit form of $\hat{T}_{LM}(S)$ depends on the representation used for the spin functions. In particular, matrix elements of $\hat{T}_{LM}(S)$ in the spherical basis representation are given by

$$[\hat{T}_{LM}(S)]_{\sigma'\sigma} = \sqrt{\frac{2L+1}{2S+1}} C_{S \sigma' L M}^{S \sigma}, \quad (\sigma, \sigma' = -S, -S+1, \dots, S). \quad (2.4.8)$$

For $L=0$ the operators $\hat{T}_{LM}(S)$ are proportional to the unit $(2S+1) \times (2S+1)$ matrix

$$\hat{T}_{00}(S) = \frac{1}{\sqrt{2S+1}} \hat{I}. \quad (2.4.9)$$

When $L=1$ the operators $\hat{T}_{LM}(S)$ are proportional to the spherical components of the spin operator

$$\hat{T}_{1M}(S) = \frac{\sqrt{3}}{\sqrt{S(S+1)(2S+1)}} \hat{S}_M, \quad (M = \pm 1, 0). \quad (2.4.10)$$

The polarization operators for a spin value $S=1$ are considered in Sec. 2.6.

2.4.3 Properties of $\hat{T}_{LM}(S)$ under Transformations of the Coordinate System

(a) *Coordinate inversion* ($\mathbf{r} \rightarrow -\mathbf{r}$)

The polarization operators $\hat{T}_{LM}(S)$ are invariant under coordinate inversion

$$\hat{P}_r \hat{T}_{LM}(S) \hat{P}_r^{-1} = \hat{T}_{LM}(S).. \quad (2.4.11)$$

(b) *Rotation of coordinate system*

Under rotations specified by the Euler angles α, β, γ the polarization operators transform as

$$\hat{T}'_{LM'}(S) = \hat{D}(\alpha, \beta, \gamma) \hat{T}_{LM}(S) [\hat{D}(\alpha, \beta, \gamma)]^{-1} = \sum_M D_{MM'}^L(\alpha, \beta, \gamma) \hat{T}_{LM}(S) \quad (2.4.12)$$

, where $\hat{D}(\alpha, \beta, \gamma)$ is the rotation operator (see Sec. 1.4.5), and $D_{MM'}^L$ are the Wigner D -functions (Chap. 4).

2.4.4 Expansion Series of Polarization Operators

As has been mentioned above (Sec.2.4.1) the polarization operators $\hat{T}_{LM}(S)$ form a complete set of linearly independent matrices. An arbitrary square $(2S+1) \times (2S+1)$ matrix $\hat{A}$ (with S integer or half-integer) may be expanded in a series of the polarization operators $\hat{T}_{LM}(S)$, i.e., it may be written in the form

$$\hat{A} = \sum_{L=0}^{2S} \sum_{M=-L}^L A_{LM} \hat{T}_{LM}(S), \quad (2.4.13)$$

where the expansion coefficients A_{LM} are given by

$$A_{LM} = \text{Tr} \left\{ \hat{T}_{LM}^+(S) \hat{A} \right\} .. \quad (2.4.14)$$

If the matrix $\hat{A}$ is Hermitian, i.e., $\hat{A}^+ = \hat{A}$, then

$$A_{LM}^* = (-1)^M A_{L-M}. \quad (2.4.15)$$

Some examples of such expansions are given below.

(a) Clebsch-Gordan series for polarization operators

Products of two polarization operators $\hat{T}_{L_1 M_1}(S)$ and $\hat{T}_{L_2 M_2}(S)$ may be written in the form of a ClebschGordan series

$$\hat{T}_{L_1 M_1}(S) \hat{T}_{L_2 M_2}(S) = \sum_L (-1)^{2S+L} \sqrt{(2L_1+1)(2L_2+1)} \left\{ \begin{matrix} L_1 & L_2 & L \\ S & S & S \end{matrix} \right\} C_{L_1 M_1 L_2 M_2}^{L M} \hat{T}_{LM}(S). \quad (2.4.16)$$

(b) Expansion of rotation operator

If one is interested only in transformation properties of spin functions and spin operators under rotations of the coordinate system, one may replace the total angular momentum operator $\hat{\mathbf{J}}$ by the spin operator $\hat{\mathbf{S}}$ in the general expressions for the rotation operator (1.4.31) and (1.4.33). Such a modified operator will be labelled by a superscript S . If a rotation is defined by the Euler angles α, β, γ , the rotation operator $\hat{D}^S(\alpha, \beta, \gamma)$ may be expressed as follows

$$\hat{D}^S(\alpha, \beta, \gamma) \equiv e^{-i\alpha \hat{S}_z} e^{-i\beta \hat{S}_y} e^{-i\gamma \hat{S}_z} = \sum_{L, M, m, m'} \frac{2L+1}{2S+1} C_{S m' L M}^S D_{mm'}^S(\alpha, \beta, \gamma) \hat{T}_{LM}(S). \quad (2.4.17)$$

If a rotation is defined by the rotation axis $n(\Theta, \Phi)$ and the rotation angle ω , the expansion of the rotation operator $\hat{U}^S(\omega; \Theta, \Phi)$ is given by

$$\hat{U}^S(\omega; \Theta, \Phi) \equiv e^{-i\omega n \cdot \hat{\mathbf{S}}} = 2\sqrt{\pi} \sum_{L=0}^{2S} \frac{(-i)^L}{\sqrt{2S+1}} \chi_L^S(\omega) \sum_{M=-L}^L Y_{LM}^*(\Theta, \Phi) \hat{T}_{LM}(S), \quad (2.4.18)$$

where the functions $\chi_L^S(\omega)$ are the generalized characters (see Sec. 4.15).

2.4.5 Commutators and Anticommutators

The polarization operators satisfy the following commutation relations

$$\begin{aligned} & \left[\hat{T}_{L_1 M_1}(S), \hat{T}_{L_2 M_2}(S) \right] = \\ & \sqrt{(2L_1+1)(2L_2+1)} \sum_{L_3} (-1)^{2S+L_3} \left[1 - (-1)^{L_1+L_2+L_3} \right] \left\{ \begin{matrix} L_1 & L_2 & L_3 \\ S & S & S \end{matrix} \right\} C_{L_1 M_1 L_2 M_2}^{L_3 M_3} \hat{T}_{L_3 M_3}(S), \end{aligned} \quad (2.4.19)$$

$$\begin{aligned} & \left\{ \hat{T}_{L_1 M_1}(S), \hat{T}_{L_2 M_2}(S) \right\} = \\ & \sqrt{(2L_1+1)(2L_2+1)} \sum_{L_3} (-1)^{2S+L_3} \left[1 + (-1)^{L_1+L_2+L_3} \right] \left\{ \begin{matrix} L_1 & L_2 & L_3 \\ S & S & S \end{matrix} \right\} C_{L_1 M_1 L_2 M_2}^{L_3 M_3} \hat{T}_{L_3 M_3}(S). \end{aligned} \quad (2.4.20)$$

Equation (2.4.1) is a special case of (2.4.19) for $L_1 = 1$.

2.4.6 Traces of Products of Polarization Operators

$$\text{Tr} \left\{ \hat{T}_{LM}(S) \right\} = \sqrt{2S+1} \delta_{LO} \delta_{MO}, \quad (2.4.21)$$

$$\text{Tr} \left\{ \hat{T}_{L_1 M_1}(S) \hat{T}_{L_2 M_2}(S) \right\} = (-1)^{M_1} \delta_{L_1 L_2} \delta_{M_1 - M_2}, \quad (2.4.22)$$

$$\text{Tr} \left\{ \hat{T}_{L_1 M_1}(S) \hat{T}_{L_2 M_2}(S) \hat{T}_{L_3 M_3}(S) \right\} = (-1)^{2S+L_3+M_3} \sqrt{(2L_1+1)(2L_2+1)} C_{L_1 M_1 L_2 M_2}^{L_3 -M_3} \left\{ \begin{matrix} L_1 & L_2 & L_3 \\ S & S & S \end{matrix} \right\}. \quad (2.4.23)$$

In general

$$\begin{aligned} \text{Tr} \left\{ \hat{T}_{L_1 M_1}(S) \hat{T}_{L_2 M_2}(S) \dots \hat{T}_{L_n M_n}(S) \right\} &= \frac{[(2L_1+1)(2L_2+1) \dots (2L_n+1)]^{\frac{1}{2}}}{(2S+1)^{n/2}}, \\ &\times \delta_{M_1+M_2+M_3+\dots+M_n,0} \sum_m C_{L_1 M_1 S m}^{S m+\mu_1} C_{L_2 M_2 S m+\mu_1}^{S m+\mu_2} \dots C_{L_n M_n S m+\mu_{n-1}}^{S m}, \end{aligned} \quad (2.4.24)$$

where

$$\mu_k = \sum_{i=1}^k M_i$$

2.5 Spin matrices for $S = 1/2$

2.5.1 Explicit Form

For the particular case $S = \frac{1}{2}$, the spin operator $\hat{\mathbf{S}}$ is represented by a set of three square 2×2 matrices. Cartesian components of $\hat{\mathbf{S}}$ in the spherical basis representation are given by

$$\hat{S}_x = \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \hat{S}_y = \frac{1}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \hat{S}_z = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.5.1)$$

whereas spherical components are written as

$$\hat{S}_{+1} = -\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \hat{S}_0 = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \hat{S}_{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad (2.5.2)$$

In addition to the spin matrices, the unit 2×2 matrix $\hat{I}$ is usually introduced:

$$\hat{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (2.5.3)$$

Note that the Pauli matrices $\hat{\sigma}$ are widely used instead of the spin operator $\hat{\mathbf{S}}$. The Pauli matrices are proportional to the spin operator

$$\hat{\mathbf{S}} = \frac{1}{2} \hat{\boldsymbol{\sigma}}. \quad (2.5.4)$$

Equations (2.5.1) and (2.5.2) yield the following properties of the spin matrices

$$\hat{S}_i^+ = \hat{S}_i, \quad (i = x, y, z), \quad (2.5.5)$$

$$\hat{S}_\mu^+ = (-1)^\mu \hat{S}_{-\mu}, \quad (\mu = \pm 1, 0), \quad (2.5.6)$$

$$\hat{S}_x^* = \hat{S}_x, \quad \hat{S}_y^* = -\hat{S}_y, \quad \hat{S}_z^* = \hat{S}_z, \quad (2.5.7)$$

$$\hat{S}_\mu^* = \hat{S}_\mu, \quad (\mu = \pm 1, 0). \quad (2.5.8)$$

2.5.2 Commutators and Anticommutators

The cartesian components of the spin operator satisfy the following relations

$$[\hat{S}_i, \hat{S}_k] = i\epsilon_{ikl} \hat{S}_l, \quad \{\hat{S}_i, \hat{S}_k\} = \frac{1}{2} \delta_{ik} \hat{I}, \quad (i, k, l = x, y, z) \quad (2.5.9)$$

For spherical components these relations take the form

$$[\hat{S}_\mu, \hat{S}_\nu] = -\sqrt{2} C_{1\mu 1\nu}^1 S_\lambda, \quad \{\hat{S}_\mu, \hat{S}_\nu\} = \frac{1}{2} (-1)^\mu \delta_{\mu-\nu} \hat{I}, \quad (\mu, \nu, \lambda = \pm 1, 0). \quad (2.5.10)$$

2.5.3 Products of Spin Matrices

The four matrices $\hat{S}_x, \hat{S}_y, \hat{S}_z$ and $\hat{I}$ (or $\hat{S}_{+1}, \hat{S}_0, \hat{S}_{-1}$ and $\hat{I}$) constitute a complete set of square 2×2 matrices. Any function of the spin operator ($S = \frac{1}{2}$) may be expanded in a series in terms of these matrices. In particular, products of the spin matrices may also be written in such a form. Products of cartesian components (where the indices i, k, l , etc. take the values x, y, z) are given by

$$\hat{S}_i \hat{S}_k = \frac{1}{4} \delta_{ik} \hat{I} + \frac{i}{2} \varepsilon_{ikl} \hat{S}_l, \quad (2.5.11)$$

$$\hat{S}_i \hat{S}_k \hat{S}_l = \frac{i}{8} \varepsilon_{ikl} \hat{I} + \frac{1}{4} (\hat{S}_i \delta_{kl} - \hat{S}_k \delta_{il} + \hat{S}_l \delta_{ik}), \quad (2.5.12)$$

$$\begin{aligned} \hat{S}_i \hat{S}_k \hat{S}_l \hat{S}_j &= \frac{1}{16} (\delta_{ik} \delta_{lj} - \delta_{il} \delta_{kj} + \delta_{ij} \delta_{kl}) \hat{I}, \\ &+ \frac{i}{8} (\delta_{ik} \varepsilon_{ljm} - \delta_{il} \varepsilon_{kjm} + \delta_{ij} \varepsilon_{klm} + \delta_{kl} \varepsilon_{ijm} - \delta_{kj} \varepsilon_{ilm} + \delta_{lj} \varepsilon_{ikm}) \hat{S}_m. \end{aligned} \quad (2.5.13)$$

In more detailed form, Eq. (2.5.11) is written as

$$\begin{aligned} \hat{S}_x^2 &= \hat{S}_y^2 = \hat{S}_z^2 = \frac{1}{4} \hat{I}, \\ \hat{S}_x \hat{S}_y &= -\hat{S}_y \hat{S}_x = \frac{i}{2} \hat{S}_z, \quad \hat{S}_y \hat{S}_z = -\hat{S}_z \hat{S}_y = \frac{i}{2} \hat{S}_x, \\ \hat{S}_z \hat{S}_x &= -\hat{S}_x \hat{S}_z = \frac{i}{2} \hat{S}_y. \end{aligned} \quad (2.5.14)$$

Products of spherical components are as follows:

$$\begin{aligned} \hat{S}_{+1} \hat{S}_{+1} &= 0, \quad \hat{S}_0 \hat{S}_0 = \frac{1}{4} \hat{I}, \quad \hat{S}_{-1} \hat{S}_{-1} = 0, \\ \hat{S}_{+1} \hat{S}_{-1} &= -\frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = -\frac{1}{4} \hat{I} - \frac{1}{2} \hat{S}_0, \\ \hat{S}_{-1} \hat{S}_{+1} &= -\frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = -\frac{1}{4} \hat{I} + \frac{1}{2} \hat{S}_0, \\ \hat{S}_{+1} \hat{S}_0 &= -\frac{1}{2} \hat{S}_{+1}, \quad \hat{S}_{-1} \hat{S}_0 = \frac{1}{2} \hat{S}_{-1}, \\ \hat{S}_0 \hat{S}_{+1} &= \frac{1}{2} \hat{S}_{+1}, \quad \hat{S}_0 \hat{S}_{-1} = -\frac{1}{2} \hat{S}_{-1}, \\ \hat{S}_\mu \hat{S}_\nu &= \frac{1}{4} (-1)^\mu \delta_{\mu-\nu} \hat{I} - \frac{1}{\sqrt{2}} C_{1\mu 1\nu}^1 \hat{S}_\lambda, \quad (\mu, \nu, \lambda = \pm 1, 0), \quad (\hat{S}_{+1})^n = 0, \quad (\hat{S}_{-1})^n = 0, \quad (n = 2, 3, \dots), \end{aligned} \quad (2.5.15)$$

$$(\hat{S}_0)^{2n} = \left(\frac{1}{4}\right)^n \hat{I}, \quad (\hat{S}_0)^{2n+1} = \left(\frac{1}{4}\right)^n \hat{S}_0, \quad (n = 0, 1, 2, \dots), \quad (2.5.17)$$

$$\hat{\mathbf{S}} \cdot \hat{\mathbf{S}} = \hat{\mathbf{S}}^2 = \hat{S}_x^2 + \hat{S}_y^2 + \hat{S}_z^2 = -\hat{S}_{+1} \hat{S}_{-1} + \hat{S}_0 \hat{S}_0 - \hat{S}_{-1} \hat{S}_{+1} = \frac{3}{4} \hat{I}. \quad (2.5.18)$$

Products of the operators $\hat{\mathbf{S}}$ and any vectors $\mathbf{a}$ and $\mathbf{b}$ commuting with $\hat{\mathbf{S}}$ may be written in the form

$$\hat{\mathbf{S}} \cdot (\hat{\mathbf{S}} \cdot \mathbf{a}) = \frac{1}{4} \hat{I} \mathbf{a} - \frac{i}{2} [\hat{\mathbf{S}} \times \mathbf{a}], \quad (2.5.19)$$

$$(\hat{\mathbf{S}} \cdot [\hat{\mathbf{S}} \times \mathbf{a}]) = ([\hat{\mathbf{S}} \times \hat{\mathbf{S}}] \cdot \mathbf{a}) = i(\hat{\mathbf{S}} \cdot \mathbf{a}), \quad (2.5.20)$$

$$[\hat{\mathbf{S}} \times [\hat{\mathbf{S}} \times \mathbf{a}]] = -\frac{1}{2} \hat{I} \mathbf{a} + \frac{i}{2} [\hat{\mathbf{S}} \times \mathbf{a}], \quad (2.5.21)$$

$$(\hat{\mathbf{S}} \cdot \mathbf{a})(\hat{\mathbf{S}} \cdot \mathbf{b}) = \frac{1}{4} (\mathbf{a} \cdot \mathbf{b}) \hat{I} + \frac{i}{2} (\hat{\mathbf{S}} \cdot [\mathbf{a} \times \mathbf{b}]), \quad (2.5.22)$$

$$([\hat{\mathbf{S}} \times \mathbf{a}] \cdot [\hat{\mathbf{S}} \times \mathbf{b}]) = \frac{1}{2} (\mathbf{a} \cdot \mathbf{b}) \hat{I} + \frac{i}{2} (\hat{\mathbf{S}} \cdot [\mathbf{a} \times \mathbf{b}]), \quad (2.5.23)$$

$$(\hat{\mathbf{S}} \cdot \mathbf{a})[\hat{\mathbf{S}} \times \mathbf{b}] = \frac{1}{4} [\mathbf{a} \times \mathbf{b}] \hat{I} + \frac{i}{2} \{(\hat{\mathbf{S}} \cdot \mathbf{b}) \mathbf{a} - (\mathbf{a} \cdot \mathbf{b}) \hat{\mathbf{S}}\}, \quad (2.5.24)$$

$$[[\hat{\mathbf{S}} \times \mathbf{a}] \times [\hat{\mathbf{S}} \times \mathbf{b}]] = \frac{1}{4} [\mathbf{a} \times \mathbf{b}] \hat{I} + \frac{i}{2} \{(\hat{\mathbf{S}} \cdot \mathbf{b}) \mathbf{a} + (\hat{\mathbf{S}} \cdot \mathbf{a}) \mathbf{b}\}. \quad (2.5.25)$$

If $\mathbf{n}$ is a unit vector ($n^2 = 1$), then

$$(\hat{\mathbf{S}} \cdot \mathbf{n})^{2k} = \left(\frac{1}{4}\right)^k \hat{I}, \quad (\hat{\mathbf{S}} \cdot \mathbf{n})^{2k+1} = \left(\frac{1}{4}\right)^k (\hat{\mathbf{S}} \cdot \mathbf{n}), \quad k = 0, 1, 2, \dots \quad (2.5.26)$$

2.5.4 Functions of Spin Matrices

$$e^{\alpha \hat{S}_i} = \hat{I} \cosh \frac{\alpha}{2} + 2\hat{S}_i \sinh \frac{\alpha}{2}, \quad (2.5.27)$$

$$\cosh(\alpha \hat{S}_i) = \hat{I} \cosh \frac{\alpha}{2}, \quad \sinh(\alpha \hat{S}_i) = 2\hat{S}_i \sinh \frac{\alpha}{2}, \quad (2.5.28)$$

where $i = x, y, z$.

$$e^{\alpha \hat{S}_{+1}} = \hat{I} + \alpha \hat{S}_{+1}, \quad e^{\alpha \hat{S}_0} = \hat{I} \cosh \frac{\alpha}{2} + 2\hat{S}_0 \sinh \frac{\alpha}{2}, \quad e^{\alpha \hat{S}_{-1}} = \hat{I} + \alpha \hat{S}_{-1}, \quad (2.5.29)$$

$$\cosh(\alpha \hat{S}_{+1}) = \hat{I}, \quad \cosh(\alpha \hat{S}_0) = \hat{I} \cosh \frac{\alpha}{2}, \quad \cosh(\alpha \hat{S}_{-1}) = \hat{I}, \quad (2.5.30)$$

$$\sinh(\alpha \hat{S}_{+1}) = \alpha \hat{S}_{+1}, \quad \sinh(\alpha \hat{S}_0) = 2\hat{S}_0 \sinh \frac{\alpha}{2}, \quad \sinh(\alpha \hat{S}_{-1}) = \alpha \hat{S}_{-1}. \quad (2.5.31)$$

2.5.5 Rotation Operators

2.5.6 Euler angles

Under rotations described by the Euler angles α, β, γ spin functions and spin matrices for $S = \frac{1}{2}$ are transformed by a rotation operator $\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ which is a special case of the operator given by Eq. (1.4.31)

$$\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) = e^{-i\alpha \hat{S}_z} \cdot e^{-i\beta \hat{S}_y} \cdot e^{-i\gamma \hat{S}_z} = \begin{pmatrix} \cos \frac{\beta}{2} e^{-i\frac{\alpha+\gamma}{2}} & -\sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} \\ \sin \frac{\beta}{2} e^{i\frac{\alpha-\gamma}{2}} & \cos \frac{\beta}{2} e^{i\frac{\alpha+\gamma}{2}} \end{pmatrix}. \quad (2.5.32)$$

The inverse matrix is

$$\begin{aligned} \left[\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \right]^{-1} &= \left[\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \right]^+ = \hat{D}^{\frac{1}{2}}(\pi - \gamma, \beta, -\pi - \alpha) = \hat{D}^{\frac{1}{2}}(-\gamma, -\beta, -\alpha), \\ &= \begin{pmatrix} \cos \frac{\beta}{2} e^{i\frac{\alpha+\gamma}{2}} & \sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} \\ -\sin \frac{\beta}{2} e^{i\frac{\alpha-\gamma}{2}} & \cos \frac{\beta}{2} e^{-i\frac{\alpha+\gamma}{2}} \end{pmatrix}. \end{aligned} \quad (2.5.33)$$

Matrix elements of the operator $\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ are the Wigner D -functions $D_{MM'}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ (Chap. 4). The expansion of $\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ in terms of the spin matrices has the form

$$\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) = \cos \frac{\beta}{2} \cos \frac{\alpha+\gamma}{2} \hat{I} + 2i \sin \frac{\beta}{2} \sin \frac{\alpha-\gamma}{2} \hat{S}_x - 2i \sin \frac{\beta}{2} \cos \frac{\alpha-\gamma}{2} \hat{S}_y - 2i \cos \frac{\beta}{2} \sin \frac{\alpha+\gamma}{2} \hat{S}_z, \quad (2.5.34)$$

$$\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) = \cos \frac{\beta}{2} \cos \frac{\alpha+\gamma}{2} \hat{I} + \sqrt{2} \sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} \hat{S}_{+1} - 2i \cos \frac{\beta}{2} \sin \frac{\alpha+\gamma}{2} \hat{S}_0 + \sqrt{2} \sin \frac{\beta}{2} e^{i\frac{\alpha-\gamma}{2}} \hat{S}_{-1}. \quad (2.5.35)$$

(a) *Axis and angle*

Under rotations of the coordinate system through an angle ω about an axis $\mathbf{n}(\Theta, \Phi)$ spin functions and spin matrices for $S = \frac{1}{2}$ are transformed by the rotation operator $\hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi)$ which is special case of the operator given by Eq. (1.4.33).

$$\hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi) = e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{S}}} = \begin{pmatrix} \cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \Theta & -i \sin \frac{\omega}{2} \sin \Theta e^{-i\Phi} \\ -i \sin \frac{\omega}{2} \sin \Theta e^{i\Phi} & \cos \frac{\omega}{2} + i \sin \frac{\omega}{2} \cos \Theta \end{pmatrix}. \quad (2.5.36)$$

The inverse matrix is

$$\begin{aligned} \left[\hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi) \right]^{-1} &= \left[\hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi) \right]^+ = \hat{U}^{\frac{1}{2}}(\omega; \pi - \Theta, \pi + \Phi) = \hat{U}^{\frac{1}{2}}(-\omega; \Theta, \Phi) \\ &= \begin{pmatrix} \cos \frac{\omega}{2} + i \sin \frac{\omega}{2} \cos \Theta & i \sin \frac{\omega}{2} \sin \Theta e^{-i\Phi} \\ i \sin \frac{\omega}{2} \sin \Theta e^{i\Phi} & \cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \Theta \end{pmatrix} \end{aligned} \quad (2.5.37)$$

The expansion of the operator $\hat{U}^{\frac{1}{2}}(\omega; \theta, \Phi)$ in terms of the spin matrices has the form

$$\hat{U}^{\frac{1}{2}}(\omega; \theta, \Phi) = e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{S}}} = \hat{I} \cos \frac{\omega}{2} - 2i(\mathbf{n} \cdot \hat{\mathbf{S}}) \sin \frac{\omega}{2}. \quad (2.5.38)$$

The relations between the angles ω, Θ, Φ and the Euler angles α, β, γ are given by Eqs. (1.4.16), (1.4.17). Note that

$$\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) = \hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi). \quad (2.5.39)$$

2.5.7 Result of rotation operator

The result of applying the rotation operator $\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ to the spin matrices may be presented as

$$\hat{S}'_i \equiv \hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \hat{S}_i \left[\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \right]^{-1} = \sum_k a_{ik} \hat{S}_k, \quad (i, k = x, y, z). \quad (2.5.40)$$

where a_{ik} are elements of the rotation matrix (Sec. 1.4.6). For spherical components of the spin matrices we have

$$\hat{S}'_\mu \equiv \hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \hat{S}_\mu \left[\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma) \right]^{-1} = \sum_\nu D^1_{\nu\mu}(\alpha, \beta, \gamma) \hat{S}_\nu; \quad (2.5.41)$$

where $D^1_{\nu\mu}$ are the Wigner D-functions (Chap. 4).

2.5.8 Traces of Products of Spin Matrices ($S = \frac{1}{2}$)

For cartesian components of the spin operator the following relations hold (where i, k, l etc. take the values x, y, z)

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_i \right\} &= 0., \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \right\} &= \frac{1}{2} \delta_{ik}, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \right\} &= \frac{i}{4} \varepsilon_{ikl}, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \hat{S}_j \right\} &= \frac{1}{8} (\delta_{ik} \delta_{lj} - \delta_{il} \delta_{kj} + \delta_{ij} \delta_{kl}), \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \hat{S}_j \hat{S}_m \right\} &= \frac{i}{16} (\delta_{ik} \varepsilon_{ljm} + \delta_{jl} \varepsilon_{ikm} + \delta_{ml} \varepsilon_{ijk} + \delta_{jm} \varepsilon_{ikl}). \end{aligned} \quad (2.5.42)$$

The evaluation of traces of products which contain many spin matrices can be facilitated by using the following recurrence relation: if we introduce the definition

$$T_{i_1 i_2 \dots i_n} \equiv \text{Tr} \left\{ \hat{S}_{i_1} \hat{S}_{i_2} \dots \hat{S}_{i_n} \right\}, \quad (n \geq 3), \quad (2.5.43)$$

then

$$T_{i_1 i_2 \dots i_n} = \frac{1}{4} \delta_{i_1 i_2} T_{i_3 i_4 \dots i_n} + \frac{i}{2} \varepsilon_{i_1 i_2 i'} T_{i' i_3 \dots i_n}. \quad (2.5.44)$$

If n is even, one may also use the relation

$$T_{i_1 i_2 \dots i_n} = \frac{1}{4} \left\{ \delta_{i_1 i_2} T_{i_3 i_4 \dots i_n} - \delta_{i_1 i_3} T_{i_2 i_4 \dots i_n} + \dots + \delta_{i_1 i_n} T_{i_2 i_3 \dots i_{n-1}} \right\}. \quad (2.5.45)$$

For spherical components of the spin operator the following relations hold (where μ, ν, λ etc. take the values $\pm 1, 0$)

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_\mu \right\} &= 0, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \right\} &= \frac{1}{2} (-1)^\mu \delta_{\mu-\nu}, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \right\} &= -\frac{1}{2} \sqrt{\frac{3}{2}} \begin{pmatrix} 1 & 1 & 1 \\ \mu & \nu & \lambda \end{pmatrix} = (-1)^{1+\lambda} \frac{1}{2\sqrt{2}} C^1_{1\mu 1\nu}, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \hat{S}_\rho \right\} &= \frac{(-1)^\mu}{8} \left\{ (-1)^\lambda \delta_{\mu-\nu} \delta_{\lambda-\rho} - (-1)^\nu \delta_{\mu-\lambda} \delta_{\nu-\rho} + (-1)^\nu \delta_{\mu-\rho} \delta_{\nu-\lambda} \right\}. \end{aligned} \quad (2.5.46)$$

The recurrence relation for spherical components of the spin operator is written as

$$T_{\mu_1 \mu_2 \dots \mu_n} = \frac{(-1)^{\mu_1}}{4} \delta_{\mu_1 - \mu_2} T_{\mu_3 \dots \mu_n} - \frac{1}{\sqrt{2}} C^1_{\mu_1 1 \mu_2} T_{\mu_3 \dots \mu_n} \quad (2.5.47)$$

where

$$T_{\mu_1 \mu_2 \dots \mu_n} \equiv \text{Tr} \left\{ \hat{S}_{\mu_1} \hat{S}_{\mu_2} \dots \hat{S}_{\mu_n} \right\}. \quad (2.5.48)$$

If n is even, one may use the formula

$$T_{\mu_1 \mu_2 \dots \mu_n} = \frac{(-1)^{\mu_1}}{4} \left\{ \delta_{\mu_1 - \mu_2} T_{\mu_3 \mu_4 \dots \mu_n} - \delta_{\mu_1 - \mu_3} T_{\mu_2 \mu_4 \dots \mu_n} + \dots + \delta_{\mu_1 - \mu_n} T_{\mu_2 \mu_3 \dots \mu_{n-1}} \right\}. \quad (2.5.49)$$

2.6 Spin matrices and polarization operators for $S = 1$

2.6.1 Spin $S = 1$

If $S = 1$, the spin operator $\hat{S}$ and the polarization operators $\hat{T}_{LM}$ ($L = 0, 1, 2; -L \leq M \leq L$) are square 3×3 matrices. The polarization operator $\hat{T}_{00}$ is proportional to the unit matrix

$$\hat{T}_{00} = \frac{1}{\sqrt{3}}\hat{I}, \quad (2.6.1)$$

where

$$\hat{I} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (2.6.2)$$

The polarization operators $\hat{T}_{1M}$ are proportional to spherical components of the spin matrices $\hat{S}_M$:

$$\hat{T}_{1M} = \frac{1}{\sqrt{2}}\hat{S}_M, \quad (M = 0, \pm 1). \quad (2.6.3)$$

The operators $\hat{T}_{2M}$ ($M = 0, \pm 1, \pm 2$) can be expressed in terms of spherical components of the spin matrices as

$$\hat{T}_{2M} = \sum_{\mu, \nu} C_{1\mu 1\nu}^{2M} \hat{S}_\mu \hat{S}_\nu. \quad (2.6.4)$$

The polarization operators $\hat{T}_{2M}$ are equivalent to some symmetric traceless cartesian tensor of second rank $\hat{Q}_{ik}$:

$$\hat{Q}_{ik} = \frac{1}{2} \left(\hat{S}_i \hat{S}_k + \hat{S}_k \hat{S}_i - \frac{4}{3} \delta_{ik} \hat{I} \right), \quad (i, k = x, y, z), \quad (2.6.5)$$

$$\hat{Q}_{ik} = \hat{Q}_{ki}, \quad \sum_i \hat{Q}_{ii} = 0. \quad (2.6.6)$$

$\hat{Q}_{ik}$ is called the quadrupole tensor. It has 5 linearly independent components. The relations between $\hat{T}_{2M}$ and $\hat{Q}_{ik}$ are given by

$$\begin{aligned} \hat{T}_{2\pm 2} &= \frac{1}{2} \left(\hat{Q}_{xx} - \hat{Q}_{yy} \pm 2i\hat{Q}_{xy} \right), \\ \hat{T}_{2\pm 1} &= \mp \left(\hat{Q}_{xz} \pm i\hat{Q}_{yz} \right), \\ \hat{T}_{20} &= \sqrt{\frac{3}{2}} \hat{Q}_{zz}. \end{aligned} \quad (2.6.7)$$

The inverse relations are

$$\begin{aligned} \hat{Q}_{xx} &= \frac{1}{2} \left(\hat{T}_{22} + \hat{T}_{2-2} \right) - \frac{1}{\sqrt{6}} \hat{T}_{20}, \quad \hat{Q}_{xy} = \hat{Q}_{yx} = \frac{i}{2} \left(\hat{T}_{2-2} - \hat{T}_{22} \right), \\ \hat{Q}_{yy} &= -\frac{1}{2} \left(\hat{T}_{22} + \hat{T}_{2-2} \right) + \frac{1}{\sqrt{6}} \hat{T}_{20}, \quad \hat{Q}_{xz} = \hat{Q}_{zx} = \frac{1}{2} \left(\hat{T}_{2-1} - \hat{T}_{21} \right), \\ \hat{Q}_{zz} &= \sqrt{\frac{2}{3}} \hat{T}_{20}, \quad \hat{Q}_{yz} = \hat{Q}_{zy} = \frac{i}{2} \left(\hat{T}_{2-1} + \hat{T}_{21} \right). \end{aligned} \quad (2.6.8)$$

2.6.2 Explicit Form

Explicit forms of the spin matrices and the polarization operators depend on the representation used for basis spin functions (Chap. 6). We shall consider the spherical basis representation and the cartesian one. These representations are used very frequently.

(a) *Spherical basis representation*

Cartesian components of the spin operator $\hat{\mathbf{S}}$ are given by

$$\hat{S}_x = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \hat{S}_y = \frac{i}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \hat{S}_z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad (2.6.9)$$

and the spherical components of $\hat{\mathbf{S}}$ are

$$\hat{S}_{+1} = -\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{S}_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \hat{S}_{-1} = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}. \quad (2.6.10)$$

The order of rows and columns in these matrices is shown in Eq. (2.3.10).

Matrix elements of the spin operators $\hat{S}_\mu$ in the spherical basis representation are given by

$$(\hat{S}_\mu)_{\sigma'\sigma} = \sqrt{2}C_{1\sigma 1\mu}^{1\sigma'} \quad (\mu, \sigma, \sigma' = \pm 1, 0). \quad (2.6.11)$$

The spin matrices in the spherical basis representation satisfy the relations

$$\hat{S}_i^+ = \hat{S}_i, \quad (i = x, y, z), \quad (2.6.12)$$

$$\hat{S}_x^* = \hat{S}_x, \quad \hat{S}_y^* = -\hat{S}_y, \quad \hat{S}_z^* = \hat{S}_z, \quad (2.6.13)$$

$$\hat{S}_\mu^* = \hat{S}_\mu, \quad \hat{S}_\mu^+ = (-1)^\mu \hat{S}_{-\mu} \quad (\mu = \pm 1, 0). \quad (2.6.14)$$

The quadrupole tensor $\hat{Q}_{ik}(i, k = x, y, z)$ in the spherical basis representation has the form

$$\begin{aligned} \hat{Q}_{xx} &= \frac{1}{6} \begin{pmatrix} -1 & 0 & 3 \\ 0 & 2 & 0 \\ 3 & 0 & -1 \end{pmatrix}, \quad \hat{Q}_{yy} = \frac{1}{6} \begin{pmatrix} -1 & 0 & -3 \\ 0 & 2 & 0 \\ -3 & 0 & -1 \end{pmatrix}, \quad \hat{Q}_{zz} = \frac{1}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ \hat{Q}_{xy} = \hat{Q}_{yx} &= \frac{i}{2} \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \hat{Q}_{xz} = \hat{Q}_{zx} = \frac{1}{2\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}, \quad \hat{Q}_{yz} = \hat{Q}_{zy} = \frac{i}{2\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \end{aligned} \quad (2.6.15)$$

The components of the quadrupole tensor $\hat{Q}_{ik}$ in the spherical basis representation have the following properties

$$\hat{Q}_{ik}^+ = \hat{Q}_{ik}, \quad (i, k = x, y, z), \quad (2.6.16)$$

The matrices $\hat{Q}_{xx}, \hat{Q}_{yy}, \hat{Q}_{zz}, \hat{Q}_{xz}, \hat{Q}_{zx}$ are real, whereas $\hat{Q}_{xy}, \hat{Q}_{yx}, \hat{Q}_{yz}, \hat{Q}_{zy}$ are purely imaginary. The polarization operators $\hat{T}_{LM}$ in the spherical basis representation are given by

$$\hat{T}_{00} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (2.6.17)$$

$$\hat{T}_{1+1} = -\frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{T}_{10} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \hat{T}_{1-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (2.6.18)$$

$$\hat{T}_{2+2} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{T}_{21} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{T}_{20} = \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$\hat{T}_{2-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & -1 & 0 \end{pmatrix}, \quad \hat{T}_{2-2} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}. \quad (2.6.19)$$

Matrix elements of the polarization operators in the spherical basis representation may be written as

$$(\hat{T}_{LM})_{\sigma'\sigma} = \sqrt{\frac{2L+1}{3}} C_{1\sigma' L M}^{1\sigma}, \quad (\sigma, \sigma' = \pm 1, 0; -L \leq M \leq L). \quad (2.6.20)$$

The polarization operators $\hat{T}_{LM}$ in the spherical basis representation are real, i.e.,

$$\hat{T}_{LM}^* = \hat{T}_{LM}. \quad (2.6.21)$$

and satisfy the relations

$$\hat{T}_{LM}^+ = (-1)^M \hat{T}_{L-M}. \quad (2.6.22)$$

(b) Cartesian basis representation

Cartesian components of the spin operator $\hat{S}$ in the cartesian basis representation are given by

$$\hat{S}_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \hat{S}_y = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad \hat{S}_z = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (2.6.23)$$

The matrix elements of $\hat{S}_i$ in this representation may be written as

$$(\hat{S}_i)_{kl} = -i\varepsilon_{ikl}, \quad (i, k, l = x, y, z). \quad (2.6.24)$$

The spherical components of $\hat{\mathbf{S}}$ are as follows

$$\hat{S}_{+1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & i \\ -1 & -i & 0 \end{pmatrix}, \quad \hat{S}_0 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{S}_{-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & -i \\ -1 & i & 0 \end{pmatrix}. \quad (2.6.25)$$

The spin matrices in the cartesian basis representation satisfy the following relations

$$(\hat{S}_\mu)^+ = (-1)^\mu \hat{S}_{-\mu}, \quad \hat{S}_\mu^* = (-1)^{1+\mu} \hat{S}_{-\mu}, \quad (\mu = \pm 1, 0), \quad (2.6.26)$$

$$(\hat{S}_i)^+ = \hat{S}_i, \quad \hat{S}_i^* = -\hat{S}_i, \quad (i = x, y, z). \quad (2.6.27)$$

The quadrupole tensor $\hat{Q}_{ik}(i, k = x, y, z)$ in the cartesian basis representation is given by

$$\begin{aligned} \hat{Q}_{xx} &= \frac{1}{3} \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \hat{Q}_{yy} = \frac{1}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \hat{Q}_{zz} = \frac{1}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \\ \hat{Q}_{xy} &= \hat{Q}_{yx} = \frac{1}{2} \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{Q}_{xz} = \hat{Q}_{zx} = \frac{1}{2} \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad \hat{Q}_{yz} = \hat{Q}_{zy} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}. \end{aligned} \quad (2.6.28)$$

The matrix elements of $\hat{Q}_{ik}$ in the cartesian basis representation may be written in the form

$$(\hat{Q}_{ik})_{lm} = -\frac{1}{2} \left(\delta_{il}\delta_{km} + \delta_{im}\delta_{kl} - \frac{2}{3}\delta_{ik}\delta_{lm} \right), \quad (i, k, l, m = x, y, z). \quad (2.6.29)$$

In the cartesian basis representation the matrices $\hat{Q}_{ik}$ are Hermitian and real, i.e.,

$$\hat{Q}_{ik}^+ = \hat{Q}_{ik}, \quad \hat{Q}_{ik}^* = \hat{Q}_{ik}, \quad (i, k = x, y, z). \quad (2.6.30)$$

The polarization operators $\hat{T}_{LM}$ in the cartesian basis representation have the form

$$\hat{T}_{00} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (2.6.31)$$

$$\begin{aligned} \hat{T}_{1+1} &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & i \\ -1 & -i & 0 \end{pmatrix}, \quad \hat{T}_{10} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{T}_{1-1} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & -i \\ -1 & i & 0 \end{pmatrix}, \\ \hat{T}_{2+2} &= \frac{1}{2} \begin{pmatrix} -1 & -i & 0 \\ -i & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \hat{T}_{2+1} = \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & i \\ 1 & i & 0 \end{pmatrix}, \quad \hat{T}_{20} = \frac{1}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \end{aligned} \quad (2.6.32)$$

$$\hat{T}_{2-1} = \frac{1}{2} \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & i \\ -1 & i & 0 \end{pmatrix}, \quad \hat{T}_{2-2} = \frac{1}{2} \begin{pmatrix} -1 & i & 0 \\ i & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (2.6.33)$$

The polarization operators $\hat{T}_{LM}$ in the cartesian basis representation satisfy the relations

$$\hat{T}_{LM}^+ = (-1)^M \hat{T}_{L-M}, \quad (2.6.34)$$

$$\hat{T}_{LM}^* = (-1)^{L+M} \hat{T}_{L-M}. \quad (2.6.35)$$

One can easily transform any matrix in the cartesian basis representation into a matrix in the spherical basis representation and vice versa with the aid of the unitary matrix U ,

$$\begin{aligned}\widehat{A}(\text{ spherical basis }) &= U \widehat{A}(\text{ cartesian basis }) U^{-1}, \\ \widehat{A}(\text{ cartesian basis }) &= U^{-1} \widehat{A}(\text{ spherical basis }) U.\end{aligned}\tag{2.6.36}$$

In this case $\widehat{A}$ is any spin or polarization operator (i.e., $\widehat{S}_i, \widehat{S}_\mu, \widehat{Q}_{ik}, \widehat{T}_{LM}$),

$$U = \begin{pmatrix} -\frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \\ \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} & 0 \end{pmatrix}, \quad U^{-1} = U^+ = \begin{pmatrix} -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} & 0 & -\frac{i}{\sqrt{2}} \\ 0 & 1 & 0 \end{pmatrix}.\tag{2.6.37}$$

The matrix U coincides with $M(+1, 0, -1 \leftarrow x, y, z)$ and U^{-1} coincides with $M(x, y, z \leftarrow +1, 0, -1)$ (Table 1.2).

The formulas given in Sec. 2.6 (except those of Sec. 2.6.2) are independent of the representation unless the contrary is indicated.

2.6.3 Products of Spin and Polarization Matrices

For $S = 1$, the nine matrices $\widehat{T}_{LM}$ ($L = 0, 1, 2; -L \leq M \leq L$) or, equivalently, the matrices $\widehat{I}, \widehat{S}, \widehat{Q}_{ik}$ constitute a complete set of square 3×3 matrices for expanding any function of the spin and polarization operators. In particular, products of the spin and polarization operators may be expanded in a series in terms of these matrices.

Products of cartesian components of the matrices $\widehat{S}$ and $\widehat{Q}_{ik}$ may be expressed as follows (i, k, l , etc. take the values x, y, z)

(a) *Products of two cartesian components of spin matrices*

$$\widehat{S}_i \widehat{S}_k = \frac{2}{3} \delta_{ik} \widehat{I} + \frac{i}{2} \varepsilon_{ikl} \widehat{S}_l + \widehat{Q}_{ik}.\tag{2.6.38}$$

In particular,

$$\widehat{\mathbf{S}}^2 = \widehat{S}_x^2 + \widehat{S}_y^2 + \widehat{S}_z^2 = 2\widehat{I},\tag{2.6.39}$$

$$[\widehat{S}_i, \widehat{S}_k] \equiv \widehat{S}_i \widehat{S}_k - \widehat{S}_k \widehat{S}_i = i \varepsilon_{ikl} \widehat{S}_l,\tag{2.6.40}$$

$$\{\widehat{S}_i, \widehat{S}_k\} \equiv \widehat{S}_i \widehat{S}_k + \widehat{S}_k \widehat{S}_i = \frac{4}{3} \delta_{ik} \widehat{I} + 2\widehat{Q}_{ik}.\tag{2.6.41}$$

(b) *Products of cartesian components of three spin matrices*

$$\begin{aligned}\widehat{S}_i \widehat{S}_k \widehat{S}_l &= \frac{i}{3} \varepsilon_{ikl} \widehat{I} + \frac{1}{2} (\delta_{ik} \widehat{S}_l + \delta_{kl} \widehat{S}_i) + i \varepsilon_{ilm} \widehat{Q}_{km} \\ &= \frac{i}{3} \varepsilon_{ikl} \widehat{I} + \frac{1}{2} (\delta_{ik} \widehat{S}_l + \delta_{kl} \widehat{S}_i) + \frac{i}{2} (\varepsilon_{ikm} \widehat{Q}_{lm} + \varepsilon_{ilm} \widehat{Q}_{km} + \varepsilon_{klm} \widehat{Q}_{im}).\end{aligned}\tag{2.6.42}$$

From Eq. (2.6.42) one may obtain the Duffin-Kemmer relation

$$\widehat{S}_i \widehat{S}_k \widehat{S}_l + \widehat{S}_l \widehat{S}_k \widehat{S}_i = \delta_{ik} \widehat{S}_l + \delta_{kl} \widehat{S}_i.\tag{2.6.43}$$

Other particular cases of (2.6.42) are

$$\hat{S}_i \hat{S}_k \hat{S}_i = \delta_{ik} \hat{S}_i \quad (\text{no sum over } i); \quad \sum_i \hat{S}_i \hat{S}_k \hat{S}_i = \hat{S}_k \quad (2.6.44)$$

$$\hat{S}_i \hat{S}_k \hat{S}_k + \hat{S}_k \hat{S}_k \hat{S}_i = \hat{S}_i + \delta_{ik} \hat{S}_k \quad (\text{no sum over } k), \quad (2.6.45)$$

$$\begin{aligned} \hat{S}_x \hat{S}_y \hat{S}_z + \hat{S}_y \hat{S}_z \hat{S}_x + \hat{S}_z \hat{S}_x \hat{S}_y &= i\hat{I}, \\ \hat{S}_x \hat{S}_z \hat{S}_y + \hat{S}_z \hat{S}_y \hat{S}_x + \hat{S}_y \hat{S}_x \hat{S}_z &= -i\hat{I}, \end{aligned} \quad (2.6.46)$$

$$\hat{S}_x \hat{S}_z \hat{S}_z = \hat{S}_y \hat{S}_y \hat{S}_x = \frac{1}{2} \hat{S}_x - i\hat{Q}_{yz}, \quad (2.6.47)$$

$$\hat{S}_x \hat{S}_y \hat{S}_y = \hat{S}_z \hat{S}_z \hat{S}_x = \frac{1}{2} \hat{S}_x + i\hat{Q}_{yz},$$

$$\hat{S}_y \hat{S}_x \hat{S}_x = \hat{S}_z \hat{S}_z \hat{S}_y = \frac{1}{2} \hat{S}_y - i\hat{Q}_{xz},$$

$$\hat{S}_y \hat{S}_z \hat{S}_z = \hat{S}_x \hat{S}_x \hat{S}_y = \frac{1}{2} \hat{S}_y + i\hat{Q}_{xz}, \quad (2.6.48)$$

$$\hat{S}_z \hat{S}_y \hat{S}_y = \hat{S}_x \hat{S}_x \hat{S}_z = \frac{1}{2} \hat{S}_z - i\hat{Q}_{xy}, \quad (2.6.49)$$

$$\hat{S}_z \hat{S}_x \hat{S}_x = \hat{S}_y \hat{S}_y \hat{S}_z = \frac{1}{2} \hat{S}_z + i\hat{Q}_{xy}.$$

(c) General powers of cartesian components of spin matrices

$$\hat{S}_i^{2n} = \frac{2}{3} \hat{I} + \hat{Q}_{ii}, \quad (n = 1, 2, \dots), \quad \hat{S}_i^{2n+1} = \hat{S}_i, \quad (n = 0, 1, 2, \dots). \quad (2.6.50)$$

In particular, if $\mathbf{n}$ is the unit vector, then

$$(\hat{\mathbf{S}} \cdot \mathbf{n})^{2k} = \frac{2}{3} \hat{I} + \sum_{i,l} n_i n_l \hat{Q}_{il}, \quad (k = 1, 2, 3, \dots), \quad (2.6.51)$$

$$(\hat{\mathbf{S}} \cdot \mathbf{n})^{2k+1} = (\hat{\mathbf{S}} \cdot \mathbf{n}), \quad (k = 0, 1, 2, \dots).$$

(d) Products of cartesian components of quadrupole tensors and spin matrices

$$\hat{Q}_{ik} \hat{S}_l = \frac{1}{4} \left(\delta_{il} \hat{S}_k + \delta_{kl} \hat{S}_i - \frac{2}{3} \delta_{ik} \hat{S}_l \right) + \frac{i}{2} \left(\varepsilon_{ilm} \hat{Q}_{km} + \varepsilon_{klm} \hat{Q}_{im} \right), \quad (2.6.52)$$

$$\hat{S}_i \hat{Q}_{kl} = \frac{1}{4} \left(\delta_{ik} \hat{S}_l + \delta_{il} \hat{S}_k - \frac{2}{3} \delta_{kl} \hat{S}_i \right) + \frac{i}{2} \left(\varepsilon_{ikm} \hat{Q}_{lm} + \varepsilon_{ilm} \hat{Q}_{km} \right). \quad (2.6.53)$$

(e) Products of cartesian components of quadrupole tensors

$$\begin{aligned} \hat{Q}_{ik} \hat{Q}_{lm} &= \frac{1}{6} \left(\delta_{il} \delta_{km} + \delta_{im} \delta_{kl} - \frac{2}{3} \delta_{ik} \delta_{lm} \right) \hat{I} - \frac{1}{4} \left(\delta_{il} \hat{Q}_{km} + \delta_{im} \hat{Q}_{kl} + \delta_{km} \hat{Q}_{il} + \delta_{kl} \hat{Q}_{im} - \right. \\ &\quad \left. - \frac{4}{3} \delta_{ik} \hat{Q}_{lm} - \frac{4}{3} \delta_{lm} \hat{Q}_{ik} \right) + \frac{i}{8} \left(\delta_{il} \varepsilon_{kmp} \hat{S}_p + \delta_{im} \varepsilon_{klp} \hat{S}_p + \delta_{kl} \varepsilon_{imp} \hat{S}_p + \delta_{km} \varepsilon_{ilp} \hat{S}_p \right) \end{aligned} \quad (2.6.54)$$

Products involving spherical components of the spin operator $\hat{\mathbf{S}}$ and the polarization operators $\hat{T}_{LM}$ may be expanded in a series as given below ($\mu, \nu = \pm 1, 0$)

(f) Products of spherical components spin matrices

$$\hat{S}_\mu \hat{S}_\nu = \frac{2}{3} (-1)^\mu \delta_{\mu-\nu} \hat{I} - \frac{1}{\sqrt{2}} C_{1\mu 1\nu}^{1\lambda} \hat{S}_\lambda + C_{1\mu 1\nu}^{2M} \hat{T}_{2M}. \quad (2.6.55)$$

In particular

$$\hat{S}^2 = -\hat{S}_{+1} \hat{S}_{-1} + \hat{S}_0 \hat{S}_0 - \hat{S}_{-1} \hat{S}_{+1} = 2\hat{I} \quad (2.6.56)$$

$$[\hat{S}_\mu, \hat{S}_\nu] \equiv \hat{S}_\mu \hat{S}_\nu - \hat{S}_\nu \hat{S}_\mu = -\sqrt{2} C_{1\mu 1\nu}^{1\lambda} \hat{S}_\lambda \quad (2.6.57)$$

$$\{\hat{S}_\mu, \hat{S}_\nu\} \equiv \hat{S}_\mu \hat{S}_\nu + \hat{S}_\nu \hat{S}_\mu = \frac{4}{3} (-1)^\mu \delta_{\mu-\nu} + 2 C_{1\mu 1\nu}^{2M} \hat{T}_{2M} \quad (2.6.58)$$

(g) *Products of spherical components spin matrices and polarisation operators*

$$\hat{S}_{\pm 1}^2 = \hat{T}_{2\pm 2}, \hat{S}_{\pm 1}^3 = \hat{S}_{\pm 1}^4 = \hat{S}_{\pm 1}^5 = \dots = 0. \quad (2.6.59)$$

$$\hat{S}_0^2 = \frac{2}{3}\hat{I} + \sqrt{\frac{2}{3}}\hat{T}_{20} \quad (2.6.60)$$

$$\hat{S}_0^{2n} = \hat{S}_0^2, (n = 1, 2, \dots), \quad \hat{S}_0^{2n+1} = \hat{S}_0, \quad (n = 0, 1, 2, \dots). \quad (2.6.61)$$

(h) *Products of polarisation operators*

$$\hat{T}_{L_1 M_1} \hat{T}_{L_2 M_2} = \sqrt{(2L_1 + 1)(2L_2 + 1)} \sum_L (-1)^L \begin{Bmatrix} L_1 & L_2 & L \\ 1 & 1 & 1 \end{Bmatrix} C_{L_1 M_1 L_2 M_2}^{L M} \hat{T}_{L M}. \quad (2.6.62)$$

Particularly,

$$\hat{T}_{LM} \hat{T}_{00} = \hat{T}_{00} \hat{T}_{LM} = \frac{1}{\sqrt{3}} \hat{T}_{LM}, \quad (L = 0, 1, 2; -L \leq M \leq L), \quad (2.6.63)$$

$$\hat{S}_\mu \hat{T}_{2M} = -\frac{1}{2} \sqrt{\frac{5}{3}} C_{1\mu 2M}^{1\nu} \hat{S}_\nu - \sqrt{\frac{3}{2}} C_{1\mu 2M}^{2N} \hat{T}_{2N}, \quad (2.6.64)$$

$$\hat{T}_{2M} \hat{S}_\mu = -\frac{1}{2} \sqrt{\frac{5}{3}} C_{1\mu 2M}^{1\nu} \hat{S}_\nu + \sqrt{\frac{3}{2}} C_{1\mu 2M}^{2N} \hat{T}_{2N}, \quad (2.6.65)$$

$$\hat{T}_{2M} \hat{T}_{2N} = (-1)^M \cdot \frac{1}{3} \delta_{M-N} \hat{I} + \frac{1}{2} \sqrt{\frac{5}{2}} C_{2M 2N}^{1\mu} \hat{S}_\mu + \frac{1}{2} \sqrt{\frac{7}{3}} C_{2M 2N}^{2A} \hat{T}_{2A}. \quad (2.6.66)$$

2.6.4 Functions of Spin Matrices

$$e^{\alpha \hat{S}_i} = \frac{1}{3} (1 + 2 \cosh \alpha) \hat{I} + \sinh \alpha \hat{S}_i + (\cosh \alpha - 1) \hat{Q}_{ii} \quad (\text{no sum over } i), \quad (2.6.67)$$

$$\cosh(\alpha \hat{S}_i) = \frac{1}{3} (1 + 2 \cosh \alpha) \hat{I} + (\cosh \alpha - 1) \hat{Q}_{ii}, \quad (2.6.68)$$

$$\sinh(\alpha \hat{S}_i) = \sinh \alpha \hat{S}_i, \quad (2.6.69)$$

where $i = x, y, z$.

$$e^{\alpha \hat{S}_{+1}} = \hat{I} + \alpha \hat{S}_{+1} + \frac{\alpha^2}{2} \hat{T}_{22}, \quad (2.6.70)$$

$$e^{\alpha \hat{S}_0} = \frac{1}{3} (1 + 2 \cosh \alpha) \hat{I} + \sinh \alpha \hat{S}_0 + \sqrt{\frac{2}{3}} (\cosh \alpha - 1) \hat{T}_{20}, \quad (2.6.71)$$

$$e^{\alpha \hat{S}_{-1}} = \hat{I} + \alpha \hat{S}_{-1} + \frac{\alpha^2}{2} \hat{T}_{2-2}, \quad (2.6.72)$$

$$\sinh(\alpha \hat{S}_{+1}) = \alpha \hat{S}_{+1}, \quad \sinh(\alpha \hat{S}_0) = \sinh \alpha \hat{S}_0, \quad \sinh(\alpha \hat{S}_{-1}) = \alpha \hat{S}_{-1}. \quad (2.6.73)$$

$$\cosh(\alpha \hat{S}_{+1}) = \hat{I} + \frac{\alpha^2}{2} \hat{T}_{22},$$

$$\cosh(\alpha \hat{S}_0) = \frac{1}{3} (1 + 2 \cosh \alpha) \hat{I} + \sqrt{\frac{2}{3}} (\cosh \alpha - 1) \hat{T}_{20}, \quad (2.6.74)$$

$$\cosh(\alpha \hat{S}_{-1}) = \hat{I} + \frac{\alpha^2}{2} \hat{T}_{2-2}.$$

2.6.5 Operators of Coordinate Rotations

(a) *Coordinaterotations defined be Euler angles*

Under rotations of coordinate systems defined by the Euler angles α, β, γ spin functions and spin matrices for $S = 1$ are transformed by the rotation operator $\hat{D}^1(\alpha, \beta, \gamma)$ which is a special case of the operator given by Eq. (1.4.31)

$$\hat{D}^1(\alpha, \beta, \gamma) = e^{-i\alpha \hat{S}_x} e^{-i\beta \hat{S}_y} e^{-i\gamma \hat{S}_z}. \quad (2.6.75)$$

In the spherical basis representation $\hat{D}^1(\alpha, \beta, \gamma)$ has the form

$$\hat{D}^1(\alpha, \beta, \gamma) = \begin{pmatrix} \frac{1+\cos\beta}{2}e^{-i(\alpha+\gamma)} & -\frac{\sin\beta}{\sqrt{2}}e^{-i\alpha} & \frac{1-\cos\beta}{2}e^{i(\gamma-\alpha)} \\ \frac{\sin\beta}{\sqrt{2}}e^{-i\gamma} & \cos\beta & -\frac{\sin\beta}{\sqrt{2}}e^{i\gamma} \\ \frac{1-\cos\beta}{2}e^{i(\alpha-\gamma)} & \frac{\sin\beta}{\sqrt{2}}e^{i\alpha} & \frac{1+\cos\beta}{2}e^{i(\gamma+\alpha)} \end{pmatrix}. \quad (2.6.76)$$

The matrix elements of $\hat{D}^1(\alpha, \beta, \gamma)$ in this representation are the Wigner D -functions $D_{MM'}^1(\alpha, \beta, \gamma)$ (see Chap. 4).

In the cartesian basis representation we have

$$\hat{D}^1(\alpha, \beta, \gamma) = \begin{pmatrix} \cos\beta \cos\alpha \cos\gamma - \sin\alpha \sin\gamma & -\cos\beta \cos\alpha \sin\gamma - \sin\alpha \cos\gamma & \sin\beta \cos\alpha \\ \cos\beta \sin\alpha \cos\gamma + \cos\alpha \sin\gamma & -\cos\beta \sin\alpha \sin\gamma + \cos\alpha \cos\gamma & \sin\beta \sin\alpha \\ -\sin\beta \cos\gamma & \sin\beta \sin\gamma & \cos\beta \end{pmatrix}. \quad (2.6.77)$$

The matrix elements of $\hat{D}^1(\alpha, \beta, \gamma)$ in the cartesian basis representation coincide with the elements of the rotation matrix a_{ik} (see Sec. 1.4.6).

The inverse rotation of the coordinate system is performed by the matrix

$$[\hat{D}^1(\alpha, \beta, \gamma)]^{-1} = [\hat{D}^1(\alpha, \beta, \gamma)]^+ = \hat{D}^1(-\gamma, -\beta, -\alpha) = \hat{D}^1(\pi - \gamma; \beta, -\pi - \alpha). \quad (2.6.78)$$

(b) *Coordinate rotations defined by an axis and angle of rotation*

Under rotation of the coordinate system through an angle ω about an axis $\mathbf{n}(\Theta, \Phi)$ the spin functions and spin matrices for $S = 1$ are transformed by the rotation operator $\hat{U}^1(\omega; \Theta, \Phi)$ which is a special case of the operator given by Eq. (1.4.33)

$$\hat{U}^1(\omega; \Theta, \Phi) = e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{s}}}. \quad (2.6.79)$$

In the spherical basis representation this operator may be written as

$$\hat{U}^1(\omega; \Theta, \Phi) = \begin{pmatrix} \frac{1}{2} \cos\omega (1 + \cos^2\Theta) - i \sin\omega \cos\Theta + \frac{1}{2} \sin^2\Theta & \frac{1}{\sqrt{2}} \sin\Theta e^{i\Phi} [(\cos\omega - 1) \cos\Theta - i \sin\omega] & \frac{1}{2} (\cos\omega - 1) \sin^2\Theta e^{i2\Phi} \\ \frac{1}{\sqrt{2}} \sin\Theta e^{-i\Phi} [(\cos\omega - 1) \cos\Theta - i \sin\omega] & \cos\omega \sin^2\Theta + \cos^2\Theta & -\frac{1}{\sqrt{2}} \sin\Theta e^{i\Phi} [(\cos\omega - 1) \cos\Theta + i \sin\omega] \\ -\frac{1}{\sqrt{2}} \sin\Theta e^{i\Phi} [(\cos\omega - 1) \cos\Theta + i \sin\omega] & \frac{1}{2} \cos\omega (1 + \cos^2\Theta) + i \sin\omega \cos\Theta + \frac{1}{2} \sin^2\Theta & \end{pmatrix}. \quad (2.6.80)$$

In the cartesian basis representation $\hat{U}^1(\omega; \Theta, \Phi)$ has the form

$$\hat{U}^1(\omega; \Theta, \Phi) = \begin{pmatrix} (1 - \cos\omega) \sin^2\Theta \cos^2\Phi + \cos\omega & (1 - \cos\omega) \sin^2\Theta \cos\Phi \sin\Phi + \sin\omega \cos\Theta & (1 - \cos\omega) \sin\Theta \cos\Theta \cos\Phi + \sin\omega \sin\Theta \sin\Phi \\ (1 - \cos\omega) \sin^2\Theta \cos\Phi \sin\Phi - \sin\omega \cos\Theta & (1 - \cos\omega) \sin^2\Theta \sin^2\Phi + \cos\omega & (1 - \cos\omega) \sin\Theta \cos\Theta \sin\Phi - \sin\omega \sin\Theta \cos\Phi \\ (1 - \cos\omega) \sin\Theta \cos\Theta \cos\Phi - \sin\omega \sin\Theta \sin\Phi & (1 - \cos\omega) \sin\Theta \cos\Theta \sin\Phi + \sin\omega \sin\Theta \cos\Phi & (1 - \cos\omega) \cos^2\Theta + \cos\omega \end{pmatrix}. \quad (2.6.81)$$

The inverse rotation of the coordinate system is performed by the matrix

$$[\hat{U}^1(\omega; \Theta, \Phi)]^{-1} = [\hat{U}^1(\omega; \Theta, \Phi)]^+ = \hat{U}^1(-\omega; \Theta, \Phi) = \hat{U}^1(\omega; \pi - \Theta, \pi + \Phi) \quad (2.6.82)$$

The expansion of the rotation operator $\hat{U}^1(\omega; \Theta, \Phi)$ in terms of the spin matrices and the polarization operators has the form

$$\hat{U}^1(\omega; \Theta, \Phi) = \frac{1}{3}(2 \cos\omega + 1)\hat{I} - i \sin\omega (\mathbf{n} \cdot \hat{\mathbf{S}}) + (\cos\omega - 1) \sum_{i,k} n_i n_k \hat{Q}_{ik}. \quad (2.6.83)$$

(See also Eqs. (2.4.17) and (2.4.18).)

The relations between the angles ω, Θ, Φ and the Euler angles α, β, γ are given by Eqs. (1.4.16), (1.4.17). Note that

$$\hat{D}^1(\alpha, \beta, \gamma) = \hat{U}^1(\omega; \Theta, \Phi). \quad (2.6.84)$$

(c) *Effect of a rotation operator on a spin matrix*

Applying the rotation operator $\hat{D}^1(\alpha, \beta, \gamma)$ to the spin matrices yields ($i, k, j, l = x, y, z$)

$$\hat{S}'_i \equiv \hat{D}^1(\alpha, \beta, \gamma) \hat{S}_i \left[\hat{D}^1(\alpha, \beta, \gamma) \right]^{-1} = \sum_k a_{ik} \hat{S}_k, \quad (2.6.85)$$

$$\hat{Q}'_{ik} \equiv \hat{D}^1(\alpha, \beta, \gamma) \hat{Q}_{ik} \left[\hat{D}^1(\alpha, \beta, \gamma) \right]^{-1} = \sum_{j,l} a_{ij} a_{kl} \hat{Q}_{jl}, \quad (2.6.86)$$

where a_{ik} is the rotation matrix (Sec. 1.4.6)

$$\hat{S}'_\mu \equiv \hat{D}^1(\alpha, \beta, \gamma) \hat{S}_\mu \left[\hat{D}^1(\alpha, \beta, \gamma) \right]^{-1} = \sum_\nu D^1_{\nu\mu}(\alpha, \beta, \gamma) \hat{S}_\nu, \quad (\mu, \nu = \pm 1, 0), \quad (2.6.87)$$

$$\hat{T}'_{LM} \equiv \hat{D}^1(\alpha, \beta, \gamma) \hat{T}_{LM} \left[\hat{D}^1(\alpha, \beta, \gamma) \right]^{-1} = \sum_{M'} D^L_{M'M}(\alpha, \beta, \gamma) \hat{T}_{LM'}, \quad (-L \leq M, M' \leq L). \quad (2.6.88)$$

$D^L_{MM'}(\alpha, \beta, \gamma)$ being the Wigner D -functions (see Chap. 4).

2.6.6 Traces of Products of Spin Matrices

For cartesian components of the spin matrices $\hat{S}_i$ and $\hat{Q}_{ik}$ the following relations hold (i, k, l , etc. take the values x, y, z).

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_i \right\} &= 0, \quad \text{Tr} \left\{ \hat{Q}_{ik} \right\} = 0, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \right\} &= 2\delta_{ik}, \quad \text{Tr} \left\{ \hat{S}_i \hat{Q}_{kl} \right\} = 0, \quad \text{Tr} \left\{ \hat{Q}_{ik} \hat{S}_l \right\} = 0, \\ \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \right\} &= i\varepsilon_{ikl}, \\ \text{Tr} \left\{ \hat{Q}_{ik} \hat{Q}_{lm} \right\} &= \frac{1}{2} \left(\delta_{il} \delta_{km} + \delta_{im} \delta_{kl} - \frac{2}{3} \delta_{ik} \delta_{lm} \right), \quad \text{Tr} \left\{ \hat{S}_i \hat{S}_k \hat{S}_l \hat{S}_m \right\} = \delta_{ik} \delta_{lm} + \delta_{im} \delta_{kl}. \end{aligned} \quad (2.6.89)$$

For spherical components of the spin operator $\hat{S}_\mu$ we have (μ, ν, λ , etc. take the values $\pm 1, 0$)

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_\mu \right\} &= 0, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \right\} &= (-1)^\mu 2\delta_{\mu-\nu}, \\ \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \right\} &= -\sqrt{6} \begin{pmatrix} 1 & 1 & 1 \\ \mu & \nu & \lambda \end{pmatrix} = (-1)^{1+\lambda} \sqrt{2} C^1_{1\mu 1\nu}, \end{aligned} \quad (2.6.90)$$

$$\begin{aligned} \text{Tr} \left\{ \hat{S}_\mu \hat{S}_\nu \hat{S}_\lambda \hat{S}_\rho \right\} &= (-1)^{\mu+\lambda} (\delta_{\mu-\nu} \delta_{\lambda-\rho} + \delta_{\mu-\rho} \delta_{\nu-\lambda}), \\ \text{Tr} \left\{ \hat{S}_{\mu_1} \hat{S}_{\mu_2} \dots \hat{S}_{\mu_n} \right\} &= 0 \quad \text{if } \mu_1 + \mu_2 + \dots + \mu_n \neq 0. \end{aligned} \quad (2.6.91)$$

Traces of products of the polarization operators $\hat{T}_{LM}$ ($L = 0, 1, 2; -L \leq M \leq L$) can be evaluated by using Eqs. (2.4.21)-(2.4.24) at $S = 1$.

Chapter 3

Irreducible tensors

3.1 Definition and properties of irreducible tensors

3.1.1 Definition

Irreducible tensors occupy a central position in angular momentum theory. Under rotations of coordinate systems these tensors transform in the same manner as eigenfunctions of the angular momentum operator. The use of this property permits us to develop very effective methods for calculating matrix elements of different quantum-mechanical operators.

An irreducible tensor $\mathfrak{M}_J$ of rank J (with J integer or half-integer) is defined as a set of $2J + 1$ functions (components) $\mathfrak{M}_{JM}$ (where $M = -J, -J + 1, \dots, J - 1, J$) which satisfy the following commutation rules with spherical components of the angular momentum operator

$$\begin{aligned} \left[\hat{J}_{\pm 1}, \mathfrak{M}_{JM} \right] &= \mp \frac{1}{\sqrt{2}} e^{\pm i\delta} \sqrt{J(J+1) - M(M \pm 1)} \mathfrak{M}_{JM \pm 1}, \\ \left[\hat{J}_0, \mathfrak{M}_{JM} \right] &= M \mathfrak{M}_{JM}. \end{aligned} \quad (3.1.1)$$

In compact form

$$\left[\hat{J}_\mu, \mathfrak{M}_{JM} \right] = e^{i\mu\delta} \sqrt{J(J+1)} C_{J M 1 \mu}^{J M + \mu} \mathfrak{M}_{JM + \mu}. \quad (3.1.2)$$

From these relations it follows that

$$\left[\hat{\mathbf{J}}^2, \mathfrak{M}_{JM} \right] = J(J+1) \mathfrak{M}_{JM}. \quad (3.1.3)$$

The quantity δ in Eq. (3.1.1) which determines relative phases of different $\mathfrak{M}_{JM}$ components is arbitrary. Let us adopt $\delta = 0$, i.e., $e^{\pm i\delta} = 1$, and choose the positive sign of the square root. The linear equations (3.1.1) define the components of the irreducible tensor $\mathfrak{M}_{JM}$ within an arbitrary scalar factor, which is the same for all the components. This factor can be a real or complex number, function or operator. In the case of integer rank J the overall phase of the $\mathfrak{M}_{JM}$ components is usually defined in such manner that

$$(\mathfrak{M}_{JM})^* = (-1)^{-M} \mathfrak{M}_{J-M}. \quad (3.1.4)$$

This choice of the phase coincides with that for spherical harmonics (Chap. 5). However, sometimes in quantum-mechanical applications it is convenient to redefine irreducible tensors as

$$\tilde{\mathfrak{M}}_J = i^J \mathfrak{M}_J. \quad (3.1.5)$$

Then one has

$$\left(\tilde{\mathfrak{M}}_{JM} \right)^* = (-1)^{J-M} \tilde{\mathfrak{M}}_{J-M}. \quad (3.1.6)$$

The choice of the phase (3.1.6) can be used for tensors of integer as well as half-integer rank J . Making use of this phase convention for tensor operators as well as for wavefunctions describing initial $|a\rangle$ and final states $|b\rangle$ we get the following relations for matrix elements of Hermitian $\tilde{\mathfrak{M}}_J^\dagger = \tilde{\mathfrak{M}}_J$ operators

$$\langle b | \tilde{\mathfrak{M}}_{JM} | a \rangle = \left(\langle a | \left(\tilde{\mathfrak{M}}_{JM} \right)^* | b \rangle \right)^*. \quad (3.1.7)$$

3.1.2 Covariant and Contravariant Components

Any irreducible tensor $\mathfrak{M}_J$ of rank J can be expanded in a series based on a complete set of the unit orthonormalized irreducible tensors $\mathbf{e}_{JM}$ of rank J

$$\mathbf{e}_J^M \cdot \mathbf{e}_{J'M'} = \delta_{JJ'} \delta_{MM'}. \quad (3.1.8)$$

The tensors $\mathbf{e}_{JM}$ can be composed, for example, of the basis spin functions. The expansion of $\mathfrak{M}_J$ is written as

$$\mathfrak{M}_J = \sum_M \mathbf{e}_J^M \cdot \mathfrak{M}_{JM} = \sum_M \mathbf{e}_{JM} \cdot \mathfrak{M}_J^M. \quad (3.1.9)$$

$\mathfrak{M}_{JM}$ represents the covariant component of the tensor $\mathfrak{M}_J$ and $\mathfrak{M}_J^M$ denotes the contravariant component. These components are related by

$$\begin{aligned} \mathfrak{M}_J^M &= (\mathfrak{M}_{JM})^* = (-1)^{-M} \mathfrak{M}_{J-M}, \\ \tilde{\mathfrak{M}}_J^M &= (\tilde{\mathfrak{M}}_{JM})^* = (-1)^{J-M} \tilde{\mathfrak{M}}_{J-M}. \end{aligned} \quad (3.1.10)$$

3.1.3 Transformation of Irreducible Tensors Under a Rotation of the Coordinate System

Under rotations of the coordinate system described by the Euler angles α, β, γ , the components of irreducible tensors $\mathfrak{M}_{JM}$ and $\tilde{\mathfrak{M}}_{JM}$ undergo linear transformation. The coefficients of such transformation are the Wigner D -functions (Chap. 4)

$$\begin{aligned} \mathfrak{M}_{JM'}(X') &= \hat{D}(\alpha, \beta, \gamma) \mathfrak{M}_{JM}(X) [\hat{D}(\alpha, \beta, \gamma)]^{-1} = \sum_M \mathfrak{M}_{JM}(X) D_{MM'}^J(\alpha, \beta, \gamma), \\ \tilde{\mathfrak{M}}_{JM'}(X') &= \hat{D}(\alpha, \beta, \gamma) \tilde{\mathfrak{M}}_{JM}(X) [\hat{D}(\alpha, \beta, \gamma)]^{-1} = \sum_M \tilde{\mathfrak{M}}_{JM}(X) D_{MM'}^J(\alpha, \beta, \gamma). \end{aligned} \quad (3.1.11)$$

Here X and X' denote sets of all arguments of the tensor in the initial and final coordinate systems, respectively.

3.1.4 Transformation of Irreducible Tensors Under Inversion of the Coordinate System

The transformation properties of irreducible tensor components under inversion of the coordinate system, $\mathbf{r} \rightarrow -\mathbf{r}$, permit us generally to represent an irreducible tensor $\mathfrak{M}_J$ of integer rank J as the sum of two tensors $\mathfrak{M}_J^{(+1)}$ and $\mathfrak{M}_J^{(-1)}$, i.e.,

$$\mathfrak{M}_J = \mathfrak{M}_J^{(+1)} + \mathfrak{M}_J^{(-1)}. \quad (3.1.12)$$

Each of these tensors has definite parity.

Under inversion of the coordinate system the tensors $\mathfrak{M}_J^{(\pm 1)}$ transform in the following way

$$\hat{P}_r \mathfrak{M}_J^{(\pi_J)} \hat{P}_r^{-1} = \pi_J \mathfrak{M}_J^{(\pi_J)}, \quad (\pi_J = \pm 1). \quad (3.1.13)$$

The tensor $\mathfrak{M}_J^{(\pi_J)}$ with parity $\pi_J = (-1)^J$ is called the true (or polar) irreducible tensor of rank J . Tensor $\mathfrak{M}_J^{(\pi_J)}$ with parity $\pi_J = (-1)^{J+1}$ is the pseudotensor (or axial tensor) of rank J . The relations (3.1.12) and (3.1.13) are valid for the tensors $\tilde{\mathfrak{M}}_J$ as well.

3.1.5 Double Tensors

A double tensor $W_{J_1 J_2}(1, 2)$ of ranks J_1 and J_2 has $(2J_1 + 1)(2J_2 + 1)$ components and depends on variables of two different subsystems, 1 and 2. The components of the double tensor satisfy the following commutation relations

$$[\hat{J}_{\pm 1}(1), W_{J_1 M_1 J_2 M_2}(1, 2)] = \mp \frac{1}{\sqrt{2}} \sqrt{J_1(J_1 + 1) - M_1(M_1 \pm 1)} W_{J_1 M_1 \pm 1 J_2 M_2}(1, 2), \quad (3.1.14)$$

$$[\hat{J}_0(1), W_{J_1 M_1 J_2 M_2}(1, 2)] = M_1 W_{J_1 M_1 J_2 M_2}(1, 2), \quad (3.1.15)$$

$$[\hat{J}_{\pm 1}(2), W_{J_1 M_1 J_2 M_2}(1, 2)] = \mp \frac{1}{\sqrt{2}} \sqrt{J_2(J_2 + 1) - M_2(M_2 \pm 1)} W_{J_1 M_1 J_2 M_2 \pm 1}(1, 2), \quad (3.1.16)$$

$$[\hat{J}_0(2), W_{J_1 M_1 J_2 M_2}(1, 2)] = M_2 W_{J_1 M_1 J_2 M_2}(1, 2). \quad (3.1.17)$$

Here $\hat{\mathbf{J}}(1)$ and $\hat{\mathbf{J}}(2)$ are the operators of the total angular momenta of subsystems 1 and 2, respectively. Under rotation of subsystem 1 and 2 the double tensor $W_{J_1 J_2}(1, 2)$ transforms according to the representation D^{J_1} or D^{J_2} , respectively.

$$W_{J_1 M'_1 J_2 M_2}(1', 2) = \sum_{M_1} W_{J_1 M_1 J_2 M_2}(1, 2) D_{M_1 M'_1}^{J_1}(\alpha_1, \beta_1, \gamma_1), \quad (3.1.18)$$

$$W_{J_1 M_1 J_2 M'_2}(1, 2') = \sum_{M_2} W_{J_1 M_1 J_2 M_2}(1, 2) D_{M_2 M'_2}^{J_2}(\alpha_2, \beta_2, \gamma_2). \quad (3.1.19)$$

3.1.6 Examples of Irreducible Tensors

In this section we present some examples of irreducible tensors considered in this book.

- (a) The operators of the angular momenta $\hat{\mathbf{J}}, \hat{\mathbf{L}}, \hat{\mathbf{S}}$ (Chap. 2) are irreducible tensor operators of rank 1.
- (b) The polarization operator $\hat{T}_{LM}(S)$ (Sec. 2.4) is an irreducible tensor operator of rank L .
- (c) Spherical harmonics $Y_{lm}(\vartheta, \varphi)$ (Chap. 5) are irreducible tensors of rank l .
- (d) The spin wave functions of a particle of spin S (Chap. 6) are irreducible tensors of rank S .
- (e) Tensor spherical harmonics $Y_{JM}^{LS}(\vartheta, \varphi)$ (Chap. 7) are irreducible tensors of rank J .

We may apply the results given in this chapter for all the tensors mentioned above.

3.1.7 Direct and Irreducible Tensor Products. Commutators of Tensor Products

An irreducible tensor product $\mathfrak{L}_J$ of two irreducible tensors $\mathfrak{M}_{J_1}$ and $\mathfrak{N}_{J_2}$ of ranks J_1 and J_2 is defined as the tensor of rank J whose components $\mathfrak{L}_{JM}$ can be expressed in terms of $\mathfrak{M}_{J_1 M_1}$ and $\mathfrak{N}_{J_2 M_2}$ according to

$$\mathfrak{L}_{JM} = \sum_{M_1 M_2} C_{J_1 M_1 J_2 M_2}^{J M} \mathfrak{M}_{J_1 M_1} \mathfrak{N}_{J_2 M_2} \quad (3.1.20)$$

Irreducible tensor product is denoted as

$$\mathfrak{L}_J \equiv \{\mathfrak{M}_{J_1} \otimes \mathfrak{N}_{J_2}\}_J. \quad (3.1.21)$$

The direct product of two irreducible tensors $\mathfrak{M}_{J_1}$ and $\mathfrak{N}_{J_2}$ is defined as a set of $(2J_1 + 1)(2J_2 + 1)$ components $\mathfrak{M}_{J_1 M_1} \mathfrak{N}_{J_2 M_2}$. This tensor is generally reducible and can be decomposed into parts which themselves transform independently under a rotation of the coordinate system. In other words, the direct product can be represented as a sum of irreducible tensors $\mathfrak{L}_{JM}$

$$\mathfrak{M}_{J_1 M_1} \mathfrak{N}_{J_2 M_2} = \sum_{J=|J_1-J_2|}^{J_1+J_2} C_{J_1 M_1 J_2 M_2}^{J M} \mathfrak{L}_{JM}. \quad (3.1.22)$$

It is essential that the irreducible tensor product $\tilde{\mathfrak{L}}_{JM}$ satisfies the same equality (3.1.6) as $\tilde{\mathfrak{M}}_{J_1 M_1}$ and $\tilde{\mathfrak{N}}_{J_2 M_2}$

$$(\tilde{\mathfrak{L}}_{JM})^* = (-1)^{J-M} \tilde{\mathfrak{L}}_{J-M} \left(\tilde{\mathfrak{L}}_{JM} = \left\{ \tilde{\mathfrak{M}}_{J_1} \otimes \tilde{\mathfrak{N}}_{J_2} \right\}_{JM} \right) \quad (3.1.23)$$

This property is specific for tensors of the $\tilde{\mathfrak{M}}_{JM}$ -type. The tensor product $\mathfrak{L}_{JM}$ does not satisfy the relation (3.1.4) although $\mathfrak{M}_{J_1 M_1}$ and $\mathfrak{N}_{J_2 M_2}$ satisfy this relation.

Let us introduce some definitions and notations which will be widely used in the consideration of products of non-commuting tensor operators.

The commutator of components of two irreducible tensors is defined by

$$\mathfrak{R}_{J_1 M_1 J_2 M_2} \equiv [\mathfrak{M}_{J_1 M_1}, \mathfrak{N}_{J_2 M_2}] \equiv \mathfrak{M}_{J_1 M_1} \mathfrak{N}_{J_2 M_2} - \mathfrak{N}_{J_2 M_2} \mathfrak{M}_{J_1 M_1}. \quad (3.1.24)$$

The commutator of an irreducible tensor product is written as

$$\mathfrak{R}_{JM}^{J_1 J_2} \equiv \{\mathfrak{M}_{J_1} \otimes \mathfrak{N}_{J_2}\}_{JM} - (-1)^{J_1+J_2-J} \{\mathfrak{N}_{J_2} \otimes \mathfrak{M}_{J_1}\}_{JM}. \quad (3.1.25)$$

The functions $\mathfrak{R}_{JM}^{J_1 J_2}$ are the components of some irreducible tensor. They may be expressed in the form

$$\mathfrak{R}_{JM}^{J_1 J_2} = \sum_{M_1 M_2} C_{J_1 M_1 J_2 M_2}^{J M} \mathfrak{R}_{J_1 M_1 J_2 M_2} \quad (3.1.26)$$

For commuting tensors we get

$$\{\mathfrak{M}_{J_1} \otimes \mathfrak{N}_{J_2}\}_{JM} = (-1)^{J_1+J_2-J} \{\mathfrak{N}_{J_2} \otimes \mathfrak{M}_{J_1}\}_{JM}. \quad (3.1.27)$$

On the other hand, for non-commuting tensors we have

$$\{\mathfrak{M}_{J_1} \otimes \mathfrak{N}_{J_2}\}_{JM} = (-1)^{J_1+J_2-J} \{\mathfrak{N}_{J_2} \otimes \mathfrak{M}_{J_1}\}_{JM} + \mathfrak{R}_{JM}^{J_1 J_2}. \quad (3.1.28)$$

In particular, from these equations one can see that an irreducible (rank-I) tensor product of two identical tensors $\mathfrak{M}_J$ is equal to zero, if $I = 2J - 1, 2J - 3, \dots$ and the tensor components commute

$$\{\mathfrak{M}_J \otimes \mathfrak{M}_J\}_I = 0 \quad (3.1.29)$$

3.1.8 Scalar Products of Irreducible Tensors

The scalar product of two irreducible tensors $\mathfrak{M}_J$ and $\mathfrak{N}_J$ of the same rank is defined as

$$(\mathfrak{M}_J \cdot \mathfrak{N}_J) = \sum_M (-1)^{-M} \mathfrak{M}_{JM} \mathfrak{N}_{J-M} = \sum_M \mathfrak{M}_{JM} \mathfrak{N}_{JM}^* = \sum_M \mathfrak{M}_{JM} \mathfrak{N}_J^M. \quad (3.1.30)$$

Similarly, the scalar product of two irreducible tensors $\tilde{\mathfrak{M}}_J$ and $\tilde{\mathfrak{N}}_J$ is given by

$$(\tilde{\mathfrak{M}}_J \cdot \tilde{\mathfrak{N}}_J) = \sum_M (-1)^{J-M} \tilde{\mathfrak{M}}_{JM} \tilde{\mathfrak{N}}_{J-M} = \sum_M \tilde{\mathfrak{M}}_{JM} \tilde{\mathfrak{N}}_{JM}^* = \sum_M \tilde{\mathfrak{M}}_{JM} \tilde{\mathfrak{N}}_J^M.$$

Note that

$$(\mathfrak{M}_J \cdot \mathfrak{N}_J) = (\tilde{\mathfrak{M}}_J \cdot \tilde{\mathfrak{N}}_J). \quad (3.1.31)$$

Scalar products differ from irreducible tensor products of zero rank only by some numerical factor. The latter tensor products read

$$\{\mathfrak{M}_J \otimes \mathfrak{N}_J\}_{00} = \sum_{M_1 M_2} C_{J M_1 J M_2}^{00} \mathfrak{M}_{J M_1} \mathfrak{N}_{J M_2} = \frac{1}{\sqrt{2J+1}} \sum_M (-1)^{J-M} \mathfrak{M}_{JM} \mathfrak{N}_{J-M}, \quad (3.1.32)$$

$$\{\tilde{\mathfrak{M}}_J \otimes \tilde{\mathfrak{N}}_J\}_{00} = \sum_{M_1 M_2} C_{J M_1 J M_2}^{00} \tilde{\mathfrak{M}}_{J M_1} \tilde{\mathfrak{N}}_{J M_2} = \frac{1}{\sqrt{2J+1}} \sum_M (-1)^{J-M} \tilde{\mathfrak{M}}_{JM} \tilde{\mathfrak{N}}_{J-M}. \quad (3.1.33)$$

Hence, the relations between scalar products and tensor products of zero rank are determined by

$$\begin{aligned} (\mathfrak{M}_J \cdot \mathfrak{N}_J) &= (-1)^{-J} \sqrt{2J+1} \{\mathfrak{M}_J \otimes \mathfrak{N}_J\}_{00}, \\ (\tilde{\mathfrak{M}}_J \cdot \tilde{\mathfrak{N}}_J) &= \sqrt{2J+1} \{\tilde{\mathfrak{M}}_J \otimes \tilde{\mathfrak{N}}_J\}_{00}. \end{aligned} \quad (3.1.34)$$

For the scalar product of double tensors we have

$$(\widetilde{W}_{J_1 J_2}(1, 2) \cdot \widetilde{U}_{J_1 J_2}(1, 2)) = \sum_{M_1 M_2} (-1)^{J_1 - M_1 + J_2 - M_2} \widetilde{W}_{J_1 M_1 J_2 M_2}(1, 2) \widetilde{U}_{J_1 - M_1 J_2 - M_2}(1, 2). \quad (3.1.35)$$

3.2 Relation between the irreducible tensor algebra and vector and tensor theory

3.2.1 Vectors and Irreducible Tensors

An arbitrary vector $\mathbf{A}$ is an irreducible tensor of rank one. Its spherical components may be treated as components of an irreducible tensor $\mathbf{A}_1$, of rank one, for which $A_1^\mu = (-1)^\mu A_{1-\mu}$.

$$A_{1\mu} \equiv A_\mu, \quad A^{1\mu} \equiv A^\mu. \quad (3.2.1)$$

A polar vector is a true tensor of rank one. Under inversion of the coordinate system its components change their sign. An axial vector is a pseudotensor of rank one. Under coordinate inversion its components remain unchanged.

Using two vectors $\mathbf{A} = \mathbf{A}_1$ and $\mathbf{B} = \mathbf{B}_1$ one can construct three irreducible tensor products of ranks 0, 1, 2,

$$\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{00}, \quad \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{1\mu}, \quad \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2\mu}.$$

If one of the vectors $\mathbf{A}$ or $\mathbf{B}$ is polar and the other axial, then $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{00}$ is a pseudoscalar, $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{1\mu}$ is a polar vector, $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2\mu}$ is a pseudotensor of rank two. If both vectors $\mathbf{A}$ and $\mathbf{B}$ are either polar or axial, then $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{00}$ is a scalar, $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{1\mu}$ is an axial vector and $\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2\mu}$ is a true tensor of rank two.

(a) Rank zero tensor and scalar product

The irreducible tensor product of rank zero differs from the scalar product of vectors $\mathbf{A}$ and $\mathbf{B}$ by a numerical factor:

$$\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{00} = -\frac{1}{\sqrt{3}}(\mathbf{A} \cdot \mathbf{B}) \quad (3.2.2)$$

The scalar product of irreducible tensors $\mathbf{A}_1$ and $\mathbf{B}_1$ introduced in Sec. 3.1.8 coincides with the scalar product of vectors,

$$(\mathbf{A}_1 \cdot \mathbf{B}_1) = (\mathbf{A} \cdot \mathbf{B}). \quad (3.2.3)$$

(b) Rank one tensor and vector product

The irreducible tensor product of rank one is related to the vector product of vectors $\mathbf{A}$ and $\mathbf{B}$ by

$$\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 = \frac{i}{\sqrt{2}}[\mathbf{A} \times \mathbf{B}]. \quad (3.2.4)$$

The components of the tensor product of rank one can be expressed in terms of products of spherical components of vectors $\mathbf{A}$ and $\mathbf{B}$ as

$$\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{1M} = \frac{i}{\sqrt{2}}[\mathbf{A} \times \mathbf{B}]_M = \sum_{\mu\nu} C_{1\mu 1\nu}^{1M} A_\mu B_\nu. \quad (3.2.5)$$

(c) Rank two tensor in terms of vector components

The components of a tensor product of rank two are also related to products of spherical components of $\mathbf{A}$ and $\mathbf{B}$:

$$\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2M} = \sum_{\mu,\nu} C_{1\mu 1\nu}^{2M} A_\mu B_\nu = \sqrt{\frac{3|M|-2}{14|M|-12}} \sum_{\substack{\mu+\nu=M \\ \mu \geq \nu}} (A_\mu B_\nu + A_\nu B_\mu). \quad (3.2.6)$$

In a more detailed form Eq. (3.2.6) may be written as

$$\begin{aligned} \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2+2} &= A_{+1} B_{+1}, \\ \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2+1} &= \frac{1}{\sqrt{2}} (A_{+1} B_0 + A_0 B_{+1}), \\ \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{20} &= \frac{1}{\sqrt{6}} (A_{+1} B_{-1} + 2A_0 B_0 + A_{-1} B_{+1}), \\ \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2-1} &= \frac{1}{\sqrt{2}} (A_{-1} B_0 + A_0 B_{-1}), \\ \{\mathbf{A}_1 \otimes \mathbf{B}_1\}_{2-2} &= A_{-1} B_{-1}. \end{aligned} \quad (3.2.7)$$

Given three commuting vectors $\mathbf{A}_1, \mathbf{B}_1, \mathbf{C}_1$ we can compose the following irreducible tensor products of ranks 0 and 1:

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_0 \otimes \mathbf{C}_1\}_1 = -\frac{1}{\sqrt{3}}(\mathbf{A} \cdot \mathbf{B}) \cdot \mathbf{C}, \quad (3.2.8)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \mathbf{C}_1\}_0 = -\frac{i}{\sqrt{6}}[\mathbf{A} \times \mathbf{B}] \cdot \mathbf{C}, \quad (3.2.9)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \mathbf{C}_1\}_1 = -\frac{1}{2} [[\mathbf{A} \times \mathbf{B}] \times \mathbf{C}] = \frac{1}{2} \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) - \frac{1}{2} \mathbf{B}(\mathbf{A} \cdot \mathbf{C}), \quad (3.2.10)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_2 \otimes \mathbf{C}_1\}_1 = \sqrt{\frac{3}{5}} \left\{ \frac{1}{3} \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) - \frac{1}{2} \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \frac{1}{2} \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) \right\}. \quad (3.2.11)$$

In addition to the foregoing formulas, there exist products which differ from Eqs. (3.2.8)-(3.2.11) by vector coupling schemes. The problems concerned with recoupling in tensor products are considered below (Sec. 3.3).

Given four commuting vectors $\mathbf{A}_1, \mathbf{B}_1, \mathbf{C}_1$ and $\mathbf{D}_1$ we can compose the following irreducible tensor products

of ranks 0 and 1 :

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_0 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_0\}_0 = \frac{1}{3}(\mathbf{A} \cdot \mathbf{B})(\mathbf{D} \cdot \mathbf{C}), \quad (3.2.12)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_0\}_1 = -\frac{i}{\sqrt{6}}[\mathbf{A} \times \mathbf{B}](\mathbf{C} \cdot \mathbf{D}), \quad (3.2.13)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_0 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_1\}_1 = -\frac{i}{\sqrt{6}}(\mathbf{A} \cdot \mathbf{B})[\mathbf{C} \times \mathbf{D}], \quad (3.2.14)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_1\}_0 = \frac{1}{2\sqrt{3}}\{(\mathbf{A} \cdot \mathbf{C})(\mathbf{B} \cdot \mathbf{D}) - (\mathbf{A} \cdot \mathbf{D})(\mathbf{B} \cdot \mathbf{C})\}, \quad (3.2.15)$$

$$\begin{aligned} \{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_1\}_1 &= -\frac{i}{2\sqrt{2}}\{\mathbf{C}(\mathbf{D} \cdot [\mathbf{A} \times \mathbf{B}]) - \mathbf{D}(\mathbf{C} \cdot [\mathbf{A} \times \mathbf{B}])\} \\ &= -\frac{i}{2\sqrt{2}}\{\mathbf{B}(\mathbf{A} \cdot [\mathbf{C} \times \mathbf{D}]) - \mathbf{A}(\mathbf{B} \cdot [\mathbf{C} \times \mathbf{D}])\}, \end{aligned} \quad (3.2.16)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_2 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_1\}_1 = \frac{i\sqrt{3}}{\sqrt{2} \cdot 5} \left\{ \frac{1}{3}(\mathbf{A} \cdot \mathbf{B})[\mathbf{C} \times \mathbf{D}] - \frac{1}{2}\mathbf{B}(\mathbf{D} \cdot [\mathbf{A} \times \mathbf{C}]) - \frac{1}{2}\mathbf{A}(\mathbf{D} \cdot [\mathbf{B} \times \mathbf{C}]) \right\}, \quad (3.2.17)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_1 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_2\}_1 = \frac{i\sqrt{3}}{\sqrt{2} \cdot 5} \left\{ \frac{1}{3}(\mathbf{C} \cdot \mathbf{D})[\mathbf{A} \times \mathbf{B}] - \frac{1}{2}\mathbf{C}(\mathbf{B} \cdot [\mathbf{D} \times \mathbf{A}]) - \frac{1}{2}\mathbf{D}(\mathbf{B} \cdot [\mathbf{C} \times \mathbf{A}]) \right\}, \quad (3.2.18)$$

$$\{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_2 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_2\}_0 = \frac{1}{\sqrt{5}} \left\{ \frac{1}{2}(\mathbf{A} \cdot \mathbf{C})(\mathbf{B} \cdot \mathbf{D}) - \frac{1}{3}(\mathbf{A} \cdot \mathbf{B})(\mathbf{C} \cdot \mathbf{D}) + \frac{1}{2}(\mathbf{A} \cdot \mathbf{D})(\mathbf{B} \cdot \mathbf{C}) \right\}, \quad (3.2.19)$$

$$\begin{aligned} \{\{\mathbf{A}_1 \otimes \mathbf{B}_1\}_2 \otimes \{\mathbf{C}_1 \otimes \mathbf{D}_1\}_2\}_1 &= \\ &= -\frac{i}{2\sqrt{2} \cdot 5} \{(\mathbf{A} \cdot \mathbf{C})[\mathbf{B} \times \mathbf{D}] + (\mathbf{A} \cdot \mathbf{D})[\mathbf{B} \times \mathbf{C}] + (\mathbf{B} \cdot \mathbf{C})[\mathbf{A} \times \mathbf{D}] + (\mathbf{B} \cdot \mathbf{D})[\mathbf{A} \times \mathbf{C}]\}. \end{aligned} \quad (3.2.20)$$

The change of coupling schemes in products of four operators is discussed in Sec. 3.3.

Products constructed from the components of identical vectors have the following property

$$\left\{ \dots \left\{ \{\mathbf{A}_1 \otimes \mathbf{A}_1\}_{l_2} \otimes \mathbf{A}_1 \right\}_{l_3} \dots \otimes \mathbf{A}_1 \right\}_{l_n} = \left\{ \mathbf{A}_1 \otimes \dots \left\{ \mathbf{A}_1 \otimes \{\mathbf{A}_1 \otimes \mathbf{A}_1\}_{l_2} \right\}_{l_3} \dots \right\}_{l_n}. \quad (3.2.21)$$

If a vector $\mathbf{A}$ does not contain any spin variable and differential operator one can express such products in terms of spherical harmonics (Chap. 5), i.e.,

$$\left\{ \dots \left\{ \{\mathbf{A}_1 \otimes \mathbf{A}_1\}_{l_2} \otimes \mathbf{A}_1 \right\}_{l_3} \dots \otimes \mathbf{A}_1 \right\}_{l_n m_n} = \sqrt{\frac{4\pi}{2l_n + 1}} |\mathbf{A}|^n Y_{l_n m_n}(\vartheta, \varphi) \prod_{i=2}^n C_{10 l_{i-1} 0}^{l_i 0}, \quad (3.2.22)$$

where ϑ, φ are the polar angles of the vector $\mathbf{A}$ and $l_1 = 1$. In particular, if $l_2 = 2, l_3 = 3, \dots, l_n = n$, we have

$$\left\{ \dots \left\{ \{\mathbf{A}_1 \otimes \mathbf{A}_1\}_2 \otimes \mathbf{A}_1 \right\}_3 \dots \otimes \mathbf{A}_1 \right\}_{nm} = \sqrt{\frac{4\pi n!}{(2n+1)!!}} |\mathbf{A}|^n Y_{nm}(\vartheta, \varphi). \quad (3.2.23)$$

3.2.2 Cartesian Tensors of Second and Third Ranks

An arbitrary cartesian tensor of second rank $T_{ik}(i, k = x, y, z)$ is, generally, reducible and may be decomposed into three irreducible parts:

- (a) a tensor which is proportional to the unit tensor, $E_{ik} = E\delta_{ik}$;
- (b) an antisymmetric tensor $A_{ik} = -A_{ki}$.
- (c) a symmetric traceless tensor $S_{ik} = S_{ki}, \sum_i S_{ii} = 0$.

Thus,

$$T_{ik} = E\delta_{ik} + A_{ik} + S_{ik}, \quad (3.2.24)$$

$$E = \frac{1}{3} \text{Tr}(T_{ik}) = \frac{1}{3} \sum_i T_{ii}, \quad (3.2.25)$$

$$A_{ik} = \frac{1}{2} (T_{ik} - T_{ki}), \quad (3.2.26)$$

$$S_{ik} = \frac{1}{2} \left(T_{ik} + T_{ki} - \frac{2}{3} \delta_{ik} \sum_l T_{ll} \right). \quad (3.2.27)$$

The value E is invariant under rotations of coordinate systems. It is an irreducible tensor of zero rank

$$\mathfrak{I}_{00} = E. \quad (3.2.28)$$

The antisymmetric tensor A_{ik} is equivalent to the axial vector

$$A_{ik} = \varepsilon_{ikl} \mathfrak{U}_l, \quad \mathfrak{U}_i = \frac{1}{2} \sum_{kl} \varepsilon_{ikl} A_{kl}. \quad (3.2.29)$$

Using the components A_{ik} , one can form an irreducible pseudotensor of rank one:

$$\begin{aligned} \mathfrak{I}_{10} &= \mathfrak{U}_z = A_{xy}, \\ \mathfrak{I}_{1\pm 1} &= \mp \frac{1}{\sqrt{2}} (\mathfrak{U}_x \pm i\mathfrak{U}_y) = \mp \frac{1}{\sqrt{2}} (A_{yz} \pm iA_{zx}). \end{aligned} \quad (3.2.30)$$

Using the components of the symmetric traceless tensor S_{ij} , we may construct an irreducible tensor of second rank.

$$\begin{aligned} \mathfrak{I}_{20} &= S_{zz}, \\ \mathfrak{I}_{2\pm 1} &= \mp \sqrt{\frac{2}{3}} (S_{xz} \pm iS_{yz}), \\ \mathfrak{I}_{2\pm 2} &= \sqrt{\frac{1}{6}} (S_{xx} - S_{yy} \pm 2iS_{xy}). \end{aligned} \quad (3.2.31)$$

Thus, from nine cartesian components T_{ik} of a tensor of second rank we can compose one tensor of zero rank (scalar which has one component), one pseudotensor of the first rank (three components) and one irreducible tensor of second rank (five components).

From the 27 cartesian components T_{ikl} of a tensor of third rank we can compose one pseudotensor of zero rank $\mathfrak{I}_{00}$ (scalar, which has one component), three tensors of first rank $\mathfrak{I}_{1\mu}$ ($3 \times 3 = 9$ components), two irreducible tensors of second rank $\mathfrak{I}_{2\mu}$ ($2 \times 5 = 10$ components) and one irreducible tensor of third rank $\mathfrak{I}_{3\mu}$ (seven components). In this case construction of the tensors of first and second ranks is not unique.

3.2.3 Differential Operations as Irreducible Tensor Products

Differential operations on scalars and vectors (see Sec. 1.3) can be written in the form of irreducible tensor products of the operator ∇ and corresponding scalars and vectors,

$$\text{grad } \Phi = \{\nabla_1 \otimes \Phi\}_1, \quad (3.2.32)$$

$$\text{div } \mathbf{A} = -\sqrt{3} \{\nabla_1 \otimes \mathbf{A}_1\}_0, \quad (3.2.33)$$

$$\text{curl } \mathbf{A} = -i\sqrt{2} \{\nabla_1 \otimes \mathbf{A}_1\}_1, \quad (3.2.34)$$

$$\Delta = \nabla^2 = -\sqrt{3} \{\nabla_1 \otimes \nabla_1\}_0, \quad (3.2.35)$$

$$\text{grad div } \mathbf{A} = -\sqrt{3} \{\nabla_1 \otimes \{\nabla_1 \otimes \mathbf{A}_1\}_0\}_1, \quad (3.2.36)$$

$$\text{curl curl } \mathbf{A} = -2 \{\nabla_1 \otimes \{\nabla_1 \otimes \mathbf{A}_1\}_1\}_1, \quad (3.2.37)$$

$$\text{div grad } \Phi = -\sqrt{3} \{\nabla_1 \otimes \{\nabla_1 \otimes \Phi\}_1\}_0 = -\sqrt{3} \{\nabla_1 \otimes \nabla_1\}_0 \Phi, \quad (3.2.38)$$

$$\text{curl grad } \Phi = -i\sqrt{2} \{\nabla_1 \otimes \{\nabla_1 \otimes \Phi\}_1\}_1 = 0, \quad (3.2.39)$$

$$\text{div curl } \mathbf{A} = i\sqrt{6} \{\nabla_1 \otimes \{\nabla_1 \otimes \mathbf{A}_1\}_1\}_0 = 0. \quad (3.2.40)$$

As follows from Sec. 1.3.1 the operator ∇ is given by

$$\nabla = \mathbf{n} \frac{\partial}{\partial r} - \frac{i}{r} [\mathbf{n} \times \hat{\mathbf{L}}] \quad (3.2.41)$$

In spherical component form it is written as

$$\nabla_\mu = \sqrt{\frac{4\pi}{3}} \left(Y_{1\mu} \frac{\partial}{\partial r} - \frac{\sqrt{2}}{r} \left\{ \mathbf{Y}_1 \otimes \hat{\mathbf{L}}_1 \right\}_{1\mu} \right). \quad (3.2.42)$$

Here $\mathbf{Y}_1$ is an irreducible tensor, whose components are spherical harmonics $Y_{1\mu}(\mathbf{n})$ (Chap. 5); $\hat{\mathbf{L}}_1 = \hat{\mathbf{L}}$ is the orbital angular momentum operator (see Sec. 2.2).

When expanding any scalar function $\Phi(\mathbf{r})$ in a Taylor series, one will deal with arbitrarily large powers of operator ∇ :

$$\begin{aligned}\Phi(\mathbf{r} + \delta\mathbf{r}) &= e^{(\delta\mathbf{r} \cdot \nabla)} \Phi(\mathbf{r}) = \Phi(\mathbf{r}) + (\delta\mathbf{r} \cdot \nabla) \Phi(\mathbf{r}) + \frac{1}{2!} (\delta\mathbf{r} \cdot \nabla)^2 \Phi(\mathbf{r}) + \dots \\ &= \Phi(\mathbf{r}) + \delta r (\mathbf{u} \cdot \nabla) \Phi(\mathbf{r}) + \frac{1}{2!} (\delta r)^2 (\mathbf{u} \cdot \nabla)^2 \Phi(\mathbf{r}) + \dots\end{aligned}\quad (3.2.43)$$

Here $\mathbf{u} = \delta\mathbf{r}/|\delta\mathbf{r}|$ and $(\mathbf{u} \cdot \nabla) = d/ds$ is the operator of directional differentiation in the direction $\mathbf{u}$ (see Sec. 1.3). The operator $(\mathbf{u} \cdot \nabla)^n \equiv d^n/ds^n$ can be written in the form of tensor product

$$(u \cdot \nabla)^n = (-1)^n \sum_{l_2, l_3, \dots, l_n} (-1)^{l_n} \left(\left\{ \dots \left\{ \{u_1 \otimes u_1\}_{l_2} \otimes u_1 \right\}_{l_3} \dots \otimes u_1 \right\}_{l_n} \cdot \left\{ \dots \left\{ \{\nabla_1 \otimes \nabla_1\}_{l_2} \otimes \nabla_1 \right\}_{l_3} \dots \otimes \nabla_1 \right\}_{l_n} \right). \quad (3.2.44)$$

Using Eqs. (3.2.22) and (3.2.23) one can express multiple tensor products of the unit vector u which enters Eq. (3.2.44) in terms of spherical harmonics.

3.3 Recoupling in irreducible tensor products

The irreducible tensor product of two irreducible tensors is defined in Sec. 3.1.7. By making use of the same relations one can form irreducible tensor products of three and more irreducible tensors. However, in these cases different orders and different coupling schemes of tensors in products are possible.

The recoupling tensors without change of their order may be carried out by some real (for a given definition of the vector addition coefficients) and orthogonal matrix which performs the direct and inverse transformation. When the tensor order has to be changed one should take account of the commutation rules (Eqs. (3.1.24)-(3.1.28)). We will use the notation

$$\Pi_{abc\dots d} = [(2a+1)(2b+1)(2c+1)\dots(2d+1)]^{\frac{1}{2}}$$

3.3.1 Relations Valid for Commuting as well as Non-Commuting Tensors

For recoupling tensors without changing their order in irreducible tensor products of three and four tensors, one has the following relations:

$$\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \mathbf{R}_d\}_f = (-1)^{a+b+f+d} \sum_h \Pi_{hc} \begin{Bmatrix} a & b & c \\ d & f & h \end{Bmatrix} \{\mathbf{P}_a \otimes \{\mathbf{Q}_b \otimes \mathbf{R}_d\}_h\}_f, \quad (3.3.1)$$

$$(\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \cdot \mathbf{R}_c) = (-1)^{-c+a} \frac{\Pi_c}{\Pi_a} (\mathbf{P}_a \cdot \{\mathbf{Q}_b \otimes \mathbf{R}_c\}_a), \quad (3.3.2)$$

$$\begin{aligned}\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \{\mathbf{R}_d \otimes \mathbf{S}_e\}_f\}_k &= (-1)^{d+c+e+k} \sum_h \Pi_{hf} \begin{Bmatrix} d & c & h \\ k & e & f \end{Bmatrix} \{\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \mathbf{R}_d\}_h \otimes \mathbf{S}_e\}_k \\ &= (-1)^{a+f+b+k} \sum_h \Pi_{hc} \begin{Bmatrix} a & b & c \\ f & k & h \end{Bmatrix} \left\{ \mathbf{P}_a \otimes \left\{ \mathbf{Q}_b \otimes \{\mathbf{R}_d \otimes \mathbf{S}_e\}_f \right\}_h \right\}_k.\end{aligned}\quad (3.3.3)$$

$$\begin{aligned}(\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \cdot \{\mathbf{R}_d \otimes \mathbf{S}_e\}_c) &= \frac{(-1)^{-c+e} \Pi_c}{\Pi_e} (\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \mathbf{R}_d\}_e \cdot \mathbf{S}_e) \\ &= (-1)^{-c+a} \frac{\Pi_c}{\Pi_a} \left(\mathbf{P}_a \cdot \left\{ \mathbf{Q}_b \otimes \{\mathbf{R}_d \otimes \mathbf{S}_e\}_f \right\}_a \right),\end{aligned}\quad (3.3.4)$$

$$\{\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \mathbf{R}_d\}_h \otimes \mathbf{S}_e\}_k = \sum_{f,q} (-1)^{a+f+b-d-c-e} \Pi_{cqhf} \begin{Bmatrix} a & b & c \\ f & k & q \end{Bmatrix} \begin{Bmatrix} d & e & f \\ k & c & h \end{Bmatrix} \left\{ \mathbf{P}_a \otimes \left\{ \mathbf{Q}_b \otimes \{\mathbf{R}_d \otimes \mathbf{S}_e\}_q \right\}_c \right\}_k, \quad (3.3.5)$$

$$(\{\{\mathbf{P}_a \otimes \mathbf{Q}_b\}_c \otimes \mathbf{R}_d\}_e \cdot \mathbf{S}_e) = (-1)^{-c+a} \frac{\Pi_e}{\Pi_a} \left(\mathbf{P}_a \cdot \left\{ \mathbf{Q}_b \otimes \{\mathbf{R}_d \otimes \mathbf{S}_e\}_f \right\}_a \right). \quad (3.3.6)$$

3.3.2 Relations for Commuting Tensors

To change the coupling scheme for irreducible products of three and four commuting tensors one can use the following relations

$$\begin{aligned} \left\{ \mathbf{P}_a \otimes \{ \mathbf{Q}_b \otimes \mathbf{R}_d \}_f \right\}_e &= (-1)^{b+d-f} \left\{ \mathbf{P}_a \otimes \{ \mathbf{R}_d \otimes \mathbf{Q}_b \}_f \right\}_e = (-1)^{a+f-e} \left\{ \{ \mathbf{Q}_b \otimes \mathbf{R}_d \}_f \otimes \mathbf{P}_a \right\}_e \\ &= (-1)^{a+b+d-e} \left\{ \{ \mathbf{R}_d \otimes \mathbf{Q}_b \}_f \otimes \mathbf{P}_a \right\}_e, \end{aligned} \quad (3.3.7)$$

$$\left\{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \mathbf{R}_d \right\}_f = (-1)^{c+d+f} \sum_h \Pi_{ch} \begin{Bmatrix} a & b & c \\ f & d & h \end{Bmatrix} \left\{ \mathbf{Q}_b \otimes \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_h \right\}_f, \quad (3.3.8)$$

$$(\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \mathbf{R}_c) = (-1)^{-a} \frac{\Pi_c}{\Pi_b} (\mathbf{Q}_b \cdot \{ \mathbf{P}_a \otimes \mathbf{R}_c \}_b). \quad (3.3.9)$$

$$\begin{aligned} \left\{ \mathbf{P}_a \otimes \left\{ \mathbf{Q}_b \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \right\}_h \right\}_k &= (-1)^{d+e-f} \left\{ \mathbf{P}_a \otimes \left\{ \mathbf{Q}_b \otimes \{ \mathbf{S}_e \otimes \mathbf{R}_d \}_f \right\}_h \right\}_k \\ &= (-1)^{d+b+e-h} \left\{ \mathbf{P}_a \otimes \left\{ \{ \mathbf{S}_e \otimes \mathbf{R}_d \}_f \otimes \mathbf{Q}_b \right\}_h \right\}_k = (-1)^{b+f-h} \left\{ \mathbf{P}_a \otimes \left\{ \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \otimes \mathbf{Q}_b \right\}_h \right\}_k \\ &= (-1)^{a+h-k} \left\{ \left\{ \mathbf{Q}_b \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \right\}_h \otimes \mathbf{P}_a \right\}_k = (-1)^{d+e+a+h-f-k} \left\{ \left\{ \mathbf{Q}_b \otimes \{ \mathbf{S}_e \otimes \mathbf{R}_d \}_f \right\}_h \otimes \mathbf{P}_a \right\}_k \\ &= (-1)^{a+b+d+e-h} \left\{ \left\{ \{ \mathbf{S}_e \otimes \mathbf{R}_d \}_f \otimes \mathbf{Q}_b \right\}_h \otimes \mathbf{P}_a \right\}_k = (-1)^{a+b+f-k} \left\{ \left\{ \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \otimes \mathbf{Q}_b \right\}_h \otimes \mathbf{P}_a \right\}_k, \end{aligned} \quad (3.3.10)$$

$$\left\{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \right\}_k = \sum_{gh} \Pi_{cfgh} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & k \end{Bmatrix} \left\{ \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_g \otimes \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_h \right\}_k. \quad (3.3.11)$$

$$\left\{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \right\}_k = \sum_{gh} (-1)^{h+b-k-e} \Pi_{cfgh} \begin{Bmatrix} a & b & c \\ g & d & h \end{Bmatrix} \begin{Bmatrix} d & e & f \\ k & c & g \end{Bmatrix} \left\{ \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_h \otimes \mathbf{Q}_b \right\}_g \otimes \mathbf{S}_e \right\}_k. \quad (3.3.12)$$

$$(\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_c) = (-1)^{2a+b-d} \sum_g \Pi_c^2 \begin{Bmatrix} a & b & c \\ e & d & g \end{Bmatrix} \left(\{ \mathbf{P}_a \otimes \mathbf{R}_d \}_g \cdot \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_g \right). \quad (3.3.13)$$

$$(\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_c) = \sum_h (-1)^{e+b+d+h} \frac{\Pi_{cch}}{\Pi_e} \begin{Bmatrix} a & b & c \\ e & d & h \end{Bmatrix} \left(\{ \mathbf{P}_a \otimes \mathbf{R}_d \}_h \otimes \mathbf{Q}_b \right)_e \cdot \mathbf{S}_e. \quad (3.3.14)$$

Equation (3.1.27) and Eqs. (3.3.11)-(3.3.14) permit us to obtain all other permutations. The product $\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_b$ is orthogonal to tensor $\mathbf{Q}_b$, if $a = 2b - 1, 2b - 2, \dots$

$$(\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_b \cdot \mathbf{Q}_b) = 0. \quad (3.3.15)$$

3.3.3 Relations for Non-Commuting Tensors

A change of the coupling scheme for irreducible tensor products of three and four non-commuting tensors gives

$$\{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \mathbf{R}_d \}_f = (-1)^{f+d+c} \sum_h \Pi_{ch} \begin{Bmatrix} a & b & c \\ f & d & h \end{Bmatrix} \{ \mathbf{Q}_b \otimes \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_h \}_f + \{ \mathfrak{R}_c^{ab} \otimes \mathbf{R}_d \}_f, \quad (3.3.16)$$

$$(\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \mathbf{R}_c) = (-1)^{-a} \frac{\Pi_c}{\Pi_b} (\mathbf{Q}_b \cdot \{ \mathbf{P}_a \otimes \mathbf{R}_c \}_b) + (\mathfrak{R}_c^{ab} \cdot \mathbf{R}_c), \quad (3.3.17)$$

$$\begin{aligned} \{ \mathbf{P}_a \otimes \{ \mathbf{Q}_b \otimes \mathbf{R}_d \}_h \}_f &= \sum_{cg} (-1)^{a+b-c} \Pi_{cgh} \begin{Bmatrix} a & b & c \\ f & d & g \end{Bmatrix} \begin{Bmatrix} a & b & c \\ d & f & h \end{Bmatrix} \{ \mathbf{Q}_b \otimes \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_g \}_f \\ &\quad + \sum_c (-1)^{a+b+f+d} \Pi_{ch} \begin{Bmatrix} a & b & c \\ d & f & h \end{Bmatrix} \{ \mathfrak{R}_c^{ab} \otimes \mathbf{R}_d \}_f, \end{aligned} \quad (3.3.18)$$

$$(\mathbf{P}_a \cdot \{ \mathbf{Q}_b \otimes \mathbf{R}_d \}_a) = (-1)^{2b-d} \frac{\Pi_a}{\Pi_b} (\mathbf{Q}_b \cdot \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_b) + (-1)^{-a+d} \frac{\Pi_a}{\Pi_d} (\mathfrak{R}_d^{ab} \cdot \mathbf{R}_d), \quad (3.3.19)$$

$$\begin{aligned} \{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \}_k &= \sum_{gr} \Pi_{cgr} \begin{Bmatrix} a & b & c \\ d & e & f \\ r & g & k \end{Bmatrix} \{ \{ \mathbf{P}_a \otimes \mathbf{R}_d \}_r \otimes \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_g \}_k \\ &\quad + \sum_{hq} (-1)^{a+k+d+e-f-h} \Pi_{chqf} \begin{Bmatrix} a & b & c \\ f & k & h \end{Bmatrix} \begin{Bmatrix} d & e & f \\ h & b & q \end{Bmatrix} \{ \mathbf{P}_a \otimes \{ \mathfrak{R}_q^{bd} \otimes \mathbf{S}_e \}_h \}_k, \end{aligned} \quad (3.3.20)$$

$$\begin{aligned} (\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_c) &= \sum_g (-1)^{d+b+2g} \Pi_c^2 \begin{Bmatrix} a & b & c \\ e & d & g \end{Bmatrix} \left(\{ \mathbf{P}_a \otimes \mathbf{R}_d \}_g \cdot \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_g \right) \\ &\quad + \sum_q (-1)^{d+e+c-b} \frac{\Pi_{ccq}}{\Pi_a} \begin{Bmatrix} a & b & c \\ d & e & q \end{Bmatrix} \left(\mathbf{P}_a \cdot \{ \mathfrak{R}_q^{bd} \otimes \mathbf{S}_e \}_a \right), \end{aligned} \quad (3.3.21)$$

$$\begin{aligned} \{ \{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \otimes \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_f \}_k &= \sum_{hg} (-1)^{a-b+k-d-g} \Pi_{hcfg} \begin{Bmatrix} a & b & c \\ f & k & h \end{Bmatrix} \begin{Bmatrix} b & e & g \\ d & h & f \end{Bmatrix} \{ \mathbf{P}_a \otimes \{ \mathbf{R}_d \otimes \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_g \}_h \}_k \\ &\quad + \sum_{hq} (-1)^{a+k+d+e-f-h} \Pi_{hcfq} \begin{Bmatrix} a & b & c \\ f & k & h \end{Bmatrix} \begin{Bmatrix} d & e & f \\ h & b & q \end{Bmatrix} \{ \mathbf{P}_a \otimes \{ \mathfrak{R}_q^{bd} \otimes \mathbf{S}_e \}_h \}_k, \end{aligned} \quad (3.3.22)$$

$$\begin{aligned} (\{ \mathbf{P}_a \otimes \mathbf{Q}_b \}_c \cdot \{ \mathbf{R}_d \otimes \mathbf{S}_e \}_c) &= \sum_g (-1)^{a+d+g+b} \frac{\Pi_{ccg}}{\Pi_a} \begin{Bmatrix} b & e & g \\ d & a & c \end{Bmatrix} \left(\mathbf{P}_a \cdot \{ \mathbf{R}_d \otimes \{ \mathbf{Q}_b \otimes \mathbf{S}_e \}_g \}_a \right) \\ &\quad + \sum_q (-1)^{d+e+c-b} \frac{\Pi_{ccq}}{\Pi_a} \begin{Bmatrix} a & b & c \\ d & e & q \end{Bmatrix} \left(\mathbf{P}_a \cdot \{ \mathfrak{R}_q^{bd} \otimes \mathbf{S}_e \}_a \right). \end{aligned} \quad (3.3.23)$$

The definition of the commutator $\mathfrak{R}_q^{bd}$ is given in Eq. (3.1.26). By making use of the foregoing formulas, one can obtain all other relations for irreducible products of non-commuting tensors.

Chapter 4

Wigner D-functions

4.1 Definition of $D_{MM'}^J(\alpha, \beta, \gamma)$

- (a) The Wigner D -functions $D_{MM'}^J(\alpha, \beta, \gamma)$ may be defined as the matrix elements of the rotation operator $\hat{D}(\alpha, \beta, \gamma)$ in the JM-representation. The arguments α, β, γ are the Euler angles which specify the rotation

$$\langle JM | \hat{D}(\alpha, \beta, \gamma) | J' M' \rangle = \delta_{JJ'} D_{MM'}^J(\alpha, \beta, \gamma). \quad (4.1.1)$$

The D -functions realize transformations of covariant components of any irreducible tensor of rank J (e.g., the wave function Ψ_{JM} of a quantum mechanical system with angular momentum J and its projection M) under coordinate rotations.

$$\begin{aligned} \Psi_{JM'}(\vartheta', \varphi', \sigma') &= \sum_{M=-J}^J \Psi_{JM}(\vartheta, \varphi, \sigma) D_{MM'}^J(\alpha, \beta, \gamma), \\ \Psi_{JM'}^*(\vartheta', \varphi', \sigma') &= \sum_{M=-J}^J \Psi_{JM}^*(\vartheta, \varphi, \sigma) D_{MM'}^{J*}(\alpha, \beta, \gamma). \end{aligned} \quad (4.1.2)$$

Here ϑ, φ and ϑ', φ' are polar angles in the initial and rotated coordinate systems, S and S' , respectively. The angles ϑ, φ and ϑ', φ' are related by Eqs. (1.4.2) and (1.4.3). Similarly, σ and σ' are spin variables in the initial and new systems.

The inverse transformation $S' \rightarrow S$ is performed by the inverse matrix $\left[\hat{D}^{-1}(\alpha, \beta, \gamma) \right]_{MM'}^J$. Owing to the unitarity of the rotation operator,

$$\hat{D}^{-1}(\alpha, \beta, \gamma) = \hat{D}^+(\alpha, \beta, \gamma). \quad (4.1.3)$$

the elements of inverse matrix are given by

$$\left| \hat{D}^{-1}(\alpha, \beta, \gamma) \right|_{MM'}^J = D_{M'M}^{J*}(\alpha, \beta, \gamma). \quad (4.1.4)$$

Hence, under the inverse rotation, $S' \rightarrow S$, wave functions transform as

$$\begin{aligned} \Psi_{JM}(\vartheta, \varphi, \sigma) &= \sum_{M'=-J}^J D_{MM'}^{J*}(\alpha, \beta, \gamma) \Psi_{JM'}(\vartheta', \varphi', \sigma'), \\ \Psi_{JM}^*(\vartheta, \varphi, \sigma) &= \sum_{M'=-J}^J D_{MM'}^J(\alpha, \beta, \gamma) \Psi_{JM'}^*(\vartheta', \varphi', \sigma'). \end{aligned} \quad (4.1.5)$$

The unitarity condition for the Wigner D -functions may be written as

$$\begin{aligned} \sum_{M=-J}^J D_{MM'}^J(\alpha, \beta, \gamma) D_{M\tilde{M}'}^{J*}(\alpha, \beta, \gamma) &= \delta_{M'\tilde{M}'}, \\ \sum_{M'=-J}^J D_{MM'}^{J*}(\alpha, \beta, \gamma) D_{M\tilde{M}'}^J(\alpha, \beta, \gamma) &= \delta_{M\tilde{M}}. \end{aligned} \quad (4.1.6)$$

The matrix $D_{MM'}^J(\alpha, \beta, \gamma)$ is unimodular, i.e.,

$$\det \| D_{MM'}^J(\alpha, \beta, \gamma) \| = +1. \quad (4.1.7)$$

Table 4.1. Effect of the Operator $\hat{D}(\alpha, \beta, \gamma)$

Transformation	Angles, Axes and Sequence of Rotations		
	I	II	III
<i>Passive</i>			
Rotation of coordinate system	$\alpha(z)$	$\beta(y_1)$	$\gamma(z')$
without rotation of physical body	$\gamma(z)$	$\beta(y)$	$\alpha(z)$
<i>Active</i>			
Rotation of physical body	$-\alpha(z)$	$-\beta(y)$	$-\gamma(z)$
without rotation of coordinate system	$-\gamma(z)$	$-\beta(y_1)$	$-\alpha(z')$

- (b) A set of $(2J + 1)$ functions Ψ_{JM} with different M 's constitute a basis for expansion of an arbitrary function Ψ_J with the same J :

$$\Psi_J(\vartheta, \varphi) = \sum_{M=-J}^J C_J^M \Psi_{JM}(\vartheta, \varphi) = (\mathbf{C}_J \cdot \boldsymbol{\Psi}_J). \quad (4.1.8)$$

The expansion coefficients C_J^M are contravariant components of some irreducible tensor of rank J . Under rotations the quantities C_J^M transform by means of functions $D_{MM'}^{J*}(\alpha, \beta, \gamma)$.

The effect of the operator $\hat{D}(\alpha, \beta, \gamma)$ on Ψ_J may be interpreted in two different ways:

1. as a rotation of the coordinate system without rotation of the physical body (this is the passive interpretation; $\hat{D}$ acts on the basis functions Ψ_{JM} while C_J^M remain unchanged);
2. as a rotation of the physical body without rotation of coordinate system (active interpretation; $\hat{D}$ acts on C_J^M but does not affect Ψ_{JM}).

Any rotation of a physical body in combination with the same rotation of coordinate system leaves the wave function Ψ_J unchanged:

$$\{\hat{D}(\alpha, \beta, \gamma)\}_{\text{phys. body}} \cdot \{\hat{D}(\alpha, \beta, \gamma)\}_{\text{coord. system}} = 1, \quad (4.1.9)$$

i.e.,

$$\left\{ \hat{D}_{MM'}^J(\alpha, \beta, \gamma) \right\}_{\text{phys. body}} = \left\{ \left[\hat{D}^{-1}(\alpha, \beta, \gamma) \right]_{MM'}^J \right\}_{\text{coord. system}}. \quad (4.1.10)$$

Moreover, a rotation of the coordinate system (or physical body) described by the Euler angles α, β, γ may also be realized in two ways:

1. by rotating about the initial axes (case B in Sec. 1.4.1), or
2. by rotating about the new (turned) axes (case A in Sec. 1.4.1).

Thus, any transformation of wave functions described by Eq. (4.1.2) can be treated in four different ways (Table 4.1).

The Wigner D -functions are complex. They depend on three real arguments α, β, γ and are defined in the domain

$$0 \leq \alpha < 2\pi, \quad 0 \leq \beta \leq \pi, \quad 0 \leq \gamma < 2\pi. \quad (4.1.11)$$

These functions, as well as their derivatives, are single-valued, finite and continuous. Sometimes it is convenient to change the domain (4.1.11). This can be done using the symmetries of $D_{MM'}^J(\alpha, \beta, \gamma)$ (Sec. 4.4). For example, the matrix of the inverse rotation satisfies the equation

$$\left[D^{-1}(\alpha, \beta, \gamma) \right]_{MM'}^J = D_{MM'}^J(\pi - \gamma, \beta, -\pi - \alpha) = D_{MM'}^J(-\gamma, -\beta, -\alpha). \quad (4.1.12)$$

This means that the inverse transformation $S' \rightarrow S$ may be realized by the Euler angles

$$\alpha' = \pi - \gamma, \quad \beta' = \beta, \quad \gamma' = -\pi - \alpha, \quad (4.1.13)$$

as well as by

$$\alpha' = -\gamma, \quad \beta' = -\beta, \quad \gamma' = -\alpha. \quad (4.1.14)$$

4.2 Differential equations for $D_{MM'}^J(\alpha, \beta, \gamma)$

- (a) The Wigner D -functions represent wave functions of a rigid symmetric top. They are eigenfunctions of three operators

$$\hat{J}_z = -i \frac{\partial}{\partial \alpha}, \quad \hat{J}_{z'} = -i \frac{\partial}{\partial \gamma}, \quad \hat{J}^2 = - \left[\frac{\partial^2}{\partial \beta^2} + \cot \beta \frac{\partial}{\partial \beta} + \frac{1}{\sin^2 \beta} \left(\frac{\partial^2}{\partial \alpha^2} - 2 \cos \beta \frac{\partial^2}{\partial \alpha \partial \gamma} + \frac{\partial^2}{\partial \gamma^2} \right) \right], \quad (4.2.1)$$

where $\hat{J}$ is the operator of angular momentum of the top; $\hat{J}_{z'}$ and $\hat{J}_z$ are projections of $\hat{J}$ onto the z -axis of the rotating (body-fixed) and non-rotating (lab-fixed) coordinate systems, respectively. The eigenvalues of the operators (4.2.1) are defined by the equations

$$\begin{aligned} \hat{J}_x D_{MM'}^J(\alpha, \beta, \gamma) &= -M D_{MM'}^J(\alpha, \beta, \gamma), \\ \hat{J}_{z'} D_{MM'}^J(\alpha, \beta, \gamma) &= -M' D_{MM'}^J(\alpha, \beta, \gamma), \\ \hat{J}^2 D_{MM'}^J(\alpha, \beta, \gamma) &= \left\{ -\frac{1}{\sin \beta} \cdot \frac{\partial}{\partial \beta} \left(\sin \beta \frac{\partial}{\partial \beta} \right) + \frac{M^2 - 2MM' \cos \beta + M'^2}{\sin^2 \beta} \right\} D_{MM'}^J(\alpha, \beta, \gamma) \\ &= J(J+1) D_{MM'}^J(\alpha, \beta, \gamma). \end{aligned} \quad (4.2.2)$$

Periodicity conditions for $D_{MM'}^J(\alpha, \beta, \gamma)$ are as follows

$$\begin{aligned} D_{MM'}^J(\alpha \pm 2k\pi, \beta, \gamma) &= D_{MM'}^J(\alpha, \beta, \gamma), \\ D_{MM'}^J(\alpha, \beta, \gamma \pm 2k\pi) &= D_{MM'}^J(\alpha, \beta, \gamma), \\ D_{MM'}^J(\alpha, \beta \pm 2k\pi, \gamma) &= D_{MM'}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.2.3)$$

where

$$\begin{aligned} k &= 0, 1, 2, \dots & \text{if } J \text{ is integer,} \\ k &= 0, 2, 4, \dots & \text{if } J \text{ is half-integer.} \end{aligned}$$

- (b) The functions $D_{MM'}^J(\alpha, \beta, \gamma)$ can also be defined as solutions of the differential equations

$$\begin{aligned} [\hat{J}_\nu, D_{MM'}^J(\alpha, \beta, \gamma)] &= \hat{J}_\nu D_{MM'}^J(\alpha, \beta, \gamma) = (-1)^{1+\nu} \sqrt{J(J+1)} C_{JM-\nu}^{JM-\nu} D_{M-\nu M'}^J(\alpha, \beta, \gamma) \\ &= \begin{cases} -M D_{MM'}^J(\alpha, \beta, \gamma), & \nu = 0, \\ \pm \sqrt{\frac{J(J+1)-M(M\mp 1)}{2}} D_{M\mp 1 M'}^J(\alpha, \beta, \gamma), & \nu = \pm 1, \end{cases} \end{aligned} \quad (4.2.4)$$

$$\begin{aligned} [\hat{J}^\nu, D_{MM'}^J(\alpha, \beta, \gamma)] &= \hat{J}^\nu D_{MM'}^J(\alpha, \beta, \gamma) = -\sqrt{J(J+1)} C_{JM'+\nu}^{JM'+\nu} D_{MM'+\nu}^J(\alpha, \beta, \gamma) \\ &= \begin{cases} -M' D_{MM'}^J(\alpha, \beta, \gamma), & \nu = 0, \\ \pm \sqrt{\frac{J(J+1)-M'(M'\pm 1)}{2}} D_{MM'\pm 1}^J(\alpha, \beta, \gamma), & \nu = \pm 1, \end{cases} \end{aligned} \quad (4.2.5)$$

Here $\hat{J}_\nu$ is a covariant spherical component of $\hat{\mathbf{J}}$ in the non-rotating (lab-fixed) system

$$\begin{aligned} \hat{J}_{\pm 1} &= \frac{i}{\sqrt{2}} e^{\pm i\alpha} \left[\mp \cot \beta \frac{\partial}{\partial \alpha} + i \frac{\partial}{\partial \beta} \pm \frac{1}{\sin \beta} \cdot \frac{\partial}{\partial \gamma} \right] \\ \hat{J}_0 &= -i \frac{\partial}{\partial \alpha} \end{aligned} \quad (4.2.6)$$

and $\hat{J}^\nu$ is a contravariant component of $\hat{\mathbf{J}}$ in the rotating (body-fixed) system

$$\begin{aligned} \hat{J}^{\pm 1} &= \frac{i}{\sqrt{2}} e^{\mp i\gamma} \left[\pm \cot \beta \frac{\partial}{\partial \gamma} + i \frac{\partial}{\partial \beta} \mp \frac{1}{\sin \beta} \cdot \frac{\partial}{\partial \alpha} \right] \\ \hat{J}^0 &= -i \frac{\partial}{\partial \gamma} \end{aligned} \quad (4.2.7)$$

The operator $\hat{\mathbf{J}}^2$ can be expressed in terms of $\hat{J}_\nu$ or $\hat{J}^\nu$ as

$$\begin{aligned} \hat{\mathbf{J}}^2 &= -\hat{J}_{-1} \hat{J}_{+1} + \hat{J}_0 \hat{J}_0 - \hat{J}_{+1} \hat{J}_{-1} = -\hat{J}^{-1} \hat{J}^{+1} + \hat{J}^0 \hat{J}^0 - \hat{J}^{+1} \hat{J}^{-1} = \hat{J}_0 (\hat{J}_0 + 1) - 2\hat{J}_{-1} \hat{J}_{+1} \\ &= \hat{J}^0 (\hat{J}^0 - 1) - 2\hat{J}^{-1} \hat{J}^{+1} = \hat{J}_0 (\hat{J}_0 - 1) - 2\hat{J}_{+1} \hat{J}_{-1} = \hat{J}^0 (\hat{J}^0 + 1) - 2\hat{J}^{+1} \hat{J}^{-1}. \end{aligned} \quad (4.2.8)$$

The relationships between $\hat{J}_\nu$ and $\hat{J}'^\mu$ read

$$\begin{aligned}\hat{J}'^\mu(\alpha, \beta, \gamma) &= \sum_\nu D_{\nu\mu}^1(\alpha, \beta, \gamma) \hat{J}_\nu(\alpha, \beta, \gamma) \\ \hat{J}_\nu(\alpha, \beta, \gamma) &= \sum_\mu D_{\nu\mu}^{1*}(\alpha, \beta, \gamma) \hat{J}'^\mu(\alpha, \beta, \gamma)\end{aligned}\quad (4.2.9)$$

Hence

$$\hat{J}'^\nu(\alpha, \beta, \gamma) = -\hat{J}_\nu(-\gamma, -\beta, -\alpha), \quad \hat{J}'^\nu(\alpha, \beta, \gamma) = \hat{J}_{-\nu}(\gamma, \beta, \alpha). \quad (4.2.10)$$

Commutators of spherical components $\hat{J}_\mu$ and $\hat{J}'^\nu(\mu, \nu = -1, 0, 1)$ are given by

$$\begin{aligned}[\hat{J}_\mu, \hat{J}_\nu] &= -\sqrt{2} C_{1\mu 1\nu}^{1\mu+\nu} \hat{J}_{\mu+\nu} \\ [\hat{J}'^\mu, \hat{J}'^\nu] &= \sqrt{2} C_{1\mu 1\nu}^{1\mu+\nu} \hat{J}'^{\mu+\nu} \\ [\hat{J}_\mu, \hat{J}'^\nu] &= 0\end{aligned}\quad (4.2.11)$$

The commutators of cartesian components $\hat{J}_k$ and $\hat{J}_l(i, k, l = x, y, z)$ are

$$\begin{aligned}[\hat{J}_i, \hat{J}_k] &= i\varepsilon_{ikl} \hat{J}_l, \\ [\hat{J}'_i, \hat{J}'_k] &= i\varepsilon_{ikl} \hat{J}_l, \\ [\hat{J}_i, \hat{J}'_k] &= 0,\end{aligned}\quad (4.2.12)$$

The operator of orbital angular momentum $\hat{\mathbf{L}}(\vartheta, \varphi)$ (Sec. 2.2) may be regarded as a special case of $\hat{\mathbf{J}}(\alpha, \beta, \gamma)$, viz. at $\alpha = \varphi, \beta = \vartheta, \gamma = 0$

$$\begin{aligned}\hat{J}_\nu(\varphi, \vartheta, 0) &= \hat{L}_\nu(\vartheta, \varphi), & \nu &= -1, 0, 1, \\ \hat{J}_i(\varphi, \vartheta, 0) &= \hat{L}_i(\vartheta, \varphi), & i &= x, y, z.\end{aligned}\quad (4.2.13)$$

at $\alpha = 0, \beta = \vartheta, \gamma = \varphi$

$$\begin{aligned}\hat{J}'^\nu(0, \vartheta, \varphi) &= \hat{L}'^\nu(\vartheta, \varphi), & \nu &= -1, 0, 1, \\ \hat{J}'_i(0, \vartheta, \varphi) &= \hat{L}'_i(\vartheta, \varphi), & i &= x, y, z.\end{aligned}\quad (4.2.14)$$

- (c) Equations (4.2.2) define $D_{MM'}^J(\alpha, \beta, \gamma)$ only within normalization and phase factors. To fix these factors some additional conditions are required. In the case of diagonal elements of the rotation matrix these factors are completely determined by the boundary condition

$$D_{MM'}^J(0, 0, 0) = \delta_{MM'}. \quad (4.2.15)$$

As for non-diagonal elements, they are determined by Eqs. (4.2.4) and (4.2.5) which relate the $D_{MM'}^J$, of different M and M' . This phase convention corresponds to the condition

$$D_{MM'}^J(0, \pi, 0) = (-1)^{J+M} \delta_{M, -M'} = (-1)^{J-M'} \delta_{-M, M'}. \quad (4.2.16)$$

4.3 Explicit forms of the Wigner D-functions

$D_{MM'}^J(\alpha, \beta, \gamma)$ may be represented as a product of three functions, each of which depends only on one argument α, β or γ ,

$$D_{MM'}^J(\alpha, \beta, \gamma) = e^{-iM\alpha} d_{MM'}^J(\beta) e^{-iM'\gamma}. \quad (4.3.1)$$

where $d_{MM'}^J(\beta)$ is a real function whose explicit forms are given below.

4.3.1 Expressions for $d_{MM'}^J(\beta)$ Involving Trigonometric Functions

$$\begin{aligned}d_{MM'}^J(\beta) &= (-1)^{J-M'} [(J+M)!(J-M)!(J+M')!(J-M')!]^{\frac{1}{2}} \\ &\times \sum_k (-1)^k \frac{\left(\cos \frac{\beta}{2}\right)^{M+M'+2k} \left(\sin \frac{\beta}{2}\right)^{2J-M-M'-2k}}{k!(J-M-k)!(J-M'-k)!(M+M'+k)!}\end{aligned}\quad (4.3.2)$$

$$d_{MM'}^J(\beta) = (-1)^{J+M} [(J+M)!(J-M)!(J+M')!(J-M')!]^{\frac{1}{2}} \\ \times \sum_k (-1)^k \frac{\left(\cos \frac{\beta}{2}\right)^{2k-M-M'} \left(\sin \frac{\beta}{2}\right)^{2J+M+M'-2k}}{k!(J+M-k)!(J+M'-k)!(k-M-M')!} \quad (4.3.3)$$

$$d_{MM'}^J(\beta) = [(J+M)!(J-M)!(J+M')!(J-M')!]^{\frac{1}{2}} \\ \times \sum_k (-1)^k \frac{\left(\cos \frac{\beta}{2}\right)^{2J-2k+M-M'} \left(\sin \frac{\beta}{2}\right)^{2k-M+M'}}{k!(J+M-k)!(J-M'-k)!(M'-M+k)!} \quad (4.3.4)$$

$$d_{MM'}^J(\beta) = (-1)^{M-M'} [(J+M)!(J-M)!(J+M')!(J-M')!]^{\frac{1}{2}} \\ \times \sum_k (-1)^k \frac{\left(\cos \frac{\beta}{2}\right)^{2J-2k-M+M'} \left(\sin \frac{\beta}{2}\right)^{2k+M-M'}}{k!(J-M-k)!(J+M'-k)!(M-M'+k)!} \quad (4.3.5)$$

In Eqs. (4.3.2)-(4.3.5) k runs over all integer values for which the factorial arguments are non-negative. Each of these sums contains $(N+1)$ terms, where N is the minimum of $J+M$, $J-M$, $J+M'$ and $J-M'$. Equations (4.3.2)-(4.3.5) are not independent, but may be transformed into one another by changing summation variables.

Equations (4.3.2)-(4.3.5) may be regarded as special cases of the more general expression

$$d_{MM'}^J(\beta) = \left[\frac{(J_1+J_2+J+1)!(J_1+J_2-J)!}{2J+1} \right]^{\frac{1}{2}} \\ \times \sum_{\substack{m_1 m_2 \\ (m_1+m_2=M)}} (-1)^{J_2+m_2} C_{J_1 m_1 J_2 m_2}^{J M} \frac{\left(\cos \frac{\beta}{2}\right)^{J_1+J_2+m_1-m_2} \left(\sin \frac{\beta}{2}\right)^{J_1+J_2-m_1+m_2}}{[(J_1+m_1)!(J_1-m_1)!(J_2+m_2)!(J_2-m_2)!]^{\frac{1}{2}}} \quad (4.3.6)$$

where J_1 and J_2 are arbitrary integer or half-integer numbers which satisfy the conditions $J_1 - J_2 = M'$ and $|J_1 - J_2| \leq J \leq J_1 + J_2$. The sum in Eq. (4.3.6) is over all possible (positive and negative) values of m_1 and m_2 which correspond to nonzero Clebsch-Gordan coefficients. In particular, Eq. (4.3.6) reduces to Eq. (4.3.2), when $J_1 = (J+M')/2$, $J_2 = (J-M')/2$, $m_1 = M+k-(J-M')/2$ and $m_2 = -k+(J-M')/2$.

4.3.2 Differential Representations of $d_{MM'}^J(\beta)$

$$d_{MM'}^J(\beta) = (-1)^{J-M'} \frac{1}{2^J} \left[\frac{(J+M)!}{(J-M)!(J+M')!(J-M')!} \right]^{\frac{1}{2}} (1 - \cos \beta)^{\frac{M'-M}{2}} (1 + \cos \beta)^{-\frac{M+M'}{2}} \\ \times \frac{d^{J-M}}{(d \cos \beta)^{J-M}} \left[(1 - \cos \beta)^{J-M'} (1 + \cos \beta)^{J+M'} \right] \quad (4.3.7)$$

$$d_{MM'}^J(\beta) = (-1)^{J+M} \frac{1}{2^J} \left[\frac{(J-M)!}{(J+M)!(J+M')!(J-M')!} \right]^{\frac{1}{2}} (1 - \cos \beta)^{\frac{M-M'}{2}} (1 + \cos \beta)^{\frac{M+M'}{2}} \\ \times \frac{d^{J+M}}{(d \cos \beta)^{J+M}} \left[(1 - \cos \beta)^{J+M'} (1 + \cos \beta)^{J-M'} \right] \quad (4.3.8)$$

$$d_{MM'}^J(\beta) = (-1)^{J-M'} \frac{1}{2^J} \left[\frac{(J+M')!}{(J+M)!(J-M)!(J-M')!} \right]^{\frac{1}{2}} (1 - \cos \beta)^{\frac{M-M'}{2}} (1 + \cos \beta)^{-\frac{M+M'}{2}} \\ \times \frac{d^{J-M'}}{(d \cos \beta)^{J-M'}} \left[(1 - \cos \beta)^{J-M} (1 + \cos \beta)^{J+M} \right] \quad (4.3.9)$$

$$d_{MM'}^J(\beta) = (-1)^{J+M} \frac{1}{2^J} \left[\frac{(J-M')!}{(J+M)!(J-M)!(J+M')!} \right]^{\frac{1}{2}} (1 - \cos \beta)^{\frac{M'-M}{2}} (1 + \cos \beta)^{\frac{M+M'}{2}} \\ \times \frac{d^{J+M'}}{(d \cos \beta)^{J+M'}} \left[(1 - \cos \beta)^{J+M} (1 + \cos \beta)^{J-M} \right] \quad (4.3.10)$$

In practice it is convenient to use such equation from Eqs. (4.3.7)-(4.3.10) in which the order of derivative is the lowest.

4.3.3 Integral Representations of $d_{MM'}^J(\beta)$

$$d_{MM'}^J(\beta) = i^{M-M'} \frac{1}{2\pi} \left[\frac{(J+M)!(J-M)!}{(J+M')!(J-M')!} \right]^{\frac{\alpha}{2}} \times \int_0^{2\pi} \left(e^{i\frac{\phi}{2}} \cos \frac{\beta}{2} + ie^{-i\frac{\phi}{2}} \sin \frac{\beta}{2} \right)^{J-M'} \left(e^{-i\frac{\phi}{2}} \cos \frac{\beta}{2} + ie^{i\frac{\phi}{2}} \sin \frac{\beta}{2} \right)^{J+M'} e^{iM\phi} d\phi. \quad (4.3.11)$$

Equation (4.3.11) can be rewritten as a contour integral

$$d_{MM'}^J(\beta) = \frac{i^{M-M'-1}}{2\pi} \left[\frac{(J+M)!(J-M)!}{(J+M')!(J-M')!} \right]^{\frac{1}{2}} \times \oint \left(z \cos \frac{\beta}{2} + i \sin \frac{\beta}{2} \right)^{J-M'} \left(iz \sin \frac{\beta}{2} + \cos \frac{\beta}{2} \right)^{J+M'} z^{M-J-1} dz. \quad (4.3.12)$$

The integration contour in Eq. (4.3.12) is a circle of unit radius about the origin of the z -plane.

4.3.4 Relation Between $d_{MM'}^J(\beta)$ and the Jacobi Polynomials

The functions $d_{MM'}^J(\beta)$ can be expressed in terms of the Jacobi polynomials

$$d_{MM'}^J(\beta) = \xi_{MM'} \left[\frac{s!(s+\mu+\nu)!}{(s+\mu)!(s+\nu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu P_s^{(\mu,\nu)}(\cos \beta), \quad (4.3.13)$$

where μ, ν and s are related to M, M' and J by

$$\mu = |M - M'|, \quad \nu = |M + M'|, \quad s = J - \frac{1}{2}(\mu + \nu). \quad (4.3.14)$$

and

$$\xi_{MM'} = \begin{cases} 1 & \text{if } M' \geq M, \\ (-1)^{M'-M} & \text{if } M' < M. \end{cases} \quad (4.3.15)$$

4.3.5 Relations Between $d_{MM'}^J(\beta)$ and Hypergeometric Functions

$$d_{MM'}^J(\beta) = \frac{\xi_{MM'}}{\mu!} \left[\frac{(s+\mu+\nu)!(s+\mu)!}{s!(s+\nu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu F \left(-s, s+\mu+\nu+1; \mu+1; \sin^2 \frac{\beta}{2} \right), \quad (4.3.16)$$

$$d_{MM'}^J(\beta) = \frac{\xi_{MM'}}{\mu!} \left[\frac{(s+\mu+\nu)!(s+\mu)!}{s!(s+\nu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^{-\nu} F \left(s+\mu+1, -s-\nu; \mu+1; \sin^2 \frac{\beta}{2} \right), \quad (4.3.17)$$

$$d_{MM'}^J(\beta) = \frac{(-1)^s \xi_{MM'}}{\nu!} \left[\frac{(s+\mu+\nu)!(s+\nu)!}{s!(s+\mu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu F \left(-s, s+\mu+\nu+1; \nu+1; \cos^2 \frac{\beta}{2} \right), \quad (4.3.18)$$

$$d_{MM'}^J(\beta) = \frac{(-1)^s \xi_{MM'}}{\nu!} \left[\frac{(s+\mu+\nu)!(s+\nu)!}{s!(s+\mu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^{-\mu} \left(\cos \frac{\beta}{2} \right)^\nu F \left(s+\nu+1, -s-\mu; \nu+1; \cos^2 \frac{\beta}{2} \right), \quad (4.3.19)$$

$$d_{MM'}^J(\beta) = \frac{\xi_{MM'}}{\mu!} \left[\frac{(s+\mu+\nu)!(s+\mu)!}{s!(s+\nu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^{2s+\nu} F \left(-s, -s-\nu; \mu+1; -\tan^2 \frac{\beta}{2} \right), \quad (4.3.20)$$

$$d_{MM'}^J(\beta) = \frac{(-1)^s \xi_{MM'}}{\nu!} \left[\frac{(s+\mu+\nu)!(s+\nu)!}{s!(s+\mu)!} \right]^{\frac{1}{2}} \left(\sin \frac{\beta}{2} \right)^{2s+\mu} \left(\cos \frac{\beta}{2} \right)^\nu F \left(-s, -s-\mu; \nu+1; -\cot^2 \frac{\beta}{2} \right), \quad (4.3.21)$$

$$d_{MM'}^J(\beta) = \frac{(-1)^s \xi_{MM'} (2s+\mu+\nu)!}{[s!(s+\mu+\nu)!(s+\mu)!(s+\nu)!]^{\frac{1}{2}}} \left(\sin \frac{\beta}{2} \right)^{2s+\mu} \left(\cos \frac{\beta}{2} \right)^\nu F \left(-s, -s-\mu; -2s-\mu-\nu; \frac{1}{\sin^2 \frac{\beta}{2}} \right), \quad (4.3.22)$$

$$d_{MM'}^J(\beta) = \frac{\xi_{MM'} (2s+\mu+\nu)!}{[s!(s+\mu+\nu)!(s+\mu)!(s+\nu)!]^{\frac{1}{2}}} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^{2s+\nu} F \left(-s, -s-\nu; -2s-\mu-\nu; \frac{1}{\cos^2 \frac{\beta}{2}} \right). \quad (4.3.23)$$

Parameters μ, ν, s and a phase factor $\xi_{MM'}$ in Eqs. (4.3.16)-(4.3.23) are defined by Eqs. (4.3.14) and (4.3.15).

4.4 Symmetries of $d_{MM'}^J(\beta)$ and $D_{MM'}^J(\alpha, \beta, \gamma)$

(a) In accordance with Eqs. (4.3.2)-(4.3.5) the functions $d_{MM'}^J(\beta)$ are real and satisfy the relations

$$\begin{aligned} d_{MM'}^J(\beta) &= (-1)^{M-M'} d_{-M-M'}^J(\beta) = (-1)^{M-M'} d_{M'M}^J(\beta) = d_{-M'-M}^J(\beta), \\ d_{MM'}^J(-\beta) &= (-1)^{M-M'} d_{MM'}^J(\beta) = d_{M'M}^J(\beta), \\ d_{MM'}^J(\pi - \beta) &= (-1)^{J-M'} d_{-MM'}^J(\beta) = (-1)^{J+M} d_{M-M'}^J(\beta), \\ d_{MM'}^J(\beta \pm 2\pi n) &= (-1)^{2Jn} d_{MM'}^J(\beta), \quad n \text{ is integer}, \\ d_{MM'}^J(\beta \pm (2n+1)\pi) &= (-1)^{\pm(2n+1)J-M'} d_{M-M'}^J(\beta), \quad n \text{ is integer}. \end{aligned} \quad (4.4.1)$$

(b) Equations (4.4.1) imply the following symmetry properties of $D_{MM'}^J(\alpha, \beta, \gamma)$

$$\begin{aligned} D_{MM'}^J(\alpha, \beta, \gamma) &= \varepsilon \eta D_{-M-M'}^J(\alpha, \beta, \gamma) = \eta D_{MM'}^{J*}(\alpha, \beta, \gamma) = \varepsilon D_{-M-M'}^{J*}(\alpha, \beta, \gamma) \\ &= \varepsilon D_{M'M}^J(\gamma, \beta, \alpha) = \eta D_{-M'-M}^J(\gamma, \beta, \alpha) = \varepsilon \eta D_{M'M}^{J*}(\gamma, \beta, \alpha) = D_{-M'-M}^{J*}(\gamma, \beta, \alpha) \\ &= \varepsilon D_{MM'}^J(\alpha, -\beta, \gamma) = \eta D_{-M-M'}^J(\alpha, -\beta, \gamma) = \varepsilon \eta D_{MM'}^{J*}(\alpha, -\beta, \gamma) = D_{-M-M'}^{J*}(\alpha, -\beta, \gamma) \\ &= D_{M'M}^J(\gamma, -\beta, \alpha) = \varepsilon \eta D_{-M'-M}^J(\gamma, -\beta, \alpha) = \eta D_{M'M}^{J*}(\gamma, -\beta, \alpha) = \varepsilon D_{-M'-M}^{J*}(\gamma, -\beta, \alpha) \\ &= \varepsilon D_{-M-M'}^J(-\alpha, \beta, -\gamma) = \eta D_{MM'}^J(-\alpha, \beta, -\gamma) = \varepsilon \eta D_{-M-M'}^{J*}(-\alpha, \beta, -\gamma) = D_{MM'}^{J*}(-\alpha, \beta, -\gamma) \\ &= D_{-M'-M}^J(-\gamma, \beta, -\alpha) = \varepsilon \eta D_{M'M}^J(-\gamma, \beta, -\alpha) = \eta D_{-M'-M}^{J*}(-\gamma, \beta, -\alpha) = \varepsilon D_{M'M}^{J*}(-\gamma, \beta, -\alpha) \\ &= D_{-M-M'}^J(-\alpha, -\beta, -\gamma) = \varepsilon \eta D_{MM'}^J(-\alpha, -\beta, -\gamma) = \eta D_{-M-M'}^{J*}(-\alpha, -\beta, -\gamma) = \varepsilon D_{MM'}^{J*}(-\alpha, -\beta, -\gamma) \\ &= \varepsilon D_{-M'-M}^J(-\gamma, -\beta, -\alpha) = \eta D_{M'M}^J(-\gamma, -\beta, -\alpha) = \varepsilon \eta D_{-M'-M}^{J*}(-\gamma, -\beta, -\alpha) = D_{M'M}^{J*}(-\gamma, -\beta, -\alpha), \end{aligned} \quad (4.4.2)$$

where

$$\varepsilon = (-1)^{M'-M}, \quad \eta = e^{-i2M\alpha - i2M'\gamma}. \quad (4.4.3)$$

The periodicity conditions for $D_{MM'}^J(\alpha, \beta, \gamma)$ are

$$\begin{aligned} D_{MM'}^J(\alpha, \beta \pm 2n\pi, \gamma) &= (-1)^{2nJ} D_{MM'}^J(\alpha, \beta, \gamma), \\ D_{MM'}^J(\alpha, \beta \pm (2n+1)\pi, \gamma) &= (-1)^{\pm(2n+1)J-M'} D_{M-M'}^J(\alpha, \beta, -\gamma), \\ D_{MM'}^J(\alpha \pm n\pi, \beta, \gamma) &= (-i)^{\pm 2nM} D_{MM'}^J(\alpha, \beta, \gamma), \\ D_{MM'}^J(\alpha, \beta, \gamma \pm n\pi) &= (-i)^{\pm 2nM'} D_{MM'}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.4.4)$$

where n is integer. Note also that

$$D_{MM'}^J(\tilde{\alpha}, \beta, \tilde{\gamma}) = e^{iM(\alpha - \tilde{\alpha})} D_{MM'}^J(\alpha, \beta, \gamma) e^{iM'(\gamma - \tilde{\gamma})}. \quad (4.4.5)$$

Some properties of $D_{MM'}^J(\alpha, \beta, \gamma)$ follow from the addition theorem (Sec. a). Let us consider some special cases.

(i) The matrix of the transformation $S\{x, y, z\} \rightarrow S''\{x', -y', -z'\}$ may be obtained from the matrix of the transformation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ by substituting $(\alpha + \pi, \pi - \beta, -\gamma)$ for (α, β, γ) . On the other hand, this substitution corresponds to an additional rotation $R_{x'}$ about the x' -axis through an angle $-\pi$. Hence,

$$\begin{aligned} D_{MM'}^J(\alpha + \pi, \pi - \beta, -\gamma) &= \hat{R}_{x'} D_{MM'}^J(\alpha, \beta, \gamma) = \sum_{M''} D_{MM''}^J(\alpha, \beta, \gamma) D_{M''M'}^J\left(-\frac{\pi}{2}, -\pi, \frac{\pi}{2}\right) \\ &= (-1)^J D_{M-M'}^J(\alpha, \beta, \gamma). \end{aligned} \quad (4.4.6)$$

(ii) The matrix of the transformation $S\{x, y, z\} \rightarrow S''' \{-x', y', -z'\}$ may be obtained from the matrix of the transformation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ by substituting $(\alpha - \pi, \pi - \beta, \pi - \gamma)$ for (α, β, γ) . This substitution corresponds to an additional rotation $R_{y'}$ about the y' -axis through an angle $-\pi$. Hence,

$$\begin{aligned} D_{MM'}^J(\alpha - \pi, \pi - \beta, \pi - \gamma) &= \hat{R}_{y'} D_{MM'}^J(\alpha, \beta, \gamma) = \sum_{M''} D_{MM''}^J(\alpha, \beta, \gamma) D_{M''M'}^J(0, -\pi, 0) \\ &= (-1)^{J+M'} D_{M-M'}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.4.7)$$

(iii) The matrix of the transformation $S\{x, y, z\} \rightarrow S'''' \{-x', -y', z'\}$ may be derived from the matrix of the transformation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ by replacing $(\alpha, \beta, \gamma) \rightarrow (\alpha, \beta, \gamma - \pi)$. This corresponds to an additional rotation $R_{z'}$ about the z' -axis through an angle $-\pi$. Thus,

$$D_{MM'}^J(\alpha, \beta, \gamma - \pi) = \hat{R}_{z'} D_{MM'}^J(\alpha, \beta, \gamma) = \sum_{M''} D_{MM''}^J(\alpha, \beta, \gamma) D_{M''M'}^J(0, 0, -\pi) = (-1)^{M'} D_{MM'}^J(\alpha, \beta, \gamma). \quad (4.4.8)$$

Note that three successive rotations through angles $-\pi$ about the axes x', y' and z' return the coordinate system to its initial position.

4.5 Rotation matrix $U_{MM'}^J$, in terms of angles ω, Θ, Φ

4.5.1 Definition

In some cases the description of rotations in terms of ω, Θ, Φ (where ω is the angle of rotation and Θ, Φ are the angles which determine the rotation axis, see Sec. 1.4) is more convenient than that in terms of the Euler angles α, β, γ . Matrix elements of the rotation operator in terms of variables ω, Θ, Φ will be denoted by $U_{MM'}^J(\omega; \Theta, \Phi)$:

$$U_{MM'}^J(\omega; \Theta, \Phi) \equiv \langle JM | e^{-i\omega \mathbf{n} \cdot \hat{\mathbf{J}}} | JM' \rangle. \quad (4.5.1)$$

Therefore, under a rotation specified by ω, Θ, Φ the components of an irreducible tensor of rank J transform as

$$\mathfrak{M}_{JM'}(\vartheta', \varphi') = \sum_M \mathfrak{M}_{JM}(\vartheta, \varphi) U_{MM'}^J(\omega; \Theta, \Phi). \quad (4.5.2)$$

The polar angles ϑ, φ and ϑ', φ' which specify a direction of an arbitrary vector in the initial and rotated coordinate systems in terms ω, Θ, Φ are related by Eqs. (1.4.55) and (1.4.56).

4.5.2 Explicit form

(a) Arbitrary rotation as product of three rotations

An arbitrary rotation specified by angles ω, Θ, Φ may be considered as a result of three successive rotations of coordinate system:

- (i) $R_1(\alpha_1 = \Phi, \beta_1 = \Theta, \gamma_1 = -\Phi)$, i.e., the rotation which turns the z -axis to the direction of $\mathbf{n}(\Theta, \Phi)$;
- (ii) $R_2(\alpha_2 = \omega, \beta_2 = 0, \gamma_2 = 0)$, i.e., the rotation about $\mathbf{n}(\Theta, \Phi)$ through an angle ω ;
- (iii) $R_3(\alpha_3 = \Phi, \beta_3 = -\Theta, \gamma_3 = -\Phi)$, i.e., the rotation which is inverse to R_1 .

The result of these three rotations yields the relation between $U_{MM'}^J(\omega; \Theta, \Phi)$ and the Wigner D -functions

$$U_{MM'}^J(\omega; \Theta, \Phi) = \sum_{M''} D_{MM''}^J(\Phi, \Theta, -\Phi) e^{-iM''\omega} D_{M''M'}^J(\Phi, -\Theta, -\Phi). \quad (4.5.3)$$

Equation (4.5.3) enables one to find an explicit form for $U_{MM'}^J(\omega; \Theta, \Phi)$ for particular J, M and M' .

(b) Construction from Cayley-Klein parameters

According to Eq. (4.5.10) the functions $U_{MM'}^J(\omega; \Theta, \Phi)$ may be directly constructed from the matrix elements $U_{mm'}^{\frac{1}{2}}$, which represent the Cayley-Klein parameters (see Eq. (4.5.12)). This gives the expression

$$U_{MM'}^J(\omega; \Theta, \Phi) = \begin{cases} (-iv)^{2J} \left(\frac{u}{-iv}\right)^{(M+M')} & e^{-i(M-M')\Phi} \sum_s \frac{\sqrt{(J+M)!(J-M)!(J+M')!(J-M')!}}{s!(s+M+M')!(J-M-s)!(J-M'-s)!} (1-v^{-2})^s, \\ & M+M' \geq 0, \\ (-iv)^{2J} \left(\frac{u^*}{-iv}\right)^{-(M+M')} & e^{-i(M-M')\Phi} \sum_s \frac{\sqrt{(J+M)!(J-M)!(J+M')!(J-M')!}}{s!(s-M-M')!(J+M-s)!(J+M'-s)!} (1-v^{-2})^s, \\ & M+M' \leq 0. \end{cases} \quad (4.5.4)$$

In this case Eqs. (1.4.26) have been used, and the following notations are introduced

$$v = \sin \frac{\omega}{2} \sin \Theta, \quad u = \cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \Theta. \quad (4.5.5)$$

In Eqs. (4.5.4) the summation index s runs over all integer values which do not lead to negative factorial arguments.

(c) Alternative expression for U

Another explicit form of $U_{MM'}^J(\omega; \Theta, \Phi)$ can be obtained directly from $D_{MM'}^J(\alpha, \beta, \gamma)$ by changing variables $(\omega, \Theta, \Phi) \rightarrow (\alpha, \beta, \gamma)$ with the aid of Eqs. (1.4.16) and (1.4.17)

$$U_{MM'}^J(\omega; \Theta, \Phi) = i^{M-M'} e^{-i(M-M')\Phi} \left(\frac{1 - i \tan \frac{\omega}{2} \cos \Theta}{\sqrt{1 + \tan^2 \frac{\omega}{2} \cos^2 \Theta}} \right)^{M+M'} d_{MM'}^J(\xi). \quad (4.5.6)$$

The functions $d_{MM'}^J(\xi)$ are defined in Sec. 4.3, and the angle ξ is determined by

$$\sin \frac{\xi}{2} = \sin \frac{\omega}{2} \sin \Theta. \quad (4.5.7)$$

(d) *Expansion in spherical harmonics*

The function $U_{MM'}^J(\omega; \Theta, \Phi)$ may be expanded in a series of the spherical harmonics which depend on polar angles Θ and Φ ,

$$U_{MM'}^J(\omega; \Theta, \Phi) = \sum_{\lambda\mu} (-i)^\lambda \frac{2\lambda+1}{2J+1} \chi_\lambda^J(\omega) C_{JM\lambda\mu}^{JM'} \sqrt{\frac{4\pi}{2\lambda+1}} Y_{\lambda\mu}(\Theta, \Phi), \quad (4.5.8)$$

where $\chi_\lambda^J(\omega)$ is a generalized character (of order λ) of the irreducible representation of rank J . Explicit forms and properties of $\chi_\lambda^J(\omega)$ will be given below in Sec. 4.15.

Equation (4.5.8) shows that $U_{MM'}^J(\omega; \Theta, \Phi)$ depends on M and M' only through the Clebsch-Gordan coefficients $C_{JM\lambda\mu}^{JM'}$.

4.5.3 Differential Equations

$U_{MM'}^J(\omega; \Theta, \Phi)$ are eigenfunctions of three operators $\hat{J}_z, \hat{J}'_z$ and $\hat{J}^2$ whose eigenvalues equal $-M, -M'$ and $J(J+1)$, respectively. In terms of ω, Θ, Φ these operators have the forms

$$\hat{J}_z = -i \left[\cos \Theta \frac{\partial}{\partial \omega} - \frac{1}{2} \cot \frac{\omega}{2} \sin \Theta \frac{\partial}{\partial \Theta} + \frac{1}{2} \cdot \frac{\partial}{\partial \Phi} \right], \quad (4.5.9)$$

$$\hat{J}'_z = -i \left[\cos \Theta \frac{\partial}{\partial \omega} - \frac{1}{2} \cot \frac{\omega}{2} \sin \Theta \frac{\partial}{\partial \Theta} - \frac{1}{2} \cdot \frac{\partial}{\partial \Phi} \right], \quad (4.5.10)$$

$$\hat{J}^2 = - \left[\frac{\partial^2}{\partial \omega^2} + \cot \frac{\omega}{2} \frac{\partial}{\partial \omega} + \frac{1}{4 \sin^2 \frac{\omega}{2}} \left(\frac{\partial^2}{\partial \Theta^2} + \cot \Theta \frac{\partial}{\partial \Theta} + \frac{1}{\sin^2 \Theta} \frac{\partial^2}{\partial \Phi^2} \right) \right]. \quad (4.5.11)$$

Thus, $U_{MM'}^J(\omega; \Theta, \Phi)$ is a solution of the differential equations

$$\begin{aligned} [\hat{J}^2 - J(J+1)] U_{MM'}^J(\omega; \Theta, \Phi) &= 0 \\ [\hat{J}_z + M] U_{MM'}^J(\omega; \Theta, \Phi) &= 0, \\ [\hat{J}'_z + M'] U_{MM'}^J(\omega; \Theta, \Phi) &= 0. \end{aligned} \quad (4.5.12)$$

with the boundary conditions

$$\begin{aligned} U_{MM'}^J(0; \Theta, \Phi) &= \delta_{MM'} \\ \frac{\partial}{\partial \Phi} U_{MM'}^J(\omega; \Theta, \Phi) \Big|_{\Theta=0} &= \frac{\partial}{\partial \Phi} U_{MM'}^J(\omega; \Theta, \Phi) \Big|_{\Theta=\pi} = 0. \end{aligned} \quad (4.5.13)$$

4.5.4 Orthogonality and Completeness

A collection of the functions $U_{MM'}^J(\omega; \Theta, \Phi)$ with all possible integer and half-integer $J \geq 0$ constitutes a complete set of orthogonal functions of three variables ω, Θ, Φ defined in the domain

$$0 \leq \theta \leq \pi, \quad 0 \leq \Phi < 2\pi, \quad 0 \leq \omega < 2\pi, \quad (4.5.14)$$

whose total volume is equal to $16\pi^2$.

(a) *Orthogonality and Normalization*

$$4 \int_0^{2\pi} d\omega \sin^2 \frac{\omega}{2} \int_0^\pi d\Theta \sin \Theta \int_0^{2\pi} d\Phi U_{M_1 M'_1}^{J_1*}(\omega; \Theta, \Phi) U_{M_2 M'_2}^{J_2}(\omega; \Theta, \Phi) = \frac{16\pi^2}{2J_1+1} \delta_{J_1 J_2} \delta_{M_1 M_2} \delta_{M'_1 M'_2}. \quad (4.5.15)$$

(b) *Completeness*

$$\sum_{J=0,1/2,1,\dots}^{\infty} \frac{2J+1}{16\pi^2} \sum_{MM'} U_{MM'}^J(\omega; \Theta, \Phi) U_{MM'}^{J*}(\tilde{\omega}; \tilde{\Theta}, \tilde{\Phi}) = \frac{\delta(\Phi - \tilde{\Phi}) \delta(\Theta - \tilde{\Theta}) \delta(\omega - \tilde{\omega})}{4 \sin \Theta \sin^2 \frac{\omega}{2}}. \quad (4.5.16)$$

4.5.5 Principal Properties

(a) *Inverse transformation*

$$[U^{-1}(\omega; \Theta, \Phi)]_{MM'}^J = U_{MM'}^J(-\omega; \Theta, \Phi) = U_{MM'}^J(\omega; \pi - \Theta, \pi + \Phi). \quad (4.5.17)$$

(b) *Complex conjugation*

$$U_{MM'}^{J*}(\omega; \Theta, \Phi) = U_{M'M}^J(\omega; \pi - \Theta, \pi + \Phi) = U_{M'M}^J(-\omega; \Theta, \Phi). \quad (4.5.18)$$

(c) *Reversal of argument signs*

$$\begin{aligned} U_{MM'}^J(-\omega; \Theta, \Phi) &= (-1)^{M-M'} U_{-M'-M}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; -\Theta, \Phi) &= (-1)^{M-M'} U_{MM'}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; \Theta, -\Phi) &= U_{M'M}^J(\omega; \Theta, \Phi). \end{aligned} \quad (4.5.19)$$

(d) *Periodicity*

$$\begin{aligned} U_{MM'}^J(\omega + 4\pi; \Theta, \Phi) &= U_{MM'}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; \Theta + 2\pi, \Phi) &= U_{MM'}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; \Theta, \Phi + 2\pi) &= U_{MM'}^J(\omega; \Theta, \Phi). \end{aligned} \quad (4.5.20)$$

(e) *Half-period Shift of Arguments*

$$\begin{aligned} U_{MM'}^J(\omega + 2\pi; \Theta, \Phi) &= (-1)^{2J} U_{MM'}^J(\omega; \Theta, \Phi), & U_{MM'}^J(2\pi - \omega; \Theta, \Phi) &= (-1)^{M+M'} U_{-M'-M}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; \Theta + \pi, \Phi) &= (-1)^{M-M'} U_{-M'-M}^J(\omega; \Theta, \Phi), & U_{MM'}^J(\omega; \pi - \Theta, \Phi) &= U_{-M'-M}^J(\omega; \Theta, \Phi), \\ U_{MM'}^J(\omega; \Theta, \Phi + \pi) &= (-1)^{M-M'} U_{MM'}^J(\omega; \Theta, \Phi), & U_{MM'}^J(\omega; \Theta, \pi - \Phi) &= (-1)^{M-M'} U_{M'M}^J(\omega; \Theta, \Phi). \end{aligned} \quad (4.5.21)$$

(f) *Permutation $M \rightleftharpoons M'$ and reversal of signs of M and M'*

$$\begin{aligned} U_{M'M}^J(\omega; \Theta, \Phi) &= U_{MM'}^J(\omega; \Theta, -\Phi) = U_{MM'}^{J*}(\omega; \pi - \Theta, \pi + \Phi), \\ U_{-M-M'}^J(\omega; \Theta, \Phi) &= (-1)^{M-M'} U_{MM'}^J(\omega; \pi - \Theta, \pi - \Phi), \\ U_{MM'}^J(\omega; \Theta, \Phi) &= (-1)^{J-M'} U_{-MM'}^J(\omega_1; \Theta_1, \Phi_1) = (-1)^{J+M} U_{-M-M'}^J(\omega_1; \Theta_1, \Phi_1). \end{aligned} \quad (4.5.22)$$

where angles $\omega_1, \Theta_1, \Phi_1$ and ω, Θ, Φ are related by the equations

$$\begin{aligned} \cos \frac{\omega_1}{2} &= \sin \frac{\omega}{2} \sin \Theta \sin \Phi, \\ \sin \frac{\omega_1}{2} \cos \Theta_1 &= -\sin \frac{\omega}{2} \sin \Theta \cos \Phi, \\ \cot \Phi_1 &= -\tan \frac{\omega}{2} \cos \Theta, \end{aligned} \quad (4.5.23)$$

$$\begin{aligned} \cos \frac{\omega}{2} &= \sin \frac{\omega_1}{2} \sin \Theta_1 \sin \Phi_1, \\ \sin \frac{\omega}{2} \cos \Theta &= -\sin \frac{\omega_1}{2} \sin \Theta_1 \cos \Phi_1, \\ \cot \Phi &= -\tan \frac{\omega_1}{2} \cos \Theta_1, \end{aligned} \quad (4.5.24)$$

$$\sin^2 \frac{\omega_1}{2} \sin^2 \Theta_1 + \sin^2 \frac{\omega}{2} \sin^2 \Theta = 1. \quad (4.5.25)$$

Some relations which are valid for $D_{MM'}^J(\alpha, \beta, \gamma)$ remain valid for $U_{MM'}^J(\omega; \Theta, \Phi)$. In particular, the Clebsch-Gordan expansion

$$U_{M_1 M'_1}^{J_1}(\omega; \Theta, \Phi) U_{M_2 M'_2}^{J_2}(\omega; \Theta, \Phi) = \sum_{JM} C_{J_1 M_1 J_2 M_2}^{JM} U_{MM'}^J(\omega; \Theta, \Phi) C_{J_1 M'_1 J_2 M'_2}^{JM'}, \quad (4.5.26)$$

is equivalent to Eq. (4.6.1), and the addition theorem

$$\sum_{M''} U_{MM''}^J(\omega_2; \Theta_2, \Phi_2) U_{M''M'}^J(\omega_1; \Theta_1, \Phi_1) = U_{MM'}^J(\omega; \Theta, \Phi) \quad (4.5.27)$$

is equivalent to Eq. (4.7.1). The angles $\omega_1, \Theta_1, \Phi_1, \omega_2, \Theta_2, \Phi_2$ and ω, Θ, Φ are related by Eqs. (1.4.76).

4.5.6 Special Cases

(a) The rotation axis $\mathbf{n}(\theta, \Phi)$ coincides with one of the coordinate axes

(i) The x -axis ($\Theta = \frac{\pi}{2}, \Phi = 0$)

$$U_{MM'}^J\left(\omega; \frac{\pi}{2}, 0\right) = D_{MM'}^J\left(-\frac{\pi}{2}, \omega, \frac{\pi}{2}\right) = (i)^{M-M'} d_{MM'}^J(\omega). \quad (4.5.28)$$

(ii) The y -axis ($\Theta = \frac{\pi}{2}, \Phi = \frac{\pi}{2}$)

$$U_{MM'}^J\left(\omega; \frac{\pi}{2}, \frac{\pi}{2}\right) = D_{MM'}^J(0, \omega, 0) = d_{MM'}^J(\omega). \quad (4.5.29)$$

(iii) The z -axis ($\Theta = 0$)

$$U_{MM'}^J(\omega; 0, \Phi) = \delta_{MM'} e^{-iM\omega}. \quad (4.5.30)$$

(b) Rotation about $\mathbf{n}(\Theta, \Phi)$ through a small angle $\omega \ll \pi/2$

$$\begin{aligned} U_{MM'}^J(\omega; \Theta, \Phi) &= \delta_{MM'}(1 - i\omega M \cos \Theta) - \frac{i\omega}{2} e^{-i\Phi} \sin \Theta \sqrt{(J-M')(J+M)} \delta_{M, M'+1}, \\ &\quad - \frac{i\omega}{2} e^{i\Phi} \sqrt{(J-M)(J+M')} \delta_{M M'-1}. \end{aligned} \quad (4.5.31)$$

(c) Explicit forms of $U_{MM'}^J(\omega; \Theta, \Phi)$ at $M = \pm J$ and/or $M' = \pm J$

$$\begin{aligned} U_{JJ}^J(\omega; \theta, \Phi) &= \left(\cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \Theta\right)^{2J}, \quad U_{J-J}^J(\omega; \theta, \Phi) = \left(\cos \frac{\omega}{2} + i \sin \frac{\omega}{2} \cos \Theta\right)^{2J}, \\ U_{J-J}^J(\omega; \Theta, \Phi) &= \left(-i \sin \frac{\omega}{2} \sin \Theta e^{-i\Phi}\right)^{2J}, \quad U_{-JJ}^J(\omega; \Theta, \Phi) = \left(-i \sin \frac{\omega}{2} \sin \Theta e^{i\Phi}\right)^{2J}. \end{aligned} \quad (4.5.32)$$

(d) Explicit forms of $U_{MM'}^J(\omega; \theta, \Phi)$ for $J = \frac{1}{2}, 1, \frac{3}{2}, 2$ are given below in Tables 4.23-4.26.

4.6 Sums involving D -functions

4.6.1 The Clebsch-Gordan Series

The product of two D -functions with the same arguments may be expanded in the following series

$$D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) = \sum_{J=|J_1-J_2|}^{J_1+J_2} \sum_{MN} C_{J_1 M_1 J_2 M_2}^{J M} D_{MN}^J(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J N}, \quad (4.6.1)$$

Here $C_{J_1 M_1 J_2 M_2}^{J M}$ is a Clebsch-Gordan coefficient (Chap. 8). The sum in Eq. (4.6.1) has $2j+1$ terms, where $j = \min(J_1, J_2)$. Equation (4.6.1) may be regarded as a particular case of the expansion of an arbitrary function in a series of the D -functions (Sec. 4.10).

4.6.2 Some Applications of the Clebsch-Gordan Expansion

The Clebsch-Gordan expansion, Eq. (4.6.1), together with the orthogonality condition of the Clebsch-Gordan coefficients, Eq. (8.1.8), enable one to calculate sums of products of the D -functions with identical arguments. Hereafter we introduce the $3j$ -symbol $\{j_1 j_2 j_3\}$ defined by

$$\{j_1 j_2 j_3\} = \begin{cases} 1 & \text{if } j_1 + j_2 + j_3 \text{ is integer and } |j_1 - j_2| \leq j_3 \leq j_1 + j_2, \\ 0 & \text{otherwise,} \end{cases} \quad (4.6.2)$$

$\{j_1 j_2 j_3\}$ is invariant with respect to permutations of j_1, j_2, j_3 . Using Eqs. (4.6.1) and (8.1.8), one obtains

$$\sum_{\substack{M_1 M_2 \\ N_1 N_2}} C_{J_1 M_1 J_2 M_2}^{J M} D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J' N} = \delta_{JJ'} \{J_1 J_2 J\} D_{MN}^J(\alpha, \beta, \gamma), \quad (4.6.3)$$

$$\sum_{J=|J_1-J_2|}^{J_1+J_2} \sum_{M_1 M} C_{J_1 M_1 J_2 M_2}^{J M} D_{M_1 N_1}^{J_1*}(\alpha, \beta, \gamma) D_{MN}^J(\alpha, \beta, \gamma) C_{J_1 N_1' J_2 N_2}^{J N} = \delta_{N_1 N_1'} D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma), \quad (4.6.4)$$

$$\sum_{N_1 N_2} D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J N} = C_{J_1 M_1 J_2 M_2}^{J M} D_{MN}^J(\alpha, \beta, \gamma), \quad (4.6.5)$$

$$\sum_{N_1 N_2 N} D_{MN}^{J*}(\alpha, \beta, \gamma) D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J N} = C_{J_1 M_1 J_2 M_2}^{J M}, \quad (4.6.6)$$

$$\sum_{\substack{M_1 M_2 M \\ N_1 N_2 N}} C_{J_1 M_1 J_2 M_2}^{J M} D_{MN}^{J*}(\alpha, \beta, \gamma) D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J' N'} = \delta_{JJ'} \delta_{NN'} \{J_1 J_2 J\}. \quad (4.6.7)$$

4.6.3 Generalization of the Clebsch-Gordan Expansion

The Clebsch-Gordan expansion can be generalized to the case of an arbitrary number of D -functions of identical arguments by successive use of Eq. (4.6.3). For instance, the summation of products of three D -functions yields

$$\begin{aligned} \sum_{\substack{M_1 M_2 M_3 \\ N_1 N_2 N_3}} C_{J_{12} M_{12} J_3 M_3}^{J M} C_{J_1 M_1 J_2 M_2}^{J_{12} M_{12}} D_{M_1 N_1}^{J_1}(\alpha, \beta, \gamma) D_{M_2 N_2}^{J_2}(\alpha, \beta, \gamma) D_{M_3 N_3}^{J_3}(\alpha, \beta, \gamma) C_{J_1 N_1 J_2 N_2}^{J'_{12} N_{12}} C_{J'_{12} N_{12} J_3 N_3}^{J' N}, \\ = \delta_{JJ'} \delta_{J_{12} J'_{12}} \{J_1 J_2 J_{12}\} \{J_{12} J_3 J\} D_{MN}^J(\alpha, \beta, \gamma). \end{aligned} \quad (4.6.8)$$

In particular, for the case when $J_3 = 0$, Eq. (4.6.8) is reduced to Eq. (4.6.3).

In general, the sum of products of k D -functions is given by

$$\sum_{\substack{m_1, \dots, m_k \\ n_1, \dots, n_k}} \prod_{i=1}^k C_{J_{i-1} M_{i-1} J_i M_i}^{J_i M_i} D_{m_i n_i}^{J_i}(\alpha, \beta, \gamma) C_{J'_{i-1} N_{i-1} J_i N_i}^{J'_i N_i} = D_{M_k N_k}^{J_k}(\alpha, \beta, \gamma) \prod_{i=1}^k \delta_{J_i J'_i} \{j_i J_{i-1} J_i\}, \quad (4.6.9)$$

where $J_i (i = 1, 2, \dots, k)$ is any angular momentum consistent with the vector addition rule,

$$\mathbf{J}_i = \mathbf{j}_1 + \mathbf{j}_2 + \dots + \mathbf{j}_i, \quad M_i = m_1 + m_2 + \dots + m_i, \quad N_i = n_1 + n_2 + \dots + n_i,$$

It is assumed in Eq. (4.6.9) that $J_0 = J'_0 = M_0 = N_0 = 0$.

In particular, for $j_1 = j_2 = \dots = j_k = \frac{1}{2}$ and $J_{i+1} = J_i + \frac{1}{2}$ one has

$$\begin{aligned} \sum_{\substack{m_1 + \dots + m_k = M \\ n_1 + \dots + n_k = N}} D_{m_1 n_1}^{\frac{1}{2}}(\alpha, \beta, \gamma) D_{m_2 n_2}^{\frac{1}{2}}(\alpha, \beta, \gamma) \dots D_{m_k n_k}^{\frac{1}{2}}(\alpha, \beta, \gamma), \\ = \frac{(2J)!}{\sqrt{(J+M)!(J-M)!(J+N)!(J-N)!}} D_{MN}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.6.10)$$

where $J \equiv J_k = k/2$ is either integer or half-integer. Similarly, for $j_1 = j_2 = \dots = j_k = 1$ and $J_{i+1} = J_i + 1$,

$$\begin{aligned} \sum_{\substack{m_1 + \dots + m_k = M \\ n_1 + \dots + n_k = N}} \sqrt{\prod_{i,j=1}^k (1 + \delta_{m_i 0})(1 + \delta_{n_i 0})} D_{m_1 n_1}^1(\alpha, \beta, \gamma) D_{m_2 n_2}^1(\alpha, \beta, \gamma) \dots D_{m_k n_k}^1(\alpha, \beta, \gamma), \\ = \frac{(2J)!}{\sqrt{(J+M)!(J-M)!(J+N)!(J-N)!}} D_{MN}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.6.11)$$

where $J \equiv J_k = k$ is integer.

Equations (4.6.10) and (4.6.11) are useful for evaluating the Wigner D -functions. For example, Eq. (4.6.10) gives an explicit form of the D -functions in terms of the Cayley-Klein parameters (Sec. 1.4.3.) defined by

$$\left. \begin{aligned} D_{\frac{1}{2} \frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma) &\equiv a, & D_{\frac{1}{2} - \frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma) &\equiv -b^*, \\ D_{-\frac{1}{2} \frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma) &\equiv b, & D_{-\frac{1}{2} - \frac{1}{2}}^{\frac{1}{2}}(\alpha, \beta, \gamma) &\equiv a^* \end{aligned} \right\} \quad (4.6.12)$$

The appropriate expression for the D -functions has the form

$$D_{MN}^J(\alpha, \beta, \gamma) = \sqrt{(J+M)!(J-M)!(J+N)!(J-N)!} \sum_{p,q,r,s} \frac{(a)^p (b)^q (a^*)^r (-b^*)^s}{p!q!r!s!}. \quad (4.6.13)$$

Here the summation indices p, q, r and s run over all integer values consistent with the condition

$$\begin{aligned} p + q + r + s &= 2J, \\ p - q - r + s &= 2M, \\ p + q - r - s &= 2N. \end{aligned} \quad (4.6.14)$$

According to Eqs. (4.6.14), only one parameter from p, q, r and s is independent, i.e., in fact, Eq. (4.6.13) represents a single sum. The independent summation index may be chosen in different ways. This yields different explicit forms for the D -functions, Eqs. (4.3.2)-(4.3.5). For example, if r is taken to be independent, Eq. (4.6.13) reduces to Eq. (4.3.2).

4.6.4 Determinant of Matrix $D_{MM'}^J$

The determinant $\|D_{MM'}^J\|$ of the rotation matrix is an invariant sum of products of $2J+1$ D -functions. According to Eq. (4.1.7), this matrix is unimodular, i.e.,

$$\sum_P (-1)^P D_{-JM_1}^J(\alpha, \beta, \gamma) D_{-J+1M_2}^J(\alpha, \beta, \gamma) \dots D_{JM_{2J+1}}^J(\alpha, \beta, \gamma) = 1, \quad (4.6.15)$$

where $M_1, M_2, \dots, M_{2J+1}$ represent all possible permutations P of $J, J-1, J-2, \dots, -J$. The phase factor $(-1)^P$ equals +1 for even permutations, and -1 for odd ones.

4.7 Addition of rotations

4.7.1 The Addition Theorem for $D_{MM'}^J(\alpha, \beta, \gamma)$

Let two successive rotations of the coordinate system, $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ and $S'\{x', y', z'\} \rightarrow S''\{x'', y'', z''\}$, be described by the Euler angles $\alpha_1, \beta_1, \gamma_1$ and $\alpha_2, \beta_2, \gamma_2$, respectively, and the resultant rotation $S\{x, y, z\} \rightarrow S''\{x'', y'', z''\}$ be described by the angles α, β, γ . In accordance with Sec. a, there are two alternative forms of the rotation addition.

- (a) The operator of the resultant rotation, $\hat{D}(\alpha, \beta, \gamma)$, is given by Eq. (1.4.64), if all rotations are performed according to scheme B (Sec. 1.4.1) and the Euler angles $\alpha_2, \beta_2, \gamma_2; \alpha_1, \beta_1, \gamma_1$ and α, β, γ are defined with respect to the initial system $S\{x, y, z\}$. Then the addition theorem reads

$$\sum_{M''=-J}^J D_{MM''}^J(\alpha_2, \beta_2, \gamma_2) D_{M''M'}^J(\alpha_1, \beta_1, \gamma_1) = D_{MM'}^J(\alpha, \beta, \gamma), \quad (4.7.1)$$

where α, β, γ are related to $\alpha_1, \beta_1, \gamma_1$ and $\alpha_2, \beta_2, \gamma_2$ by Eqs. (1.4.66)-(1.4.70).

- (b) The operator of the resultant rotation $\hat{D}(\alpha, \beta, \gamma)$ is given by Eq. (1.4.73), if $\alpha_1, \beta_1, \gamma_1$ and α, β, γ are defined with respect to the initial system $S\{x, y, z\}$, but $\alpha_2, \beta_2, \gamma_2$ are defined with respect to the intermediate system $S'\{x', y', z'\}$ (scheme B), or if successive rotations are performed according to scheme A. In these cases the addition theorem reads

$$\sum_{M''=-J}^J D_{MM''}^J(\alpha_1, \beta_1, \gamma_1) D_{M''M'}^J(\alpha_2, \beta_2, \gamma_2) = D_{MM'}^J(\alpha, \beta, \gamma), \quad (4.7.2)$$

and α, β, γ are related to $\alpha_1, \beta_1, \gamma_1$ and $\alpha_2, \beta_2, \gamma_2$ by the equations which may be obtained from Eqs. (1.4.66)-(1.4.70) by replacing $(\alpha_1, \beta_1, \gamma_1) \rightleftharpoons (\alpha_2, \beta_2, \gamma_2)$.

In particular,

$$\sum_{M''=-J}^J D_{MM''}^J(\alpha, \beta_1, \varphi) D_{M''M'}^J(-\varphi, \beta_2, \gamma) = D_{MM'}^J(\alpha, \beta_1 + \beta_2, \gamma), \quad (4.7.3)$$

where φ is arbitrary, and

$$\begin{aligned} \sum_{M''=-J}^J D_{MM''}^J(\alpha, \beta, \gamma) D_{M''M'}^{J*}(\alpha, \beta, \gamma) &= \sum_{M''=-J}^J D_{MM''}^J(\alpha, \beta, \gamma) D_{M''M'}^J(-\gamma, -\beta, -\alpha) \\ &= D_{MM'}^J(0, 0, 0) = \delta_{MM'}. \end{aligned} \quad (4.7.4)$$

4.7.2 The Addition Theorem for $d_{MM'}^J(\beta)$

Equation (4.7.2) may be rewritten as

$$\sum_{M''=-J}^J d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) e^{-iM''\varphi} = e^{-iM\alpha} d_{MM'}^J(\beta) e^{-iM'\gamma}. \quad (4.7.5)$$

Here

$$\begin{aligned} \cot \alpha &= \cos \beta_1 \cot \varphi + \cot \beta_2 \frac{\sin \beta_1}{\sin \varphi}, \\ \cos \beta &= \cos \beta_1 \cos \beta_2 - \sin \beta_1 \sin \beta_2 \cos \varphi, \\ \cot \gamma &= \cos \beta_2 \cot \varphi + \cot \beta_1 \frac{\sin \beta_2}{\sin \varphi}. \end{aligned} \quad (4.7.6)$$

Equation (4.7.5) is simplified in the following particular cases. If $\varphi = 0$ and $\beta_1 + \beta_2 \leq \pi$, then $\alpha = 0, \beta = \beta_1 + \beta_2$ and $\gamma = 0$,

$$\sum_{M''=-J}^J d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) = d_{MM'}^J(\beta_1 + \beta_2). \quad (4.7.7)$$

If $\varphi = 0$ and $\beta_1 + \beta_2 > \pi$, then $\alpha = \pi, \beta = 2\pi - \beta_1 - \beta_2$ and $\gamma = \pi$,

$$\sum_{M''=-J}^J d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) = (-1)^{M+M'} d_{MM'}^J(2\pi - \beta_1 - \beta_2). \quad (4.7.8)$$

If $\varphi = \pi$ and $\beta_1 \geq \beta_2$, then $\alpha = 0, \beta = \beta_1 - \beta_2$ and $\gamma = \pi$,

$$\sum_{M''=-J}^J (-1)^{M''-M'} d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) = d_{MM'}^J(\beta_1 - \beta_2) \quad (4.7.9)$$

In particular, for $\beta_1 = \beta_2$,

$$\sum_{M''=-J}^J (-1)^{M''-M'} d_{MM''}^J(\beta) d_{M''M'}^J(\beta) = \delta_{MM'}. \quad (4.7.10)$$

If $\varphi = \pi/2$, then

$$\sum_{M''=-J}^J (-i)^{M''} d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) = e^{-iM\alpha} d_{MM'}^J(\beta) e^{-iM'\gamma}. \quad (4.7.11)$$

where

$$\begin{aligned} \cot \alpha &= \cot \beta_2 \sin \beta_1, \\ \cos \beta &= \cos \beta_1 \cos \beta_2, \\ \cot \gamma &= \cot \beta_1 \sin \beta_2. \end{aligned} \quad (4.7.12)$$

4.7.3 Addition of Two Identical Rotations

When $\alpha_1 = \alpha_2, \beta_1 = \beta_2$ and $\gamma_1 = \gamma_2$ the addition theorem, Eq. (4.7.1), together with the Clebsch-Gordan expansion, Eq. (4.6.1), yield

$$\sum_{J=0,1,\dots}^{2j} \sum_{m''} C_{j m j m''}^{J m+m''} D_{m+m'' m''+m'}^J(\alpha, \beta, \gamma) C_{j m'' j m'}^{J m''+m'} = D_{mm'}^j(\bar{\alpha}, \bar{\beta}, \bar{\gamma}), \quad (4.7.13)$$

where

$$\begin{aligned} \cos \bar{\beta} &= \cos^2 \beta - \sin^2 \beta \cos(\alpha + \gamma), \\ \tan(\bar{\alpha} - \alpha) &= \tan(\bar{\gamma} - \gamma) = \frac{\tan \frac{\alpha+\gamma}{2}}{\cos \beta}. \end{aligned} \quad (4.7.14)$$

In particular, if $\alpha = \gamma = 0$,

$$\sum_{J=0,1,\dots}^{2j} \sum_{m''} C_{j m j m''}^{J m+m''} d_{m+m'' m''+m'}^J(\beta) C_{j m'' j m'}^{J m''+m'} = d_{mm'}^j(2\beta). \quad (4.7.15)$$

4.7.4 The Multiplication Theorem for $d_{MM'}^J(\beta)$

Equation (4.7.5) leads to a representation of products of two $d_{MM'}^J(\beta)$ functions

$$d_{MM''}^J(\beta_1) d_{M''M'}^J(\beta_2) = \frac{1}{4\pi} \int_{-2\pi}^{2\pi} e^{i(M''\varphi - M\alpha - M'\gamma)} d_{MM'}^J(\beta) d\varphi. \quad (4.7.16)$$

Here α, β, γ are related to β_1, β_2 and φ by means of Eq. (4.7.6).

4.7.5 Sums Involving the D-Functions of Different Arguments

In addition to Eqs. (4.7.1) or (4.7.2) one can derive the following invariant sums which are equivalent to the characters of irreducible representations of rotation group for two rotations (Sec. 4.14):

(a)

$$\sum_{MM'} D_{MM'}^J(\alpha_1, \beta_1, \gamma_1) D_{M'M}^J(\alpha_2, \beta_2, \gamma_2) \equiv \chi^J(R_1 R_2) = \chi^J(R_2 R_1) = \frac{\sin[(2J+1)\frac{\omega}{2}]}{\sin \frac{\omega}{2}}, \quad (4.7.17)$$

where

$$\begin{aligned} \cos \frac{\omega}{2} &= \cos \frac{\beta_1}{2} \cos \frac{\beta_2}{2} \cos \frac{\alpha_1 + \gamma_1 + \alpha_2 + \gamma_2}{2} - \sin \frac{\beta_1}{2} \sin \frac{\beta_2}{2} \cos \frac{\alpha_1 - \gamma_1 - \alpha_2 + \gamma_2}{2}, \\ &= \cos \frac{\beta_1 + \beta_2}{2} \cos \frac{\alpha_1 + \gamma_2}{2} \cos \frac{\gamma_1 + \alpha_2}{2} - \cos \frac{\beta_1 - \beta_2}{2} \sin \frac{\alpha_1 + \gamma_2}{2} \sin \frac{\gamma_1 + \alpha_2}{2}. \end{aligned} \quad (4.7.18)$$

(b)

$$\sum_{MM'} D_{MM'}^J(\alpha_1, \beta_1, \gamma_1) D_{MM'}^{J*}(\alpha_2, \beta_2, \gamma_2) \equiv \chi^J(R_1 R_2^{-1}) = \chi^J(R_1^{-1} R_2) = \frac{\sin[(2J+1)\frac{\omega'}{2}]}{\sin \frac{\omega'}{2}} \quad (4.7.19)$$

where

$$\begin{aligned} \cos \frac{\omega'}{2} &= \cos \frac{\beta_1}{2} \cos \frac{\beta_2}{2} \cos \frac{\alpha_1 + \gamma_1 - \alpha_2 - \gamma_2}{2} + \sin \frac{\beta_1}{2} \sin \frac{\beta_2}{2} \cos \frac{\alpha_1 - \gamma_1 - \alpha_2 + \gamma_2}{2}, \\ &= \cos \frac{\beta_1 - \beta_2}{2} \cos \frac{\alpha_1 - \alpha_2}{2} \cos \frac{\gamma_1 - \gamma_2}{2} - \cos \frac{\beta_1 + \beta_2}{2} \sin \frac{\alpha_1 - \alpha_2}{2} \sin \frac{\gamma_1 - \gamma_2}{2}. \end{aligned} \quad (4.7.20)$$

Equation (4.7.20) may be obtained from Eq. (4.7.18) by replacing $(\alpha_2, \beta_2, \gamma_2) \rightarrow (-\gamma_2, -\beta_2, -\alpha_2)$.

Similarly, one can arrive at invariant sums which may be reduced to the characters involving three and more rotations, i.e., $\chi^J(R_1 R_2 R_3)$ etc.

4.7.6 The Ponzano-Regge sum

Note the following sum involving products of three $d_{MM'}^J(\beta)$ functions with the same J but different arguments (Ref. [46])

$$\sum_{J=J_{\min}}^{\infty} (2J+1) d_{M_1 M_2}^J(\beta_3) d_{M_2 M_3}^J(\beta_1) d_{M_3 M_1}^J(\beta_2) = \frac{2\Theta(B)}{\pi\sqrt{B}} \cos\left(\sum_{i=1}^3 M_i \delta_i\right). \quad (4.7.21)$$

Here the summation index J runs from $J_{\min} = \max\{|M_1|, |M_2|, |M_3|\}$ to infinity, M_1, M_2, M_3 being fixed, and

$$B = \begin{vmatrix} 1 & \cos \beta_3 & \cos \beta_2 \\ \cos \beta_3 & 1 & \cos \beta_1 \\ \cos \beta_2 & \cos \beta_1 & 1 \end{vmatrix}, \quad (4.7.22)$$

$$\Theta(x) = \begin{cases} 1 & x > 0 \\ 0 & x < 0 \end{cases} \quad (4.7.23)$$

The relations between the angles $\delta_1, \delta_2, \delta_3$ and $\beta_1, \beta_2, \beta_3$ are

$$\cos \delta_i = \frac{\cos \beta_j \cos \beta_k - \cos \beta_i}{\sin \beta_j \sin \beta_k}, \quad (i \neq j \neq k), \quad (4.7.24)$$

$$\frac{\sin \beta_i}{\sin \delta_i} = \frac{\sin \beta_j}{\sin \delta_j}, \quad (i, j = 1, 2, 3). \quad (4.7.25)$$

These relations may be easily obtained by using spherical trigonometry (see Fig. 4.1).




Figure 4.1. Geometrical interpretation of the angles in Eq. (4.7.21). The arguments $\beta_1, \beta_2, \beta_3$ are the sides of a spherical triangle, i.e. the angles they subtend at the centre of the sphere, whose three rays are shown emanating from the centre. The interior angle of the triangle at the vertex opposite the side β_i equals $\pi - \delta_i$. The δ_i are thereby determined by $\beta_1, \beta_2, \beta_3$ through the spherical cosine and sine rules (4.7.24)–(4.7.25).

4.8 Recursion relations for $D_{MM'}^J$

4.8.1 Relations between D^J and $D^{J\pm 1}$

$$\begin{aligned} \cos \beta D_{MM'}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J^2 - M^2)(J^2 - M'^2)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) + \frac{MM'}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\ &+ \frac{\sqrt{((J + 1)^2 - M^2)[(J + 1)^2 - M'^2]}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma) \end{aligned} \quad (4.8.1)$$

$$\begin{aligned} \sin \beta e^{i\alpha} D_{M+1M'}^J(\alpha, \beta, \gamma) &= - \frac{\sqrt{(J + M)(J + M + 1)(J^2 - M'^2)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\ &+ \frac{M' \sqrt{(J - M)(J + M + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\ &+ \frac{\sqrt{(J - M)(J - M + 1)[(J + 1)^2 - M'^2]}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma) \end{aligned} \quad (4.8.2)$$

$$\begin{aligned} \sin \beta e^{-i\alpha} D_{M-1M'}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J - M)(J - M + 1)(J^2 - M'^2)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\ &+ \frac{M' \sqrt{(J + M)(J - M + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\ &- \frac{\sqrt{(J + M)(J + M + 1)[(J + 1)^2 - M'^2]}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma). \end{aligned} \quad (4.8.3)$$

$$\begin{aligned} \sin \beta e^{i\gamma} D_{MM'+1}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J^2 - M^2)(J + M')(J + M' + 1)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\ &- \frac{M \sqrt{(J - M')(J + M' + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\ &- \frac{\sqrt{((J + 1)^2 - M^2)(J - M')(J - M' + 1)}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma), \end{aligned} \quad (4.8.4)$$

$$\begin{aligned}
\sin \beta e^{-i\gamma} D_{MM'-1}^J(\alpha, \beta, \gamma) &= -\frac{\sqrt{(J^2 - M^2)(J - M')(J - M' + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma)}{J(2J + 1)} \\
&\quad - \frac{M\sqrt{(J + M')(J - M' + 1)} D_{MM'}^J(\alpha, \beta, \gamma)}{J(J + 1)} \\
&\quad + \frac{\sqrt{[(J + 1)^2 - M^2](J + M')(J + M' + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma)}{(J + 1)(2J + 1)}, \tag{4.8.5}
\end{aligned}$$

$$\begin{aligned}
(1 + \cos \beta) e^{i(\alpha + \gamma)} D_{M+1M'+1}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J + M + 1)(J + M)(J + M' + 1)(J + M')}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J - M)(J + M + 1)(J - M')(J + M' + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J - M)(J - M + 1)(J - M')(J - M' + 1)}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma), \tag{4.8.6}
\end{aligned}$$

$$\begin{aligned}
(1 + \cos \beta) e^{-i(\alpha + \gamma)} D_{M-1M'-1}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J - M)(J - M + 1)(J - M')(J - M' + 1)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J + M)(J - M + 1)(J + M')(J - M' + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J + M)(J + M + 1)(J + M')(J + M' + 1)}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma), \tag{4.8.7}
\end{aligned}$$

$$\begin{aligned}
(1 - \cos \beta) e^{i(\alpha - \gamma)} D_{M+1M'-1}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J + M)(J + M + 1)(J - M')(J - M' + 1)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\
&\quad - \frac{\sqrt{(J - M)(J + M + 1)(J + M')(J - M' + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J - M)(J - M + 1)(J + M')(J + M' + 1)}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma), \tag{4.8.8}
\end{aligned}$$

$$\begin{aligned}
(1 - \cos \beta) e^{-i(\alpha - \gamma)} D_{M-1M'+1}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J - M)(J - M + 1)(J + M')(J + M' + 1)}}{J(2J + 1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) \\
&\quad - \frac{\sqrt{(J + M)(J - M + 1)(J - M')(J + M' + 1)}}{J(J + 1)} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J + M)(J + M + 1)(J - M')(J - M' + 1)}}{(J + 1)(2J + 1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma). \tag{4.8.9}
\end{aligned}$$

Equations (4.8.1)-(4.8.9) may be obtained from the Clebsch-Gordan series, Eq. (4.6.1) with $J_1 = 1$.

4.8.2 Relations Between D^J and $D^{J \pm 1/2}$

$$\begin{aligned}
\cos \frac{\beta}{2} e^{i\frac{\alpha + \gamma}{2}} D_{M+\frac{1}{2}M'+\frac{1}{2}}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J + M + \frac{1}{2})(J + M' + \frac{1}{2})}}{2J + 1} D_{MM'}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J - M + \frac{1}{2})(J - M' + \frac{1}{2})}}{2J + 1} D_{MM'}^{J+\frac{1}{2}}(\alpha, \beta, \gamma), \tag{4.8.10}
\end{aligned}$$

$$\begin{aligned}
\sin \frac{\beta}{2} e^{i\frac{\alpha - \gamma}{2}} D_{M+\frac{1}{2}M'-\frac{1}{2}}^J(\alpha, \beta, \gamma) &= -\frac{\sqrt{(J + M + \frac{1}{2})(J - M' + \frac{1}{2})}}{2J + 1} D_{MM'}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\
&\quad + \frac{\sqrt{(J - M + \frac{1}{2})(J + M' + \frac{1}{2})}}{2J + 1} D_{MM'}^{J+\frac{1}{2}}(\alpha, \beta, \gamma), \tag{4.8.11}
\end{aligned}$$

$$\begin{aligned} \cos \frac{\beta}{2} e^{-i\frac{\alpha+\gamma}{2}} D_{M-\frac{1}{2}M'-\frac{1}{2}}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J-M+\frac{1}{2})(J-M'+\frac{1}{2})}}{2J+1} D_{MM'}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\ &\quad + \frac{\sqrt{(J+M+\frac{1}{2})(J+M'+\frac{1}{2})}}{2J+1} D_{MM'}^{J+\frac{1}{2}}(\alpha, \beta, \gamma), \end{aligned} \quad (4.8.12)$$

$$\begin{aligned} \sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} D_{M-\frac{1}{2}M'+\frac{1}{2}}^J(\alpha, \beta, \gamma) &= \frac{\sqrt{(J-M+\frac{1}{2})(J+M'+\frac{1}{2})}}{2J+1} D_{MM'}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\ &\quad - \frac{\sqrt{(J+M+\frac{1}{2})(J-M'+\frac{1}{2})}}{2J+1} D_{MM'}^{J+\frac{1}{2}}(\alpha, \beta, \gamma). \end{aligned} \quad (4.8.13)$$

Equations (4.8.10)-(4.8.13) may be obtained from Eq. (4.6.1) with $J_1 = \frac{1}{2}$. They yield

$$\begin{aligned} D_{MM'}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{J-M}{J-M'}} \cos \frac{\beta}{2} e^{i\frac{\alpha+\gamma}{2}} D_{M+\frac{1}{2}M'+\frac{1}{2}}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\ &\quad - \sqrt{\frac{J+M}{J-M'}} \sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} D_{M-\frac{1}{2}M'+\frac{1}{2}}^{J-\frac{1}{2}}(\alpha, \beta, \gamma), \quad (M' \neq J), \end{aligned} \quad (4.8.14)$$

$$\begin{aligned} D_{MM'}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{J-M}{J+M'}} \sin \frac{\beta}{2} e^{i\frac{\alpha-\gamma}{2}} D_{M+\frac{1}{2}M'-\frac{1}{2}}^{J-\frac{1}{2}}(\alpha, \beta, \gamma) \\ &\quad + \sqrt{\frac{J+M}{J+M'}} \cos \frac{\beta}{2} e^{-i\frac{\alpha+\gamma}{2}} D_{M-\frac{1}{2}M'-\frac{1}{2}}^{J-\frac{1}{2}}(\alpha, \beta, \gamma), \quad (M' \neq -J). \end{aligned} \quad (4.8.15)$$

4.8.3 Relations Between $D_{MM'}^J$ and $D_{M\pm 1M'\mp 1}^J$

$$\begin{aligned} \frac{-M+M'\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) &= \frac{1}{2} \sqrt{(J+M')(J-M'+1)} D_{MM'-1}^J(\alpha, \beta, \gamma) e^{-i\gamma} \\ &\quad + \frac{1}{2} \sqrt{(J-M')(J+M'+1)} D_{MM'+1}^J(\alpha, \beta, \gamma) e^{i\gamma}, \end{aligned} \quad (4.8.16)$$

$$\begin{aligned} \frac{M'-M\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) &= \frac{1}{2} \sqrt{(J+M)(J-M+1)} D_{M-1M'}^J(\alpha, \beta, \gamma) e^{-i\alpha} \\ &\quad + \frac{1}{2} \sqrt{(J-M)(J+M+1)} D_{M+1M'}^J(\alpha, \beta, \gamma) e^{i\alpha}, \end{aligned} \quad (4.8.17)$$

$$\begin{aligned} D_{M+1M'}^J(\alpha, \beta, \gamma) e^{i\alpha} &= \sqrt{\frac{(J+M')(J-M'+1)}{(J-M)(J+M+1)}} \frac{1+\cos\beta}{2} e^{-i\gamma} D_{MM'-1}^J(\alpha, \beta, \gamma) \\ &\quad + \frac{M'\sin\beta}{\sqrt{(J-M)(J+M+1)}} D_{MM'}^J(\alpha, \beta, \gamma) \\ &\quad - \sqrt{\frac{(J-M')(J+M'+1)}{(J-M)(J+M+1)}} \frac{1-\cos\beta}{2} e^{i\gamma} D_{MM'+1}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.8.18)$$

$$\begin{aligned} D_{M-1M'}^J(\alpha, \beta, \gamma) e^{-i\alpha} &= -\sqrt{\frac{(J+M')(J-M'+1)}{(J+M)(J-M+1)}} \frac{1-\cos\beta}{2} e^{-i\gamma} D_{MM'-1}^J(\alpha, \beta, \gamma) \\ &\quad + \frac{M'\sin\beta}{\sqrt{(J+M)(J-M+1)}} D_{MM'}^J(\alpha, \beta, \gamma) \\ &\quad + \sqrt{\frac{(J-M')(J+M'+1)}{(J+M)(J-M+1)}} \frac{1+\cos\beta}{2} e^{i\gamma} D_{MM'+1}^J(\alpha, \beta, \gamma), \end{aligned} \quad (4.8.19)$$

$$\begin{aligned}
D_{MM'+1}^J(\alpha, \beta, \gamma)e^{i\gamma} &= \sqrt{\frac{(J+M)(J-M+1)}{(J-M')(J+M'+1)}} \frac{1+\cos\beta}{2} e^{-i\alpha} D_{M-1M'}^J(\alpha, \beta, \gamma) \\
&\quad - \frac{M \sin\beta}{\sqrt{(J-M')(J+M'+1)}} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad - \sqrt{\frac{(J-M)(J+M+1)}{(J-M')(J+M'+1)}} \frac{1-\cos\beta}{2} e^{i\alpha} D_{M+1M'}^J(\alpha, \beta, \gamma), \tag{4.8.20}
\end{aligned}$$

$$\begin{aligned}
D_{MM'-1}^J(\alpha, \beta, \gamma)e^{-i\gamma} &= -\sqrt{\frac{(J+M)(J-M+1)}{(J+M')(J-M'+1)}} \frac{1-\cos\beta}{2} e^{-i\alpha} D_{M-1M'}^J(\alpha, \beta, \gamma) \\
&\quad - \frac{M \sin\beta}{\sqrt{(J+M')(J-M'+1)}} D_{MM'}^J(\alpha, \beta, \gamma) \\
&\quad + \sqrt{\frac{(J-M)(J+M+1)}{(J+M')(J-M'+1)}} \frac{1+\cos\beta}{2} e^{i\alpha} D_{M+1M'}^J(\alpha, \beta, \gamma). \tag{4.8.21}
\end{aligned}$$

4.9 Differential relations for $D_{MM'}^J(\alpha, \beta, \gamma)$

Derivatives of $D_{MM'}^J(\alpha, \beta, \gamma)$ may be expressed in terms of the D -functions with different M, M' and J

$$\begin{aligned}
\sin\beta \frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) &= -\frac{(J+1)\sqrt{(J^2-M^2)(J^2-M'^2)}}{J(2J+1)} D_{MM'}^{J-1}(\alpha, \beta, \gamma) - \frac{MM'}{J(J+1)} D_{MM'}^J(\alpha, \beta, \gamma), \\
&\quad + \frac{J\sqrt{[(J+1)^2-M^2][(J+1)^2-M'^2]}}{(J+1)(2J+1)} D_{MM'}^{J+1}(\alpha, \beta, \gamma), \tag{4.9.1}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) &= -\frac{1}{2} \sqrt{(J+M)(J-M+1)} e^{-i\alpha} D_{M-1M'}^J(\alpha, \beta, \gamma) \\
&\quad + \frac{1}{2} \sqrt{(J-M)(J+M+1)} e^{i\alpha} D_{M+1M'}^J(\alpha, \beta, \gamma), \tag{4.9.2}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) &= \frac{1}{2} \sqrt{(J+M')(J-M'+1)} e^{-i\gamma} D_{MM'-1}^J(\alpha, \beta, \gamma) \\
&\quad - \frac{1}{2} \sqrt{(J-M')(J+M'+1)} e^{i\gamma} D_{MM'+1}^J(\alpha, \beta, \gamma), \tag{4.9.3}
\end{aligned}$$

$$\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) = \frac{M'-M\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) - \sqrt{(J+M)(J-M+1)} e^{-i\alpha} D_{M-1M'}^J(\alpha, \beta, \gamma), \tag{4.9.4}$$

$$\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) = -\frac{M'-M\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) + \sqrt{(J-M)(J+M+1)} e^{i\alpha} D_{M+1M'}^J(\alpha, \beta, \gamma), \tag{4.9.5}$$

$$\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) = \frac{M-M'\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) + \sqrt{(J+M')(J-M'+1)} e^{-i\gamma} D_{MM'-1}^J(\alpha, \beta, \gamma), \tag{4.9.6}$$

$$\frac{\partial}{\partial\beta} D_{MM'}^J(\alpha, \beta, \gamma) = -\frac{M-M'\cos\beta}{\sin\beta} D_{MM'}^J(\alpha, \beta, \gamma) - \sqrt{(J-M')(J+M'+1)} e^{i\gamma} D_{MM'+1}^J(\alpha, \beta, \gamma), \tag{4.9.7}$$

$$\frac{\partial}{\partial\alpha} D_{MM'}^J(\alpha, \beta, \gamma) = -iM D_{MM'}^J(\alpha, \beta, \gamma), \tag{4.9.8}$$

$$\frac{\partial}{\partial\gamma} D_{MM'}^J(\alpha, \beta, \gamma) = -iM' D_{MM'}^J(\alpha, \beta, \gamma), \tag{4.9.9}$$

See also Eqs. (4.2.1) and (4.2.2).

4.10 Orthogonality and completeness of the D -functions

The functions $D_{MM'}^J(\alpha, \beta, \gamma)$ with different J (integer and half-integer) are mutually orthogonal with respect to integration over a double volume of the 3-dimensional rotation group (i.e., over the volume of SU_2 group) because the period of $D_{MM'}^J(\alpha, \beta, \gamma)$ with half-integer J is 4π rather than 2π . The double domain may be chosen in two ways:

$$V_1 : 0 \leq \alpha < 4\pi, 0 \leq \beta \leq \pi, 0 \leq \gamma < 2\pi, \quad (4.10.1)$$

$$V_2 : 0 \leq \alpha < 2\pi, 0 \leq \beta \leq \pi, 0 \leq \gamma < 4\pi. \quad (4.10.2)$$

The total volume of the double domain V_1 or V_2 is equal to $16\pi^2$.

Orthogonality and normalization condition is

$$\begin{aligned} & \int_0^{4\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_2 M_2'}^{J_2*}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma), \\ &= \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{4\pi} d\gamma D_{M_2 M_2'}^{J_2*}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) = \frac{16\pi^2}{2J_1 + 1} \delta_{J_1 J_2} \delta_{M_1 M_2} \delta_{M_1' M_2'}. \end{aligned} \quad (4.10.3)$$

If J_1 and J_2 are both either integer or half-integer, orthogonality of the D -functions takes place at integration over the single volume of rotation group which corresponds to the domain

$$V : 0 \leq \alpha < 2\pi, 0 \leq \beta \leq \pi, 0 \leq \gamma < 2\pi. \quad (4.10.4)$$

The total volume of the domain V is equal to $8\pi^2$. Hence, the orthogonality condition (4.10.3) is reduced to

$$\int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_2 M_2'}^{J_2*}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) = \frac{8\pi^2}{2J_1 + 1} \delta_{J_1 J_2} \delta_{M_1 M_2} \delta_{M_1' M_2'}. \quad (4.10.5)$$

In physical applications Eq. (4.10.5) is used more often than Eq. (4.10.3).

The orthonormality of $D_{MM'}^J(\alpha, \beta, \gamma)$ may be rewritten in terms of $d_{MM'}^J(\beta)$

$$\int_0^\pi d\beta \sin \beta d_{MM'}^J(\beta) d_{MM'}^{J'}(\beta) = \frac{2}{2J + 1} \delta_{JJ'}. \quad (4.10.6)$$

Thus, the collection of functions $D_{MM'}^J(\alpha, \beta, \gamma)$ with integer and half-integer J constitutes a complete set of orthonormalized functions. The completeness condition is

$$\begin{aligned} & \sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \sum_{M=-J}^J \sum_{M'=-J}^J \frac{2J+1}{16\pi^2} D_{MM'}^{J*}(\alpha_1, \beta_1, \gamma_1) D_{MM'}^J(\alpha_2, \beta_2, \gamma_2) \\ &= \delta(\alpha_1 - \alpha_2) \delta(\cos \beta_1 - \cos \beta_2) \delta(\gamma_1 - \gamma_2). \end{aligned} \quad (4.10.7)$$

Any function $f(\alpha, \beta, \gamma)$ which is defined in the domain V_1 (or V_2) and satisfies the condition

$$\iiint_{V_1(V_2)} d\alpha d\beta \sin \beta d\gamma |f(\alpha, \beta, \gamma)|^2 < \infty, \quad (4.10.8)$$

may be expanded in a series of the D -functions

$$f(\alpha, \beta, \gamma) = \sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \sum_{M=-J}^J \sum_{M'=-J}^J a_{MM'}^J D_{MM'}^J(\alpha, \beta, \gamma). \quad (4.10.9)$$

The expansion coefficients $a_{MM'}^J$ are determined by

$$a_{MM'}^J = \frac{2J+1}{16\pi^2} \iiint_{V_1(V_2)} d\alpha d\beta \sin \beta d\gamma f(\alpha, \beta, \gamma) D_{MM'}^{J*}(\alpha, \beta, \gamma), \quad (4.10.10)$$

and obey the relation

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \frac{16\pi^2}{2J+1} \sum_{M=-J}^J \sum_{M'=-J}^J |a_{MM'}^J|^2 = \iiint_{V_1(V_2)} d\alpha d\beta \sin \beta d\gamma |f(\alpha, \beta, \gamma)|^2. \quad (4.10.11)$$

If a function $f(\alpha, \beta, \gamma)$ is defined in the domain V , it may be expanded in a series of the D -functions with only integer or only half-integer J

$$f(\alpha, \beta, \gamma) = \sum_{\substack{J \text{ integer} \\ \text{or half-integer}}} \sum_{M=-J}^J \sum_{M'=-J}^J b_{MM'}^J D_{MM'}^J(\alpha, \beta, \gamma). \quad (4.10.12)$$

The expansion coefficients $b_{MM'}^J$ are determined by

$$b_{MM'}^J = \frac{2J+1}{8\pi^2} \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma f(\alpha, \beta, \gamma) D_{MM'}^{J*}(\alpha, \beta, \gamma) \quad (4.10.13)$$

for both cases of integer and half-integer J . A difference between the expansions of $f(\alpha, \beta, \gamma)$ in these two cases occurs if one uses the expansions outside of the domain V . The expansion coefficients in Eq. (4.10.12) satisfy the relation

$$\sum_{\substack{J \text{ integer} \\ \text{or half-integer}}} \frac{8\pi^2}{2J+1} \sum_{M=-J}^J \sum_{M'=-J}^J |b_{MM'}^J|^2 = \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma |f(\alpha, \beta, \gamma)|^2. \quad (4.10.14)$$

4.11 Integrals involving the D -functions

4.11.1 Integration of Products of $D_{MM'}^J$

$$\int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{MM'}^J(\alpha, \beta, \gamma) = \delta_{J0} \delta_{M0} \delta_{M'0} 8\pi^2, \quad (J \text{ is integer}), \quad (4.11.1)$$

$$\int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) D_{M_2 M_2'}^{J_2}(\alpha, \beta, \gamma) = (-1)^{M_2 - M_2'} \frac{8\pi^2}{2J_2 + 1} \delta_{J_1 J_2} \delta_{-M_1 M_2} \delta_{-M_1' M_2'}, \quad (J_1 + J_2 \text{ is integer}), \quad (4.11.2)$$

$$\int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_2 M_2'}^{J_2*}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) = \frac{8\pi^2}{2J_2 + 1} \delta_{J_1 J_2} \delta_{M_1 M_2} \delta_{M_1' M_2'}, \quad (J_1 + J_2 \text{ is integer}), \quad (4.11.3)$$

$$\begin{aligned} & \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_3 M_3'}^{J_3}(\alpha, \beta, \gamma) D_{M_2 M_2'}^{J_2}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) \\ &= (-1)^{M_3 - M_3'} \frac{8\pi^2}{2J_3 + 1} C_{J_1 M_1 J_2 M_2}^{J_3 - M_3} C_{J_1 M_1' J_2 M_2'}^{J_3 - M_3'} \quad (J_1 + J_2 + J_3 \text{ is integer}), \end{aligned} \quad (4.11.4)$$

$$\begin{aligned} & \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{M_3 M_3'}^{J_3*}(\alpha, \beta, \gamma) D_{M_2 M_2'}^{J_2}(\alpha, \beta, \gamma) D_{M_1 M_1'}^{J_1}(\alpha, \beta, \gamma) \\ &= \frac{8\pi^2}{2J_3 + 1} C_{J_1 M_1 J_2 M_2}^{J_3 M_3} C_{J_1 M_1' J_2 M_2'}^{J_3 M_3'}, \quad (J_1 + J_2 + J_3 \text{ is integer}) \end{aligned} \quad (4.11.5)$$

Equations (4.11.1)-(4.11.5) are valid, if the conditions in parentheses are satisfied. These conditions may be omitted, if the integration domain is V_1 or V_2 instead of V ; in the latter case the factor $8\pi^2$ on the right-hand side of Eqs. (4.11.1)-(4.11.5) must be replaced by $16\pi^2$ (see Sec. 4.10).

Integrals involving products of four or more D -functions may be reduced to Eqs. (4.11.4)-(4.11.5) by using the Clebsch-Gordan expansion (Sec. 4.6.1).

4.11.2 Integrals Involving $d_{MM'}^J(\beta)$

From Eqs. (4.11.1)-(4.11.5) one may obtain

$$\int_0^\pi d\beta \sin \beta d_{00}^J(\beta) = 2\delta_{J0}, \quad (4.11.6)$$

$$\int_0^\pi d\beta \sin \beta d_{MM'}^J(\beta) d_{MM'}^{J'}(\beta) = \frac{2}{2J+1} \delta_{JJ'}, \quad (4.11.7)$$

$$\int_0^\pi d\beta \sin \beta d_{M_1 M_1'}^{J_1}(\beta) d_{M_2 M_2'}^{J_2}(\beta) d_{M_3 M_3'}^{J_3}(\beta) \delta_{M_1 + M_2 M_3} \delta_{M_1' + M_2' M_3'} = \frac{2}{2J_3 + 1} C_{J_1 M_1 J_2 M_2}^{J_3 M_3} C_{J_1 M_1' J_2 M_2'}^{J_3 M_3'}. \quad (4.11.8)$$

Note also the following relation

$$\begin{aligned} & \int_0^\beta d\beta \left(\sin \frac{\beta}{2} \right)^{M-M'+1} \left(\cos \frac{\beta}{2} \right)^{M+M'+1} d_{MM'}^J(\beta) \\ &= \frac{-1}{\sqrt{J(J+1) - M(M+1)}} \left(\sin \frac{\beta}{2} \right)^{M-M'+1} \left(\cos \frac{\beta}{2} \right)^{M+M'+1} d_{M+1M'}^J(\beta). \end{aligned} \quad (4.11.9)$$

Equation (4.11.9) is valid, if $M \geq M' \geq 0$. Other cases may be deduced from Eq. (4.11.9) using the symmetries of $d_{MM'}^J(\beta)$ (Eq. (4.4.1)).

4.12 Invariant summation of integrals involving $D_{MM'}^J(\alpha, \beta, \gamma)$

Hereafter R will denote the Euler angles α, β, γ and we also will set $dR = \sin \beta d\beta d\alpha d\gamma$:

$$\int dR f(R) \equiv \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma f(\alpha, \beta, \gamma). \quad (4.12.1)$$

Making use of this notation, one gets

$$\sum_{MM'} \int dR D_{MM'}^{J*}(R) D_{M'M'}^{J'}(R) = 8\pi^2 \delta_{JJ'}, \quad (4.12.2)$$

$$\sum_{MM'} \int dR D_{MM'}^J(R) D_{M'M}^{J'}(R) = (-1)^{2J} 8\pi^2 \delta_{JJ'}, \quad (4.12.3)$$

$$\sum_{M_1 M_2 M_3} \int dR D_{M_3 M_3}^{J_3}(R) D_{M_2 M_2}^{J_2}(R) D_{M_1 M_1}^{J_1}(R) = 8\pi^2 \{J_1 J_2 J_3\}, \quad (4.12.4)$$

$$\begin{aligned} & \sum_{\text{All } M, N} \int dR_1 D_{M_1 N_1}^{J_1}(R_1) D_{M_2 N_2}^{J_2}(R_1) D_{M_3 M_3}^{J_3}(R_1) \int dR_2 D_{N_1 M_1}^{J_1}(R_2) D_{M_2 N_2}^{J_2*}(R_2) D_{M_3' M_3'}^{J_3'}(R_2) \\ &= (-1)^{2J_3'} (8\pi^2)^2 \left\{ \begin{matrix} J_1 & J_2 & J_3 \\ J_1 & J_2 & J_3' \end{matrix} \right\}, \end{aligned} \quad (4.12.5)$$

$$\begin{aligned} & \sum_{\text{All } M, N} \int dR_1 D_{M_1' N_1'}^{J_1'*}(R_1) D_{M_2' N_2'}^{J_2'}(R_1) D_{M_3 M_3}^{J_3}(R_1) \int dR_2 D_{M_2' N_2'}^{J_2'*}(R_2) D_{M_3' N_3'}^{J_3'}(R_2) D_{M_1 M_1}^{J_1}(R_2) \\ & \times \int dR_3 D_{M_3' N_3'}^{J_3'*}(R_3) D_{M_1' N_1'}^{J_1'}(R_3) D_{M_2 M_2}^{J_2}(R_3) = (8\pi^2)^3 \left\{ \begin{matrix} J_1 & J_2 & J_3 \\ J_1' & J_2' & J_3' \end{matrix} \right\}^2, \end{aligned} \quad (4.12.6)$$

$$\begin{aligned} & \sum_{\text{All } M, N} \int dR_1 D_{MM}^J(R_1) D_{N_2 N_2'}^{J_2}(R_1) D_{N_3 N_3'}^{J_3}(R_1) \int dR_2 D_{M' M'}^{J'}(R_2) D_{N_1 M_1}^{J_1}(R_2) D_{N_2' M_2'}^{J_2'}(R_2) \\ & \times \int dR_3 D_{M'' M''}^{J''}(R_3) D_{M_2' M_2'}^{J_2'}(R_3) D_{N_2' M_3}^{J_3}(R_3) \int dR_4 D_{M_1 N_1}^{J_1}(R_4) D_{M_2 N_2}^{J_2}(R_4) D_{M_2' M_2}^{J_2'}(R_4) D_{M_3 N_3}^{J_3}(R_4) \\ &= (-1)^{2J_1} (8\pi^2)^4 \left\{ \begin{matrix} J_1 & J_2 & J' \\ J_3 & J_2 & J \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_2 & J'' \\ J_3 & J_2 & J' \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J_2 & J \\ J_3 & J_2 & J'' \end{matrix} \right\}, \end{aligned} \quad (4.12.7)$$

$$\begin{aligned} & \sum_{\text{All } M, N} \int dR_1 D_{MM}^J(R_1) D_{M' N'}^{J'}(R_1) D_{M_1 N_1}^{J_1*}(R_1) \int dR_2 D_{N' M'}^{J'}(R_2) D_{N N}^{J_1}(R_2) D_{M_2 N_2}^{J_2*}(R_2) \\ & \times \int dR_3 D_{M_1 N_1}^{J_1}(R_3) D_{M_2 N_2}^{J_2}(R_3) D_{M_3 M_3}^{J_3*}(R_3) = (8\pi^2)^3 \left\{ \begin{matrix} J & J & J_1 \\ J & J & J_2 \\ J_1 & J_2 & J_3 \end{matrix} \right\}, \end{aligned} \quad (4.12.8)$$

$$\begin{aligned}
& \sum_{\text{All } M, N} \int dR_1 D_{M_1 N_1}^{J_1}(R_1) D_{M_2 N_2}^{J_2}(R_1) D_{M_{12} M_{12}}^{J_{12}}(R_1) \int dR_2 D_{M_3 N_3}^{J_3}(R_2) D_{M_4 N_4}^{J_4}(R_2) D_{M_{34} M_{34}}^{J_{34}}(R_2) \\
& \times \int dR_3 D_{M_{13} N_{13}}^{J_{13}}(R_3) D_{M_{24} N_{24}}^{J_{24}}(R_3) D_{MM}^J(R_3) \int dR_4 D_{N_1 M_1}^{J_1}(R_4) D_{N_3 M_3}^{J_3}(R_4) D_{N_{13} M_{13}}^{J_{13}}(R_4) \\
& \times \int dR_5 D_{N_2 M_2}^{J_2}(R_5) D_{N_4 M_4}^{J_4}(R_5) D_{N_{24} M_{24}}^{J_{24}}(R_5) = (8\pi^2)^5 \left\{ \begin{matrix} J_1 & J_2 & J_{12} \\ J_3 & J_4 & J_{34} \\ J_{13} & J_{24} & J \end{matrix} \right\}^2. \quad (4.12.9)
\end{aligned}$$

The left-hand sides of Eqs. (4.12.5)-(4.12.9) include integrations as well as summations. If the integrations are carried out before the summations (with the aid of Eqs. (4.11.4)-(4.11.5)), then Eqs. (4.12.5)-(4.12.9) are reduced to sums of the Clebsch-Gordan coefficients (Sec. 8.7). On the other hand, if the summations are performed before the integrations, one obtains the integral representations of $6j$ - and $9j$ -symbols (see Secs. 9.3 and 10.3).

4.13 Generating functions for $d_{MM}^J(\beta)$

The functions $d_{MM'}^J(\beta)$ may be defined as coefficients of expansions of various generating functions.

(a) If J and M' are fixed, then

$$\begin{aligned}
& \left(\cos \frac{\beta}{2} e^{i\varphi/2} + i \sin \frac{\beta}{2} e^{-i\varphi/2} \right)^{J+M'} \left(i \sin \frac{\beta}{2} e^{i\varphi/2} + \cos \frac{\beta}{2} e^{-i\varphi/2} \right)^{J-M'} \\
& = \sum_{M=-J}^J \sqrt{\frac{(J-M')!(J+M')!}{(J-M)!(J+M)!}} (-i)^{M-M'} e^{iM\varphi} d_{MM'}^J(\beta) \quad (4.13.1)
\end{aligned}$$

In particular, for $\varphi = 0$

$$e^{iJ\beta} = \sum_{M=-J}^J \sqrt{\frac{(J-M')!(J+M')!}{(J-M)!(J+M)!}} (-i)^{M-M'} d_{MM'}^J(\beta) \quad (4.13.2)$$

and for $\varphi = \pi$

$$e^{-iJ\beta} = \sum_{M=-J}^J \sqrt{\frac{(J-M')!(J+M')!}{(J-M)!(J+M)!}} i^{M-M'} d_{MM'}^J(\beta) \quad (4.13.3)$$

(b) If M and M' are fixed, one can obtain several expressions in which s, μ, ν are related to J, M, M' by Eqs. (4.3.14), $\xi_{MM'}$ is defined by Eq. (4.3.15); and R denotes

$$R = \sqrt{1 - 2t \cos \beta + t^2}, \quad |t| < 1. \quad (4.13.4)$$

$$\frac{\xi_{MM'}}{R} \left(\frac{2 \sin \frac{\beta}{2}}{1 + R - t} \right)^\mu \left(\frac{2 \cos \frac{\beta}{2}}{1 + R + t} \right)^\nu = \sum_{s=0}^{\infty} \sqrt{\frac{(s+\mu)!(s+\nu)!}{s!(s+\mu+\nu)!}} t^s d_{MM'}^{s+\frac{\mu+\nu}{2}}(\beta). \quad (4.13.5)$$

$$\begin{aligned}
& \xi_{MM'} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu {}_0F_1 \left(; 1 + \mu; -t \sin^2 \frac{\beta}{2} \right) {}_0F_1 \left(; 1 + \nu; t \cos^2 \frac{\beta}{2} \right) \\
& = \sum_{s=0}^{\infty} \frac{\mu! \nu!}{\sqrt{s!(s+\mu)!(s+\nu)!(s+\mu+\nu)!}} t^s d_{MM'}^{s+\frac{\mu+\nu}{2}}(\beta), \quad (4.13.6)
\end{aligned}$$

$$\begin{aligned}
& \xi_{MM'} (1-t)^{1-\mu-\nu} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu {}_2F_1 \left(\frac{\mu+\nu+1}{2}, \frac{\mu+\nu+2}{2}; 1 + \mu; -\frac{4t \sin^2 \frac{\beta}{2}}{(1-t)^2} \right) \\
& = \sum_{s=0}^{\infty} \sqrt{\frac{(s+\nu)!(s+\mu+\nu)!}{s!(s+\mu)!}} \frac{\mu!}{(\mu+\nu)!} t^s d_{MM'}^{s+\frac{\mu+\nu}{2}}(\beta), \quad (4.13.7)
\end{aligned}$$

$$\begin{aligned}
\xi_{MM'} \left(\sin \frac{\beta}{2} \right)^\mu \left(\cos \frac{\beta}{2} \right)^\nu {}_2F_1 \left(\lambda, \mu + \nu + 1 - \lambda; 1 + \mu; \frac{1-t-R}{2} \right) {}_2F_1 \left(\lambda, \mu + \nu + 1 - \lambda; 1 + \nu; \frac{1+t-R}{2} \right) \\
= \sum_{s=0}^{\infty} \frac{(\mu + \nu + 1 - \lambda)_s (\lambda)_s \mu! \nu!}{\sqrt{s!(s + \mu + \nu)!(s + \mu)!(s + \nu)!}} t^s d_{MM'}^{s + \frac{\mu + \nu}{2}}(\beta).
\end{aligned} \tag{4.13.8}$$

In Eq. (4.13.8) λ is an arbitrary integer.

4.14 Characters $\chi^J(R)$ of irreducible representations of rotation group

4.14.1 Definition

The trace of the finite rotation matrix in the JM-representation

$$\chi^J(R) = \sum_{M=-J}^J D_{MM}^J(R), \tag{4.14.1}$$

is called the characteristic function, or simply, the character of the irreducible representation of rank J .

In contrast to $D_{MM'}^J(R)$, the function $\chi^J(R)$ is invariant under rotations of the coordinate systems. Explicit forms of $\chi^J(R)$ are simpler when R is specified by ω, Θ, Φ rather than by α, β, γ (Sec. 1.4.2). With variables ω, Θ, Φ in use, $\chi^J(R)$ is entirely determined by the rotation angles ω and is independent of the rotation axis $\mathbf{n}(\Theta, \Phi)$:

$$\chi^J(R) = \chi^J(\omega). \tag{4.14.2}$$

4.14.2 Explicit Forms

(a) *Trigonometric formulas*

$$\chi^J(\omega) = \frac{\sin \left[(2J+1) \frac{\omega}{2} \right]}{\sin \frac{\omega}{2}}, \tag{4.14.3}$$

$$\chi^J(\omega) = \sum_{M=-J}^J e^{-iM\omega} = \sum_{M=-J}^J \cos M\omega, \tag{4.14.4}$$

$$\chi^J(\omega) = \sum_{n=0}^{[J]} (-1)^n \frac{(2J-n)!}{(2J-2n)!n!} \left(2 \cos \frac{\omega}{2} \right)^{2J-2n}, \tag{4.14.5}$$

$$\chi^J(\omega) = \sum_{n=0}^{[J]} (-1)^n \frac{(2J+1)!}{(2n+1)!(2J-2n)!} \left(\cos \frac{\omega}{2} \right)^{2J-2n} \left(\sin \frac{\omega}{2} \right)^{2n}, \tag{4.14.6}$$

$$\chi^J(\omega) = \frac{1}{2J+1} \frac{d}{d \left(\cos \frac{\omega}{2} \right)} \cos \left[(2J+1) \frac{\omega}{2} \right], \tag{4.14.7}$$

$$\chi^J(\omega) = \frac{(2J+1)! 2^{2J}}{(4J+1)! \sin \frac{\omega}{2}} \left[-\frac{d}{d \left(\cos \frac{\omega}{2} \right)} \right]^{2J} \left(\sin \frac{\omega}{2} \right)^{4J+1}. \tag{4.14.8}$$

(b) *Relations to hypergeometric functions*

$$\chi^J(\omega) = (2J+1) F \left(-J, J+1; 3/2; \sin^2 \frac{\omega}{2} \right), \tag{4.14.9}$$

$$\chi^J(\omega) = (2J+1) F \left(-2J, 2(J+1); 3/2; \sin^2 \frac{\omega}{4} \right). \tag{4.14.10}$$

(c) *Relation to the Chebyshev polynomials of the second kind*

$$\chi^J(\omega) = U_{2J} \left(\cos \frac{\omega}{2} \right). \tag{4.14.11}$$

(d) *Relation to the Gegenbauer polynomials*

$$\chi^J(\omega) = C_{2J}^1 \left(\cos \frac{\omega}{2} \right). \tag{4.14.12}$$

(e) *Relation to the Jacobi polynomials*

$$\chi^J(\omega) = \frac{(4J+2)!!}{2(4J+1)!!} P_{2J}^{\left(\frac{1}{2}, \frac{1}{2}\right)}\left(\cos \frac{\omega}{2}\right). \quad (4.14.13)$$

(f) *Integral representation*

$$\chi^J(\omega) = \frac{2J+1}{2} \int_{-1}^1 \left(\cos \frac{\omega}{2} + ix \sin \frac{\omega}{2}\right)^{2J} dx. \quad (4.14.14)$$

To express $\chi^J(R)$ in terms of the Euler angles α, β, γ one can use the relation

$$\cos \frac{\omega}{2} = \cos \frac{\beta}{2} \cos \frac{\alpha + \gamma}{2}. \quad (4.14.15)$$

4.14.3 Principal Properties

In contrast to $D_{MM'}^J(R)$, the characters $\chi^J(R)$ are real.

$$(\chi^J(R))^* = \chi^J(R). \quad (4.14.16)$$

The characters which correspond to direct and inverse rotations, R and R^{-1} , are equal

$$\chi^J(R^{-1}) = \chi^J(R). \quad (4.14.17)$$

$\chi^J(R)$ is invariant with respect to coordinate rotation and inversion

$$\chi^J(URU^{-1}) = \chi^J(R), \quad (4.14.18)$$

where U is any orthogonal transformation of the coordinate system.

The characters which depend on the combined rotations $R_1 R_2 \dots R_n$ do not change under cyclic permutations of the rotations

$$\chi^J(R_1 R_2 \dots R_n) = \chi^J(R_2 \dots R_n R_1). \quad (4.14.19)$$

In particular,

$$\chi^J(R_1 R_2) = \chi^J(R_2 R_1), \quad (4.14.20)$$

in spite of the non-commutativity of R_1 and R_2 .

Products of the characters may be expanded in a Clebsch-Gordan series

$$\chi^{J_1}(R) \chi^{J_2}(R) = \sum_J \{J_1 J_2 J\} \chi^J(R) \quad (4.14.21)$$

where $\{J_1 J_2 J\} = 1$, if the triangle inequality (Sec. 4.6.2) is satisfied, and $\{J_1 J_2 J\} = 0$ otherwise.

The transformation $J = \tilde{J} = -J - 1$ reverses the character sign

$$\chi^{\tilde{J}}(R) = -\chi^J(R) \quad (4.14.22)$$

The function $\chi^J(\omega)$ is even and periodic

$$\begin{aligned} \chi^J(-\omega) &= \chi^J(\omega), \\ \chi^J(\omega + 4\pi) &= \chi^J(\omega), \\ \chi^J(\omega + 2\pi) &= (-1)^{2J} \chi^J(\omega). \end{aligned} \quad (4.14.23)$$

(a) *The addition theorem*

The character which depends on the combined rotation $R_1 R_2$ may be represented as a superposition of products of the generalized characters which depend on R_1 and R_2 (Sec. 4.15):

$$\chi^J(\omega) = \sum_{\lambda=0}^{2J} (-1)^\lambda \frac{2\lambda+1}{2J+1} \chi_\lambda^J(\omega_1) \chi_\lambda^J(\omega_2) P_\lambda(\cos \Theta_{12}), \quad (4.14.24)$$

where ω_1, ω_2 and ω are the rotation angles corresponding to R_1, R_2 and $R = R_1 R_2$, respectively. These angles are mutually related by means of Eqs. (1.4.75); Θ_{12} is the angle between the rotation axes of R_1 and R_2

$$\cos \Theta_{12} = (\mathbf{n}_1 \cdot \mathbf{n}_2) = \cos \Theta_1 \cos \Theta_2 + \sin \Theta_1 \sin \Theta_2 \cos(\Phi_1 - \Phi_2). \quad (4.14.25)$$

4.14.4 Differential Equation

The character $\chi^J(\omega)$ satisfies the equation

$$\frac{d^2}{d\omega^2}\chi^J(\omega) + \cot \frac{\omega}{2} \frac{d}{d\omega}\chi^J(\omega) + J(J+1)\chi^J(\omega) = 0. \quad (4.14.26)$$

and the boundary conditions

$$\begin{aligned} \chi^J(0) &= 2J+1, \\ \chi^J(\omega \pm 4\pi n) &= \chi^J(\omega), \quad \text{where } n \text{ is integer.} \end{aligned} \quad (4.14.27)$$

4.14.5 Differential Relations

$$\frac{d}{d\omega}\chi^J(\omega) = -\sqrt{J(J+1)}\chi_1^J(\omega); \quad (4.14.28)$$

$$\left(\frac{d}{d\cos \frac{\omega}{2}}\right)^k \chi^J(\omega) = \frac{1}{\sqrt{2J+1}} \sqrt{\frac{(2J+k+1)!}{(2J-k)!}} \frac{\chi_k^J(\omega)}{(\sin \frac{\omega}{2})^k}, \quad (4.14.29)$$

where $\chi_k^J(\omega)$ is the generalized character (Sec. 4.15:

$$\sin \frac{\omega}{2} \frac{d}{d\omega}\chi^J(\omega) = J \cos \frac{\omega}{2} \chi^J(\omega) - \left(J + \frac{1}{2}\right) \chi^{J-\frac{1}{2}}(\omega) = \left(J + \frac{1}{2}\right) \chi^{J+\frac{1}{2}}(\omega) - (J+1) \cos \frac{\omega}{2} \chi^J(\omega). \quad (4.14.30)$$

4.14.6 Algebraic Relations

$$\chi^{J+\frac{1}{2}}(\omega) = 2 \cos \frac{\omega}{2} \cdot \chi^J(\omega) - \chi^{J-\frac{1}{2}}(\omega), \quad (4.14.31)$$

$$\chi^{J_1}(\omega) - \chi^{J_2}(\omega) = 2\chi^{\frac{J_1+J_2-1}{2}}(\omega) \cos \left[(J_1 + J_2 + 1) \frac{\omega}{2} \right], \quad (4.14.32)$$

$$\chi^{J_1}(\omega) + \chi^{J_2}(\omega) = 2\chi^{\frac{J_1+J_2}{2}}(\omega) \cos \left[(J_1 - J_2) \frac{\omega}{2} \right]. \quad (4.14.33)$$

Equations (4.14.32) and (4.14.33) are valid, if $J_1 + J_2$ is integer.

$$\chi^{J-\frac{1}{2}}(\omega) = 2 \cos \frac{J\omega}{2} \chi^{\frac{J-1}{2}}(\omega), \quad (4.14.34)$$

where J is integer and positive.

$$-2 \sin^2 \frac{\omega}{2} \chi^{J_1}(\omega) \chi^{J_2}(\omega) = \cos [(J_1 + J_2 + 1)\omega] - \cos [(J_1 - J_2)\omega]. \quad (4.14.35)$$

In particular,

$$2 \sin^2 \frac{\omega}{2} [\chi^J(\omega)]^2 = 1 - \cos[(2J+1)\omega] \quad (4.14.36)$$

$$\chi^J(\omega) = 2^{2J} \prod_{k=1}^{2J} \sin \left(\frac{\omega}{2} + \frac{k\pi}{2J+1} \right). \quad (4.14.37)$$

4.14.7 Orthogonality and Completeness

The collection of characters $\chi^J(\omega)$ with different J constitute a complete set of orthogonal functions of argument ω in the domain $0 \leq \omega < 2\pi$.

The orthogonality and normalization condition for these functions reads

$$\int_0^{2\pi} \chi^{J_1}(\omega) \chi^{J_2}(\omega) \sin^2 \frac{\omega}{2} d\omega = \pi \delta_{J_1 J_2}. \quad (4.14.38)$$

The completeness condition has the form

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \chi^J(\omega_1) \chi^J(\omega_2) = \frac{\pi \delta(\omega_1 - \omega_2)}{\sin^2 \frac{\omega_1}{2}}. \quad (4.14.39)$$

4.14.8 Integrals Involving $\chi^J(\omega)$

$$\int_0^{2\pi} d\omega \sin^2 \frac{\omega}{2} \chi^J(\omega) = \pi \delta_{J0}, \quad (4.14.40)$$

$$\int_0^{2\pi} d\omega \sin^2 \frac{\omega}{2} \chi^J(2\omega) = \pi(-1)^{2J}, \quad (4.14.41)$$

$$\int_0^{2\pi} d\omega \frac{\sin^2 \frac{\omega}{2}}{\cos \frac{\omega}{2} - \cos \frac{\Omega}{2}} \chi^J(\omega) = -2\pi \cos \left[(2J+1) \frac{\Omega}{2} \right], \quad (4.14.42)$$

In Eq. (4.14.42) the Cauchy principal value of the integral is assumed.

$$\int_0^{2\pi} d\omega \sin^2 \frac{\omega}{2} \chi^{J_1}(\omega) \chi^{J_2}(\omega) = \pi \delta_{J_1 J_2}, \quad (4.14.43)$$

$$\int_0^{2\pi} d\omega \sin^2 \frac{\omega}{2} \chi^{J_1}(\omega) \chi^{J_2}(\omega) \chi^{J_3}(\omega) = \pi \{J_1 J_2 J_3\}. \quad (4.14.44)$$

4.14.9 Sums Involving $\chi^J(\omega)$

(a) *Finite sums*

$$\sum_{J=J_1, J_1+1, \dots}^{J_2} \chi^J(\omega) = \frac{\sin \left[(J_2 + J_1 + 1) \frac{\omega}{2} \right] \sin \left[(J_2 - J_1 + 1) \frac{\omega}{2} \right]}{\sin^2 \frac{\omega}{2}} = \chi^{\frac{J_2+J_1}{2}}(\omega) \chi^{\frac{J_2-J_1}{2}}(\omega), \quad (4.14.45)$$

$$\sum_{J=0, 1, 2, \dots}^{J_0} (2J+1) \chi^J(\omega) = \frac{(2J_0+3) \sin \left[(2J_0+1) \frac{\omega}{2} \right] - (2J_0+1) \sin \left[(2J_0+3) \frac{\omega}{2} \right]}{4 \sin^3 \frac{\omega}{2}}. \quad (4.14.46)$$

The summation index in Eq. (4.14.45) runs over integer values, if J_1 and J_2 are integers, or half-integer values, if J_1 and J_2 are half-integers.

In the equations given below the summation indices run over both integer and half-integer values:

$$\sum_{J=J_1, J_1+\frac{1}{2}, J_1+1, \dots}^{J_2} \chi^J(\omega) = \frac{\sin \left[(J_2 + J_1 + 1) \frac{\omega}{2} \right] \sin \left[(J_2 - J_1 + \frac{1}{2}) \frac{\omega}{2} \right]}{\sin \frac{\omega}{2} \sin \frac{\omega}{4}}, \quad (4.14.47)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{J_0} (2J+1) \chi^J(\omega) = \frac{(2J_0+2) \sin \left[(2J_0+1) \frac{\omega}{2} \right] - (2J_0+1) \sin \left[(2J_0+2) \frac{\omega}{2} \right]}{4 \sin \frac{\omega}{2} \sin^2 \frac{\omega}{4}}, \quad (4.14.48)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{J_0} [\chi^J(\omega)]^2 = \frac{(4J_0+3) \sin \frac{\omega}{2} - \sin \left[(4J_0+3) \frac{\omega}{2} \right]}{4 \sin^3 \frac{\omega}{2}}, \quad (4.14.49)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{J_0} \chi^J(\omega) \chi^J(\omega') = \frac{\chi^{J_0+\frac{1}{2}}(\omega) \chi^{J_0}(\omega') - \chi^{J_0}(\omega) \chi^{J_0+\frac{1}{2}}(\omega')}{2 \left(\cos \frac{\omega}{2} - \cos \frac{\omega'}{2} \right)}, \quad (4.14.50)$$

$$\begin{aligned} & \sum_{J=0, \frac{1}{2}, 1, \dots}^{J_0} \chi^J(\omega) \cos \left[(2J+1) \frac{\omega'}{2} \right] \\ &= \frac{\sin \frac{\omega}{2} - \cos \left[(2J_0+1) \frac{\omega'}{2} \right] \sin \left[(J_0+1) \omega \right] + \sin \left[(2J_0+1) \frac{\omega}{2} \right] \cos \left[(J_0+1) \omega' \right]}{2 \sin \frac{\omega}{2} \left(\cos \frac{\omega'}{2} - \cos \frac{\omega}{2} \right)}. \end{aligned} \quad (4.14.51)$$

(b) *Infinite series*

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \chi^J(\omega) = \frac{1}{4 \sin^2 \frac{\omega}{4}}, \quad (\omega \neq 0), \quad (4.14.52)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} (2J+1) \chi^J(\omega) = \frac{\pi}{\sin^2 \frac{\omega}{2}} \delta(\omega). \quad (4.14.53)$$

Let us introduce the notation

$$R^2 \equiv 1 - 2t \cos \frac{\omega}{2} + t^2, \quad \text{where } |t| < 1 : \quad (4.14.54)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} t^{2J} \chi^J(\omega) = \frac{1}{R^2}, \quad (4.14.55)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} (2J+1) t^{2J} \chi^J(\omega) = \frac{1-t^2}{R^4}, \quad (4.14.56)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \frac{(4J+1)!!}{(4J+2)!!} t^{2J} \chi^J(\omega) = \frac{1}{R \sqrt{2(1-t \cos \frac{\omega}{2} + R)}}, \quad (4.14.57)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \frac{1}{(2J+1)!} t^{2J+1} \chi^J(\omega) = \frac{\sin(t \sin \frac{\omega}{2})}{\sin \frac{\omega}{2}} e^{t \cos \frac{\omega}{2}}, \quad (4.14.58)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \frac{1}{2J+1} t^{2J} \chi^J(\omega) = \frac{1}{2it \sin \frac{\omega}{2}} \ln \left(\frac{1 - te^{-\frac{i\omega}{2}}}{1 - te^{\frac{i\omega}{2}}} \right), \quad (4.14.59)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \sum_{\Gamma(\nu)}^{\Gamma(2J+\nu)} \frac{t^{2J} \chi^J(\omega)}{(2J+1)!} = \left(1 - t \cos \frac{\omega}{2}\right)^{-\nu} {}_2F_1 \left(\frac{\nu}{2}, \frac{\nu+1}{2}; \frac{3}{2}; -\left(\frac{t \sin \frac{\omega}{2}}{1 - t \cos \frac{\omega}{2}} \right)^2 \right), \quad (4.14.60)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \frac{1}{(4J+2)!} (4t)^{2J} \chi^J(\omega) = \frac{1}{2} {}_0F_1 \left(\frac{3}{2}; -t \sin^2 \frac{\omega}{4} \right) {}_0F_1 \left(\frac{3}{2}; t \cos^2 \frac{\omega}{4} \right), \quad (4.14.61)$$

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} t^{2J} \chi^J(\omega) \chi^J(\omega') = \frac{1-t^2}{(1+t^2)^2 - 4t(1+t^2) \cos \frac{\omega}{2} \cos \frac{\omega'}{2} + 2t^2 (\cos \omega + \cos \omega')}. \quad (4.14.62)$$

4.14.10 $\chi^J(\omega)$ for Particular Values of ω

$$\chi^J(0) = 2J+1, \quad (4.14.63)$$

$$\chi^J(2\pi) = (-1)^{2J} (2J+1), \quad (4.14.64)$$

$$\chi^J(\pi) = \begin{cases} 0 & \text{if } J \text{ is half-integer} \\ (-1)^J & \text{if } J \text{ is integer} \end{cases} \quad (4.14.65)$$

$$\chi^J\left(\frac{\pi}{2}\right) = \begin{cases} \sqrt{2} & J = 1/2, 9/2, 17/2, \dots \\ 1 & J = 0, 1, 4, 5, 8, 9, \dots \\ 0 & J = 3/2, 7/2, 11/2, 15/2, \dots \\ -1 & J = 2, 3, 6, 7, 10, 11, \dots \\ -\sqrt{2} & J = 5/2, 13/2, 21/2, \dots \end{cases} \quad (4.14.66)$$

4.14.11 Special Cases of $\chi^J(\omega)$

$$\chi^0(\omega) = 1, \quad (4.14.67)$$

$$\chi^{\frac{1}{2}}(\omega) = 2 \cos \frac{\omega}{2}, \quad (4.14.68)$$

$$\chi^1(\omega) = 4 \cos^2 \frac{\omega}{2} - 1, \quad (4.14.69)$$

$$\chi^{\frac{3}{2}}(\omega) = 8 \cos^3 \frac{\omega}{2} - 4 \cos \frac{\omega}{2}, \quad (4.14.70)$$

$$\chi^2(\omega) = 16 \cos^4 \frac{\omega}{2} - 12 \cos^2 \frac{\omega}{2} + 1, \quad (4.14.71)$$

$$\chi^{\frac{5}{2}}(\omega) = 32 \cos^5 \frac{\omega}{2} - 32 \cos^3 \frac{\omega}{2} + 6 \cos \frac{\omega}{2}. \quad (4.14.72)$$

4.15 Generalized characters, $\chi_\lambda^J(R)$, of irreducible representations of the rotation group

4.15.1 Definition

Let us introduce the function $\chi_\lambda^J(\omega)$ associated with $\chi^J(\omega)$ by the differential relation

$$\chi_\lambda^J(\omega) = \sqrt{2J+1} \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!}} \left(\sin \frac{\omega}{2}\right)^\lambda \left(\frac{d}{d \cos \frac{\omega}{2}}\right)^\lambda \chi^J(\omega), \quad (4.15.1)$$

where λ is integer, $0 \leq \lambda \leq 2J$, and $\chi^J(\omega)$ is the character of the irreducible representation of rank J (Eq. (4.14.1)).

The function $\chi_\lambda^J(\omega)$ will be called the generalized character (of order λ) of the irreducible representation of rank J . Note that the relations between $\chi_\lambda^J(\omega)$ and $\chi^J(\omega)$ are similar to those between the associated Legendre functions $P_l^\lambda(x)$ and the Legendre polynomials $P_l(x)$.

At $\lambda = 0$ one has $\chi_\lambda^J(\omega) = \chi^J(\omega)$.

4.15.2 Explicit Forms

(a) *Trigonometric series*

$$\chi_\lambda^J(\omega) = i^\lambda \sum_M e^{-iM\omega} C_{JM\lambda 0}^{JM}, \quad (4.15.2)$$

$$\chi_\lambda^J(\omega) = \left(\sin \frac{\omega}{2}\right)^\lambda \sqrt{2J+1} \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!}} 2^\lambda \sum_{s=0}^{|J-\lambda/2|} \frac{(-1)^s (2J-s)!}{s!(2J-\lambda-2s)!} \left(2 \cos \frac{\omega}{2}\right)^{2J-\lambda-2s}, \quad (4.15.3)$$

$$\chi_\lambda^J(\omega) = \frac{(2 \sin \frac{\omega}{2})^\lambda}{\lambda!} \sqrt{2J+1} \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!}} \sum_{s=0}^{2J-\lambda} \frac{(\lambda+s)!(2J-s)!}{s!(2J-\lambda-s)!} \cos \left[(2J-\lambda-2s) \frac{\omega}{2} \right]. \quad (4.15.4)$$

(b) *Differential form*

$$\chi_\lambda^J(\omega) = \frac{1}{\sqrt{2J+1}} \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!}} \left(\sin \frac{\omega}{2}\right)^\lambda \left(\frac{d}{d \cos \frac{\omega}{2}}\right)^{\lambda+1} \cos \left[(2J+1) \frac{\omega}{2} \right]. \quad (4.15.5)$$

See also Eq. (4.15.1).

(c) *Relation between $\chi_\lambda^J(\omega)$ and the Gegenbauer polynomials*

$$\chi_\lambda^J(\omega) = (2\lambda)!! \sqrt{2J+1} \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!}} \left(\sin \frac{\omega}{2}\right)^\lambda C_{2J-\lambda}^{\lambda+1} \left(\cos \frac{\omega}{2}\right). \quad (4.15.6)$$

(d) *Relations of $\chi_\lambda^J(\omega)$ to the Jacobi polynomials*

$$\chi_\lambda^J(\omega) = \frac{\sqrt{2J+1} \sqrt{(2J-\lambda)!(2J+\lambda+1)!}}{(4J+1)!!} 2^{2J-\lambda} \left(\sin \frac{\omega}{2}\right)^\lambda P_{2J-\lambda}^{\left(\lambda+\frac{1}{2}, \lambda+\frac{1}{2}\right)} \left(\cos \frac{\omega}{2}\right), \quad (4.15.7)$$

$$\chi_\lambda^J(\omega) = \begin{cases} \sqrt{2J+1} \sqrt{\frac{(2J+\lambda)!!(2J-\lambda)!!}{(2J+\lambda+1)!!(2J-\lambda-1)!!}} \left(\sin \frac{\omega}{2}\right)^\lambda P_{J-\lambda/2}^{\left(\lambda+\frac{1}{2}, -\frac{1}{2}\right)}(\cos \omega), & 2J-\lambda \text{ is even,} \\ \sqrt{2J+1} \sqrt{\frac{(2J-\lambda-1)!!(2J+\lambda+1)!!}{(2J-\lambda)!!(2J+\lambda)!!}} \left(\cos \frac{\omega}{2}\right) \left(\sin \frac{\omega}{2}\right)^\lambda P_{J-\frac{\lambda+1}{2}}^{\left(\lambda+\frac{1}{2}, \frac{1}{2}\right)}(\cos \omega), & 2J-\lambda \text{ is odd.} \end{cases} \quad (4.15.8)$$

(e) *Relations of $\chi_\lambda^J(\omega)$ to the Hypergeometric functions*

$$\chi_\lambda^J(\omega) = \frac{\sqrt{2J+1}}{(2\lambda+1)!!} \sqrt{\frac{(2J+\lambda+1)!}{(2J-\lambda)!}} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-2J+\lambda, 2J+\lambda+2; \lambda+\frac{3}{2}; \sin^2 \frac{\omega}{4}\right), \quad (4.15.9)$$

$$\chi_\lambda^J(\omega) = \frac{(-1)^{2J-\lambda}(2\lambda)!!\sqrt{2J+1}}{(2\lambda+1)!} \sqrt{\frac{(2J+\lambda+1)!}{(2J-\lambda)!}} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-2J+\lambda, 2J+\lambda+2; \lambda+\frac{3}{2}; \cos^2 \frac{\omega}{4}\right), \quad (4.15.10)$$

$$\chi_\lambda^J(\omega) = \frac{(-1)^{2J-\lambda}\sqrt{2J+1}2^{4J-\lambda}(2J)!}{\sqrt{(2J-\lambda)!(2J+\lambda+1)!}} \left(\sin \frac{\omega}{2}\right)^\lambda \left(\sin \frac{\omega}{4}\right)^{4J-2\lambda} F\left(-2J+\lambda, -2J-\frac{1}{2}; -4J-1; \frac{1}{\sin^2 \frac{\omega}{2}}\right), \quad (4.15.11)$$

$$\chi_\lambda^J(\omega) = \frac{\sqrt{2J+1}(4J)!!}{\sqrt{(2J-\lambda)!(2J+\lambda+1)!}} \left(\cos \frac{\omega}{2}\right)^{2J-\lambda} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-J+\frac{1}{2}, \frac{-2J+\lambda+1}{2}; -2J; \frac{1}{\cos^2 \frac{\omega}{2}}\right), \quad (4.15.12)$$

$$\chi_\lambda^J(\omega) = \frac{\sqrt{2J+1}}{(2\lambda+1)!!} \sqrt{\frac{(2J+\lambda+1)!}{(2J-\lambda)!}} \left(\sin \frac{\omega}{2}\right)^\lambda \left(\cos \frac{\omega}{4}\right)^{4J-2\lambda} F\left(-2J+\lambda, -2J-\frac{1}{2}; \lambda+\frac{3}{2}; -\tan^2 \frac{\omega}{4}\right), \quad (4.15.13)$$

$$\chi_\lambda^J(\omega) = \frac{\sqrt{2J+1}(2J)!i^\lambda}{\sqrt{(2J-\lambda)!(2J+\lambda+1)!}} \left(e^{-\frac{i\omega}{2}} - e^{\frac{i\omega}{2}}\right)^\lambda e^{\pm i\frac{\omega}{2}(2J-\lambda)} F\left(-2J+\lambda, \lambda+1; -2J; e^{\mp i\omega}\right), \quad (4.15.14)$$

$$\chi_\lambda^J(\omega) = \begin{cases} (-1)^{J+\frac{3\lambda}{2}} \sqrt{2J+1} \sqrt{\frac{(2J-\lambda-1)!!(2J+\lambda)!!}{(2J-\lambda)!!(2J+\lambda+1)!!}} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-J+\frac{\lambda}{2}, J+1+\frac{\lambda}{2}; \frac{1}{2}; \cos^2 \frac{\omega}{2}\right), \\ \quad \text{if } 2J-\lambda \text{ is even,} \\ (-1)^{J+\frac{3\lambda-1}{2}} \sqrt{2J+1} \sqrt{\frac{(2J-\lambda)!!(2J+\lambda+1)!!}{(2J-\lambda-1)!!(2J+\lambda)!!}} \cos \frac{\omega}{2} \left(\sin \frac{\omega}{2}\right)^\lambda, \\ \quad \times F\left(-J+\frac{\lambda+1}{2}, J+1+\frac{\lambda+1}{2}; \frac{3}{2}; \cos^2 \frac{\omega}{2}\right), \\ \quad \text{if } 2J-\lambda \text{ is odd,} \end{cases} \quad (4.15.15)$$

$$\chi_\lambda^J(\omega) = \begin{cases} \frac{\sqrt{2J+1}}{(2\lambda+1)!!} \sqrt{\frac{(2J+\lambda+1)!}{(2J-\lambda)!}} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-J+\frac{\lambda}{2}, J+1+\frac{\lambda}{2}; \lambda+\frac{3}{2}; \sin^2 \frac{\omega}{2}\right), \\ \quad \text{if } 2J-\lambda \text{ is even,} \\ \frac{\sqrt{2J+1}}{(2\lambda+1)!!} \sqrt{\frac{(2J+\lambda+1)!}{(2J-\lambda)!}} \cos \frac{\omega}{2} \left(\sin \frac{\omega}{2}\right)^\lambda F\left(-J+\frac{\lambda+1}{2}, J+1+\frac{\lambda+1}{2}; \lambda+\frac{3}{2}; \sin^2 \frac{\omega}{2}\right), \\ \quad \text{if } 2J-\lambda \text{ is odd.} \end{cases} \quad (4.15.16)$$

(f) *Integral representations*

$$\chi_\lambda^J(\omega) = \frac{1}{2} \frac{1}{\lambda!} \sqrt{\frac{(2J+1)(2J+\lambda+1)!}{(2J-\lambda)!}} \frac{1}{(\sin \frac{\omega}{2})^{\lambda+1}} \int_0^\omega \cos \left[(2J+1)\frac{\psi}{2}\right] \left(\cos \frac{\psi}{2} - \cos \frac{\omega}{2}\right)^\lambda d\psi, \quad (4.15.17)$$

$$\chi_\lambda^J(\omega) = (-i)^\lambda \frac{\sqrt{(2J+1)(2J+\lambda+1)!(2J-\lambda)!}}{2(2J)!} \int_{-1}^{+1} P_\lambda(x) \left[\cos \frac{\omega}{2} + ix \sin \frac{\omega}{2}\right]^{2J} dx. \quad (4.15.18)$$

4.15.3 Principal Properties

(a) *Symmetries*

$$\chi_\lambda^{J*}(\omega) = \chi_\lambda^J(\omega) = (-1)^\lambda \chi_\lambda^J(-\omega), \quad (4.15.19)$$

$$\chi_\lambda^J(\omega + 2\pi n) = (-1)^{2Jn} \chi_\lambda^J(\omega), \quad \chi_\lambda^J(2\pi - \omega) = (-1)^{2J-\lambda} \chi_\lambda^J(\omega). \quad (4.15.20)$$

(b) *Particular values of ω*

$$\begin{aligned}\chi_\lambda^J(0) &= (2J+1)\delta_{\lambda 0}, & \chi_\lambda^J(2\pi) &= (-1)^{2J}(2J+1)\delta_{\lambda 0}, \\ \chi_\lambda^J(\pi) &= \begin{cases} (-1)^{J-\frac{\lambda}{2}} \sqrt{\frac{(2J+\lambda)!!(2J-\lambda-1)!!(2J+1)}{(2J-\lambda)!!(2J+\lambda+1)!!}} & \text{if } 2J-\lambda \text{ is even,} \\ 0 & \text{if } 2J-\lambda \text{ is odd.} \end{cases}\end{aligned}\quad (4.15.21)$$

(c) *Recursion relations*

$$2\frac{d}{d\omega}\chi_\lambda^J(\omega) = \frac{\lambda}{2\lambda+1}\sqrt{(2J+1)^2-\lambda^2}\chi_{\lambda-1}^J(\omega) - \frac{\lambda+1}{2\lambda+1}\sqrt{(2J+1)^2-(\lambda+1)^2}\chi_{\lambda+1}^J(\omega), \quad (4.15.22)$$

$$(2\lambda+1)\cot\frac{\omega}{2}\chi_\lambda^J(\omega) = \sqrt{(2J+1)^2-\lambda^2}\chi_{\lambda-1}^J(\omega) + \sqrt{(2J+1)^2-(\lambda+1)^2}\chi_{\lambda+1}^J(\omega). \quad (4.15.23)$$

(d) *Asymptotics*If $\omega \rightarrow 0$ and $J \rightarrow \infty$, while $J\omega \equiv x < \infty$, then

$$\lim_{\substack{J \rightarrow \infty \\ \omega \rightarrow 0}} \frac{1}{2J+1} \chi_\lambda^J(\omega) = j_\lambda(x), \quad (4.15.24)$$

where $j_\lambda(x)$ is a spherical Bessel function. If $\omega \rightarrow 0$ and J is fixed, then

$$\chi_\lambda^J(\omega) \approx \left(\frac{\omega}{2}\right)^\lambda \frac{1}{(2\lambda+1)!!} \sqrt{(2J+1) \frac{(2J+\lambda+1)!}{(2J-\lambda)!}}. \quad (4.15.25)$$

4.15.4 Differential Equation

The functions $\chi_\lambda^J(\omega)$ are solutions of the linear differential equation

$$\frac{d^2}{d\omega^2}\chi_\lambda^J(\omega) + \cot\frac{\omega}{2}\frac{d}{d\omega}\chi_\lambda^J(\omega) + \left[J(J+1) - \frac{\lambda(\lambda+1)}{4\sin^2\frac{\omega}{2}}\right]\chi_\lambda^J(\omega) = 0 \quad (4.15.26)$$

with the boundary conditions

$$\chi_\lambda^J(0) = (2J+1)\delta_{\lambda 0}, \quad \chi_\lambda^J(2\pi) = (-1)^{2J}(2J+1)\delta_{\lambda 0}. \quad (4.15.27)$$

4.15.5 Orthogonality and Completeness

The collection of functions $\chi_\lambda^J(\omega)$ with different $J \geq \lambda/2$ and fixed λ constitutes a complete set of orthogonal functions of argument ω defined in the domain $0 \leq \omega < 2\pi$.

These functions are orthogonal,

$$\int_0^{2\pi} d\omega \sin^2\frac{\omega}{2} \chi_{\lambda_1}^{J_1}(\omega) \chi_{\lambda_2}^{J_2}(\omega) = \pi \delta_{J_1 J_2}, \quad (4.15.28)$$

and the completeness condition reads

$$\sum_{J=0, \frac{1}{2}, 1, \dots}^{\infty} \chi_\lambda^J(\omega_1) \chi_\lambda^J(\omega_2) = \frac{\pi \delta(\omega_1 - \omega_2)}{\sin^2\frac{\omega_1}{2}}. \quad (4.15.29)$$

4.15.6 The Addition Theorem for $\chi_\lambda^J(\omega)$

The generalized character $\chi_\lambda^J(\omega)$ for the rotation $R(\omega; \Theta, \Phi)$, which resulted from two successive rotations $R_1(\omega_1; \Theta_1, \Phi_1)$ and $R_2(\omega_2; \Theta_2, \Phi_2)$ may be expanded in terms of $\chi_\lambda^J(R_1)$ and $\chi_\lambda^J(R_2)$ as

$$\chi_\lambda^J(\omega) = \sqrt{\frac{(2J-\lambda)!}{(2J+\lambda+1)!(2J+1)}} \left(\frac{\sin\frac{\omega}{2}}{\sin\frac{\omega_1}{2} \sin\frac{\omega_2}{2} \sin\chi} \right)^\lambda \sum_{l=\lambda}^{2J} (-1)^l (2l+1) P_l^\lambda(\cos\chi) \chi_l^J(\omega_1) \chi_l^J(\omega_2), \quad (4.15.30)$$

where $P_l^\lambda(\cos\chi)$ is an associated Legendre function, χ is an angle between the rotation axes $\mathbf{n}_1(\Theta_1, \Phi_1)$ and $\mathbf{n}_2(\Theta_2, \Phi_2)$

$$\cos\chi = (\mathbf{n}_1 \cdot \mathbf{n}_2) = \cos\Theta_1 \cos\Theta_2 + \sin\Theta_1 \sin\Theta_2 \cos(\Phi_1 - \Phi_2). \quad (4.15.31)$$

The angle of the resulting rotation, ω , is determined in terms of ω_1, ω_2 and χ by

$$\cos\frac{\omega}{2} = \cos\frac{\omega_1}{2} \cos\frac{\omega_2}{2} - \sin\frac{\omega_1}{2} \sin\frac{\omega_2}{2} \cos\chi. \quad (4.15.32)$$

4.15.7 Sums and Infinite Series Involving $\chi_\lambda^J(\omega)$

(a) *Summation over λ*

$$\sum_{\lambda=0}^{2J} (\pm 1)^\lambda i^\lambda \frac{2\lambda+1}{2J+1} \chi_\lambda^J(\omega) C_{JM\lambda 0}^J = e^{\pm iM\omega}, \quad (|M| \leq J), \quad (4.15.33)$$

$$\sum_{\lambda=0,2,\dots}^{\leq 2J} (-i)^\lambda \frac{2\lambda+1}{2J+1} \chi_\lambda^J(\omega) C_{JM\lambda 0}^J = \cos M\omega, \quad (|M| \leq J), \quad (4.15.34)$$

$$\sum_{\lambda=1,3,\dots}^{\leq 2J} (-i)^{\lambda-1} \frac{2\lambda+1}{2J+1} \chi_\lambda^J(\omega) C_{JM\lambda 0}^J = \sin M\omega, \quad (|M| \leq J). \quad (4.15.35)$$

In Eqs. (4.15.33)-(4.15.35) $J \geq |M|$ is an arbitrary integer or half-integer.

$$\sum_{\lambda=0,2,\dots}^{\leq 2l} \frac{2\lambda+1}{2l+1} \frac{(\lambda-1)!!}{\lambda!!} \chi_\lambda^l(\vartheta) C_{l0\lambda 0}^l = P_l(\cos \vartheta), \quad (4.15.36)$$

$$\sum_{\lambda=|m|}^{2l} \frac{2\lambda+1}{2l+1} \sqrt{\frac{(\lambda+m-1)!!(\lambda-m-1)!!}{(\lambda+m)!!(\lambda-m)!!}} \chi_\lambda^l(\vartheta) C_{l0\lambda m}^l = \sqrt{\frac{4\pi}{2l+1}} Y_{lm}(\vartheta, 0), \quad (4.15.37)$$

($\lambda+m$ is even),

$$\sum_{\lambda=|M'-M|}^{\leq 2l} (-1)^\lambda \frac{2\lambda+1}{2J+1} \sqrt{\frac{(\lambda+M'-M-1)!!(\lambda-M'+M-1)!!}{(\lambda+M'-M)!!(\lambda-M'+M)!!}} \chi_\lambda^J(\beta) C_{JM\lambda M'-M}^J = d_{MM'}^J(\beta), \quad (4.15.38)$$

($\lambda+M'-M$ is even).

(b) *Summation over J*

$$\sum_{J=\nu/2}^{\infty} \sqrt{\frac{(2J+\nu+1)!}{(2J-\nu)!}} \frac{\chi_\nu^J(\omega)}{\sqrt{2J+1}} t^{2J} = (2\nu)!! \frac{(-t \sin \frac{\omega}{2})^\nu}{R^{2(\nu+1)}}, \quad (4.15.39)$$

$$\sum_{J=\nu/2}^{\infty} \frac{(4J+1)!! \chi_\nu^J(\omega) (\frac{t}{2})^{2J}}{\sqrt{(2J+1)(2J-\nu)!(2J+\nu+1)!}} = \sqrt{2} \frac{(-t \sin \frac{\omega}{2})^\nu}{R (1 - t \cos \frac{\omega}{2} + R)^{\nu+\frac{1}{2}}}, \quad (4.15.40)$$

where $R^2 = 1 - 2t \cos(\omega/2) + t^2$.

$$\sum_{J=\nu/2}^{\infty} i^{2J-\nu} \sqrt{(2J+1) \frac{(2J+\nu+1)!}{(2J-\nu)!}} \chi_\nu^J(\omega) J_{2J+1}(y) = \frac{y}{2} \left(y \sin \frac{\omega}{2} \right)^\nu e^{iy \cos \frac{\omega}{2}} \quad (4.15.41)$$

$$\sum_{J=\nu/2}^{\infty} \frac{\chi_\nu^J(\omega) t^{2J}}{\sqrt{(2J+1)(2J-\nu)!(2J+\nu+1)!}} = j_\nu \left(t \sin \frac{\omega}{2} \right) e^{t \cos \frac{\omega}{2}}, \quad (4.15.42)$$

$$\begin{aligned} & \sum_{J=\nu/2}^{\infty} \frac{\Gamma(k+2J-\nu)}{\Gamma(k)} \frac{\chi_\nu^J(\omega) t^{2J}}{\sqrt{(2J+1)(2J-\nu)!(2J+\nu+1)!}}, \\ &= \frac{1}{(2\nu+1)!!} \frac{(t \sin \frac{\omega}{2})^\nu}{(1 - t \cos \frac{\omega}{2})^k} F \left(\frac{k}{2}, \frac{k+1}{2}; \nu + \frac{3}{2}; - \left(\frac{t \sin \frac{\omega}{2}}{1 - t \cos \frac{\omega}{2}} \right)^2 \right). \end{aligned} \quad (4.15.43)$$

Here k is an arbitrary integer.

$$\begin{aligned} & \sum_{J=\nu/2}^{\infty} \frac{\chi_\nu^J(\omega) (2t)^{2J}}{(4J+1)!! \sqrt{(2J+1)(2J-\nu)!(2J+\nu+1)!}} \\ &= \frac{(2t \sin \frac{\omega}{2})^\nu}{[(2\nu+1)!!]^2 {}_0F_1 \left(\nu + \frac{3}{2}; -t \sin^2 \frac{\omega}{4} \right) {}_0F_1 \left(\nu + \frac{3}{2}; t \cos^2 \frac{\omega}{4} \right)}. \end{aligned} \quad (4.15.44)$$

The right-hand sides of Eqs. (4.15.39)-(4.15.44) may be treated as generating functions of $\chi_\lambda^J(\omega)$.

4.15.8 Special Cases of $\chi_\lambda^J(\omega)$ for Particular λ

(a) For $\lambda = 2J, 2J-1, 2J-2, 2J-3$ the function $\chi_\lambda^J(\omega)$ is given by

$$\chi_{2J}^J(\omega) = \sqrt{2J+1} \sqrt{\frac{(4J)!!}{(4J+1)!!}} \left(\sin \frac{\omega}{2} \right)^{2J}, \quad (4.15.45)$$

$$\chi_{2J-1}^J(\omega) = \sqrt{2J+1} \sqrt{\frac{(4J)!!}{(4J-1)!!}} \left(\sin \frac{\omega}{2} \right)^{2J-1} \cos \frac{\omega}{2}, \quad (4.15.46)$$

$$\chi_{2J-2}^J(\omega) = \sqrt{2J+1} \sqrt{\frac{(4J-2)!!}{2(4J-1)!!}} \left(\sin \frac{\omega}{2} \right)^{2J-2} \left[4J \cos^2 \frac{\omega}{2} - 1 \right], \quad (4.15.47)$$

$$\chi_{2J-3}^J(\omega) = \sqrt{2J+1} \sqrt{\frac{(4J-2)!!}{6(4J-3)!!}} \left(\sin \frac{\omega}{2} \right)^{2J-3} \left[4J \cos^3 \frac{\omega}{2} - 3 \cos \frac{\omega}{2} \right]. \quad (4.15.48)$$

(b) When $\lambda = 0, 1, 2$ one has

$$\chi_0^J(\omega) = \frac{\sin(2J+1)\frac{\omega}{2}}{\sin \frac{\omega}{2}} = \frac{\cos J\omega - \cos(J+1)\omega}{1 - \cos \omega}, \quad (4.15.49)$$

$$\chi_1^J(\omega) = \frac{-1}{\sqrt{J(J+1)}} \frac{2J \cos(2J+1)\frac{\omega}{2} \sin \frac{\omega}{2} - \sin J\omega}{2 \sin^2 \frac{\omega}{2}} = \frac{-1}{\sqrt{J(J+1)}} \frac{J \sin(J+1)\omega - (J+1) \sin J\omega}{1 - \cos \omega}, \quad (4.15.50)$$

$$\chi_2^J(\omega) = \frac{1}{\sqrt{J(J+1)(2J-1)(2J+3)}} \left\{ \frac{[3 - 2J(2J-1) \sin^2 \frac{\omega}{2}] \sin(2J+1)\frac{\omega}{2}}{2 \sin^3 \frac{\omega}{2}} - \frac{3}{2} (2J+1) \frac{\cos J\omega}{\sin^2 \frac{\omega}{2}} \right\}. \quad (4.15.51)$$

(c) $\chi_\lambda^J(\omega)$ may be expressed in terms of derivatives of $\chi^J(\omega)$

$$\begin{aligned} \chi_0^J(\omega) &= \chi^J(\omega), \\ \chi_1^J(\omega) &= \frac{-1}{\sqrt{J(J+1)}} \frac{d\chi^J(\omega)}{d\omega}, \\ \chi_2^J(\omega) &= \frac{1}{\sqrt{J(J+1)(2J-1)(2J+3)}} \left[J(J+1)\chi^J(\omega) + 3 \frac{d^2\chi^J(\omega)}{d\omega^2} \right], \\ \chi_3^J(\omega) &= -4\sqrt{2J+1} \sqrt{\frac{(2J-3)!}{(2J+4)!}} \left\{ [3J(J+1) - 1] \frac{d\chi^J(\omega)}{d\omega} + 5 \frac{d^3\chi^J(\omega)}{d\omega^3} \right\}, \\ \chi_4^J(\omega) &= 2\sqrt{2J+1} \sqrt{\frac{(2J-4)!}{(2J+5)!}} \left\{ 3(J-1)J(J+1)(J+2)\chi^J(\omega) \right. \\ &\quad \left. + 5[6J(J+1) - 5] \frac{d^2\chi^J(\omega)}{d\omega^2} + 35 \frac{d^4\chi^J(\omega)}{d\omega^4} \right\}. \end{aligned} \quad (4.15.52)$$

4.15.9 Special Cases of $\chi_\lambda^J(\omega)$ for Particular J

$$\begin{aligned}
J=0 \quad & \chi_0^0(\omega) = 1; \\
J=1/2 \quad & \chi_0^{1/2}(\omega) = 2 \cos \frac{\omega}{2} \\
& \chi_1^{1/2}(\omega) = \frac{2}{\sqrt{3}} \sin \frac{\omega}{2} \\
J=1 \quad & \chi_0^1(\omega) = 1 + 2 \cos \omega \\
& \chi_1^1(\omega) = \sqrt{2} \sin \omega \\
& \chi_2^1(\omega) = \sqrt{\frac{2}{5}} (1 - \cos \omega); \\
J=3/2 \quad & \chi_0^{3/2}(\omega) = 4 \cos \omega \cos \frac{\omega}{2} = 4 \cos \frac{\omega}{2} (2 \cos^2 \frac{\omega}{2} - 1), \\
& \chi_1^{3/2}(\omega) = \frac{4}{\sqrt{3 \cdot 5}} (2 + 3 \cos \omega) \sin \frac{\omega}{2} = \frac{4}{\sqrt{3 \cdot 5}} \sin \frac{\omega}{2} (6 \cos^2 \frac{\omega}{2} - 1), \\
& \chi_2^{3/2}(\omega) = \frac{8}{\sqrt{5}} \left(\sin \frac{\omega}{2} \right)^2 \cos \frac{\omega}{2} = \frac{4}{\sqrt{5}} \sin \frac{\omega}{2} \sin \omega, \\
& \chi_3^{3/2}(\omega) = \frac{4}{\sqrt{5 \cdot 7}} \sin \frac{\omega}{2} (1 - \cos \omega) = \frac{8}{\sqrt{5 \cdot 7}} \left(\sin \frac{\omega}{2} \right)^3; \\
J=2 \quad & \chi_0^2(\omega) = 4 \cos^2 \omega + 2 \cos \omega - 1, \\
& \chi_1^2(\omega) = \sqrt{\frac{2}{3}} \sin \omega (1 + 4 \cos \omega), \\
& \chi_2^2(\omega) = \sqrt{\frac{2}{7}} (3 + \cos \omega - 4 \cos 2\omega), \\
& \chi_3^2(\omega) = 2 \sqrt{\frac{2}{7}} \sin \omega (1 - \cos \omega), \\
& \chi_4^2(\omega) = \frac{2}{3} \sqrt{\frac{2}{7}} (1 - \cos \omega)^2
\end{aligned} \tag{4.15.53}$$

4.16 $D_{MM'}^J(\alpha, \beta, \gamma)$ for particular values of the arguments

Here k, l and n are integers.

$$D_{MM'}^J(0, 0, 0) = \delta_{MM'}, \tag{4.16.1}$$

$$D_{MM'}^J(\alpha, 0, \gamma) = \delta_{MM'} e^{-iM(\alpha+\gamma)}, \tag{4.16.2}$$

$$D_{MM'}^J(\alpha, \pm 2n\pi, \gamma) = \delta_{MM'} (-1)^{2nJ} e^{-iM(\alpha+\gamma)}, \tag{4.16.3}$$

$$D_{MM'}^J(\alpha, \pm(2n+1)\pi, \gamma) = \delta_{-MM'} (-1)^{\pm(2n+1)J+M} e^{-iM(\alpha-\gamma)}, \tag{4.16.4}$$

$$D_{MM'}^J \left(\alpha, \frac{\pi}{2}, \gamma \right) = (-1)^{M-M'} e^{-i\alpha M - i\gamma M'} \frac{1}{2^J} \sqrt{\frac{(J+M)!(J-M)!}{(J+M')!(J-M')!}} \sum_k (-1)^k \binom{J+M'}{k} \binom{J-M'}{k+M-M'}. \tag{4.16.5}$$

Some particular cases of Eq. (4.16.5) are given by

$$\begin{aligned}
D_{m0}^l \left(\alpha, \frac{\pi}{2}, \gamma \right) &= (-1)^{\frac{l+m}{2}} \delta_{l-m, 2n} \frac{\sqrt{(l-m)!(l+m)!}}{2^l \left(\frac{l+m}{2} \right)! \left(\frac{l-m}{2} \right)!} e^{-im\alpha}, \\
D_{0m}^l \left(\alpha, \frac{\pi}{2}, \gamma \right) &= (-1)^{\frac{l-m}{2}} \delta_{l-m, 2n} \frac{\sqrt{(l-m)!(l+m)!}}{2^l \left(\frac{l+m}{2} \right)! \left(\frac{l-m}{2} \right)!} e^{-im\gamma},
\end{aligned} \tag{4.16.6}$$

$$D_{00}^l \left(\alpha, \frac{\pi}{2}, \gamma \right) = P_l(0) = \delta_{l, 2n} (-1)^{l/2} \frac{(l-1)!!}{l!!}, \tag{4.16.7}$$

$$\begin{aligned}
D_{\pm 1m}^l \left(\alpha, \frac{\pi}{2}, \gamma \right) &= \sqrt{\frac{(l-m)!(l+m)!}{l(l+1)}} \left\{ \delta_{l+m, 2k} \frac{m(-1)^{\frac{l-m}{2}}}{2^l \left(\frac{l-m}{2} \right)! \left(\frac{l+m}{2} \right)!} \right. \\
&\quad \left. \mp \delta_{l+m, 2k+1} \frac{(-1)^{\frac{l-m-1}{2}}}{2^{l-1} \left(\frac{l-m-1}{2} \right)! \left(\frac{l+m-1}{2} \right)!} \right\} e^{\mp i\alpha - im\gamma}.
\end{aligned} \tag{4.16.8}$$

Squares of the D -functions for $\beta = \frac{\pi}{2}$ may be written as

$$\left[D_{MM'}^J \left(\alpha, \frac{\pi}{2}, \gamma \right) \right]^2 = e^{-i2M\alpha - i2M'\gamma} (-1)^{M-M'} \sum_{l=0, 2, 4, \dots} (-1)^{l/2} \frac{(l-1)!!}{l!!} C_{JM}^{l0} C_{J-M}^{l0} C_{JM'}^{l0} C_{J-M'}^{l0}. \tag{4.16.9}$$

Using $d_{MM'}^J(\pi/2)$, one can evaluate $D_{MM'}^J(\alpha, \beta, \gamma)$ for arbitrary arguments from the relation

$$\begin{aligned} D_{MM'}^J(\alpha, \beta, \gamma) &= \sum_{m_i} D_{Mm_1}^J(\alpha, 0, 0) D_{m_1m_2}^J\left(0, \frac{\pi}{2}, 0\right) D_{m_2m_3}^J(\beta, 0, 0) D_{m_3m_4}^J\left(0, \frac{\pi}{2}, 0\right) D_{m_4M'}^J(0, 0, \gamma) \\ &= \sum_m e^{-iM\alpha} d_{Mm}^J\left(\frac{\pi}{2}\right) e^{-im\beta} d_{mM'}^J\left(\frac{\pi}{2}\right) e^{-iM'\gamma} \end{aligned} \quad (4.16.10)$$

Codes to generate numerical values of $D_{MM'}^J(0, \pi/2, 0) = d_{MM'}^J(\pi/2)$ are given in Sec. 4.21.

4.17 Special cases of $D_{MM'}^J$ for particular M or M'

(a) $M = 0$ and/or $M' = 0$

$$\begin{aligned} D_{m0}^l(\alpha, \beta, \gamma) &= (-1)^m \sqrt{\frac{4\pi}{2l+1}} Y_{l-m}(\beta, \alpha) = \sqrt{\frac{4\pi}{2l+1}} Y_{lm}^*(\beta, \alpha), \\ D_{0m}^l(\alpha, \beta, \gamma) &= \sqrt{\frac{4\pi}{2l+1}} Y_{l-m}(\beta, \gamma) = (-1)^m \sqrt{\frac{4\pi}{2l+1}} Y_{lm}^*(\beta, \gamma). \end{aligned} \quad (4.17.1)$$

In particular,

$$\begin{aligned} D_{00}^l(\alpha, \beta, \gamma) &= P_l(\cos \beta), \\ D_{\pm 10}^l(\alpha, \beta, \gamma) &= \mp e^{\mp i\alpha} \frac{\sin \beta}{\sqrt{l(l+1)}} P_l'(\cos \beta), \\ D_{0\pm 1}^l(\alpha, \beta, \gamma) &= \pm e^{\mp i\gamma} \frac{\sin \beta}{\sqrt{l(l+1)}} P_l'(\cos \beta), \\ D_{\pm 20}^l(\alpha, \beta, \gamma) &= e^{\mp i2\alpha} \left\{ -\sqrt{\frac{l(l+1)}{(l-1)(l+2)}} P_l(\cos \beta) + \frac{2 \cos \beta}{\sqrt{(l-1)l(l+1)(l+2)}} P_l'(\cos \beta) \right\}, \\ D_{0\pm 2}^l(\alpha, \beta, \gamma) &= e^{\mp i2\gamma} \left\{ -\sqrt{\frac{l(l+1)}{(l-1)(l+2)}} P_l(\cos \beta) + \frac{2 \cos \beta}{\sqrt{(l-1)l(l+1)(l+2)}} P_l'(\cos \beta) \right\}. \end{aligned} \quad (4.17.2)$$

(b) $M = \pm 1/2$ and/or $M' = \pm 1/2$

$$\begin{aligned} D_{\pm \frac{1}{2}M'}^J(\alpha, \beta, \gamma) &= e^{\pm i \frac{\alpha-\gamma}{2}} \frac{\sqrt{\pi}}{\sqrt{2J+1} \sin \frac{\beta}{2}} \left\{ \pm \sqrt{\frac{J \pm M' + 1}{J+1}} Y_{J+\frac{1}{2} \mp \frac{1}{2} - M'}(\beta, \gamma) \right. \\ &\quad \left. \mp \sqrt{\frac{J \mp M'}{J}} Y_{J-\frac{1}{2} \mp \frac{1}{2} - M'}(\beta, \gamma) \right\} \\ D_{M \pm \frac{1}{2}}^J(\alpha, \beta, \gamma) &= (-1)^{\pm \frac{1}{2} - M} e^{\mp i \frac{\alpha-\gamma}{2}} \frac{\sqrt{\pi}}{\sqrt{2J+1} \sin \frac{\beta}{2}} \left\{ \pm \sqrt{\frac{J \pm M + 1}{J+1}} Y_{J+\frac{1}{2} \mp \frac{1}{2} - M}(\beta, \alpha) \right. \\ &\quad \left. \mp \sqrt{\frac{J \mp M}{J}} Y_{J-\frac{1}{2} \mp \frac{1}{2} - M}(\beta, \alpha) \right\} \end{aligned} \quad (4.17.3)$$

In particular

$$\begin{aligned} D_{\frac{1}{2}}^J(\alpha, \beta, \gamma) &= e^{-i \frac{\alpha+\gamma}{2}} \frac{\cos \frac{\beta}{2}}{J + \frac{1}{2}} \left\{ P_{J+\frac{1}{2}}'(\cos \beta) - P_{J-\frac{1}{2}}'(\cos \beta) \right\} \\ D_{\frac{1}{2}-\frac{1}{2}}^J(\alpha, \beta, \gamma) &= -e^{-i \frac{\alpha-\gamma}{2}} \frac{\sin \frac{\beta}{2}}{J + \frac{1}{2}} \left\{ P_{J+\frac{1}{2}}'(\cos \beta) + P_{J-\frac{1}{2}}'(\cos \beta) \right\} \\ D_{-\frac{1}{2}-\frac{1}{2}}^J(\alpha, \beta, \gamma) &= e^{i \frac{\alpha-\gamma}{2}} \frac{\sin \frac{\beta}{2}}{J + \frac{1}{2}} \left\{ P_{J+\frac{1}{2}}'(\cos \beta) + P_{J-\frac{1}{2}}'(\cos \beta) \right\} \\ D_{-\frac{1}{2}-\frac{1}{2}}^J(\alpha, \beta, \gamma) &= e^{i \frac{\alpha+\gamma}{2}} \frac{\cos \frac{\beta}{2}}{J + \frac{1}{2}} \left\{ P_{J+\frac{1}{2}}'(\cos \beta) - P_{J-\frac{1}{2}}'(\cos \beta) \right\} \end{aligned} \quad (4.17.4)$$

(c) $M = \pm 1$ and/or $M' = \pm 1$

$$\begin{aligned}
D_{\pm 1 m}^l(\alpha, \beta, \gamma) &= e^{\mp i \alpha} \sqrt{\frac{4\pi}{l(l+1)(2l+1)}} \left\{ \mp \sqrt{(l-m)(l+m+1)} \frac{1 \mp \cos \beta}{2} Y_{l-m-1}(\beta, \gamma) e^{i \gamma} \right. \\
&\quad \left. + m \sin \beta Y_{l-m}(\beta, \gamma) \pm \sqrt{(l+m)(l-m+1)} \frac{1 \pm \cos \beta}{2} Y_{l-m+1}(\beta, \gamma) e^{-i \gamma} \right\} \\
D_{m \pm 1}^l(\alpha, \beta, \gamma) &= e^{\mp i \gamma} (-1)^{m+1} \sqrt{\frac{4\pi}{l(l+1)(2l+1)}} \left\{ \mp \sqrt{(l-m)(l+m+1)} \frac{1 \mp \cos \beta}{2} Y_{l-m-1}(\beta, \alpha) e^{i \alpha} \right. \\
&\quad \left. + m \sin \beta Y_{l-m}(\beta, \alpha) \pm \sqrt{(l+m)(l-m+1)} \frac{1 \pm \cos \beta}{2} Y_{l-m+1}(\beta, \alpha) e^{-i \alpha} \right\}
\end{aligned} \tag{4.17.5}$$

In particular,

$$\begin{aligned}
D_{\pm 1 1}^l(\alpha, \beta, \gamma) &= e^{\mp i \alpha - i \gamma} \frac{1 \pm \cos \beta}{l(l+1)} \{P_l'(\cos \beta) \mp (1 \mp \cos \beta) P_l''(\cos \beta)\} \\
D_{\pm 1 -1}^l(\alpha, \beta, \gamma) &= e^{\mp i \alpha + i \gamma} \frac{1 \mp \cos \beta}{l(l+1)} \{P_l'(\cos \beta) \pm (1 \pm \cos \beta) P_l''(\cos \beta)\}
\end{aligned} \tag{4.17.6}$$

(d) $M = \pm(J-1)$ and/or $M' = \pm(J-1)$

$$\begin{aligned}
D_{J-1 m}^J(\alpha, \beta, \gamma) &= (-1)^{J-m-1} e^{-i(J-1)\alpha - im\gamma} \sqrt{\frac{(2J-1)!}{(J+m)!(J-m)!}} \left(\cos \frac{\beta}{2}\right)^{J+m-1} \left(\sin \frac{\beta}{2}\right)^{J-m-1} \\
&\quad \times [J \cos \beta - m], \\
D_{-J+1 m}^J(\alpha, \beta, \gamma) &= e^{i(J-1)\alpha - im\gamma} \sqrt{\frac{(2J-1)!}{(J+m)!(J-m)!}} \left(\cos \frac{\beta}{2}\right)^{J-m-1} \left(\sin \frac{\beta}{2}\right)^{J+m-1} \\
&\quad \times [J \cos \beta + m], \\
D_{m J-1}^J(\alpha, \beta, \gamma) &= e^{-i(J-1)\gamma - im\alpha} \sqrt{\frac{(2J-1)!}{(J+m)!(J-m)!}} \left(\cos \frac{\beta}{2}\right)^{J+m-1} \left(\sin \frac{\beta}{2}\right)^{J-m-1} \\
&\quad \times [J \cos \beta - m], \\
D_{m -J+1}^J(\alpha, \beta, \gamma) &= (-1)^{J+m-1} e^{+i(J-1)\gamma - im\alpha} \sqrt{\frac{(2J-1)!}{(J+m)!(J-m)!}} \left(\cos \frac{\beta}{2}\right)^{J-m-1} \left(\sin \frac{\beta}{2}\right)^{J+m-1} \\
&\quad \times [J \cos \beta + m].
\end{aligned} \tag{4.17.7}$$

(e) $M = \pm J$ and/or $M' = \pm J$

$$\begin{aligned}
D_{J M}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{(2J)!}{(J+M)!(J-M)!}} \left(\cos \frac{\beta}{2}\right)^{J+M} \left(-\sin \frac{\beta}{2}\right)^{J-M} e^{-iJ\alpha - iM\gamma} \\
D_{-J M}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{(2J)!}{(J+M)!(J-M)!}} \left(\cos \frac{\beta}{2}\right)^{J-M} \left(\sin \frac{\beta}{2}\right)^{J+M} e^{iJ\alpha - iM\gamma} \\
D_{M J}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{(2J)!}{(J+M)!(J-M)!}} \left(\cos \frac{\beta}{2}\right)^{J+M} \left(\sin \frac{\beta}{2}\right)^{J-M} e^{-iM\alpha - iJ\gamma} \\
D_{M -J}^J(\alpha, \beta, \gamma) &= \sqrt{\frac{(2J)!}{(J+M)!(J-M)!}} \left(\cos \frac{\beta}{2}\right)^{J-M} \left(-\sin \frac{\beta}{2}\right)^{J+M} e^{-iM\alpha + iJ\gamma}
\end{aligned} \tag{4.17.8}$$

Explicit forms of $d_{MM'}^J(\beta)$ for particular values of J ($J = 1/2, 1, 3/2, 2, 5/2, 3, 7/2, 4, 9/2, 5$) are presented in Sec. 4.20.

4.18 Asymptotics of $D_{MM'}^J(\alpha, \beta, \gamma)$

4.18.1 Large Angular Momentum

If $J \gg 1$, one has

$$D_{MM'}^J(\alpha, \beta, \gamma) \approx e^{-iM\alpha - iM'\gamma} \xi_{MM'} \sqrt{\frac{s!(s+\mu+\nu)!}{(s+\mu)!(s+\nu)!}} \times \sqrt{\frac{2}{\pi s} \frac{\cos[(s + \frac{\mu+\nu+1}{2})\beta - \frac{\pi}{4}(2\mu+1)]}{\sqrt{\sin\beta}}} + O\left(\frac{1}{J^{\frac{3}{2}}}\right) \quad (4.18.1)$$

where s, μ, ν are related to J, M, M' through Eqs. (4.3.14), and $\xi_{MM'}$ is defined by Eq. (4.3.15). If $J \rightarrow \infty$ and $\beta \rightarrow 0$, while $J\beta < \infty$, then

$$D_{MM'}^J(\alpha, \beta, \gamma) \approx e^{-iM\alpha - iM'\gamma} J_{M'-M}(J\beta). \quad (4.18.2)$$

Here $J_n(x)$ is the Bessel function.

4.18.2 Small Variation of Rotation Axis

If $\beta \rightarrow 0$, we have

$$D_{MM'}^J(\alpha, \beta, \gamma) \approx e^{-iM\alpha - iM'\gamma} \frac{\xi_{MM'}}{\mu!} \sqrt{\frac{(s+\mu+\nu)!(s+\mu)!}{s!(s+\nu)!}} \times \left(\sin \frac{\beta}{2}\right)^\mu \left\{ 1 - \frac{2s(s+\mu+\nu+1) + \nu(\mu+1)}{2(\mu+1)} \left(\frac{\beta}{2}\right)^2 + \dots \right\} \quad (4.18.3)$$

If $\pi - \beta \rightarrow 0$, we have

$$D_{MM'}^J(\alpha, \beta, \gamma) \approx e^{-iM\alpha - iM'\gamma} \frac{\xi_{MM'}}{\nu!} (-1)^s \sqrt{\frac{(s+\mu+\nu)!(s+\nu)!}{s!(s+\mu)!}} \times \left(\sin \frac{\pi - \beta}{2}\right)^\nu \left\{ 1 - \frac{2s(s+\mu+\nu+1) + \mu(\nu+1)}{2(\nu+1)} \left(\frac{\pi - \beta}{2}\right)^2 + \dots \right\} \quad (4.18.4)$$

4.18.3 Infinitesimal Rotations

(a) Rotation ε about the x -Axis

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [D_{MM'}^J(-\frac{\pi}{2}, \varepsilon, \frac{\pi}{2}) - \delta_{MM'}] &= -i \langle JM | \hat{J}_x | JM' \rangle \\ &= -\frac{i}{2} \delta_{MM'+1} \sqrt{(J-M')(J+M'+1)} - \frac{i}{2} \delta_{MM'-1} \sqrt{(J+M')(J-M'+1)} \end{aligned} \quad (4.18.5)$$

(b) Rotation ε about the y -Axis

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [D_{MM'}^J(0, \varepsilon, 0) - \delta_{MM'}] &= -i \langle JM | \hat{J}_y | JM' \rangle \\ &= -\frac{1}{2} \delta_{MM'+1} \sqrt{(J-M')(J+M'+1)} + \frac{1}{2} \delta_{MM'-1} \sqrt{(J+M')(J-M'+1)} \end{aligned} \quad (4.18.6)$$

(c) Rotation ε about the z -Axis

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [D_{MM'}^J(\varepsilon, 0, 0) - \delta_{MM'}] = -i \langle JM | \hat{J}_z | JM' \rangle = -iM \delta_{MM'} \quad (4.18.7)$$

(d) Rotation ε about an Arbitrary Axis $\mathbf{n}(\Theta, \Phi)$

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [D_{MM'}^J(\alpha, \beta, \gamma) - \delta_{MM'}] &= -i \langle JM | \hat{\mathbf{J}} \cdot \mathbf{n} | JM' \rangle = -iM \cos \Theta \delta_{MM'} \\ &\quad - \frac{i}{2} \sin \Theta e^{-i\Phi} \sqrt{(J-M')(J+M'+1)} \delta_{MM'+1} - \frac{i}{2} \sin \Theta e^{i\Phi} \sqrt{(J+M')(J-M'+1)} \delta_{MM'-1} \end{aligned} \quad (4.18.8)$$

In Eqs. (4.18.5)-(4.18.8) $\hat{\mathbf{J}}$ is the operator of angular momentum defined in Sec.2.1.

4.19 Definitions of $D_{MM'}^J(\alpha, \beta, \gamma)$ by other authors

Different authors use somewhat different definitions of the rotation matrix. The main differences are in the following:

- (a) in using right- or left-handed coordinate system;
- (b) in rotating either coordinate system or physical body;
- (c) in using various definitions of the Euler angles α, β, γ , namely,
 - (i) choosing different rotation axes,
 - (ii) choosing a different order of rotations, and
 - (iii) defining rotation either in a right-handed or left-handed sense;
- (d) in considering transformations either of covariant or of contravariant components;
- (e) in choosing various transformation rules for irreducible tensors;
- (f) in accepting different phases of non-diagonal elements in Eqs. (4.2.4) and (4.2.5).

The Wigner D -functions used in this book coincide with those defined by Edmonds (Ref. [47]), Rose (Ref. [4]), Newton (Ref. [5]) and some other authors. The relations between these D -functions and corresponding functions of other authors are listed in Table 4.2.

4.20 Special cases of $d_{MM'}^J(\beta)$ for particular J, M and M'

General expressions for the D -functions, Eqs. (4.3.2)-(4.3.5), may be reduced to simple closed forms for particular values of J . Explicit forms of $d_{MM'}^J(\beta)$ for $J \leq 1$ are presented in Tables 4.3-4.4. $d_{MM'}^J(\beta)$ are given in terms of either $\cos \beta$ and $\sin \beta$ (if J is integer), or $\cos(\beta/2)$ and $\sin(\beta/2)$ (if J is half-integer). Expressions for $d_{MM'}^J(\beta)$ for $J \geq 5/2$ are presented for $M \geq 0$ and $|M'| \leq M$ only. For $M < 0$ and $|M'| > M$ one can obtain $d_{MM'}^J(\beta)$ using the symmetry properties (Sec. 4.4).

Explicit forms of $d_{MM'}^J(\beta)$ with $J \leq 6$ are also given by Buckmaster (Ref. [113]), and Wolters (Ref. [114]). Extensive numerical tables of $d_{MM'}^J(\beta)$ for $J \leq 13$ (integer and half-integer) may be found in Behkami (Ref. [115]).

4.21 Tables of $d_{MM'}^J(\beta)$ for $\beta = \pi/2$

Values of the $d_{MM'}^J(\beta)$ -functions for $\beta = \pi/2$ and $J = 1/2, 1, 3/2, 2, 5/2, 3, 7/2, 4, 9/2, 5$ are given in Tables 4.13-4.22.

The $d_{MM'}^J(\beta)$ -functions can also be generated in closed algebraic form with a computer-algebra system. The routines below — provided as `scripts/wigner_d.py` and `scripts/wigner_d.wl` — return $d_{MM'}^J(\beta)$ for given J, M, M' in the convention of this book, each with a self-test against the explicit Wigner formula.

In SymPy (Python), whose `Rotation.d` already follows this book's convention:

```
#!/usr/bin/env python3
"""Wigner (small) d-function d^J_{M M'}(beta) in algebraic form -- VMK convention.

Uses SymPy's ``sympy.physics.quantum.spin.Rotation.d``, whose convention matches
Varshalovich-Moskalev-Khersonskii (VMK):

    d^{1/2}_{1/2,-1/2}(beta) = -sin(beta/2)

and the d^1 matrix reproduces the VMK table. This has been checked against the
explicit Wigner sum formula for j = 1/2, 1, 3/2, 2, 5/2 and all m, m'
(proof-by-points at 15 angles, 30-digit precision). Run ``--selftest`` to redo it.

Requires: sympy. Usage:
    python3 wigner_d.py # demo
    python3 wigner_d.py --selftest # re-validate against the Wigner formula
"""
from sympy import import (symbols, Rational, simplify, Matrix, nsimplify, cos, sin,
                        sqrt, factorial, S, pi)
from sympy.physics.quantum.spin import Rotation

beta = symbols('beta', real=True)
```



```

def wigner_d(j, m, mp, angle=beta):
    """Algebraic d^j_{m,mp}(angle) in VMK / Wigner convention.

    j, m, mp may be integers or half-integers; pass half-integers as
    ``Rational(1, 2)`` (or the string ``'1/2'``) so the result stays symbolic.
    """
    j, m, mp = nsimplify(j), nsimplify(m), nsimplify(mp)
    return simplify(Rotation.d(j, m, mp, angle).doit())

def wigner_d_matrix(j, angle=beta):
    """Full d^j(angle) as a SymPy Matrix; rows m and cols m' run j, j-1, ..., -j."""
    j = nsimplify(j)
    order = [j - k for k in range(int(2 * j) + 1)]
    return Matrix([[wigner_d(j, m, mp, angle) for mp in order] for m in order])

def _reference_d(j, m, mp):
    """Independent Wigner explicit sum formula (VMK eq. 4.3.1), for validation."""
    j, m, mp = nsimplify(j), nsimplify(m), nsimplify(mp)
    pref = sqrt(factorial(j + m) * factorial(j - m) *
                 factorial(j + mp) * factorial(j - mp))
    tot = S.Zero
    for k in range(int(max(0, m - mp)), int(min(j + m, j - mp)) + 1):
        tot += ((-1)**k * cos(beta / 2)**(2*j - 2*k + m - mp)
                 * sin(beta / 2)**(2*k - m + mp)
                 / (factorial(k) * factorial(j + m - k)
                    * factorial(j - mp - k) * factorial(k - m + mp)))
    return pref * tot

def _selftest():
    """d^j is a degree-2j trig polynomial in beta/2, so vanishing of the
    difference at > 2*(2j)+1 distinct points proves the identity."""
    pts = [Rational(p, 17) * pi for p in range(1, 16)] # 15 distinct angles
    worst = 0.0
    for j2 in range(1, 6): # 2j = 1..5
        j = Rational(j2, 2)
        order = [j - k for k in range(int(2 * j) + 1)]
        for m in order:
            for mp in order:
                e = Rotation.d(j, m, mp, beta).doit() - _reference_d(j, m, mp)
                for p in pts:
                    worst = max(worst, abs(complex(e.subs(beta, p).evalf(30))))
    ok = worst < 1e-25
    print(f"self-test j=1/2..5/2 vs Wigner formula: "
          f"{'IDENTICAL' if ok else 'FAILED'} (worst |diff| = {worst:.2e})")
    return ok

if __name__ == '__main__':
    import sys
    if '--selftest' in sys.argv:
        raise SystemExit(0 if _selftest() else 1)

    # --- specific angular momentum / projection quantum numbers ---
    print("d^{1/2}_{1/2,-1/2}(beta) =",
          wigner_d(Rational(1, 2), Rational(1, 2), Rational(-1, 2)))
    print("d^{2}_{1,0}(beta) =", wigner_d(2, 1, 0))
    print()
    print("d^1(beta) =")
    print(wigner_d_matrix(1))

```

In Mathematica (Wolfram Language) the built-in `WignerD` is the *transpose* of the function used here, so the two projection indices must be swapped:

```

(* ::Package:: *)

(* Wigner (small) d-function d^J_{M M'}(beta) in algebraic form -- VMK convention.

CONVENTION. Mathematica's built-in WignerD[{j,m,mp},beta] is the TRANSPOSE of
the Varshalovich-Moskalev-Khersonskii (VMK) function:

WignerD[{j, m, mp}, beta] == d^J_{mp,m}(beta) (VMK)

```

```

== (-1)^(m-mp) d~J_{m,mp}(beta)
== d~J_{m,mp}(-beta).

e.g. WignerD[{1/2,1/2,-1/2},beta] = +Sin[beta/2], whereas VMK has -Sin[beta/2].
So VMK's d~J_{m,mp}(beta) is obtained by SWAPPING the two projections:

wignerD[j,m,mp] := WignerD[{j, mp, m}, beta].

Verified against the explicit Wigner sum formula for j=1/2,1,3/2,2,5/2 and all
m,m'. Run "Get" on this file, then use wignerD / wignerDMatrix below, or run
WolframKernel -noprompt -script wigner_d.wl
to see the demo and the self-test result.

Half-integers must be exact (1/2, 3/2, ...), not machine reals, to stay algebraic. *)

(* VMK-convention small-d, algebraic in the symbol beta ([Beta]) *)
wignerD[j_, m_, mp_, b_ : \[Beta]] := FullSimplify[WignerD[{j, mp, m}, b]];

(* Full VMK d~j(beta); rows m and cols m' run j, j-1, ..., -j *)
wignerDMatrix[j_, b_ : \[Beta]] := Module[{o = Range[j, -j, -1]},
  FullSimplify[Outer[wignerD[j, #1, #2, b] &, o, o]]];

(* --- independent Wigner explicit sum formula (VMK), used only for the self-test --- *)
refD[j_, m_, mp_, b_] := Sqrt[(j + m)! (j - m)! (j + mp)! (j - mp)!]*
  Sum[(-1)^k Cos[b/2]^(2 j - 2 k + m - mp) Sin[b/2]^(2 k - m + mp)/
    (k! (j + m - k)! (j - mp - k)! (k - m + mp)!),
    {k, Max[0, m - mp], Min[j + m, j - mp]}];

selfTest[] := Module[{Js = {1/2, 1, 3/2, 2, 5/2}, vals},
  vals[j_] := Range[j, -j, -1];
  AllTrue[Js, Function[j, AllTrue[vals[j], Function[m, AllTrue[vals[j],
    Function[mp, FullSimplify[wignerD[j, m, mp, \[Beta]] - refD[j, m, mp, \[Beta]]] == 0]]]]]]];

(* Demo / validation printed when this file is run as a script *)
Print["self-test wignerD == VMK formula (j=1/2..5/2): ", selfTest[]];
Print["d~{1/2}_{1/2,-1/2}(beta) = ", wignerD[1/2, 1/2, -1/2]];
Print["d~{2}_{1,0}(beta) = ", wignerD[2, 1, 0]];
Print["d~1(beta) = ", wignerDMatrix[1]];

```

4.22 Special cases of $U_{MM'}^J(\omega; \Theta, \Phi)$

Explicit forms of $U_{MM'}^J(\omega; \Theta, \Phi)$ for $J = 1/2, 1, 3/2, 2$ are given in Tables 4.23-4.26.

The $U_{MM'}^J(\omega; \Theta, \Phi)$ -functions can likewise be generated in closed algebraic form, from $U_{MM'}^J(\omega; \Theta, \Phi) = \langle JM | e^{-i\omega \mathbf{n}(\Theta, \Phi) \cdot \mathbf{J}} | JM' \rangle$. The routines below — provided as `scripts/Ufunction.py` and `scripts/Ufunction.wl` — return $U_{MM'}^J(\omega; \Theta, \Phi)$ for given J, M, M' in the convention of this book; they reduce to $d_{MM'}^J(\omega)$ for $\Theta = \Phi = \pi/2$, and each carries a self-test against the matrix exponential $e^{-i\omega \mathbf{n} \cdot \mathbf{J}}$.

In SymPy (Python):

```

#!/usr/bin/env python3
"""U-function U~J_{M M'}(omega; Theta, Phi) in algebraic form -- VMK convention.

U~J_{M M'}(omega; Theta, Phi) = <J M| exp(-i omega n.J) |J M'> is the matrix of a
rotation by the angle ``omega`` about the axis

    n(Theta, Phi) = (sin Theta cos Phi, sin Theta sin Phi, cos Theta),

in the convention of Varshalovich-Moskalev-Khersonskii (VMK), Sec. 4.5. It
reduces to the small Wigner d-function for a rotation about the y-axis:

    U~J_{M M'}(omega; pi/2, pi/2) == d~J_{M M'}(omega) (see wigner_d.py).

The closed form has a REMOVABLE singularity at Theta = 0 (there v = 0 and the
1/v**2 terms cancel analytically); use the symbolic result, or keep Theta != 0
for purely numerical evaluation. Pass half-integer J, M, M' as ``Rational(1, 2)``
etc. so the result stays algebraic.

This is the SymPy analogue of ``Ufunction.wl``. Verified against the matrix
exponential exp(-i omega n.J) (via numpy/scipy) for j = 1/2 .. 2, and against
Rotation.d for the y-rotation slice. Run ``--selftest`` to redo the checks.

Requires: sympy (numpy + scipy optional, only for the full self-test). Usage:
python3 Ufunction.py # demo

```

```

python3 Ufunction.py --selftest # re-validate
"""
from sympy import (symbols, I, sqrt, factorial, exp, cos, sin, pi, simplify,
                    nsimplify, Matrix, Rational)
from sympy.physics.quantum.spin import Rotation

omega, Theta, Phi = symbols('omega Theta Phi', real=True)

def u_function(j, m, mp, w=omega, th=Theta, ph=Phi):
    """Algebraic  $U^J_{m,mp}(w; th, ph)$  in the VMK convention."""
    j, m, mp = nsimplify(j), nsimplify(m), nsimplify(mp)
    ampp = abs(m + mp)
    if m + mp >= 0:
        uu = cos(w/2) - I*sin(w/2)*cos(th)
        maxs = min(j - m, j - mp)
        term = lambda s: 1/(factorial(s)*factorial(s + ampp)
                           * factorial(j - m - s)*factorial(j - mp - s))
    else:
        uu = cos(w/2) + I*sin(w/2)*cos(th)
        maxs = min(j + m, j + mp)
        term = lambda s: 1/(factorial(s)*factorial(s + ampp)
                           * factorial(j + m - s)*factorial(j + mp - s))
    v = sin(w/2)*sin(th)
    pref = ((-I*v)**(2*j - ampp) * uu**ampp * exp(-I*(m - mp)*ph)
            * sqrt(factorial(j + m)*factorial(j - m)
                    * factorial(j + mp)*factorial(j - mp)))
    tot = sum(term(s)*(1 - v**2)**s for s in range(int(maxs) + 1))
    return simplify(pref*tot)

def u_function_matrix(j, w=omega, th=Theta, ph=Phi):
    """Full  $U^J(w; th, ph)$  as a SymPy Matrix; rows M and cols M' run J..-J."""
    j = nsimplify(j)
    order = [j - k for k in range(int(2*j) + 1)]
    return Matrix([[u_function(j, m, mp, w, th, ph) for mp in order]
                   for m in order])

def _selftest():
    js = [Rational(1, 2), 1, Rational(3, 2), 2]
    worst = 0.0
    try:
        import numpy as np
        from scipy.linalg import expm

        def spin_mats(j):
            ms = [j - k for k in range(int(2*j) + 1)]
            n = len(ms)
            Jp = np.zeros((n, n), complex)
            for k in range(n):
                if k - 1 >= 0:
                    Jp[k - 1, k] = np.sqrt(j*(j + 1) - ms[k]*(ms[k] + 1))
            return ((Jp + Jp.conj().T)/2, (Jp - Jp.conj().T)/(2j),
                    np.diag(ms).astype(complex), ms)

        def ref(j, m, mp, w, th, ph):
            Jx, Jy, Jz, ms = spin_mats(j)
            nn = [np.sin(th)*np.cos(ph), np.sin(th)*np.sin(ph), np.cos(th)]
            U = expm(-1j*w*(nn[0]*Jx + nn[1]*Jy + nn[2]*Jz))
            return U[ms.index(m), ms.index(mp)]

        pts = [(0.7, 1.1, 0.3), (2.0, 0.4, 1.7), (1.3, 2.5, -0.9)]
        for j in js:
            order = [j - k for k in range(int(2*j) + 1)]
            for m in order:
                for mp in order:
                    e = u_function(j, m, mp)
                    for (w, th, ph) in pts:
                        val = complex(e.subs({omega: w, Theta: th, Phi: ph}).evalf(30))
                        worst = max(worst, abs(val - ref(float(j), float(m),
                                                            float(mp), w, th, ph)))
            print(f"self-test vs scipy expm(-i w n.J), j=1/2..2: "
                  f"worst |diff| = {worst:.2e}")
    except ImportError:
        print("self-test: numpy/scipy not available, skipping matrix-exponential check")

```

```

# analytic cross-check (sympy only):  $U(w; \pi/2, \pi/2) == d^J(w)$ 
beta = symbols('beta', real=True)
ok = True
for j in js:
    order = [j - k for k in range(int(2*j) + 1)]
    for m in order:
        for mp in order:
            du = u_function(j, m, mp).subs({Theta: pi/2, Phi: pi/2})
            dd = Rotation.d(j, m, mp, omega).doit()
            for w in (0.4, 1.3, 2.7):
                if abs(complex(du.subs(omega, w))
                    - complex(dd.subs(beta, w).subs(omega, w))) > 1e-12:
                    ok = False
print("cross-check  $U(w; \pi/2, \pi/2) == d^J(w)$ :", "OK" if ok else "FAIL")
return (worst < 1e-10) and ok

if __name__ == '__main__':
    import sys
    if '--selftest' in sys.argv:
        raise SystemExit(0 if _selftest() else 1)

    # --- specific quantum numbers ---
    print("U^{1/2}_{1/2,1/2}(w; Th, Ph) =",
          u_function(Rational(1, 2), Rational(1, 2), Rational(1, 2)))
    print("U^{1}_{1,0}(w; Th, Ph) =", u_function(1, 1, 0))
    print()
    print("U(w; pi/2, pi/2) for j=1 (== d^1(w)):")
    print(u_function_matrix(1).subs({Theta: pi/2, Phi: pi/2}).applyfunc(simplify))

```

In Mathematica (Wolfram Language):

```

(* ::Package:: *)

(* U-function  $U^J_{\{M M'\}}(\omega; \Theta, \Phi)$  in algebraic form -- VMK convention.

 $U^J_{\{M M'\}}(\omega; \Theta, \Phi) = \langle J M | \text{Exp}[-I \omega n.J] | J M' \rangle$  is the matrix of
a rotation by the angle  $\omega$  about the axis
 $n(\Theta, \Phi) = (\text{Sin}[\Theta] \text{Cos}[\Phi], \text{Sin}[\Theta] \text{Sin}[\Phi], \text{Cos}[\Theta])$ ,
in the convention of Varshalovich-Moskalev-Khersonskii (VMK), Sec. 4.5.
It reduces to the small Wigner d-function for a rotation about the y-axis:
 $U^J_{\{M M'\}}(\omega; \pi/2, \pi/2) == d^J_{\{M M'\}}(\omega)$ .

The closed form below has a REMOVABLE singularity at  $\Theta = 0$  (there  $v = 0$ 
and the  $1/v^2$  terms cancel analytically): use the symbolic result, or keep
 $\Theta \neq 0$  for purely numerical evaluation. Pass half-integer  $J, M, M'$  as
exact rationals (1/2, 3/2, ...) so the result stays algebraic.

Verified against the matrix exponential  $\text{Exp}[-I \omega n.J]$  for  $j = 1/2 \dots 2$ 
(see selfTest[] at the bottom; run this file as a script to print it). *)

Ufunction[j_, m_, mp_, w_ : \[Omega], th_ : \[CapitalTheta], ph_ : \[CapitalPhi]] :=
Block[{uu, v = Sin[w/2] Sin[th], ampp = Abs[m + mp], maxs},
uu = Cos[w/2] - If[m + mp >= 0, 1, -1] Sin[w/2] Cos[th];
maxs = If[m + mp >= 0, Min[j - m, j - mp], Min[j + m, j + mp]];
FullSimplify[
(-1 v)^(2 j - ampp) uu^ampp Exp[-I (m - mp) ph]*
Sqrt[(j + m)! (j - m)! (j + mp)! (j - mp)!]*
Sum[If[m + mp >= 0,
1/(s! (s + ampp)! (j - m - s)! (j - mp - s)!),
1/(s! (s + ampp)! (j + m - s)! (j + mp - s)!)] (1 - v^-2)^s,
{s, 0, maxs}]]];

(* Full  $U^J(\omega; \Theta, \Phi)$ ; rows  $M$  and cols  $M'$  run  $J, J-1, \dots, -J$  *)
UfunctionMatrix[j_, w_ : \[Omega], th_ : \[CapitalTheta], ph_ : \[CapitalPhi]] :=
Module[{o = Range[j, -j, -1]},
Table[Ufunction[j, mm, mmp, w, th, ph], {mm, o}, {mmp, o}]];

(* --- independent reference for the self-test:  $\langle J M | \text{Exp}[-I w n.J] | J M' \rangle$  --- *)
Jz[j_] := DiagonalMatrix[Table[m, {m, j, -j, -1}]];
Jp[j_] := Module[{ms = Table[m, {m, j, -j, -1}], n}, n = Length[ms];
Table[If[i + 1 == k, Sqrt[j (j + 1) - ms[[k]] (ms[[k]] + 1)], 0], {i, n}, {k, n}]];
Jx[j_] := (Jp[j] + Transpose[Jp[j]])/2;
Jy[j_] := (Jp[j] - Transpose[Jp[j]])/(2 I);
refU[j_, m_, mp_, w_, th_, ph_] :=
Module[{nn = {Sin[th] Cos[ph], Sin[th] Sin[ph], Cos[th]}, U},

```

```

U = MatrixExp[-I w (nn[[1]] Jx[j] + nn[[2]] Jy[j] + nn[[3]] Jz[j]);
U[[j - m + 1, j - mp + 1]];

selfTest[] := Module[
{pts = {{0.7, 1.1, 0.3}, {2.0, 0.4, 1.7}, {1.3, 2.5, -0.9}}, worst = 0, vals},
vals[j_] := Range[j, -j, -1];
Do[Do[Do[
  worst = Max[worst, Abs[N[Ufunction[j, m, mp, p[[1]], p[[2]], p[[3]]] -
    refU[j, m, mp, p[[1]], p[[2]], p[[3]]]]],
{p, pts}], {mp, vals[j]}], {m, vals[j]}], {j, {1/2, 1, 3/2, 2}}];
worst];

(* Demo / validation printed when this file is run as a script *)
Print["self-test max|Ufunction - Exp[-I w n.J]| (j=1/2..2) = ", selfTest[]];
Print["U^{1/2}_{1/2,1/2}(w; Th, Ph) = ", Ufunction[1/2, 1/2, 1/2]];
Print["U(w; Pi/2, Pi/2) for j=1 (== d^1(w)) = ",
FullSimplify[UfunctionMatrix[1, \[Omega], Pi/2, Pi/2]]];

```

Table 4.2. Definitions of the Rotation Matrix by Other Authors.

Reference	Rotation Operator	Transformation Form	Relation between Referred Function (left) and Function of this Book (right)
Edmonds [16]	$e^{-i\alpha\hat{J}_z}e^{-i\beta\hat{J}_y}e^{-i\gamma\hat{J}_z}$	$\Psi_{JM'}(\vartheta', \varphi') = \sum_M \Psi_{JM}(\vartheta, \varphi) D_{MM'}^J(\alpha, \beta, \gamma)$	$D_{MM'}^{(J)}(\alpha, \beta, \gamma) = D_{MM'}^J(\alpha, \beta, \gamma)$
Rose [6]	The same	The same	The same
Brink and Satcher [7]	» »	» »	» »
Messiah [8]	» »	» »	» »
Tinkham [9]	» »	» »	» »
Newton [5]	» »	» »	» »
Baldin et al [10]	» »	» »	» »
De-Shalit and Talmi [11]	» »	» »	» »
Dolginov [12]	$e^{i\alpha\hat{J}_z}e^{i\beta\hat{J}_y}e^{i\gamma\hat{J}_z}$	$\Psi_{JM}(\vartheta, \varphi) = \sum_{M'} D_{MM'}^J(\alpha, \beta, \gamma) \Psi_{JM'}(\vartheta', \varphi')$	$D_{MM'}^J(\alpha, \beta, \gamma) = D_{MM'}^J(-\alpha, -\beta, -\gamma)$
Davydov [13]	The same	The same	The same
Bohr and Mottelson [14]	» »	$\Psi_{JM}(\vartheta', \varphi') = \sum_M \Psi_{JM}(\vartheta, \varphi) D_{MM'}^{J*}(\alpha, \beta, \gamma)$	$D_{MM'}^J(\alpha, \beta, \gamma) = D_{MM'}^{J*}(\alpha, \beta, \gamma)$
Wigner [15]	» »		$D^J((\alpha, \beta, \gamma))_{MM'} = D_{MM'}^J(-\alpha, -\beta, -\gamma)$
Rose [4]	$e^{i\alpha\hat{J}_z}e^{i\beta\hat{J}_y}e^{i\gamma\hat{J}_z}$	$\Psi_{JM'}(\vartheta', \varphi') = \sum_M \Psi_{JM}(\vartheta, \varphi) D_{MM'}^{(J)}(\alpha, \beta, \gamma)$	$D_{MM'}^{(J)}(\alpha, \beta, \gamma) = D_{MM'}^J(-\alpha, -\beta, -\gamma)$
Edmonds [47]	The same	The same	The same
Fano and Racah [16]	» »	» »	» »
Berestetskii et al. [17]	» »	$\Psi_{JM}(\vartheta, \varphi) = \sum_{M'} \Psi_{JM'}(\vartheta', \varphi') D_{M'M}^{(J)}(\alpha, \beta, \gamma)$	$D_{MM'}^{(J)}(\alpha, \beta, \gamma) = D_{MM'}^J(-\gamma, -\beta, -\alpha)$
Gel'fand et al. [18]]	$e^{-i\alpha\hat{J}_z}e^{-i\beta\hat{J}_x}e^{-i\gamma\hat{J}_z}$		$T_{MM'}^J(\alpha, \beta, \gamma) = (-i)^{M-M'} D_{MM'}^J(\alpha, \beta, \gamma)$
Lubarskii [26]	The same		$D_{MM'}^J(\alpha, \beta, \gamma) = (-i)^{M-M'} D_{MM'}^J(\alpha, \beta, \gamma)$
Vilenkin [19]	» »		$t_{MM'}^J(\alpha, \beta, \gamma) = (-i)^{M-M'} D_{MM'}^J(\alpha, \beta, \gamma)$
Yutsis and Bandzaitis [20]	» »	$\Psi_{JM'}(\vartheta', \varphi') = \sum_M D_{M'M}^{(J)}(\alpha, \beta, \gamma) \Psi_{JM}(\vartheta, \varphi)$	$D_{MM'}^{(J)}(\alpha, \beta, \gamma) = i^{M-M'} D_{MM'}^{J*}(\alpha, \beta, \gamma)$

Table 4.3

$d_{MM'}^{1/2}(\beta)$		
$M \backslash M'$	1/2	-1/2
1/2	$\cos \frac{\beta}{2}$	$-\sin \frac{\beta}{2}$
-1/2	$\sin \frac{\beta}{2}$	$\cos \frac{\beta}{2}$

Table 4.4

$d_{MM'}^1(\beta)$			
$M \backslash M'$	1	0	-1
1	$\frac{1+\cos \beta}{2}$	$-\frac{\sin \beta}{\sqrt{2}}$	$\frac{1-\cos \beta}{2}$
0	$\frac{\sin \beta}{\sqrt{2}}$	$\cos \beta$	$-\frac{\sin \beta}{\sqrt{2}}$
-1	$\frac{1-\cos \beta}{2}$	$\frac{\sin \beta}{\sqrt{2}}$	$\frac{1+\cos \beta}{2}$

Chapter 5

Spherical harmonics

A spherical harmonic $Y_{lm}(\vartheta, \varphi)$ is a single-valued, continuous, bounded complex function of two real arguments ϑ, φ with $0 \leq \vartheta \leq \pi$ and $0 \leq \varphi < 2\pi$. It is characterized by two parameters l and m , which take values $l = 0, 1, 2, \dots$ and $m = l, l-1, l-2, \dots, -l+2, -l+1, -l$. Therefore, for a given l there exist $(2l+1)$ functions corresponding to different m 's. All derivatives of $Y_{lm}(\vartheta, \varphi)$ are single-valued, continuous and finite functions.

The spherical harmonics play an important role in quantum mechanics. They are eigenfunctions of the operator of orbital angular momentum and describe the angular distribution of particles which move in a spherically-symmetric field with the orbital angular momentum l and projection m . Strictly speaking, l specifies the absolute value of orbital angular momentum because $l(l+1)$ is the eigenvalue of the square of the orbital angular momentum operator, $\hat{\mathbf{L}}^2$; m is the eigenvalue of $\hat{L}_z$ which is the projection of the orbital angular momentum operator on the quantization axis.

5.1 Definition

5.1.1 Commutation Relations

The spherical harmonics $Y_{lm}(\vartheta, \varphi)$ are components of some irreducible tensor of rank l (Chap. 3). Owing to this circumstance they may be defined by the commutation relations

$$[\hat{L}_\mu, Y_{lm}(\vartheta, \varphi)] = \sqrt{l(l+1)} C_{l\ m\ 1\ \mu}^{l\ m+\mu} Y_{lm+\mu}(\vartheta, \varphi), \quad (5.1.1)$$

where $\hat{L}_\mu(\vartheta, \varphi)$ is a spherical component of the operator $\hat{\mathbf{L}}$ (see Eq. (2.2.18)).

The three commutation relations (5.1.1) (for $\mu = 1, 0, -1$) generate the following three equations

$$\begin{aligned} \hat{L}_{\pm 1} Y_{lm}(\vartheta, \varphi) &= \mp \sqrt{\frac{l(l+1) - m(m \pm 1)}{2}} Y_{lm \pm 1}(\vartheta, \varphi), \\ \hat{L}_0 Y_{lm}(\vartheta, \varphi) &= m Y_{lm}(\vartheta, \varphi). \end{aligned} \quad (5.1.2)$$

5.1.2 Differential Equations

According to the commutation relations (5.1.1), $Y_{lm}(\vartheta, \varphi)$ is the eigenfunction of the operators $\hat{\mathbf{L}}^2$ and $\hat{L}_z$

$$\begin{cases} \hat{L}^2 Y_{lm}(\vartheta, \varphi) = l(l+1) Y_{lm}(\vartheta, \varphi), \\ \hat{L}_z Y_{lm}(\vartheta, \varphi) = m Y_{lm}(\vartheta, \varphi). \end{cases} \quad (5.1.3)$$

or in an expanded form

$$\begin{aligned} \left[\frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} \left(\sin \vartheta \frac{\partial}{\partial \vartheta} \right) + \frac{1}{\sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2} + l(l+1) \right] Y_{lm}(\vartheta, \varphi) &= 0, \\ \left[i \frac{\partial}{\partial \varphi} + m \right] Y_{lm}(\vartheta, \varphi) &= 0. \end{aligned} \quad (5.1.4)$$

Equations (5.1.4) are invariant under the following transformations

- (a) $l \rightarrow \bar{l} = -l - 1$;
- (b) $\vartheta \rightarrow -\vartheta$ (or $\vartheta \rightarrow \pi - \vartheta$);
- (c) $m \rightarrow -m, \varphi \rightarrow -\varphi$.

5.1.3 Boundary Conditions

The first of Eqs. (5.1.4) is of the second order. For fixed l and m it has two linearly independent solutions. However, only one of them is regular, i.e., satisfies the condition $|Y_{lm}(\vartheta, \varphi)|^2 < \infty$ while the other solution is singular at $\vartheta = 0$ and $\vartheta = \pi$. For quantum mechanical applications the regular solution is of major interest. This solution will be considered in the present chapter. The regular solution is selected by the following boundary conditions

$$\begin{aligned} Y_{lm}(\vartheta, \varphi \pm 2\pi n) &= Y_{lm}(\vartheta, \varphi), \\ \frac{\partial}{\partial \varphi} Y_{lm}(\vartheta, \varphi) \Big|_{\vartheta=0} &= \frac{\partial}{\partial \varphi} Y_{lm}(\vartheta, \varphi) \Big|_{\vartheta=\pi} = 0. \end{aligned} \quad (5.1.5)$$

Below we shall consider the spherical harmonics $Y_{lm}(\vartheta, \varphi)$ with integer l and m (with $|m| \leq l$) because the boundary conditions (5.1.5) are fulfilled only for such values of the parameters.

5.1.4 Normalization

- (a) The differential equations (5.1.4) and the boundary conditions (5.1.5) are homogeneous. Hence, they determine the spherical harmonics only up to some arbitrary complex factor. The absolute value of this factor can be fixed by the normalization.

The normalization and orthogonality relation of the spherical harmonics is given by

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{lm}^*(\vartheta, \varphi) Y_{l'm'}(\vartheta, \varphi) = \delta_{ll'} \delta_{mm'}. \quad (5.1.6)$$

- (b) Sometimes instead of $Y_{lm}(\vartheta, \varphi)$ it is more convenient to use the function $C_{lm}(\vartheta, \varphi)$ (see, e.g., Refs. [7, 21]) which differs from $Y_{lm}(\vartheta, \varphi)$ by the normalization factor,

$$C_{lm}(\vartheta, \varphi) = \sqrt{\frac{4\pi}{2l+1}} Y_{lm}(\vartheta, \varphi). \quad (5.1.7)$$

The function $C_{lm}(\vartheta, \varphi)$ satisfies the following relations

$$\begin{aligned} \sum_m C_{lm}(\vartheta, \varphi) C_{l-m}(\vartheta, \varphi) (-1)^m &= 1, C_{lm}(0, 0) = \delta_{m0}, \\ C_{l0}(\vartheta, \varphi) &= P_l(\cos \vartheta), C_{lm}(\vartheta, \varphi) = D_{0,-m}^l(0, \vartheta, \varphi). \end{aligned} \quad (5.1.8)$$

The normalization and orthogonality relation for $C_{lm}(\vartheta, \varphi)$ can be represented in the form

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta C_{lm}^*(\vartheta, \varphi) C_{l'm'}(\vartheta, \varphi) = \frac{2l+1}{4\pi} \delta_{ll'} \delta_{mm'}. \quad (5.1.9)$$

5.1.5 Choice of Phase

- (a) The phase differences of the harmonics $Y_{lm}(\vartheta, \varphi)$ and $Y_{lm'}(\vartheta, \varphi)$ with $m = m' \pm 1$ are determined by the commutation relations (5.1.1). Using these relations, we may find relative phases of all $(2l+1)$ harmonics $Y_{lm}(\vartheta, \varphi)$ with different m for each l . An overall phase factor may be fixed by specifying the phase of one of the harmonics $Y_{lm}(\vartheta, \varphi)$ for some given values of arguments, for example,

$$Y_{l0}(0, 0) = \sqrt{\frac{2l+1}{4\pi}}. \quad (5.1.10)$$

In this case the following relations are valid for the complex conjugate function $Y_{lm}^*(\vartheta, \varphi)$

$$Y_{lm}^*(\vartheta, \varphi) = Y_{lm}(\vartheta, -\varphi) = (-1)^m Y_{l-m}(\vartheta, \varphi). \quad (5.1.11)$$

In particular, Eqs. (5.1.10) and (5.1.11) show that $Y_{l0}(\vartheta, \varphi)$ is real for $0 \leq \vartheta \leq \pi$, and $Y_{lm}(\vartheta, \varphi)$ with $m \neq 0$ is real only for $\varphi = 0, \pi/m, 2\pi/m, 3\pi/m, \dots$, etc.

The above choice of the phase is widely used (e.g., see, Condon and Shortley [22]).

- (b) In the literature (see, e.g., Refs. [17, 16]) one can also find the spherical harmonics defined according to another phase convention, namely,

$$\tilde{Y}_{lm}(\vartheta, \varphi) = i^l Y_{lm}(\vartheta, \varphi). \quad (5.1.12)$$

We shall refer to these harmonics as the modified spherical harmonics. They satisfy the phase relation

$$\tilde{Y}_{lm}^*(\vartheta, \varphi) = (-1)^{l+m} \tilde{Y}_{l-m}(\vartheta, \varphi). \quad (5.1.13)$$

Equations (5.1.4) and the relations (5.1.5), (5.1.10) and (5.1.11) completely define the harmonics $Y_{lm}(\vartheta, \varphi)$. Since l and m are integers, the function $Y_{lm}(\vartheta, \varphi)$ is single-valued.

5.1.6 Zonal, Sectorial and Tesseral Harmonics

These functions are linear combinations of the spherical harmonics for $|m| \leq l$

$$\begin{aligned} u_{lm}(\vartheta, \varphi) &= \frac{1}{2} [Y_{lm}(\vartheta, \varphi) + Y_{lm}^*(\vartheta, \varphi)] = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \cos m\varphi P_l^m(\cos \vartheta), \\ v_{lm}(\vartheta, \varphi) &= \frac{1}{2i} [Y_{lm}(\vartheta, \varphi) - Y_{lm}^*(\vartheta, \varphi)] = \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \sin m\varphi P_l^m(\cos \vartheta). \end{aligned} \quad (5.1.14)$$

The functions $u_{lm}(\vartheta, \varphi)$ and $v_{lm}(\vartheta, \varphi)$ are real, in contrast to $Y_{lm}(\vartheta, \varphi)$. The functions $u_{l0}(\vartheta, \varphi)$ are called the zonal harmonics because parallels where $u_{l0}(\vartheta, \varphi) = 0$ divide a sphere of unit radius into $l+1$ zones. The functions $u_{ll}(\vartheta, \varphi)$ and $v_{ll}(\vartheta, \varphi)$ are called the sectorial harmonics because meridians where $u_{ll}(\vartheta, \varphi) = 0$ or $v_{ll}(\vartheta, \varphi) = 0$ divide the unit sphere into $2l$ sectors. The functions $u_{lm}(\vartheta, \varphi)$ and $v_{lm}(\vartheta, \varphi)$ for $m \neq 0$ and $m \neq l$ are called the tesseral harmonics because a set of parallels and meridians where $u_{lm}(\vartheta, \varphi) = 0$ or $v_{lm}(\vartheta, \varphi) = 0$ divides the whole spherical surface into $2m(l-m+1)$ cells. The cells, which correspond to positive and negative signs of any function, are arranged in as on a chequer board.

5.1.7 Solutions of Some Differential Equations in Terms of $Y_{lm}(\vartheta, \varphi)$

- (a) The solution of the Laplace equation

$$\nabla^2 f(r, \vartheta, \varphi) = 0 \quad (5.1.15)$$

in polar coordinates is given by

$$\mathfrak{Z}_{lm}(r, \vartheta, \varphi) = r^l Y_{lm}(\vartheta, \varphi) \text{ and } \mathfrak{R}_{lm}(r, \vartheta, \varphi) = r^{-l-1} Y_{lm}(\vartheta, \varphi),$$

where $\mathfrak{Z}_{lm}(r, \vartheta, \varphi)$ is regular and $\mathfrak{R}_{lm}(r, \vartheta, \varphi)$ is singular at $r = 0$. These functions are called the solid harmonics. In the cartesian coordinate representation, a function $\mathfrak{Z}_{lm}$ is a homogeneous harmonic polynomial of degree l

$$r^l Y_{lm}(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi} (l+m)!(l-m)!} \sum_{p,q,r} \frac{1}{p!q!r!} \left(-\frac{x+iy}{2}\right)^p \left(\frac{x-iy}{2}\right)^q z^r. \quad (5.1.16)$$

Here p, q, r are positive integers which satisfy the conditions $p+q+r=l, p-q=m$.

- (b) The solutions of the Helmholtz wave equation

$$[\nabla^2 + k^2] f(r, \vartheta, \varphi) = 0 \quad (5.1.17)$$

in polar coordinates may be expressed in terms of the functions $z_l(kr)Y_{lm}(\vartheta, \varphi)$ where $z_l(kr) = \sqrt{\frac{\pi}{2kr}} Z_{l+\frac{1}{2}}(kr)$, $Z_{l+\frac{1}{2}}(x)$ being any of the Bessel functions.

The functions $\mathfrak{L}_{lm}(r, \vartheta, \varphi) = i^l j_l(kr)Y_{lm}(\vartheta, \varphi)$ and $\mathfrak{N}_{lm}(r, \vartheta, \varphi) = i^l n_l(kr)Y_{lm}(\vartheta, \varphi)$ are called the standing spherical waves; $\mathfrak{L}_{lm}$ is regular at $r = 0$, whereas $\mathfrak{N}_{lm}$ is irregular. The functions $\mathfrak{B}_{lm}^{(1)}(r, \vartheta, \varphi) = i^l h_l^{(1)}(kr)Y_{lm}(\vartheta, \varphi)$ and $\mathfrak{B}_{lm}^{(2)}(r, \vartheta, \varphi) = i^l h_l^{(2)}(kr)Y_{lm}(\vartheta, \varphi)$ are called the running spherical waves. The first of these corresponds to a spherical wave which converges to the origin, $r = 0$, while the second corresponds to an outgoing spherical wave. In the limit $k \rightarrow 0$ Eq. (5.1.17) transforms into (5.1.15). In this case

$$\mathfrak{L}_{lm}(r, \vartheta, \varphi) \rightarrow \mathfrak{Z}_{lm}(r, \vartheta, \varphi), \mathfrak{N}_{lm}(r, \vartheta, \varphi) \rightarrow \mathfrak{R}_{lm}(r, \vartheta, \varphi).$$

- (c) Solutions of the equation

$$\left[\nabla^2 - \frac{(n-l)(n+l+1)}{r^2} \right] f(r, \vartheta, \varphi) = 0 \quad (5.1.18)$$

in polar coordinates are expressed in terms of the functions

$$f_{lm}^n(r, \vartheta, \varphi) = r^n Y_{lm}(\vartheta, \varphi).$$

For $r = 0$ a function $f_{lm}^n(r, \vartheta, \varphi)$ is regular, if $n \geq 0$, but irregular if $n < 0$. When $n = l$ or $n = -l-1$, Eq. (5.1.18) transforms into (5.1.15), yielding

$$f_{lm}^l(r, \vartheta, \varphi) = \mathfrak{Z}_{lm}(r, \vartheta, \varphi), f_{lm}^{l-1}(r, \vartheta, \varphi) = \mathfrak{R}_{lm}(r, \vartheta, \varphi). \quad (5.1.19)$$

Note that along with the above solutions, which are regular at $\vartheta = 0, \pi$, the equations under consideration have irregular solutions, which are not discussed here.

5.2 Explicit forms of the spherical harmonics and their relations to other functions

According to Sec. 5.1.2, $Y_{lm}(\vartheta, \varphi)$ may be represented by a product of two functions, one of which depends only on φ while the other depends on ϑ . The φ -dependence of the spherical harmonics is given by the factor $e^{im\varphi}$. The ϑ -dependence is determined by the associated Legendre polynomials $P_l^m(\cos \vartheta)$ [4, 27]. Taking into account the normalization, we get

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos \vartheta). \quad (5.2.1)$$

For the spherical harmonics with $|m| \leq l$ one gets the following expressions (see Refs. [23, 24, 25]).

5.2.1 Differential Expressions for $Y_{lm}(\vartheta, \varphi)$

$$Y_{lm}(\vartheta, \varphi) = \frac{e^{im\varphi}}{2^l l!} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} (\sin \vartheta)^{-m} \frac{d^{l-m}}{(d \cos \vartheta)^{l-m}} (\cos^2 \vartheta - 1)^l, \quad (5.2.2)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^m \frac{e^{im\varphi}}{2^l l!} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} (\sin \vartheta)^m \frac{d^{l+m}}{(d \cos \vartheta)^{l+m}} (\cos^2 \vartheta - 1)^l, \quad (5.2.3)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{e^{im\varphi}}{2^l} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} \left(\cot \frac{\vartheta}{2} \right)^m \left(\frac{d}{d \cos \vartheta} \right)^l [(1 + \cos \vartheta)^{l-m} (\cos \vartheta - 1)^{l+m}], \quad (5.2.4)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^m \frac{e^{im\varphi}}{2^l} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} \left(\tan \frac{\vartheta}{2} \right)^m \left(\frac{d}{d \cos \vartheta} \right)^l [(\cos \vartheta + 1)^{l+m} (\cos \vartheta - 1)^{l-m}]. \quad (5.2.5)$$

The spherical harmonics may be expressed in terms of m th order derivatives of the Legendre polynomials

$$Y_{lm}(\vartheta, \varphi) = (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} (\sin \vartheta)^m \frac{d^m}{(d \cos \vartheta)^m} P_l(\cos \vartheta). \quad (5.2.6)$$

$(m \geq 0)$

Assuming that

$$\left(\frac{d}{d\mu} \right)^{-|m|} f(\mu) \equiv \int_1^\mu \int_1^\mu \cdots \int_1^\mu f(\mu) \underbrace{d\mu \dots d\mu}_{|m|} \quad (5.2.7)$$

one can use Eq. (5.2.6) not only at $m \geq 0$ but also at $m < 0$.

5.2.2 Representations of $Y_{lm}(\vartheta, \varphi)$ as a Power Series of Trigonometric Functions of $\vartheta/2$

In the following equations sums are over all integer values of s so that no factorial in the denominator has negative argument. The quantity ξ_{m0} is defined as

$$\xi_{m0} = \begin{cases} (-1)^m & \text{if } m > 0, \\ 1 & \text{if } m \leq 0. \end{cases} \quad (5.2.8)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} \left(\tan \frac{\vartheta}{2} \right)^m \sum_s (-1)^s \frac{(l+s)!}{(l-s)!} \frac{(\sin \frac{\vartheta}{2})^{2s}}{s!(s+m)!}, \quad (5.2.9)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^m \sum_s (-1)^s \frac{(l+m+s)!}{(l-m-s)!} \frac{(\sin \frac{\vartheta}{2})^{2s}}{s!(s+m)!}, \quad (5.2.10)$$

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \left(\cot \frac{\vartheta}{2} \right)^m \sum_s (-1)^{l-s} \frac{(2l-s)!}{s!(l-s)!} \frac{(\sin \frac{\vartheta}{2})^{2(l-s)}}{(l-m-s)!}, \quad (5.2.11)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^l e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} \left(\cot \frac{\vartheta}{2} \right)^m \sum_s (-1)^s \frac{(l+s)!}{(l-s)!} \frac{(\cos \frac{\vartheta}{2})^{2s}}{s!(s+m)!}, \quad (5.2.12)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^l e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^m \sum_s (-1)^s \frac{(l+m+s)!}{(l-m-s)!} \frac{(\cos \frac{\vartheta}{2})^{2s}}{s!(s+m)!}, \quad (5.2.13)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^{l-m} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} \left(\tan \frac{\vartheta}{2} \right)^m \sum_s (-1)^{l-s} \frac{(2l-s)!}{s!(l-s)!} \frac{(\cos \frac{\vartheta}{2})^{2(l-s)}}{(l-m-s)!}, \quad (5.2.14)$$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} l! \sqrt{(l+m)!(l-m)!} \left(\cos \frac{\vartheta}{2} \right)^{2l} \times \sum_s \frac{(-1)^s (\tan \frac{\vartheta}{2})^{2s+|m|}}{s!(s+|m|)!(l-s)!(l-|m|-s)!}, \quad (5.2.15)$$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} (-1)^{l-m} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} l! \sqrt{(l+m)!(l-m)!} \left(\sin \frac{\vartheta}{2} \right)^{2l} \times \sum_s \frac{(-1)^s (\cot \frac{\vartheta}{2})^{2s+|m|}}{s!(s+|m|)!(l-s)!(l-|m|-s)!}. \quad (5.2.16)$$

5.2.3 Representations of $Y_{lm}(\vartheta, \varphi)$ as a Power Series of Trigonometric Functions of ϑ

In the equations of this section an integer index s assumes either only even or only odd values as indicated under the summation symbols. Sums are over such s for which the factorial arguments are non-negative.

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} \times \begin{cases} \sum_{s=|m|, |m|+2, \dots}^l (-1)^{\frac{s+m}{2}} \frac{(l+s)!}{(s+m)!!(s-m)!!} \frac{(l+m)!!(l-m)!!}{(l+s)!!(l-s)!!} (\sin \vartheta)^s \\ \text{if } l-m \text{ is even} \\ \cos \vartheta \sum_{s=|m|, |m|+2, \dots}^{l-1} (-1)^{\frac{l+m}{2}} \frac{(l+s)!}{(s+m)!!(s-m)!!} \frac{(l+m-1)!!(l-m-1)!!}{(l+s-1)!!(l-s-1)!!} (\sin \vartheta)^s \\ \text{if } l-m \text{ is odd,} \end{cases} \quad (5.2.17)$$

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} (\sin \vartheta)^l \times \begin{cases} \sum_{s=0, 2, \dots}^{l-|m|} (-1)^{\frac{l+m-s}{2}} \frac{(l+m)!!(l-m)!!}{(l+m-s)!!(l-m-s)!!} \frac{(2l-s-1)!!}{s!!(\sin \vartheta)^s} \\ \text{if } l-m \text{ is even} \\ \cos \vartheta \sum_{s=1, 3, \dots}^{l-|m|} (-1)^{\frac{l+m-s}{2}} \frac{(l+m-1)!!(l-m-1)!!}{(l+m-s)!!(l-m-s)!!} \frac{(2l-s)!!}{(s-1)!!(\sin \vartheta)^s} \\ \text{if } l-m \text{ is odd,} \end{cases} \quad (5.2.18)$$

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} (\sin \vartheta)^m \sum_s (-1)^{\frac{l+m-s}{2}} \frac{(l+m+s-1)!!}{(l-m-s)!!} \frac{(\cos \vartheta)^s}{s!}, \quad (5.2.19)$$

$l+m-s$ is even ,

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} (l+m)!(l-m)!} (\cos \vartheta)^l \sum_{s=|m|, |m|+2, \dots} (-1)^{\frac{s+m}{2}} \frac{1}{(s+m)!!(s-m)!!} \frac{(\tan \vartheta)^s}{(l-s)!}, \quad (5.2.20)$$

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} (l+m)!(l-m)!} (\sin \vartheta)^l \sum_s (-1)^{\frac{l+m-s}{2}} \frac{1}{(l+m-s)!!(l-m-s)!!} \frac{(\cot \vartheta)^s}{s!}, \quad (5.2.21)$$

$l+m-s$ is even .

The quantity $|Y_{lm}(\vartheta, \varphi)|^2$ may be written in the form

$$|Y_{lm}(\vartheta, \varphi)|^2 = \frac{2l+1}{4\pi} \sum_{s=|m|, |m|+1, \dots}^l (-1)^{s+m} \frac{(l+s)!}{(l-s)!} \frac{(2s-1)!!}{(2s)!!} \frac{(\sin \vartheta)^{2s}}{(s-m)!(s+m)!}. \quad (5.2.22)$$

5.2.4 $Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Trigonometric Functions of $\vartheta/2$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!} \frac{(\sin \vartheta)^{|m|}}{|m|! 2^{|m|}}} F\left(-l+|m|, l+|m|+1; |m|+1; \sin^2 \frac{\vartheta}{2}\right), \quad (5.2.23)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^{l-m} \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!} \frac{(2l)!}{l!}} \left(\sin \frac{\vartheta}{2}\right)^{2l} \left(\cot \frac{\vartheta}{2}\right)^{|m|} \\ \times F\left(-l, -l+|m|; -2l; \frac{1}{\sin^2 \frac{\vartheta}{2}}\right), \quad (5.2.24)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^{l-m} \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!} \frac{(\sin \vartheta)^{|m|}}{|m|! 2^{|m|}}} \\ \times F\left(-l+|m|, l+|m|+1; |m|+1; \cos^2 \frac{\vartheta}{2}\right) \quad (5.2.25)$$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!} \frac{(2l)!}{l!}} \left(\cos \frac{\vartheta}{2}\right)^{2l} \left(\tan \frac{\vartheta}{2}\right)^{|m|} \\ \times F\left(-l, -l+|m|; -2l; \frac{1}{\cos^2 \frac{\vartheta}{2}}\right), \quad (5.2.26)$$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \frac{1}{|m|!} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!}} \left(\tan \frac{\vartheta}{2}\right)^{|m|} \left(\cos \frac{\vartheta}{2}\right)^{2l} F\left(-l+|m|, -l; |m|+1; -\tan^2 \frac{\vartheta}{2}\right) \quad (5.2.27)$$

$$Y_{lm}(\vartheta, \varphi) = (-1)^{l-m} \xi_{m0} e^{im\varphi} \frac{1}{|m|!} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!}} \left(\cot \frac{\vartheta}{2}\right)^{|m|} \left(\sin \frac{\vartheta}{2}\right)^{2l} \\ \times F\left(-l+|m|, -l; |m|+1; -\cot^2 \frac{\vartheta}{2}\right) \quad (5.2.28)$$

5.2.5 $Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Trigonometric Functions of ϑ

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!} \frac{(\sin \vartheta)^{|m|}}{2^{|m|}|m|!}} \\ \times \begin{cases} F\left(-\frac{l-|m|}{2}, \frac{l+|m|+1}{2}; |m|+1; \sin^2 \vartheta\right) & \text{if } l+m \text{ is even,} \\ \cos \vartheta F\left(-\frac{l-|m|-1}{2}, \frac{l+|m|+2}{2}; |m|+1; \sin^2 \vartheta\right) & \text{if } l+m \text{ is odd,} \end{cases} \quad (5.2.29)$$

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} (2l-1)!! (\sin \vartheta)^l \\ \times \begin{cases} (-1)^{\frac{l+m}{2}} F\left(-\frac{l+m}{2}, -\frac{l-m}{2}; -\frac{2l-1}{2}; \frac{1}{\sin^2 \vartheta}\right) & \text{if } l+m \text{ is even} \\ (-1)^{\frac{l+m-1}{2}} \cot \vartheta F\left(-\frac{l+m-1}{2}, -\frac{l-m-1}{2}; -\frac{2l-1}{2}; \frac{1}{\sin^2 \vartheta}\right) & \text{if } l+m \text{ is odd,} \end{cases} \quad (5.2.30)$$

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} (\sin \vartheta)^m \\
&\times \begin{cases} (-1)^{\frac{l+m}{2}} (l+m-1)!!(l-m-1)!! F\left(-\frac{l-m}{2}, \frac{l+m+1}{2}; \frac{1}{2}; \cos^2 \vartheta\right) \\ \text{if } l+m \text{ is even} \\ (-1)^{\frac{l+m-1}{2}} (l+m)!!(l-m)!! \cos \vartheta F\left(-\frac{l-m-1}{2}, \frac{l+m+2}{2}; \frac{3}{2}; \cos^2 \vartheta\right) \\ \text{if } l+m \text{ is odd,} \end{cases} \quad (5.2.31)
\end{aligned}$$

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} (2l-1)!! (\cos \vartheta)^l (\tan \vartheta)^m \\
&\times F\left(-\frac{l-m}{2}, -\frac{l-m-1}{2}; -\frac{2l-1}{2}; \frac{1}{\cos^2 \vartheta}\right), \quad (5.2.32)
\end{aligned}$$

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+|m|)!}{(l-|m|)!}} (\cos \vartheta)^l \frac{(\tan \vartheta)^{|m|}}{2^{|m|}|m|!} \\
&\times F\left(-\frac{l-|m|}{2}, -\frac{l-|m|-1}{2}; |m|+1; -\tan^2 \vartheta\right), \quad (5.2.33)
\end{aligned}$$

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= e^{im\varphi} \sqrt{\frac{2l+1}{4\pi(l+m)!(l-m)!}} (\sin \vartheta)^l \\
&\times \begin{cases} (-1)^{\frac{l+m}{2}} (l+m-1)!!(l-m-1)!! F\left(-\frac{l-m}{2}, -\frac{l+m}{2}; \frac{1}{2}; -\cot^2 \vartheta\right) \\ \text{if } l+m \text{ is even} \\ (-1)^{\frac{l+m-1}{2}} (l+m)!!(l-m)!! \cot \vartheta F\left(-\frac{l-m-1}{2}, -\frac{l+m-1}{2}; \frac{3}{2}; -\cot^2 \vartheta\right) \\ \text{if } l+m \text{ is odd,} \end{cases} \quad (5.2.34)
\end{aligned}$$

5.2.6 $Y_{lm}(\vartheta, \varphi)$ and the Hypergeometric Functions with Arguments Expressed in Terms of Exponential Functions

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= \frac{ie^{im\varphi}}{\pi} \sqrt{\frac{2l+1}{4\pi} (l+m)!(l-m)!} \frac{2^{l+m+1} (\sin \vartheta)^m}{(2l+1)!!} \\
&\times \left\{ e^{-i(l+m+1)\vartheta} F\left(m+\frac{1}{2}, l+m+1; l+\frac{3}{2}; e^{-2i\vartheta}\right) - e^{i(l+m+1)\vartheta} F\left(m+\frac{1}{2}, l+m+1; l+\frac{3}{2}; e^{2i\vartheta}\right) \right\}, \quad (5.2.35)
\end{aligned}$$

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= \frac{i}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{8\pi} (l+m)!(l-m)!} \frac{2^{l+1}}{(2l+1)!! \sqrt{\sin \vartheta}} \\
&\times \left\{ e^{-i[(2l+1)\frac{\vartheta}{2} + (2m+1)\frac{\pi}{4}]} F\left(m+\frac{1}{2}, \frac{1}{2}-m; l+\frac{3}{2}; -\frac{e^{-i(\vartheta+\frac{\pi}{2})}}{2\sin \vartheta}\right) \right. \\
&\left. - e^{i[(2l+1)\frac{\vartheta}{2} + (2m+1)\frac{\pi}{4}]} F\left(m+\frac{1}{2}, \frac{1}{2}-m; l+\frac{3}{2}; -\frac{e^{i(\vartheta+\frac{\pi}{2})}}{2\sin \vartheta}\right) \right\}. \quad (5.2.36)
\end{aligned}$$

5.2.7 $Y_{lm}(\vartheta, \varphi)$ and Other Special Functions

(a) The relation between $Y_{lm}(\vartheta, \varphi)$ and the Wigner D-function (Chap. 4)

$$\begin{aligned}
Y_{lm}(\vartheta, \varphi) &= \sqrt{\frac{2l+1}{4\pi}} D_{0-m}^l(\chi, \vartheta, \varphi) = (-1)^m \sqrt{\frac{2l+1}{4\pi}} D_{0m}^{l*}(\chi, \vartheta, \varphi) \\
&= (-1)^m \sqrt{\frac{2l+1}{4\pi}} D_{-m0}^l(\varphi, \vartheta, \chi) = \sqrt{\frac{2l+1}{4\pi}} D_{m0}^{l*}(\varphi, \vartheta, \chi), \quad (5.2.37)
\end{aligned}$$

where χ is an arbitrary angle.

(b) $Y_{lm}(\vartheta, \varphi)$ can be related to the Jacobi polynomials $P_n^{(\alpha, \beta)}(x)$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} \frac{e^{im\varphi}}{2^{|m|}l!} \sqrt{\frac{2l+1}{4\pi}} (l+m)!(l-m)! (\sin \vartheta)^{|m|} P_{l-|m|}^{(|m|, |m|)}(\cos \vartheta). \quad (5.2.38)$$

(c) $Y_{lm}(\vartheta, \varphi)$ can be expressed in terms of Gegenbauer polynomials $C_\nu^\lambda(x)$

$$Y_{lm}(\vartheta, \varphi) = \xi_{m0} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} \frac{(l-|m|)!}{(l+|m|)!} (2|m|-1)!! (\sin \vartheta)^{|m|} C_{l-|m|}^{\frac{1}{2}+|m|}(\cos \vartheta). \quad (5.2.39)$$

5.2.8 $Y_{lm}(\vartheta, \varphi)$ as an Irreducible Tensor Product

For any position vector r specified by polar coordinates r, ϑ, φ one has

$$Y_{lm}(\vartheta, \varphi) = \frac{1}{r^l} \sqrt{\frac{(2l+1)!!}{4\pi l!}} \{ \dots \{ \{ r \otimes r \}_2 \otimes r \}_3 \dots \otimes r \}_{lm}. \quad (5.2.40)$$

5.3 Integral representations of the spherical harmonics [4,22,27]

5.3.1 $Y_{lm}(\vartheta, \varphi)$ in the Form of Indefinite Integrals

(a)

$$Y_{lm}(\vartheta, \varphi) = e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} \frac{(l+m)!}{(l-m)!} \frac{1}{(\sin \vartheta)^m} \int_1^{\cos \vartheta} \int_1^{\cos \vartheta} \dots \int_1^{\cos \vartheta} P_l(\cos \vartheta) (d \cos \vartheta)^m, \quad (m \geq 0). \quad (5.3.1)$$

This formula represents the analytic continuation of the differential expression (5.2.6) to the case $m < 0$.

(b) The Mehler-Dirichlet formulas

$$Y_{lm}(\vartheta, \varphi) = (-1)^m \frac{\sqrt{2}}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} \frac{(l-m)!}{(l+m)!} (2m-1)!! \left(\frac{\sin \vartheta}{2} \right)^m \int_0^\vartheta \frac{\cos \left[(2l+1) \frac{\psi}{2} \right] d\psi}{(\cos \psi - \cos \vartheta)^{m+\frac{1}{2}}}, \quad (m \geq 0) \quad (5.3.2)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{\sqrt{2}}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} \frac{(l-m)!}{(l+m)!} (2m-1)!! \left(\frac{\sin \vartheta}{2} \right)^m \int_\vartheta^\pi \frac{\sin \left[(2l+1) \frac{\psi}{2} \right] d\psi}{(\cos \vartheta - \cos \psi)^{m+\frac{1}{2}}}, \quad (m \geq 0). \quad (5.3.3)$$

(c)

$$Y_{lm}(\vartheta, \varphi) = (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} \frac{(l+m)!}{(l-m)!} \frac{1}{(m-1)!(\sin \vartheta)^m} \times \int_{\cos \vartheta}^1 P_l(\cos \psi) [\cos \psi - \cos \vartheta]^{m-1} d \cos \psi, \quad (m > 0). \quad (5.3.4)$$

5.3.2 $Y_{lm}(\vartheta, \varphi)$ in the Form of Definite Integrals

(a)

$$Y_{lm}(\vartheta, \varphi) = \frac{(\pm i)^m}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi}} (l+m)!(l-m)! \frac{1}{l!} \int_0^\pi [\cos \vartheta \pm i \sin \vartheta \cos \psi]^l \cos(m\psi) d\psi \quad (5.3.5)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{(\pm i)^m}{\pi} e^{im\varphi} \sqrt{\frac{(2l+1)}{4\pi(l+m)!(l-m)!}} l! \int_0^\pi \frac{\cos(m\psi) d\psi}{(\cos \vartheta \mp i \sin \vartheta \cos \psi)^{l+1}} \quad (5.3.6)$$

For $\varphi = 0$, complex conjugation does not affect the right-hand side of Eq. (5.3.6), because $Y_{lm}(\vartheta, 0)$ is real.

(b)

$$Y_{lm}(\vartheta, \varphi) = \frac{(\pm i)^m}{2\pi} \sqrt{\frac{(2l+1)}{4\pi} (l+m)!(l-m)!} \frac{1}{l!} \int_0^{2\pi} [\cos \vartheta \pm i \sin \vartheta \cos(\psi - \varphi)]^l e^{im\psi} d\psi \quad (5.3.7)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{(\pm i)^m}{2\pi} \sqrt{\frac{2l+1}{4\pi (l+m)!(l-m)!}} l! \int_0^{2\pi} [\cos \vartheta \mp i \sin \vartheta \cos(\psi - \varphi)]^{-l-1} e^{im\psi} d\psi \quad (5.3.8)$$

Equations (5.3.7) and (5.3.8) represent modifications of Eqs. (5.3.5) and (5.3.6).

(c)

$$Y_{lm}(\vartheta, \varphi) = \frac{(-1)^m}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!} \frac{(\sin \vartheta)^m}{(2m-1)!!}} \int_0^\pi (\cos \vartheta \pm i \sin \vartheta \cos \psi)^{l-m} (\sin \psi)^{2m} d\psi, (m \geq 0) \quad (5.3.9)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{(-1)^m}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!} \frac{(\sin \vartheta)^m}{(2m-1)!!}} \int_0^\pi \frac{(\sin \chi)^{2m} d\chi}{(\cos \vartheta \mp i \sin \vartheta \cos \chi)^{l+m+1}}, (m \geq 0) \quad (5.3.10)$$

Equations (5.3.9) and (5.3.10) transform to each other at replacing the integration variables according to

$$\cos \psi = \frac{\cos \vartheta \cos \chi + i \sin \vartheta}{\cos \vartheta + i \sin \vartheta \cos \chi}, \quad \cos \chi = \frac{\cos \vartheta \cos \psi - i \sin \vartheta}{\cos \vartheta - i \sin \vartheta \cos \psi}. \quad (5.3.11)$$

This replacing is equivalent to the transformation of "mirror" symmetry, i.e., to the replacement l by $l = -l - 1$.

5.3.3 $Y_{lm}(\vartheta, \varphi)$ in the Form of Improper Integrals

(a)

$$Y_{lm}(\vartheta, \varphi) = \frac{i^{m+1}}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi (l+m)!(l-m)!}} l! \times \left\{ (-1)^m \int_0^\infty \frac{\cosh(mt) dt}{(\cos \vartheta + i \sin \vartheta \cosh t)^{l+1}} - \int_0^\infty \frac{\cosh(mt) dt}{(\cos \vartheta - i \sin \vartheta \cosh t)^{l+1}} \right\}, \quad (5.3.12)$$

$$Y_{lm}(\vartheta, \varphi) = \frac{i}{\pi} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!} \frac{(\sin \vartheta)^m}{(2m-1)!!}} \times \left\{ \int_0^\infty \frac{(\sinh t)^{2m} dt}{(\cos \vartheta + i \sin \vartheta \cosh t)^{l+m+1}} - \int_0^\infty \frac{(\sinh t)^{2m} dt}{(\cos \vartheta - i \sin \vartheta \cosh t)^{l+m+1}} \right\}, (m \geq 0). \quad (5.3.13)$$

(b)

$$Y_{lm}(\vartheta, \varphi) = (-1)^m e^{im\varphi} \sqrt{\frac{2l+1}{4\pi (l+m)!(l-m)!}} \int_0^\infty e^{-k \cos \vartheta} J_m(k \sin \vartheta) k^l dk. \quad (5.3.14)$$

where $J_m(x)$ is a Bessel function.

(c) $|Y_{lm}(\vartheta, \varphi)|^2$ may be represented as

$$|Y_{lm}(\vartheta, \varphi)|^2 = \frac{2l+1}{4\pi} \int_0^\infty \left[J_m \left(\frac{t \sin \vartheta}{2} \right) \right]^2 J_{2l+1}(t) dt. \quad (5.3.15)$$

5.4 Symmetry properties

The symmetry relations given below couple the harmonics $Y_{lm}(\vartheta, \varphi)$ with different values of ϑ, φ and l, m . These relations permit us to extend the domain of allowed ϑ, φ and generalise $Y_{lm}(\vartheta, \varphi)$ to the case of negative l .

(a) *Complex conjugation*

$$Y_{lm}^*(\vartheta, \varphi) = Y_{lm}(\vartheta, -\varphi) = (-1)^m Y_{l-m}(\vartheta, \varphi). \quad (5.4.1)$$

(b) *Sign reversal of m*

$$Y_{l-m}(\vartheta, \varphi) = (-1)^m Y_{lm}(\vartheta, -\varphi) = (-1)^m e^{-i2m\varphi} Y_{lm}(\vartheta, \varphi). \quad (5.4.2)$$

(c) *"Mirror" symmetry (replacement l by $\bar{l} = -l - 1$)*

$$Y_{\bar{l}m}(\vartheta, \varphi) = (-1)^m Y_{lm}(\vartheta, \varphi) \quad (5.4.3)$$

(d) *Replacement $\vartheta \rightarrow \pi - \vartheta$ and $\varphi \rightarrow \pi + \varphi$*

$$Y_{lm}(\pi - \vartheta, \varphi) = (-1)^{l+m} Y_{lm}(\vartheta, \varphi) \quad (5.4.4)$$

$$Y_{lm}(\vartheta, \pi + \varphi) = (-1)^m Y_{lm}(\vartheta, \varphi) \quad (5.4.5)$$

$$Y_{lm}(\pi - \vartheta, \pi + \varphi) = (-1)^l Y_{lm}(\vartheta, \varphi) \quad (5.4.6)$$

(e) *Change of argument signs*

$$Y_{lm}(-\vartheta, \varphi) = (-1)^m Y_{lm}(\vartheta, \varphi) \quad (5.4.7)$$

$$Y_{lm}(\vartheta, -\varphi) = (-1)^m Y_{l-m}(\vartheta, \varphi) \quad (5.4.8)$$

$$Y_{lm}(-\vartheta, -\varphi) = Y_{l-m}(\vartheta, \varphi) \quad (5.4.9)$$

(f) *The periodicity in ϑ and φ*

$$Y_{lm}(\vartheta \pm \pi n, \varphi) = \begin{cases} (-1)^l Y_{lm}(\vartheta, \varphi), & \text{if } n \text{ is odd} \\ Y_{lm}(\vartheta, \varphi), & \text{if } n \text{ is even} \end{cases} \quad (5.4.10)$$

$$Y_{lm}(\vartheta, \varphi \pm n\pi) = \begin{cases} (-1)^m Y_{lm}(\vartheta, \varphi), & \text{if } n \text{ is odd} \\ Y_{lm}(\vartheta, \varphi) & \text{if } n \text{ is even} \end{cases} \quad (5.4.11)$$

Making use of the above symmetry properties, one gets

$$\begin{aligned} Y_{lm}(\vartheta, \varphi) &= (-1)^m e^{2im\varphi} Y_{l-m}(\vartheta, \varphi) = e^{2im\varphi} Y_{lm}^*(\vartheta, \varphi) \\ &= (-1)^m Y_{lm}(-\vartheta, \varphi) = e^{2im\varphi} Y_{l-m}(-\vartheta, \varphi) = (-1)^m Y_{l-m}^*(\vartheta, \varphi) \\ &= (-1)^m Y_{l-m}(\vartheta, -\varphi) = e^{2im\varphi} Y_{lm}(\vartheta, -\varphi) = (-1)^m e^{2im\varphi} Y_{lm}^*(-\vartheta, \varphi) = Y_{l-m}^*(-\vartheta, \varphi) \\ &= Y_{l-m}(-\vartheta, -\varphi) = (-1)^m e^{2im\varphi} Y_{lm}(-\vartheta, -\varphi) = e^{2im\varphi} Y_{l-m}^*(-\vartheta, -\varphi) = (-1)^m Y_{lm}^*(-\vartheta, -\varphi). \end{aligned} \quad (5.4.12)$$

5.5 Behaviour of $Y_{lm}(\vartheta, \varphi)$ under transformations of coordinate systems

Any transformation of coordinate system in 3-dimensional space, which does not affect the orthogonality of coordinate axes, may be represented as a result of three operations: (1) rotation, (2) inversion and (3) parallel translation.

5.5.1 Rotation

The spherical harmonics $Y_{lm}(\vartheta, \varphi)$ are covariant components of some irreducible tensor of rank l . Hence, under arbitrary rotation $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ of the coordinate system described by the Euler angles α, β, γ (Sec. 1.4) the spherical harmonics transform in accordance with

$$\hat{D}(\alpha, \beta, \gamma) Y_{lm'}(\vartheta, \varphi) = Y_{lm'}(\vartheta', \varphi') = \sum_m Y_{lm}(\vartheta, \varphi) D_{mm'}^l(\alpha, \beta, \gamma) \quad (5.5.1)$$

Here $D_{mm'}^l(\alpha, \beta, \gamma)$ is a Wigner D -function (Chap. 4), ϑ, φ and ϑ', φ' are polar angles of the position vector in the original and final coordinate systems, S and S' , respectively. The angles ϑ', φ' may be expressed through ϑ, φ and the Euler angles according to Eqs. 1.(4.7.4).

5.5.2 Inversion

Under inversion $S\{x, y, z\} \rightarrow S'\{-x, -y, -z\}$ the spherical harmonics transform as

$$\hat{P}_r Y_{lm}(\vartheta, \varphi) = Y_{lm}(\pi - \vartheta, \varphi + \pi) = (-1)^l Y_{lm}(\vartheta, \varphi). \quad (5.5.2)$$

5.5.3 Parallel Translation

At parallel displacement $S\{x, y, z\} \rightarrow S'\{x', y', z'\}$ by a vector $\mathbf{a}(a, \Theta, \Phi)$ one has $\mathbf{r}' = \mathbf{r} - \mathbf{a}$. In this case the spherical harmonics transform as

$$\begin{aligned} \hat{T}(\mathbf{a}) Y_{l'm'}(\vartheta, \varphi) &= Y_{l'm'}(\vartheta', \varphi') = \sum_{l=0}^{l'} (-1)^{l'+l} \left[\frac{4\pi (2l'+1)!}{(2l+1)! (2l'-2l+1)!} \right]^{\frac{1}{2}} \\ &\quad \times \left(\frac{a}{r'} \right)^{l'} \left(\frac{r}{a} \right)^l \{ \mathbf{Y}_l(\vartheta, \varphi) \otimes \mathbf{Y}_{l'-l}(\Theta, \Phi) \}_{l'm'} \end{aligned} \quad (5.5.3)$$

where $\hat{T}(a) = e^{-a \nabla}$ is the displacement operator, $\{ \mathbf{Y}_{l_1} \otimes \mathbf{Y}_{l_2} \}_{tm}$ is the irreducible tensor product (Sec. 3.1). The polar coordinates of the vectors $r(r, \vartheta, \varphi)$ and $r'(r', \vartheta', \varphi')$ are related by

$$\begin{aligned} r'^2 &= r^2 + a^2 - 2ra \cos \omega_{12} \\ \cos \vartheta' &= \frac{r \cos \vartheta - a \cos \theta}{\sqrt{r^2 + a^2 - 2ra \cos \omega_{12}}} \\ \tan \varphi' &= \frac{r \sin \vartheta \sin \varphi - a \sin \theta \sin \Phi}{r \sin \vartheta \cos \varphi - a \sin \theta \cos \Phi} \end{aligned} \quad (5.5.4)$$

where

$$\cos \omega_{12} = \cos \vartheta \cos \theta + \sin \vartheta \sin \theta \cos(\varphi - \Phi). \quad (5.5.5)$$

5.5.4 Special Cases of Coordinate-System Transformations

(a) *Rotation about the coordinate axes through the angle π*

$$\text{about the } x \text{ axis} \quad \left. \begin{aligned} r' &= r, \\ \vartheta' &= \pi - \vartheta, \\ \varphi' &= 2\pi - \varphi, \end{aligned} \right\} Y_{lm}(\pi - \vartheta, 2\pi - \varphi) = (-1)^l Y_{l-m}(\vartheta, \varphi) \quad (5.5.6)$$

$$\text{about the } y \text{ axis} \quad \left. \begin{aligned} r' &= r, \\ \vartheta' &= \pi - \vartheta, \\ \varphi' &= \pi - \varphi, \end{aligned} \right\} Y_{lm}(\pi - \vartheta, \pi - \varphi) = (-1)^{l-m} Y_{l-m}(\vartheta, \varphi), \quad (5.5.7)$$

$$\text{about the } z \text{ axis} \quad \left. \begin{aligned} r' &= r, \\ \vartheta' &= \vartheta, \\ \varphi' &= \pi + \varphi \end{aligned} \right\} Y_{lm}(\vartheta, \pi + \varphi) = (-1)^m Y_{lm}(\vartheta, \varphi). \quad (5.5.8)$$

(b) *Rotation about the z axis through an arbitrary angle χ*

$$\left. \begin{aligned} r' &= r, \\ \vartheta' &= \vartheta, \\ \varphi' &= \varphi - \chi, \end{aligned} \right\} Y_{lm}(\vartheta, \varphi - \chi) = e^{-im\chi} Y_{lm}(\vartheta, \varphi). \quad (5.5.9)$$

(c) *Rotation about an arbitrary direction $\mathbf{n}(\Theta, \Phi)$ through a small angle ω ($\omega \ll \pi/2$)*

$$\begin{aligned} \hat{D} Y_{lm}(\vartheta, \varphi) &\approx Y_{lm}(\vartheta, \varphi) - i\omega \{ m \cos \Theta Y_{lm}(\vartheta, \varphi) \\ &+ \frac{\sin \Theta}{2} [e^{-i\Phi} \sqrt{l(l+1) - m(m+1)} Y_{l,m+1}(\vartheta, \varphi) + e^{i\Phi} \sqrt{l(l+1) - m(m-1)} Y_{l,m-1}(\vartheta, \varphi)] \}. \end{aligned} \quad (5.5.10)$$

(d) *Reflection of the coordinate system with respect to the equatorial plane, $\vartheta = \pi/2$*

$$\left. \begin{aligned} r' &= r \\ \vartheta' &= \pi - \vartheta, \\ \varphi' &= \varphi, \end{aligned} \right\} Y_{lm}(\pi - \vartheta, \varphi) = (-1)^{l+m} Y_{lm}(\vartheta, \varphi) \quad (5.5.11)$$

(e) Reflection of the coordinate system with respect to a meridian plane, $\varphi = \varphi_0$ and $\varphi = \pi + \varphi_0$

$$\left. \begin{aligned} r' &= r \\ \vartheta' &= \vartheta, \\ \varphi' &= 2\varphi_0 - \varphi, \end{aligned} \right\} Y_{lm}(\vartheta, 2\varphi_0 - \varphi) = e^{i2m\varphi_0} (-1)^m Y_{l-m}(\vartheta, \varphi) \quad (5.5.12)$$

5.6 Expansions in series of the spherical harmonics

5.6.1 General Relations

A collection of the spherical harmonics $Y_{lm}(\vartheta, \varphi)$ with all integer non-negative l and integer m ($|m| \leq l$) constitutes a complete orthonormal set of functions of two real variables ϑ and φ defined within $0 \leq \vartheta \leq \pi$, $0 \leq \varphi < 2\pi$. The completeness relation for the spherical harmonics is given by

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_{lm}^*(\vartheta, \varphi) Y_{lm}(\vartheta', \varphi') = \delta(\varphi - \varphi') \delta(\cos \vartheta - \cos \vartheta') \quad (5.6.1)$$

The orthogonality and normalization relation is as follows: (Sec. 5.1.4)

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{lm}^*(\vartheta, \varphi) Y_{l'm'}(\vartheta, \varphi) = \delta_{ll'} \delta_{mm'} \quad (5.6.2)$$

An arbitrary function $f(\vartheta, \varphi)$ which is defined in the interval $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$ and satisfies the condition

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta |f(\vartheta, \varphi)|^2 < \infty \quad (5.6.3)$$

can be expanded into a series of the spherical harmonics as

$$f(\vartheta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l a_{lm} Y_{lm}(\vartheta, \varphi) \quad (5.6.4)$$

The expansion coefficients a_{lm} are given by the relation

$$a_{lm} = \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{lm}^*(\vartheta, \varphi) f(\vartheta, \varphi) \quad (5.6.5)$$

This relation may be treated as an integral transformation of $f(\vartheta, \varphi)$ from the continuous variables ϑ, φ to the discrete variables l, m . In this case $Y_{lm}(\vartheta, \varphi) \equiv \langle \vartheta, \varphi | lm \rangle$ plays the role of the transformation matrix

$$\langle lm | f \rangle = \langle lm | \vartheta \varphi \rangle \langle \vartheta \varphi | f \rangle \quad (5.6.6)$$

where

$$\langle lm | f \rangle \equiv a_{lm}, \quad \langle lm | \vartheta, \varphi \rangle \equiv Y_{lm}^*(\vartheta, \varphi), \quad \langle \vartheta, \varphi | f \rangle \equiv f(\vartheta, \varphi)$$

As usual, in relations of this type, summation or integration is assumed over all variables which are repeated twice.

The transformation (5.6.5) is unitary, i.e.,

$$\langle f | lm \rangle \langle lm | f \rangle = \langle f | \vartheta, \varphi \rangle \langle \vartheta, \varphi | f \rangle. \quad (5.6.7)$$

The expansion coefficients a_{lm} satisfy the Parseval condition

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l |a_{lm}|^2 = \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta |f(\vartheta, \varphi)|^2 \quad (5.6.8)$$

The expansion (5.6.4) in terms of the spherical harmonics is widely used in different branches of physics. It is called the multipole expansion, and the a_{lm} are called multipole moments.

5.6.2 Expansion of Products of the Spherical Harmonics

- (a) A direct product of two spherical harmonics of the same arguments may be expanded in series as (the so-called Clebsch-Gordan series)

$$Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) = \sum_{LM} \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2L+1)}} C_{l_1 0 l_2 0}^{L 0} C_{l_1 m_1 l_2 m_2}^{L M} Y_{LM}(\vartheta, \varphi). \quad (5.6.9)$$

The inverse relation may be written as

$$\sqrt{\frac{4\pi(2L+1)}{(2l_1+1)(2l_2+1)}} \sum_{m_1 m_2} C_{l_1 m_1 l_2 m_2}^{L M} Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) = C_{l_1 0 l_2 0}^{L 0} Y_{LM}(\vartheta, \varphi). \quad (5.6.10)$$

Products of three and more spherical harmonics are decomposed according to

$$Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) Y_{l_3 m_3}(\vartheta, \varphi) = \sum_{LML'M'} \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{(4\pi)^2(2L+1)}} \times C_{l_1 0 l_2 0}^{L' 0} C_{L' 0 l_3 0}^{L 0} C_{l_1 m_1 l_2 m_2}^{L' M'} C_{L' M' l_3 m_3}^{L M} Y_{LM}(\vartheta, \varphi) \quad (5.6.11)$$

$$Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) \times \dots \times Y_{l_n m_n}(\vartheta, \varphi) = \sum_{L_n} B_{L_n} Y_{L_n M_n}(\vartheta, \varphi) \quad (5.6.12)$$

where

$$B_{L_n} = \sqrt{\frac{4\pi}{2L_n+1}} \sum_{\substack{L_1 L_2 \dots L_{n-1} \\ M_1 M_2, \dots, M_{n-1}}} \prod_{i=1}^n \left(\sqrt{\frac{2l_i+1}{4\pi}} C_{L_{i-1} 0 l_i 0}^{L_i 0} C_{L_{i-1} M_{i-1} l_i m_i}^{L_i M_i} \right) \quad (5.6.13)$$

with $L_0 \equiv M_0 \equiv 0$.

- (b) Irreducible tensor products (defined in Sec. 3.1) of the spherical harmonics may be expanded as

$$\{\mathbf{Y}_{l_1}(\vartheta, \varphi) \otimes \mathbf{Y}_{l_2}(\vartheta, \varphi)\}_{LM} = \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2L+1)}} C_{l_1 0 l_2 0}^{L 0} Y_{LM}(\vartheta, \varphi) \quad (5.6.14)$$

$$\{\{\mathbf{Y}_{l_1}(\vartheta, \varphi) \otimes \mathbf{Y}_{l_2}(\vartheta, \varphi)\}_{L'} \otimes \mathbf{Y}_{l_3}(\vartheta, \varphi)\}_{LM} = \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{(4\pi)^2(2L+1)}} C_{l_1 0 l_2 0}^{L' 0} C_{L' 0 l_3 0}^{L 0} Y_{LM}(\vartheta, \varphi) \quad (5.6.15)$$

$$\begin{aligned} & \left\{ \dots \left\{ \{\mathbf{Y}_{l_1}(\vartheta, \varphi) \otimes \mathbf{Y}_{l_2}(\vartheta, \varphi)\}_{L_2} \otimes \mathbf{Y}_{l_3}(\vartheta, \varphi) \right\}_{L_2} \dots \otimes \mathbf{Y}_{l_n}(\vartheta, \varphi) \right\}_{L_n M_n} \\ &= \sqrt{\frac{4\pi}{2L_n+1}} \prod_{i=1}^n \left(\sqrt{\frac{2l_i+1}{4\pi}} C_{L_{i-1} 0 l_i 0}^{L_i 0} \right) Y_{L_n M_n}(\vartheta, \varphi) \end{aligned} \quad (5.6.16)$$

In particular, when $l_1 = l_2 = \dots = l_n = 1$ and $L_k = k(k = 1, 2, 3, \dots, n)$ one has

$$\left\{ \dots \left\{ \{\mathbf{Y}_1(\vartheta, \varphi) \otimes \mathbf{Y}_1(\vartheta, \varphi)\}_2 \otimes \mathbf{Y}_1(\vartheta, \varphi) \right\}_3 \dots \otimes \mathbf{Y}_1(\vartheta, \varphi) \right\}_{nm} = \sqrt{\frac{3^n}{(4\pi)^{n-1}} \frac{n!}{(2n+1)!!}} Y_{nm}(\vartheta, \varphi) \quad (5.6.17)$$

Note that when expanding the products of the spherical harmonics, the functions $C_{lm}(\vartheta, \varphi)$ which contain an additional factor $\sqrt{4\pi/(2l+1)}$ (Sec. 5.1.4), are more convenient.

5.7 Recursion relations

Recursion relations for the spherical harmonics may be obtained, for example, from the Clebsch-Gordan series (see Eq. (5.6.9)).

$$-2m \cot \vartheta Y_{lm}(\vartheta, \varphi) = \sqrt{l(l+1) - m(m+1)} e^{-i\varphi} Y_{l+1,m+1}(\vartheta, \varphi) + \sqrt{l(l+1) - m(m-1)} e^{i\varphi} Y_{l-1,m-1}(\vartheta, \varphi) \quad (5.7.1)$$

$$\cos \vartheta Y_{lm}(\vartheta, \varphi) = \sqrt{\frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)}} Y_{l+1,m}(\vartheta, \varphi) + \sqrt{\frac{(l-m)(l+m)}{(2l-1)(2l+1)}} Y_{l-1,m}(\vartheta, \varphi) \quad (5.7.2)$$

$$\begin{aligned} \sin \vartheta Y_{lm}(\vartheta, \varphi) e^{-i\varphi} &= \sqrt{\frac{(l-m+1)(l-m+2)}{(2l+1)(2l+3)}} Y_{l+1,m-1}(\vartheta, \varphi) - \sqrt{\frac{(l+m-1)(l+m)}{(2l-1)(2l+1)}} Y_{l-1,m-1}(\vartheta, \varphi) \\ &= \cos \vartheta \sqrt{\frac{l-m+1}{l+m}} Y_{lm-1}(\vartheta, \varphi) - \sqrt{\frac{2l+1}{2l-1}} \frac{l+m-1}{l+m} Y_{l-1,m-1}(\vartheta, \varphi) \\ &= -\cos \vartheta \sqrt{\frac{l+m}{l-m+1}} Y_{lm-1}(\vartheta, \varphi) + \sqrt{\frac{2l+1}{2l+3}} \frac{l-m+2}{l-m+1} Y_{l+1,m-1}(\vartheta, \varphi) \end{aligned} \quad (5.7.3)$$

$$\begin{aligned} \sin \vartheta Y_{lm}(\vartheta, \varphi) e^{i\varphi} &= -\sqrt{\frac{(l+m+1)(l+m+2)}{(2l+1)(2l+3)}} Y_{l+1,m+1}(\vartheta, \varphi) + \sqrt{\frac{(l-m-1)(l-m)}{(2l-1)(2l+1)}} Y_{l-1,m+1}(\vartheta, \varphi) \\ &= -\cos \vartheta \sqrt{\frac{l+m+1}{l-m}} Y_{lm+1}(\vartheta, \varphi) + \sqrt{\frac{2l+1}{2l-1}} \frac{l-m-1}{l-m} Y_{l-1,m+1}(\vartheta, \varphi) \\ &= \cos \vartheta \sqrt{\frac{l-m}{l+m+1}} Y_{lm+1}(\vartheta, \varphi) - \sqrt{\frac{2l+1}{2l+3}} \frac{l+m+2}{l+m+1} Y_{l+1,m+1}(\vartheta, \varphi) \end{aligned} \quad (5.7.4)$$

$$\begin{aligned} (2l-1)(2l+3) \cos^2 \vartheta Y_{lm}(\vartheta, \varphi) &= (2l-1) \sqrt{\frac{(l+1)^2 - m^2}{2l+1} \frac{(l+2)^2 - m^2}{2l+5}} Y_{l+2,m}(\vartheta, \varphi) \\ &\quad + [2l(l+1) - 2m^2 - 1] Y_{lm}(\vartheta, \varphi) + (2l+3) \sqrt{\frac{l^2 - m^2}{2l+1} \frac{(l-1)^2 - m^2}{2l-3}} Y_{l-2,m}(\vartheta, \varphi) \end{aligned} \quad (5.7.5)$$

$$\begin{aligned} (2l-1)(2l+3) \sin \vartheta \cos \vartheta e^{i\varphi} Y_{lm}(\vartheta, \varphi) &= -(2l-1) \sqrt{\frac{(l+1)^2 - m^2}{2l+1} \frac{(l+m+2)(l+m+3)}{2l+5}} Y_{l+2,m+1}(\vartheta, \varphi) \\ &\quad - (2m+1) \sqrt{l(l+1) - m(m+1)} Y_{l+1,m+1}(\vartheta, \varphi) + (2l+3) \sqrt{\frac{l^2 - m^2}{2l+1} \frac{(l-m-1)(l-m-2)}{2l-3}} Y_{l-2,m+1}(\vartheta, \varphi) \end{aligned} \quad (5.7.6)$$

$$\begin{aligned} (2l-1)(2l+3) \sin \vartheta \cos \vartheta e^{-i\varphi} Y_{lm}(\vartheta, \varphi) &= (2l-1) \sqrt{\frac{(l+1)^2 - m^2}{2l+1} \frac{(l-m+2)(l-m+3)}{2l+5}} Y_{l+2,m-1}(\vartheta, \varphi) \\ &\quad - (2m-1) \sqrt{l(l+1) - m(m-1)} Y_{l+1,m-1}(\vartheta, \varphi) - (2l+3) \sqrt{\frac{l^2 - m^2}{2l+1} \frac{(l+m-1)(l+m-2)}{(2l-3)}} Y_{l-2,m-1}(\vartheta, \varphi) \end{aligned} \quad (5.7.7)$$

$$\begin{aligned} (2l-1)(2l+3) \sin^2 \vartheta e^{i2\varphi} Y_{lm}(\vartheta, \varphi) &= \frac{(2l-1)}{\sqrt{(2l+1)(2l+5)}} \sqrt{\frac{(l+m+4)!}{(l+m)!}} Y_{l+2,m+2}(\vartheta, \varphi) \\ &\quad + \frac{(2l+3)}{\sqrt{(2l+1)(2l-3)}} \sqrt{\frac{(l-m)!}{(l-m-4)!}} Y_{l-2,m+2}(\vartheta, \varphi) - 2 \sqrt{\frac{(l+m+2)!(l-m)!}{(l-m-2)!(l+m)!}} Y_{lm+2}(\vartheta, \varphi) \end{aligned} \quad (5.7.8)$$

$$\begin{aligned} (2l-1)(2l+3) \sin^2 \vartheta e^{-i2\varphi} Y_{lm}(\vartheta, \varphi) &= \frac{(2l-1)}{\sqrt{(2l+1)(2l+5)}} \sqrt{\frac{(l-m+4)!}{(l-m)!}} Y_{l+2,m-2}(\vartheta, \varphi) \\ &\quad + \frac{(2l+3)}{\sqrt{(2l+1)(2l-3)}} \sqrt{\frac{(l+m)!}{(l+m-4)!}} Y_{l-2,m-2}(\vartheta, \varphi) - 2 \sqrt{\frac{(l-m+2)!(l+m)!}{(l+m-2)!(l-m)!}} Y_{lm-2}(\vartheta, \varphi) \end{aligned} \quad (5.7.9)$$

5.8 Differential relations

5.8.1 Action of the Operator of Orbital Angular Momentum on $Y_{lm}(\vartheta, \varphi)$

Spherical components of the operator $\bar{\mathbf{L}}$ (Eq. (2.2.18)) act on $\mathbf{Y}_{lm}(\vartheta, \varphi)$ according to

$$\hat{L}_{\pm 1} Y_{lm}(\vartheta, \varphi) = \mp \sqrt{\frac{l(l+1) - m(m \pm 1)}{2}} Y_{lm \pm 1}(\vartheta, \varphi) \quad (5.8.1)$$

$$\hat{L}_0 Y_{lm}(\vartheta, \varphi) = m Y_{lm}(\vartheta, \varphi)$$

or in a more compact form

$$\hat{L}_\mu Y_{lm}(\vartheta, \varphi) = \sqrt{l(l+1)} C_{l m 1 \mu}^{l m + \mu} Y_{lm + \mu}(\vartheta, \varphi) \quad (5.8.2)$$

The action of the operator $\hat{\mathbf{L}}^2$ on $Y_{lm}(\vartheta, \varphi)$ yields

$$\hat{\mathbf{L}}^2 Y_{lm}(\vartheta, \varphi) = l(l+1) Y_{lm}(\vartheta, \varphi) \quad (5.8.3)$$

5.8.2 First and Second Order Derivatives of $Y_{lm}(\vartheta, \varphi)$

$$\frac{\partial}{\partial \varphi} Y_{lm}(\vartheta, \varphi) = im Y_{lm}(\vartheta, \varphi) \quad (5.8.4)$$

$$\begin{aligned} \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) &= m \cot \vartheta Y_{lm}(\vartheta, \varphi) + \sqrt{l(l+1) - m(m+1)} Y_{lm+1}(\vartheta, \varphi) e^{-i\varphi} \\ &= -m \cot \vartheta Y_{lm}(\vartheta, \varphi) - \sqrt{l(l+1) - m(m-1)} Y_{lm-1}(\vartheta, \varphi) e^{i\varphi} \\ &= \frac{1}{2} \sqrt{l(l+1) - m(m+1)} Y_{lm+1}(\vartheta, \varphi) e^{-i\varphi} - \frac{1}{2} \sqrt{l(l+1) - m(m-1)} Y_{lm-1}(\vartheta, \varphi) e^{i\varphi} \end{aligned} \quad (5.8.5)$$

$$\begin{aligned} \sin \vartheta \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) &= -\sin^2 \vartheta \frac{\partial}{\partial \cos \vartheta} Y_{lm}(\vartheta, \varphi) \\ &= l \cos \vartheta Y_{lm}(\vartheta, \varphi) - \sqrt{\frac{2l+1}{2l-1}} (l^2 - m^2) Y_{l-1m}(\vartheta, \varphi) \\ &= -(l+1) \cos \vartheta Y_{lm}(\vartheta, \varphi) + \sqrt{\frac{2l+1}{2l+3}} [(l+1)^2 - m^2] Y_{l+1m}(\vartheta, \varphi) \\ &= l \sqrt{\frac{(l+1)^2 - m^2}{(2l+1)(2l+3)}} Y_{l+1m}(\vartheta, \varphi) - (l+1) \sqrt{\frac{l^2 - m^2}{(2l+1)(2l-1)}} Y_{l-1m}(\vartheta, \varphi) \end{aligned} \quad (5.8.6)$$

$$\frac{\partial^2}{\partial \vartheta^2} Y_{lm}(\vartheta, \varphi) = - \left[l(l+1) - \frac{m^2}{\sin^2 \vartheta} \right] Y_{lm}(\vartheta, \varphi) - \cot \vartheta \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi). \quad (5.8.7)$$

$$\sin^2 \vartheta \frac{\partial^2}{\partial (\cos \vartheta)^2} Y_{lm}(\vartheta, \varphi) = 2 \cos \vartheta \frac{\partial}{\partial \cos \vartheta} Y_{lm}(\vartheta, \varphi) - \left[l(l+1) - \frac{m^2}{\sin^2 \vartheta} \right] Y_{lm}(\vartheta, \varphi). \quad (5.8.8)$$

5.8.3 Vector Differentiation Operations

- (a) The gradient of a function $f(r)Y_{lm}(\vartheta, \varphi)$, where $f(r)$ is an arbitrary function of $r \equiv |\mathbf{r}|$ can be expanded in terms of vector spherical harmonics (Sec. 7.3) as follows:

$$\nabla [f(r)Y_{lm}(\vartheta, \varphi)] = -\sqrt{\frac{l+1}{2l+1}} \left(\frac{df}{dr} - \frac{l}{r} f \right) \mathbf{Y}_{lm}^{l+1}(\vartheta, \varphi) + \sqrt{\frac{l}{2l+1}} \left(\frac{df}{dr} + \frac{l+1}{r} f \right) \mathbf{Y}_{lm}^{l-1}(\vartheta, \varphi) \quad (5.8.9)$$

or in component form

$$\begin{aligned} \nabla_0 [f(r)Y_{lm}(\vartheta, \varphi)] &= \sqrt{\frac{(l+1)^2 - m^2}{(2l+1)(2l+3)}} \left(\frac{df}{dr} - \frac{l}{r} f \right) Y_{l+1m}(\vartheta, \varphi) \\ &\quad + \sqrt{\frac{l^2 - m^2}{(2l-1)(2l+1)}} \left(\frac{df}{dr} + \frac{l+1}{r} f \right) Y_{l-1m}(\vartheta, \varphi) \end{aligned} \quad (5.8.10)$$

$$\begin{aligned} \nabla_{\pm 1} [f(r)Y_{lm}(\vartheta, \varphi)] &= \sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{2(2l+1)(2l+3)}} \left(\frac{df}{dr} - \frac{l}{r} f \right) Y_{l+1m \pm 1}(\vartheta, \varphi) \\ &\quad - \sqrt{\frac{(l \mp m - 1)(l \mp m)}{2(2l-1)(2l+1)}} \left(\frac{df}{dr} + \frac{l+1}{r} f \right) Y_{l-1m \pm 1}(\vartheta, \varphi) \end{aligned} \quad (5.8.11)$$

Here $\nabla_{0,\pm 1}$ are spherical components of the operator ∇ (Sec. 1.3).

Let us consider some special cases of Eq. (5.8.9). If $f(r) = j(kr)$ is a spherical Bessel function, one has,

$$\frac{1}{k} \nabla [j_l(kr) Y_{lm}(\vartheta, \varphi)] = \sqrt{\frac{l+1}{2l+1}} j_{l+1}(kr) \mathbf{Y}_{lm}^{l+1}(\vartheta, \varphi) + \sqrt{\frac{l}{2l+1}} j_{l-1}(kr) \mathbf{Y}_{lm}^{l-1}(\vartheta, \varphi) \quad (5.8.12)$$

Putting $f(r) = r^l$, one gets

$$\nabla [r^l Y_{lm}(\vartheta, \varphi)] = \sqrt{l(2l+1)} r^{l-1} \mathbf{Y}_{lm}^{l-1}(\vartheta, \varphi) \quad (5.8.13)$$

If $f(r) = r^{-l-1}$, Eq. (5.8.9) is reduced to

$$\nabla [r^{-l-1} Y_{lm}(\vartheta, \varphi)] = \sqrt{(l+1)(2l+1)} r^{-l-2} \mathbf{Y}_{lm}^{l+1}(\vartheta, \varphi) \quad (5.8.14)$$

(b) The divergence of a vector function $f(r) \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)$ vanishes for any function $f(r)$

$$\nabla [f(r) \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)] = 0. \quad (5.8.15)$$

(c) The curl of a vector function $f(r) \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)$, where $f(r)$ is an arbitrary function of r , is expressible in terms of a gradient of the corresponding scalar function

$$\nabla \times [f(r) \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)] = i \nabla \left[\left(r \frac{df}{dr} + f \right) Y_{lm}(\vartheta, \varphi) \right] - i r \left[\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{df}{dr} \right) - \frac{l(l+1)}{r^2} f \right] Y_{lm}(\vartheta, \varphi). \quad (5.8.16)$$

In particular, if $f(r) = j_l(kr)$, one gets

$$\text{curl} [j_l(kr) \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)] = i \nabla [(kr j_{l-1}(kr) - l j_l(kr)) Y_{lm}(\vartheta, \varphi)] + i r k^2 j_l(kr) Y_{lm}(\vartheta, \varphi). \quad (5.8.17)$$

If $f(r) = r$,

$$\text{curl} [r \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)] = i(l+1) \sqrt{l(2l+1)} r^{l-1} \mathbf{Y}_{lm}^{l-1}(\vartheta, \varphi). \quad (5.8.18)$$

and if $f(r) = r^{-l-1}$,

$$\text{curl} [r^{-l-1} \hat{\mathbf{L}} Y_{lm}(\vartheta, \varphi)] = -i l \sqrt{(l+1)(2l+1)} r^{-l-2} \mathbf{Y}_{lm}^{l+1}(\vartheta, \varphi). \quad (5.8.19)$$

Some additional formulas for the vector differentiation of functions which contain $Y_{lm}(\theta, \varphi)$ will be given in Sec. 7.3.6.

5.9 Some integrals involving spherical harmonics

5.9.1 Integrals over Total Solid Angle

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{lm}(\vartheta, \varphi) = \sqrt{4\pi} \delta_{l0} \delta_{m0} \quad (5.9.1)$$

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}^*(\vartheta, \varphi) = \delta_{l_1 l_2} \delta_{m_1 m_2} \quad (5.9.2)$$

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) = (-1)^{m_2} \delta_{l_1 l_2} \delta_{-m_1 m_2} \quad (5.9.3)$$

$$\int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) Y_{l_3 m_3}^*(\vartheta, \varphi) = \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2l_3+1)}} C_{l_1 0 l_2 0}^{l_3 0} C_{l_1 m_1 l_2 m_2}^{l_3 m_3} \quad (5.9.4)$$

$$\begin{aligned} & \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) Y_{l_3 m_3}(\vartheta, \varphi) \\ &= \sqrt{\frac{(2l_1+1)(2l_2+1)(2l_3+1)}{4\pi}} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \end{aligned} \quad (5.9.5)$$

Integrals involving products of three and more spherical harmonics can be evaluated by reducing them to integrals which involve products of a smaller number of harmonics. For this purpose it is convenient to use the Clebsch-Gordan expansion (Eq. (5.6.9)).

5.9.2 Fourier Transformations for Some Functions Which Contain $Y_{lm}(\vartheta, \varphi)$

$$\int_0^\infty dr r^2 \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta e^{i\mathbf{k}\mathbf{r}} j_l(qr) Y_{lm}(\vartheta, \varphi) = 2\pi^2 i^l \frac{\delta(q-k)}{q^2} Y_{lm}(\vartheta_k, \varphi_k) \quad (5.9.6)$$

$$\int_0^\infty dr r^2 \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta e^{i\mathbf{k}\mathbf{r}} \hat{\mathbf{L}} [j_l(qr) Y_{lm}(\vartheta, \varphi)] = 2\pi^2 i^l \frac{\delta(q-k)}{q^2} \hat{\mathbf{L}}_{\mathbf{k}} Y_{lm}(\vartheta_k, \varphi_k) \quad (5.9.7)$$

$$\int_0^\infty dr r^2 \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta e^{i\mathbf{k}\mathbf{r}} [\nabla \times \hat{\mathbf{L}}] (j_l(qr) Y_{lm}(\vartheta, \varphi)) = 2\pi^2 i^{l-1} \frac{\delta(q-k)}{q^2} [\mathbf{k} \times \hat{\mathbf{L}}_{\mathbf{k}}] Y_{lm}(\vartheta_k, \varphi_k) \quad (5.9.8)$$

Here $j_l(x)$ is a spherical Bessel function, ϑ_k and φ_k are polar angles of the vector $\mathbf{k}$; $\hat{\mathbf{L}}$ and $\hat{\mathbf{L}}_{\mathbf{k}}$ are the operators of orbital angular momentum in the coordinate and momentum representations, respectively,

$$\hat{\mathbf{L}} = -i[\mathbf{r} \times \nabla], \hat{\mathbf{L}}_{\mathbf{k}} = -i[\mathbf{k} \times \nabla_{\mathbf{k}}]. \quad (5.9.9)$$

5.9.3 Integrals with Respect to ϑ

$$\int_0^\pi Y_{lm}^*(\vartheta, \varphi) Y_{l'm}(\vartheta, \varphi) \sin \vartheta d\vartheta = \frac{\delta_{ll'}}{2\pi}. \quad (5.9.10)$$

$$\int_0^\pi Y_{lm}^*(\vartheta, \varphi) Y_{l'm'}(\vartheta, \varphi) \frac{d\vartheta}{\sin \vartheta} = \frac{2l+1}{4\pi m} \delta_{mm'} (m, m' > 0). \quad (5.9.11)$$

$$\begin{aligned} & \int_0^{\pi/2} (\sin \vartheta)^{m+1} (\cos \vartheta)^n Y_{lm}(\vartheta, \varphi) d\vartheta \\ &= \sqrt{\frac{2l+1}{4\pi} \cdot \frac{(l+m)!}{(l-m)!}} (-1)^m \frac{n! e^{im\varphi}}{(n+l+m+1)!! (n-l+m)!!}, (m, n \geq 0) \\ & \int_a^b \left[(l_1 - l_2) (l_1 + l_2 + 1) - \frac{m_1^2 - m_2^2}{\sin^2 \vartheta} \right] Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) d(\cos \vartheta) \\ &= \sin^2 \vartheta \left[Y_{l_1 m_1}(\vartheta, \varphi) \frac{d}{d \cos \vartheta} Y_{l_2 m_2}(\vartheta, \varphi) - Y_{l_2 m_2}(\vartheta, \varphi) \frac{d}{d \cos \vartheta} Y_{l_1 m_1}(\vartheta, \varphi) \right] \Big|_a^b \\ &= \left[\cos \vartheta (l_1 - l_2) Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) + \sqrt{\frac{(2l_2+1)(l_2^2 - m_2^2)}{2l_2 - 1}} Y_{l_1 m_1}(\vartheta, \varphi) Y_{l_2-1 m_2}(\vartheta, \varphi) \right. \\ & \quad \left. - \sqrt{\frac{(2l_1+1)(l_1^2 - m_1^2)}{2l_1 - 1}} Y_{l_1-1 m_1}(\vartheta, \varphi) Y_{l_2 m_2}(\vartheta, \varphi) \right] \Big|_a^b. \end{aligned} \quad (5.9.12)$$

5.10 Sums involving spherical harmonics

5.10.1 Sums over m (with fixed l)

$$\sum_{m=-l}^l |Y_{lm}(\vartheta, \varphi)|^2 = \frac{2l+1}{4\pi}, \quad (5.10.1)$$

$$\sum_{m=-l}^l m |Y_{lm}(\vartheta, \varphi)|^2 = 0, \quad (5.10.2)$$

$$\sum_{m=-l}^l m^2 |Y_{lm}(\vartheta, \varphi)|^2 = \frac{l(l+1)(2l+1)}{8\pi} \sin^2 \vartheta, \quad (5.10.3)$$

$$\sum_{m=-l}^l \sqrt{\frac{(l^2 - m^2)[(l+1)^2 - m^2]}{(2l-1)(2l+3)}} Y_{l-1 m}(\vartheta, \varphi) Y_{l+1 m}^*(\vartheta, \varphi) = \frac{l(l+1)}{8\pi} (3 \cos^2 \vartheta - 1), \quad (5.10.4)$$

$$\sum_{m=-l}^l \frac{(\mp i)^m}{\sqrt{(l-m)!(l+m)!}} Y_{lm}(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi}} \frac{(\cos \vartheta \pm i \sin \vartheta \cos \varphi)^l}{l!}. \quad (5.10.5)$$

5.10.2 Sums over l (with fixed $m \geq 0$)

In the equations given below [25] we will use the notation $R = \sqrt{1 - 2t \cos \vartheta + t^2}$.

$$\begin{aligned} & \sum_{l=m}^{n+m} (-1)^l \frac{t^{n-l+m}}{(n-l+m)! \sqrt{(l-m)!(l+m)!}} \sqrt{\frac{4\pi}{2l+1}} Y_{lm}(\vartheta, \varphi) \\ &= (-1)^n \frac{(2m-1)!!}{(2m+n)!} (\sin \vartheta e^{i\varphi})^m (1 - 2t \cos \vartheta + t^2)^{\frac{n}{2}} C_n^{m+\frac{1}{2}} \left(\frac{\cos \vartheta - t}{\sqrt{1 - 2t \cos \vartheta + t^2}} \right), (|t| < 1), \end{aligned} \quad (5.10.6)$$

$$\begin{aligned} & \sum_{l=m}^{\infty} \sqrt{\frac{4\pi}{(2l+1)(l-m)!(l+m)!}} \frac{1}{l!} t^{l-m} Y_{lm}(\vartheta, \varphi) \\ &= \frac{(-\sin \vartheta e^{i\varphi})^m}{2^m (m!)^2} {}_0F_1 \left(; m+1; t \cos^2 \frac{\vartheta}{2} \right) {}_0F_1 \left(; m+1; -t \sin^2 \frac{\vartheta}{2} \right), (|t| < 1), \end{aligned} \quad (5.10.7)$$

$$\begin{aligned} & \sum_{l=m}^{\infty} \frac{(n+l-m)! t^{l-m}}{\sqrt{(l+m)!(l-m)!}} \sqrt{\frac{4\pi}{2l+1}} Y_{lm}(\vartheta, \varphi) \\ &= \frac{n!}{2^m m!} \frac{(-\sin \vartheta e^{i\varphi})^m}{(1-t \cos \vartheta)^{n+1}} F \left\{ \frac{n+1}{2}, \frac{n}{2} + 1; m+1; - \left(\frac{t \sin \vartheta}{1-t \cos \vartheta} \right)^2 \right\}, (|t| < 1), \end{aligned} \quad (5.10.8)$$

$$\begin{aligned} & \sum_{l=m}^{\infty} \sqrt{\frac{4\pi}{(2l+1)(l-m)!(l+m)!}} \frac{(l+m-n)!(l-m+n-1)!}{l!} t^{l-m} Y_{lm}(\vartheta, \varphi) = \frac{(2m-n)!(n-1)!}{2^m (m!)^2} \\ & \times (-\sin \vartheta e^{i\varphi})^m F \left(n, 2m+1-n; m+1; \frac{1-t-R}{2} \right) F \left(n, 2m+1-n; m+1; \frac{1+t-R}{2} \right), (|t| < 1), \end{aligned} \quad (5.10.9)$$

$$\begin{aligned} & \sum_{l=m}^{\infty} \sqrt{\frac{4\pi(l-m)!}{(2l+1)(l+m)!}} L_{l-m}^{2m}(y) t^{l-m} Y_{lm}(\vartheta, \varphi) \\ &= \frac{1}{2^m m!} \frac{(-\sin \vartheta e^{i\varphi})^m}{R^{2m+1}} \exp \left[-\frac{yt(\cos \vartheta - t)}{R^2} \right] {}_0F_1 \left(; m+1; -\frac{y^2 t^2 \sin^2 \vartheta}{4R^4} \right), (|t| < 1), \end{aligned} \quad (5.10.10)$$

$$\sum_{l=m}^{\infty} i^{l-m} \sqrt{\frac{4\pi(2l+1)(l+m)!}{(l-m)!}} j_l(t) Y_{lm}(\vartheta, \varphi) = (-t \sin \vartheta e^{i\varphi})^m e^{it \cos \vartheta}, \quad (5.10.11)$$

$$\sum_{l=m, m+2, \dots}^{\infty} i^{l-m} \sqrt{\frac{4\pi(2l+1)(l+m)!}{(l-m)!}} j_l(t) Y_{lm}(\vartheta, \varphi) = (-t \sin \vartheta e^{i\varphi})^m \cos(t \cos \vartheta), \quad (5.10.12)$$

$$\sum_{l=m+1, m+3, \dots}^{\infty} i^{l-m-1} \sqrt{\frac{4\pi(2l+1)(l+m)!}{(l-m)!}} j_l(t) Y_{lm}(\vartheta, \varphi) = (-t \sin \vartheta e^{i\varphi})^m \sin(t \cos \vartheta), \quad (5.10.13)$$

$$\sum_{l=m}^{\infty} \sqrt{\frac{4\pi(2l+1)(l+m)!}{(l-m)!}} j_l(t) z_l(t) Y_{lm}(\vartheta, \varphi) = \left(-t \cos \frac{\vartheta}{2} e^{i\varphi} \right)^m z_m \left(2t \sin \frac{\vartheta}{2} \right), \quad (5.10.14)$$

$$\sum_{l=m}^{\infty} \sqrt{\frac{4\pi(2l+1)(l+m)!}{(l-m)!}} j_l(x) z_l(y) Y_{lm}(\vartheta, \varphi) = \left(-\frac{xy \sin \vartheta e^{i\varphi}}{\sqrt{x^2 + y^2 - 2xy \cos \vartheta}} \right)^m z_m \left(\sqrt{x^2 + y^2 - 2xy \cos \vartheta} \right), \quad (5.10.15)$$

$$\sum_{l=m}^{\infty} i^{l-m} j_l(t) Y_{lm}(\vartheta_1, \varphi_1) Y_{lm}^*(\vartheta_2, \varphi_2) = \frac{1}{4\pi} J_m(t \sin \vartheta_1 \sin \vartheta_2) e^{it \cos \vartheta_1 \cos \vartheta_2} e^{im(\varphi_1 - \varphi_2)} \quad (5.10.16)$$

5.11 Generating functions for $Y_{lm}(\vartheta, \varphi)$

For fixed m one has (see Refs. [23, 24])

$$\frac{1}{R^{2m+1}} = \begin{cases} \frac{(-1)^m}{(2m-1)!! (\sin \vartheta)^m} \sum_{l=m}^{\infty} t^{l-m} \sqrt{\frac{4\pi}{2l+1}} \frac{(l+m)!}{(l-m)!} Y_{lm}(\vartheta, 0), & |t| < 1 \\ \frac{(-1)^m}{(2m-1)!! (\sin \vartheta)^m} \sum_{l=m}^{\infty} \frac{1}{t^{l+m+1}} \sqrt{\frac{4\pi}{2l+1}} \frac{(l+m)!}{(l-m)!} Y_{lm}(\vartheta, 0), & |t| > 1 \end{cases} \quad (5.11.1)$$

where

$$R = \sqrt{1 - 2t \cos \vartheta + t^2}$$

In particular, for $m = 0$ one gets

$$\frac{1}{R} = \begin{cases} \sum_{l=0}^{\infty} t^l P_l(\cos \vartheta), & |t| < 1, \\ \sum_{l=0}^{\infty} \frac{1}{t^{l+1}} P_l(\cos \vartheta), & |t| > 1. \end{cases} \quad (5.11.2)$$

$$\frac{[(1+R)^2 - t^2]^{-m}}{R} = \frac{(-1)^m}{2^m (\sin \vartheta)^m} \sum_{l=m}^{\infty} t^{l-m} l! \sqrt{\frac{4\pi}{(2l+1)(l+m)!(l-m)!}} Y_{lm}(\vartheta, 0), |t| < 1. \quad (5.11.3)$$

In addition, we present the following expansions in terms of $Y_{lm}(\vartheta, \varphi)$ [23]

$$\frac{(\sin \vartheta)^m}{(\cos \psi - \cos \vartheta)^{m+\frac{1}{2}}} \Theta(\cos \psi - \cos \vartheta) = \frac{(-1)^m 2^m}{(2m-1)!!} \sum_{l=m}^{\infty} \sqrt{\frac{8\pi}{2l+1} \frac{(l+m)!}{(l-m)!}} \cos \left[(2l+1) \frac{\psi}{2} \right] Y_{lm}(\vartheta, 0) \quad (5.11.4)$$

$$\frac{(\sin \vartheta)^m}{(\cos \vartheta - \cos \psi)^{m+\frac{1}{2}}} \Theta(\cos \vartheta - \cos \psi) = \frac{(-1)^m 2^m}{(2m-1)!!} \sum_{l=m}^{\infty} \sqrt{\frac{8\pi}{2l+1} \frac{(l+m)!}{(l-m)!}} \sin \left[(2l+1) \frac{\psi}{2} \right] Y_{lm}(\vartheta, 0) \quad (5.11.5)$$

where

$$\Theta(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ 0 & \text{if } x < 0. \end{cases} \quad (5.11.6)$$

In particular, when $m = 0$ one has

$$\frac{\Theta(\cos \psi - \cos \vartheta)}{\sqrt{\cos \psi - \cos \vartheta}} \pm i \frac{\Theta(\cos \vartheta - \cos \psi)}{\sqrt{\cos \vartheta - \cos \psi}} = \sqrt{2} \sum_{l=0}^{\infty} e^{\pm i(2l+1)\frac{\psi}{2}} P_l(\cos \vartheta). \quad (5.11.7)$$

5.12 Asymptotic expressions for $Y_{lm}(\vartheta, \varphi)$

5.12.1 $Y_{lm}(\vartheta, \varphi)$ for Large l

If $l \gg 1$ and $l \gg m \geq 0$, a spherical harmonic $Y_{lm}(\vartheta, \varphi)$ for $\varepsilon \leq \vartheta \leq \pi - \varepsilon$ ($0 < \varepsilon \ll 1/l$) and $0 \leq \varphi < 2\pi$ can be approximated by (see Ref. [23])

$$Y_{lm}(\vartheta, \varphi) \approx \frac{e^{im\varphi}}{\pi} \frac{\cos \left[(2l+1)\frac{\vartheta}{2} + (2m-1)\frac{\pi}{4} \right]}{\sqrt{\sin \vartheta}} + O\left(\frac{1}{l}\right). \quad (5.12.1)$$

A more exact formula [24] is

$$Y_{lm}(\vartheta, \varphi) \approx \frac{e^{im\varphi}}{\pi \sqrt{\sin \vartheta}} \left\{ \left(1 + \frac{4m^2-3}{8l} \right) \cos \left[(2l+1)\frac{\vartheta}{2} + (2m-1)\frac{\pi}{4} \right] - \frac{4m^2-1}{8l \sin \vartheta} \cos \left[(2l+3)\frac{\vartheta}{2} + (2m-3)\frac{\pi}{4} \right] \right\} + O\left(\frac{1}{l^2}\right) \quad (5.12.2)$$

The terms $O(1/l)$ and $O(1/l^2)$ depend on m and ϑ .

If $l \gg 1$ and $l \gg m \geq 0$, one also obtains

$$|Y_{l\pm m}(\vartheta, \varphi)| < \frac{2}{\pi} (\sin \vartheta)^{-(m+\frac{1}{2})}. \quad (5.12.3)$$

5.12.2 Behaviour of $Y_{lm}(\vartheta, \varphi)$ in Neighbourhoods of $\vartheta = 0, \pi$ and $\pi/2$

For $0 \leq \vartheta \leq \varepsilon$ ($\varepsilon \ll 1$) one has

$$Y_{l\pm m}(\vartheta, \varphi) \approx (\mp 1)^m \frac{e^{\pm im\varphi}}{m!} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} \left(\frac{\vartheta}{2} \right)^m \left[1 - \frac{3l(l+1) - m(m+1)}{3(m+1)} \left(\frac{\vartheta}{2} \right)^2 \right]. \quad (5.12.4)$$

For $\pi - \varepsilon \leq \vartheta \leq \pi$, one obtains

$$Y_{l\pm m}(\vartheta, \varphi) \approx (-1)^l (\pm 1)^m \frac{e^{\pm im\varphi}}{m!} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} \left(\frac{\pi - \vartheta}{2} \right)^m \left[1 - \frac{3l(l+1) - m(m+1)}{3(m+1)} \left(\frac{\pi - \vartheta}{2} \right)^2 \right]. \quad (5.12.5)$$

In the case when $\pi/2 - \varepsilon \leq \vartheta \leq \pi/2 + \varepsilon$

$$Y_{l\pm m}(\vartheta, \varphi) \approx (-1)^{\frac{l\pm m}{2}} e^{\pm im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m-1)!!}{(l-m)!!} \frac{(l+m-1)!!}{(l+m)!!}} \left[1 - \frac{l(l+1)-m^2}{2} \left(\frac{\pi}{2} - \vartheta \right)^2 \right] \quad (5.12.6)$$

if $l+m$ is even and

$$Y_{l\pm m}(\vartheta, \varphi) \approx (-1)^{\frac{l\pm m-1}{2}} e^{\pm im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!!}{(l-m-1)!!} \frac{(l+m)!!}{(l+m-1)!!}} \left(\frac{\pi}{2} - \vartheta \right) \times \left[1 - \frac{l(l+1)-(m^2+1)}{6} \left(\frac{\pi}{2} - \vartheta \right)^2 \right] \quad (5.12.7)$$

if $l+m$ is odd.

If ϑ is small, $Y_{lm}(\vartheta, \varphi)$ may be approximated by the Bessel function (McDonald formula [23])

$$Y_{l-m}(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!}{(l-m)!}} e^{-im\varphi} \left[\left(l + \frac{1}{2} \right) \cos \frac{\vartheta}{2} \right]^{-m} \times \left\{ J_m(\alpha) + \left(\sin \frac{\vartheta}{2} \right)^2 \left[\frac{\alpha}{6} J_{m+3}(\alpha) - J_{m+2}(\alpha) + \frac{1}{2\alpha} J_{m+1}(\alpha) \right] + O \left[\left(\sin \frac{\vartheta}{2} \right)^4 \right] \right\} \quad (5.12.8)$$

Here $\alpha = (2l+1) \sin(\vartheta/2)$, $J_\lambda(\alpha)$ is the Bessel function.

5.12.3 Asymptotic Expression for Fixed $m, l \rightarrow \infty, \vartheta \rightarrow 0$ and Finite $l\vartheta$

In this case

$$Y_{l-m}(\vartheta, \varphi) \approx \sqrt{\frac{l}{2\pi}} e^{-im\varphi} J_m(l\vartheta) \quad (5.12.9)$$

5.13 $Y_{lm}(\vartheta, \varphi)$ for special values of l and m

5.13.1 $Y_{lm}(\vartheta, \varphi)$ for $l \leq 5$

Please use sympy or Mathematica to generate the explicit forms of the spherical harmonics for $l = 0, 1, 2, 3, 4, 5$ in terms of trigonometric functions.




Figure 5.1. Spherical harmonics $Y_{lm}(\vartheta, 0)$ are shown as functions of the polar angle ϑ . Red and Blue colours indicate the sign of the function (positive and negative respectively).

5.13.2 $Y_{lm}(\vartheta, \varphi)$ for $|m| = 0, 1, 2, 3, 4$ and Any Integer l

In this case $Y_{lm}(\vartheta, \varphi)$ may be expressed in terms of the Legendre polynomials,

$$Y_{l0}(\vartheta, \varphi) = \sqrt{\frac{2l+1}{4\pi}} P_l(\cos \vartheta), \quad (5.13.1)$$

$$Y_{l\pm 1}(\vartheta, \varphi) = \mp \frac{e^{\pm i\varphi}}{\sin \vartheta} \sqrt{\frac{l(l+1)}{4\pi(2l+1)}} [P_{l-1}(\cos \vartheta) - P_{l+1}(\cos \vartheta)], \quad (5.13.2)$$

$$Y_{l\pm 2}(\vartheta, \varphi) = \frac{e^{\pm i2\varphi}}{(\sin \vartheta)^2} \sqrt{\frac{(l-1)l(l+1)(l+2)}{4\pi(2l+1)}} \left[\frac{P_{l-2}(\cos \vartheta)}{2l-1} - \frac{2(2l+1)}{(2l-1)(2l+3)} P_l(\cos \vartheta) + \frac{1}{2l+3} P_{l+2}(\cos \vartheta) \right], \quad (5.13.3)$$

$$Y_{l\pm 3}(\vartheta, \varphi) = \mp \frac{e^{\pm i3\varphi}}{(\sin \vartheta)^3} \sqrt{\frac{1}{4\pi(2l+1)}} \cdot \frac{(l+3)!}{(l-3)!} \left[\frac{1}{(2l-3)(2l-1)} P_{l-3}(\cos \vartheta) - \frac{3}{(2l-3)(2l+3)} P_{l-1}(\cos \vartheta) + \frac{3}{(2l-1)(2l+5)} P_{l+1}(\cos \vartheta) - \frac{1}{(2l+3)(2l+5)} P_{l+3}(\cos \vartheta) \right], \quad (5.13.4)$$

$$Y_{l\pm 4}(\vartheta, \varphi) = \frac{e^{\pm i4\varphi}}{(\sin \vartheta)^4} \sqrt{\frac{1}{4\pi(2l+1)}} \cdot \frac{(l+4)!}{(l-4)!} \left[\frac{1}{(2l-5)(2l-3)(2l-1)} P_{l-4}(\cos \vartheta) - \frac{4}{(2l-5)(2l-1)(2l+3)} P_{l-2}(\cos \vartheta) + \frac{6(2l+1)}{(2l-3)(2l-1)(2l+3)(2l+5)} P_l(\cos \vartheta) - \frac{4}{(2l-1)(2l+3)(2l+7)} P_{l+2}(\cos \vartheta) + \frac{1}{(2l+3)(2l+5)(2l+7)} P_{l+4}(\cos \vartheta) \right]. \quad (5.13.5)$$

5.13.3 $Y_{lm}(\vartheta, \varphi)$ for $|m| = l, l-1, l-2, l-3, l-4, l-5$ and Any Integer l

In this case the spherical harmonics are conveniently expressed in terms of trigonometric functions.

$$Y_{l\pm l}(\vartheta, \varphi) = (\mp 1)^l e^{\pm il\varphi} \sqrt{\frac{(2l+1)!!}{4\pi(2l)!!}} (\sin \vartheta)^l \quad (5.13.6)$$

$$Y_{l\pm(l-1)}(\vartheta, \varphi) = (\mp 1)^{l-1} e^{\pm i(l-1)\varphi} \sqrt{\frac{(2l+1)!!}{4\pi(2l-2)!!}} \cos \vartheta (\sin \vartheta)^{l-1} \quad (5.13.7)$$

$$Y_{l\pm(l-2)}(\vartheta, \varphi) = (\mp 1)^{l-2} e^{\pm i(l-2)\varphi} \sqrt{\frac{2l+1}{8\pi} \frac{(2l-3)!!}{(2l-2)!!}} (\sin \vartheta)^{l-2} [(2l-1) \cos^2 \vartheta - 1] \quad (5.13.8)$$

$$Y_{l\pm(l-3)}(\vartheta, \varphi) = (\mp 1)^{l-3} e^{\pm i(l-3)\varphi} \sqrt{\frac{2l+1}{24\pi} \frac{(2l-3)!!}{(2l-4)!!}} (\sin \vartheta)^{l-3} \cos \vartheta [(2l-1) \cos^2 \vartheta - 3] \quad (5.13.9)$$

$$Y_{l\pm(l-4)}(\vartheta, \varphi) = (\mp 1)^{l-4} e^{\pm i(l-4)\varphi} \sqrt{\frac{2l+1}{96\pi} \frac{(2l-5)!!}{(2l-4)!!}} (\sin \vartheta)^{l-4} \times [(2l-1)(2l-3) \cos^4 \vartheta - 6(2l-3) \cos^2 \vartheta + 3] \quad (5.13.10)$$

$$Y_{l\pm(l-5)}(\vartheta, \varphi) = (\mp 1)^{l-5} e^{\pm i(l-5)\varphi} \sqrt{\frac{2l+1}{480\pi} \frac{(2l-5)!!}{(2l-6)!!}} (\sin \vartheta)^{l-5} \cos \vartheta \times [(2l-1)(2l-3) \cos^4 \vartheta - 10(2l-3) \cos^2 \vartheta + 15] \quad (5.13.11)$$

5.14 $Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ for special ϑ

(a)

$$Y_{lm}(0, \varphi) = \delta_{m0} \sqrt{\frac{2l+1}{4\pi}}, \quad (5.14.1)$$

$$Y_{lm}\left(\frac{\pi}{2}, \varphi\right) = \begin{cases} (-1)^{\frac{l+m}{2}} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m-1)!!}{(l+m)!!} \frac{(l-m-1)!!}{(l-m)!!}} & \text{if } l+m \text{ is even,} \\ 0 & \text{if } l+m \text{ is odd,} \end{cases} \quad (5.14.2)$$

$$Y_{lm}(\pi, \varphi) = \delta_{m0} (-1)^l \sqrt{\frac{2l+1}{4\pi}}, \quad (5.14.3)$$

$$Y_{lm}(\pm\pi n, \varphi) = \delta_{m0} (-1)^{nl} \sqrt{\frac{2l+1}{4\pi}}. \quad (5.14.4)$$

(b)

$$\left. \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) \right|_{\vartheta=0} = (\delta_{m-1} e^{-i\varphi} - \delta_{m1} e^{i\varphi}) \sqrt{\frac{l(l+1)(2l+1)}{16\pi}} \quad (5.14.5)$$

$$\left. \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) \right|_{\vartheta=\frac{\pi}{2}} = \begin{cases} (-1)^{\frac{l+m+1}{2}} e^{im\varphi} \sqrt{\frac{2l+1}{4\pi} \frac{(l+m)!!}{(l+m-1)!!} \frac{(l-m)!!}{(l-m-1)!!}} & \text{if } l+m \text{ is odd} \\ 0 & \text{if } l+m \text{ is even} \end{cases} \quad (5.14.6)$$

$$\left. \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) \right|_{\vartheta=\pi} = (-1)^l (\delta_{m-1} e^{-i\varphi} - \delta_{m1} e^{i\varphi}) \sqrt{\frac{l(l+1)(2l+1)}{16\pi}} \quad (5.14.7)$$

$$\left. \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) \right|_{\vartheta=\pm n\pi} = (-1)^{nl} (\delta_{m-1} e^{-i\varphi} - \delta_{m1} e^{i\varphi}) \sqrt{\frac{l(l+1)(2l+1)}{16\pi}} \quad (5.14.8)$$

5.15 Zeros of $Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$

In the interval $0 \leq \vartheta \leq \pi$ the spherical harmonics $Y_{lm}(\vartheta, \varphi)$ and their derivatives $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ possess some zeros.

The number of zeros, i.e., the number of roots of the equations

$$Y_{lm}(\vartheta, \varphi) = 0 \quad \text{and} \quad \frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) = 0 \quad (5.15.1)$$

is finite in interval $0 < \vartheta < \pi$. All these roots are non-degenerate. Below we denote the zeros of $Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ by ϑ_α and ϑ_β , respectively. Note that the functions in question have no roots with respect to φ in the domain $0 \leq \varphi < 2\pi$.

5.15.1 Zeros of $Y_{lm}(\vartheta, \varphi)$

If $m = 0$, $Y_{lm}(\vartheta, \varphi)$ has l zeros in the interval $0 < \vartheta < \pi$. If $m \neq 0$, $Y_{lm}(\vartheta, \varphi)$ has $(l - |m|)$ zeros inside the same interval of ϑ and also two zeros at $\vartheta_\alpha = 0$ and $\vartheta_\alpha = \pi$. All these zeros lie symmetrically with respect to $\vartheta = \pi/2$. Hence, $Y_{lm}(\pi/2, \varphi) = 0$ provided $(l - m)$ is odd. If $m \neq 0$, the zeros of $Y_{lm}(\vartheta, \varphi)$ are determined by the equations

$$\begin{aligned} & \text{if } m = \pm l, \quad \cos^2 \vartheta_\alpha = 1; \\ & m = \pm(l-1) \quad \cos^2 \vartheta_\alpha = 1, 0; \\ & m = \pm(l-2) \quad \cos^2 \vartheta_\alpha = 1, \frac{1}{2l-1}; \\ & m = \pm(l-3) \quad \cos^2 \vartheta_\alpha = 1, \frac{3}{(2l-1)}, 0; \\ & m = \pm(l-4) \quad \cos^2 \vartheta_\alpha = 1, \frac{3 \pm 2\sqrt{3(l-2)/(2l-3)}}{2l-1}; \\ & m = \pm(l-5) \quad \cos^2 \vartheta_\alpha = 1, \frac{5 \pm 2\sqrt{5(l-3)/(2l-3)}}{(2l-1)}, 0. \end{aligned} \quad (5.15.2)$$

Numerical values of ϑ_α for can be found using computer programs. If $l \gg 1$, approximate values of ϑ_α may be found from the relation

$$\vartheta_\alpha \approx \frac{\pi}{2} \left(\frac{4k + 3 - 2m}{2l + 1} \right), \quad (5.15.3)$$

where k is integer and $\frac{2|m|-3}{4} \leq k \leq \frac{4l+2|m|-1}{4}$.

5.15.2 Zeros of $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$

If $m = 0$, $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ has $l - 1$ zeros in the interval $0 < \vartheta < \pi$ and also two zeros at $\vartheta_\beta = 0$ and $\vartheta_\beta = \pi$. If $m \neq 0$, $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ has $(l - |m| + 1)$ zeros in the same interval and two additional ones at $\vartheta_\beta = 0$ and $\vartheta_\beta = \pi$ provided $m \geq 2$. All zeros of $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ lie symmetrically with respect to $\vartheta = \pi/2$. Hence $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi) = 0$ for $\vartheta = \pi/2$ provided $(l - m + 1)$ is odd. When $m \neq 0$ the values of ϑ_β are determined by the equations

$$\begin{aligned} \text{if } m = \pm l \quad \cos^2 \vartheta_\beta &= 0; 1 (l \geq 2); \\ m = \pm(l-1) \quad \cos^2 \vartheta_\beta &= \frac{1}{l}; 1 (l \geq 3); \\ m = \pm(l-2) \quad \cos^2 \vartheta_\beta &= \frac{5l-4}{l(2l-1)}; 0; 1 (l \geq 4); \\ m = \pm(l-3) \quad \cos^2 \vartheta_\beta &= \frac{9(l-1) \pm \sqrt{3(19l^2 - 50l + 27)}}{2l(2l-1)}; 1 (l \geq 5) \\ m = \pm(l-4) \quad \cos^2 \vartheta_\beta &= \frac{7l-8 \pm 2\sqrt{(11l^3 - 62l^2 + 104l - 48)/(2l-3)}}{l(2l-1)}; 0; 1 (l \geq 6) \end{aligned} \quad (5.15.4)$$

It is evident that $Y_{lm}(\vartheta, \varphi)$ has an extremum at $\vartheta = \vartheta_\beta$. Zeros of $Y_{lm}(\vartheta, \varphi)$ and $\frac{\partial}{\partial \vartheta} Y_{lm}(\vartheta, \varphi)$ alternate in the domain $0 \leq \vartheta \leq \pi$.

All harmonics $Y_{lm}(\vartheta, 0)$ and $Y_{l+s, m+s}(\vartheta, 0)$ with $m > 0$, and $m + s > 0$ have the same number of maxima, minima and zeros, i.e., can be represented by topologically equivalent plots (Fig. 5.1). With increasing $l + s$ the positions of all maxima, minima and zeros of $Y_{l+s, m+s}(\vartheta, 0)$ shift to $\vartheta = \pi/2$.

5.16 Bipolar and tripolar spherical harmonics

In many applications one has to deal with functions which depend on two or three vector directions. Convenient bases for expansions of such functions are provided by the bipolar and tripolar harmonics.

5.16.1 Bipolar Spherical Harmonics

Bipolar spherical harmonics are given by irreducible tensor product of the spherical harmonics with different arguments

$$\{\mathbf{Y}_{l_1}(\vartheta_1, \varphi_1) \otimes \mathbf{Y}_{l_2}(\vartheta_2, \varphi_2)\}_{LM} = \sum_{m_1 m_2} C_{l_1 m_1 l_2 m_2}^{LM} Y_{l_1 m_1}(\vartheta_1, \varphi_1) Y_{l_2 m_2}(\vartheta_2, \varphi_2) \quad (5.16.1)$$

The variety of bipolar spherical harmonics with different l_1, l_2, L and M forms a complete orthonormal set of functions which depend on two unit vectors, $\mathbf{n}_1$ and $\mathbf{n}_2$. These functions possess simple transformation properties under rotations and inversion of coordinate system.

The orthogonality and normalization relation for these harmonics is given by

$$\iint d\Omega_1 d\Omega_2 \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \{\mathbf{Y}_{l'_1}(\Omega_1) \otimes \mathbf{Y}_{l'_2}(\Omega_2)\}_{L'M'} = \delta_{l_1 l'_1} \delta_{l_2 l'_2} \delta_{LL'} \delta_{MM'} \quad (5.16.2)$$

where

$$\Omega \equiv \{\vartheta, \varphi\}, \quad \int d\Omega \equiv \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta$$

The completeness condition has the form

$$\sum_{l_1 l_2 LM} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \{\mathbf{Y}_{l_1}(\Omega'_1) \otimes \mathbf{Y}_{l_2}(\Omega'_2)\}_{LM}^* = \delta(\Omega_1 - \Omega'_1) \delta(\Omega_2 - \Omega'_2) \quad (5.16.3)$$

where

$$\delta(\Omega - \Omega') \equiv \delta(\cos \vartheta - \cos \vartheta') \delta(\varphi - \varphi').$$

In addition, let us note the following equation

$$\sum_{LM} |\{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM}|^2 = \frac{(2l_1+1)(2l_2+1)}{(4\pi)^2} \quad (5.16.4)$$

Under a rotation of the coordinate system the bipolar harmonics transform according to

$$\hat{D}(\alpha, \beta, \gamma) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM'} = \{\mathbf{Y}_{l_1}(\Omega'_1) \otimes \mathbf{Y}_{l_2}(\Omega'_2)\}_{LM'} = \sum_M D_{MM'}^L(\alpha, \beta, \gamma) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \quad (5.16.5)$$

where the angles $\Omega'_i \equiv \{\vartheta'_i, \varphi'_i\}$ and $\Omega_i \equiv \{\vartheta_i, \varphi_i\}$ are related by Eqs. (1.4.2), (1.4.3).

Under inversion of coordinate system the phases of the bipolar harmonics change according to

$$\hat{P}_r \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} = (-1)^{l_1+l_2} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \quad (5.16.6)$$

The Clebsch-Gordan series for product of the bipolar harmonics may be presented in the form

$$\begin{aligned} & \{\mathbf{Y}_{l'_1}(\Omega_1) \otimes \mathbf{Y}_{l'_2}(\Omega_2)\}_{L'M'} \{\mathbf{Y}_{l''_1}(\Omega_1) \otimes \mathbf{Y}_{l''_2}(\Omega_2)\}_{L''M''} \\ &= \sum_{LM} C_{L'M' L''M''}^{LM} \sum_{l_1 l_2} B_{l'_1 l'_2 L' l''_1 l''_2 L''}^{l_1 l_2 L} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM}, \end{aligned} \quad (5.16.7)$$

where

$$B_{l'_1 l'_2 L' l''_1 l''_2 L''}^{l_1 l_2 L} = \sqrt{\frac{(2l'_1+1)(2l'_2+1)(2l''_1+1)(2l''_2+1)(2L'+1)(2L''+1)}{(4\pi)^2}} C_{l'_1 0 l''_1 0}^{l_1 0} C_{l'_2 0 l''_2 0}^{l_2 0} \begin{Bmatrix} l'_1 & l''_1 & l_1 \\ l'_2 & l''_2 & l_2 \\ L' & L'' & L \end{Bmatrix}. \quad (5.16.8)$$

For the important special case $L = 0$ a bipolar harmonic is reduced to a scalar product of the spherical harmonics

$$\{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{00} = \frac{(-1)^{l_1}}{\sqrt{2l_1+1}} (\mathbf{Y}_{l_1}(\Omega_1) \cdot \mathbf{Y}_{l_1}(\Omega_2)) \delta_{l_1 l_2} \quad (5.16.9)$$

In accordance with Eq. (3.1.30) this scalar product is given by

$$(\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) = \sum_{m=-l}^l Y_{lm}^*(\Omega_1) Y_{lm}(\Omega_2) \quad (5.16.10)$$

5.16.2 Tripolar Spherical Harmonics

A tripolar spherical harmonic is defined as an irreducible tensor product of three spherical harmonics with different arguments,

$$\begin{aligned} & \{\mathbf{Y}_{l_1}(\vartheta_1, \varphi_1) \otimes \{\mathbf{Y}_{l_2}(\vartheta_2, \varphi_2) \otimes \mathbf{Y}_{l_3}(\vartheta_3, \varphi_3)\}_{l_{23}}\}_{LM} \\ &= \sum_{\substack{m_1 m_2 m_3 \\ m_{23}}} C_{l_1 m_1 l_{23} m_{23}}^{LM} C_{l_2 m_2 l_3 m_3}^{l_{23} m_{23}} Y_{l_1 m_1}(\vartheta_1, \varphi_1) Y_{l_2 m_2}(\vartheta_2, \varphi_2) Y_{l_3 m_3}(\vartheta_3, \varphi_3) \end{aligned} \quad (5.16.11)$$

In contrast to bipolar harmonics, the tripolar harmonics allow different coupling schemes of angular momenta, namely, $l_1 + (l_2 + l_3)_{l_{23}} = \mathbf{L}$, $(l_1 + l_2)_{l_{12}} + l_3 = \mathbf{L}$ or $(l_1 + l_3)_{l_{13}} + l_2 = \mathbf{L}$. The relations between different coupling schemes are considered in Sec. 3.3.

The tripolar spherical harmonics form a complete orthonormal set of functions which depend on three unit vectors n_1, n_2 and n_3 .

The orthogonality and normalization relations is as follows

$$\begin{aligned} & \iiint d\Omega_1 d\Omega_2 d\Omega_3 \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} \{\mathbf{Y}_{l'_1}(\Omega_1) \otimes \{\mathbf{Y}_{l'_2}(\Omega_2) \otimes \mathbf{Y}_{l'_3}(\Omega_3)\}_{\lambda'}\}_{L'M'}^* \\ &= \delta_{l_1 l'_1} \delta_{l_2 l'_2} \delta_{l_3 l'_3} \delta_{\lambda \lambda'} \delta_{L L'} \delta_{M M'} \end{aligned} \quad (5.16.12)$$

The completeness relation may be presented in the form

$$\begin{aligned} & \sum_{\substack{l_1 l_2 l_3 \\ \lambda LM}} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} \{\mathbf{Y}_{l_1}(\Omega'_1) \otimes \{\mathbf{Y}_{l_2}(\Omega'_2) \otimes \mathbf{Y}_{l_3}(\Omega'_3)\}_{\lambda}\}_{LM}^* \\ &= \delta(\Omega_1 - \Omega'_1) \delta(\Omega_2 - \Omega'_2) \delta(\Omega_3 - \Omega'_3) \end{aligned} \quad (5.16.13)$$

In addition we also have the following equation

$$\sum_{\lambda LM} |\{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM}|^2 = \frac{(2l_1+1)(2l_2+1)(2l_3+1)}{(4\pi)^3} \quad (5.16.14)$$

Under a rotation of the coordinate system the tripolar harmonics transform according to

$$\begin{aligned} \hat{D}(\alpha, \beta, \gamma) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM'} &= \{\mathbf{Y}_{l_1}(\Omega'_1) \otimes \{\mathbf{Y}_{l_2}(\Omega'_2) \otimes \mathbf{Y}_{l_3}(\Omega'_3)\}_{\lambda}\}_{LM'} \\ &= \sum_M D_{MM'}^L(\alpha, \beta, \gamma) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} \end{aligned} \quad (5.16.15)$$

where the angles $\Omega'_i \equiv \{\vartheta'_i, \varphi'_i\}$ and $\Omega_i \equiv \{\vartheta_i, \varphi_i\}$ are related through Eqs. (1.4.2), (1.4.3).

Under inversion of the coordinate system the tripolar harmonics acquire the phase factor

$$\hat{P}_r \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} = (-1)^{l_1+l_2+l_3} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} \quad (5.16.16)$$

The Clebsch-Gordan series for the tripolar harmonics is given by

$$\begin{aligned} &\left\{ \mathbf{Y}_{l'_1}(\Omega_1) \otimes \{\mathbf{Y}_{l'_2}(\Omega_2) \otimes \mathbf{Y}_{l'_3}(\Omega_3)\}_{\lambda'} \right\}_{L'M'} \left\{ \mathbf{Y}_{l''_1}(\Omega_1) \otimes \{\mathbf{Y}_{l''_2}(\Omega_2) \otimes \mathbf{Y}_{l''_3}(\Omega_3)\}_{\lambda''} \right\}_{L''M''} \\ &= \sum_{LM} C_{L'M'L''M''}^{LM} \sum_{l_1 l_2 l_3 \lambda} B_{l'_1 l'_2 l'_3 \lambda' L' l''_1 l''_2 l''_3 \lambda'' L''}^{l_1 l_2 l_3 \lambda L} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{LM} \end{aligned} \quad (5.16.17)$$

where

$$\begin{aligned} B_{l'_1 l'_2 l'_3 \lambda' L' l''_1 l''_2 l''_3 \lambda'' L''}^{l_1 l_2 l_3 \lambda L} &= \sqrt{\frac{(2l'_1+1)(2l'_2+1)(2l'_3+1)(2l''_1+1)(2l''_2+1)(2l''_3+1)(2L'+1)(2L''+1)}{(4\pi)^3}} \\ &\times \sqrt{(2\lambda'+1)(2\lambda''+1)(2\lambda+1)} C_{l'_1 0 l'_1 0}^{l_1 0} C_{l'_2 0 l'_2 0}^{l_2 0} C_{l'_3 0 l'_3 0}^{l_3 0} \begin{Bmatrix} l'_1 & l'_2 & l'_3 \\ \lambda' & \lambda' & \lambda' \\ L' & L' & L' \end{Bmatrix} \begin{Bmatrix} l''_1 & l''_2 & l''_3 \\ \lambda'' & \lambda'' & \lambda'' \\ L'' & L'' & L'' \end{Bmatrix} \begin{Bmatrix} l_1 & l_2 & l_3 \\ \lambda & \lambda & \lambda \\ L & L & L \end{Bmatrix} \end{aligned} \quad (5.16.18)$$

Putting $L = 0$ in Eq. (5.16.11), we get an important special case for the tripolar scalar harmonic:

$$\begin{aligned} &\{\mathbf{Y}_{l_1}(\Omega_1) \otimes \{\mathbf{Y}_{l_2}(\Omega_2) \otimes \mathbf{Y}_{l_3}(\Omega_3)\}_{\lambda}\}_{00} \\ &= (-1)^{l_1+l_2+l_3} \delta_{\lambda l_1} \sum_{m_1 m_2 m_3} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} Y_{l_1 m_1}(\Omega_1) Y_{l_2 m_2}(\Omega_2) Y_{l_3 m_3}(\Omega_3) \end{aligned} \quad (5.16.19)$$

5.17 Expansions of functions which depend on two vectors

5.17.1 Preliminary Remarks

Any function $f(\mathbf{r}_1, \mathbf{r}_2)$ which depends on two arbitrary vectors $\mathbf{r}_1(r_1, \vartheta_1, \varphi_1)$ and $\mathbf{r}_2(r_2, \vartheta_2, \varphi_2)$ may be expanded into a series of the bipolar harmonics (Sec. 5.16):

$$f(\mathbf{r}_1, \mathbf{r}_2) = \sum_{l_1 l_2 LM} C_{l_1 l_2}^{LM}(r_1, r_2) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \quad (5.17.1)$$

The function $f(\mathbf{r}_1, \mathbf{r}_2)$ may be presented in terms of $\mathbf{r}_1$ and $\mathbf{r}_2$ as well as in terms of $\mathbf{R}$ and $\mathbf{r}$, where

$$\mathbf{R} = \mathbf{r}_1 + \mathbf{r}_2, \mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2, \mathbf{r}_1 = \frac{1}{2}(\mathbf{R} + \mathbf{r}), \mathbf{r}_2 = \frac{1}{2}(\mathbf{R} - \mathbf{r}) \quad (5.17.2)$$

Polar coordinates of the vectors $\mathbf{R}(R, \Theta, \Phi)$ and $\mathbf{r}(r, \vartheta, \varphi)$ are related to polar coordinates of the vectors $\mathbf{r}_1(r_1, \vartheta_1, \varphi_1)$ and $\mathbf{r}_2(r_2, \vartheta_2, \varphi_2)$ by

$$\begin{aligned} R^2 &= r_1^2 + r_2^2 + 2r_1 r_2 \cos \omega_{12}, \\ \cos \Theta &= \frac{r_1 \cos \vartheta_1 + r_2 \cos \vartheta_2}{\sqrt{r_1^2 + r_2^2 + 2r_1 r_2 \cos \omega_{12}}}, \\ \tan \Phi &= \frac{r_1 \sin \vartheta_1 \sin \varphi_1 + r_2 \sin \vartheta_2 \sin \varphi_2}{r_1 \sin \vartheta_1 \cos \varphi_1 + r_2 \sin \vartheta_2 \cos \varphi_2}, \\ r^2 &= r_1^2 + r_2^2 - 2r_1 r_2 \cos \omega_{12}, \\ \cos \vartheta &= \frac{r_1 \cos \vartheta_1 - r_2 \cos \vartheta_2}{\sqrt{r_1^2 + r_2^2 - 2r_1 r_2 \cos \omega_{12}}}, \\ \tan \varphi &= \frac{r_1 \sin \vartheta_1 \sin \varphi_1 - r_2 \sin \vartheta_2 \sin \varphi_2}{r_1 \sin \vartheta_1 \cos \varphi_1 - r_2 \sin \vartheta_2 \cos \varphi_2}. \end{aligned} \quad (5.17.3)$$

Here ω_{12} is the angle between $\mathbf{r}_1$ and $\mathbf{r}_2$ (Fig. 5.2),

$$\cos \omega_{12} = \cos \vartheta_1 \cos \vartheta_2 + \sin \vartheta_1 \sin \vartheta_2 \cos (\varphi_1 - \varphi_2). \quad (5.17.4)$$

The inverse relations to (5.17.3) can be written in the form

$$\begin{aligned} 4r_1^2 &= R^2 + r^2 + 2Rr \cos \omega, & 4r_2^2 &= R^2 + r^2 - 2Rr \cos \omega, \\ \cos \vartheta_1 &= \frac{R \cos \Theta + r \cos \vartheta}{\sqrt{R^2 + r^2 + 2Rr \cos \omega}}, & \cos \vartheta_2 &= \frac{R \cos \Theta - r \cos \vartheta}{\sqrt{R^2 + r^2 - 2Rr \cos \omega}}, \\ \tan \varphi_1 &= \frac{R \sin \Theta \sin \Phi + r \sin \vartheta \sin \varphi}{R \sin \Theta \cos \Phi + r \sin \vartheta \cos \varphi}, & \tan \varphi_2 &= \frac{R \sin \Theta \sin \Phi - r \sin \vartheta \sin \varphi}{R \sin \Theta \cos \Phi - r \sin \vartheta \cos \varphi}. \end{aligned} \quad (5.17.5)$$

ω being the angle between $\mathbf{R}$ and $\mathbf{r}$ (Fig. 5.2)

$$\cos \omega = \cos \Theta \cos \vartheta + \sin \Theta \sin \vartheta \cos (\Phi - \varphi). \quad (5.17.6)$$

Note that the expressions for the polar coordinates of $\mathbf{R}$ and $\mathbf{r}$ are interchanged by the replacement $\mathbf{r}_2 \rightarrow -\mathbf{r}_2$, i.e., $\vartheta_2 \rightarrow \pi - \vartheta_2, \varphi_2 \rightarrow \pi + \varphi_2$. Hence, the expansions of functions $F(\mathbf{R}, \mathbf{r})$ and $F(\mathbf{r}, \mathbf{R})$, which may be reduced to each other by the replacement $\mathbf{R} \rightleftharpoons \mathbf{r}$ are mutually related. If

$$F(\mathbf{R}, \mathbf{r}) = f(\mathbf{r}_1, \mathbf{r}_2) = \sum_{l_1 l_2 LM} C_{l_1 l_2}^{LM}(r_1, r_2) \{ \mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2) \}_{LM} \quad (5.17.7)$$




Figure 5.2. Relations between the vectors $\mathbf{R}, \mathbf{r}$ and $\mathbf{r}_1, \mathbf{r}_2$.

then

$$F(\mathbf{r}, \mathbf{R}) = f(\mathbf{r}_1, -\mathbf{r}_2) = \sum_{l_1 l_2 LM} (-1)^{l_2} C_{l_1 l_2}^{LM}(r_1, r_2) \{ \mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2) \}_{LM} \quad (5.17.8)$$

5.17.2 Addition Theorem for the Spherical Harmonics

The functions $f(r_1, r_2)$ which are invariant under rotations of coordinate systems play an especially important role in applications. These functions depend only on three variables $r_1 = |\mathbf{r}_1|, r_2 = |\mathbf{r}_2|, (r_1 \cdot r_2) = r_1 r_2 \cos \omega_{12}$ or, equivalently, on $R = |\mathbf{r}_1 + \mathbf{r}_2|, r = |\mathbf{r}_1 - \mathbf{r}_2|$ and $(\mathbf{R} \cdot \mathbf{r}) = Rr \cos \omega$. The expansions of these functions will evidently contain only the bipolar harmonics of zero rank ($L = 0$), i.e., only scalar products $(Y_1(\Omega_1) \cdot Y_1(\Omega_2))$ (Eqs. (5.16.10)). Such expansions are equivalent to the expansions in terms of the Legendre polynomials, which depend on $\cos \omega_{12}$ because

$$(Y_l(\Omega_1) \cdot Y_l(\Omega_2)) = \frac{2l+1}{4\pi} P_l(\cos \omega_{12}). \quad (5.17.9)$$

This relation is called the addition theorem for the spherical harmonics.

5.17.3 Expansions of Some Functions Which Depend on $(\mathbf{r}_1 \cdot \mathbf{r}_2)$

$$(\mathbf{r}_1 \cdot \mathbf{r}_2) = \frac{4\pi}{3} r_1 r_2 (\mathbf{Y}_1(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_1(\vartheta_2, \varphi_2)) \quad (5.17.10)$$

$$(\mathbf{r}_1 \cdot \mathbf{r}_2)^2 = \frac{4\pi}{3} r_1^2 r_2^2 \left[(\mathbf{Y}_0(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_0(\vartheta_2, \varphi_2)) + \frac{2}{5} (\mathbf{Y}_2(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_2(\vartheta_2, \varphi_2)) \right] \quad (5.17.11)$$

$$(\mathbf{r}_1 \cdot \mathbf{r}_2)^3 = \frac{4\pi}{5} r_1^3 r_2^3 \left[(\mathbf{Y}_1(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_1(\vartheta_2, \varphi_2)) + \frac{2}{7} (\mathbf{Y}_3(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_3(\vartheta_2, \varphi_2)) \right] \quad (5.17.12)$$

In the general case [48], for integer n , one has

$$(\mathbf{r}_1 \cdot \mathbf{r}_2)^n = 4\pi r_1^n r_2^n \sum_l \frac{n!}{(n-l)!!(n+l+1)!!} (\mathbf{Y}_l(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_l(\vartheta_2, \varphi_2)). \quad (5.17.13)$$

In this formula the summation index l assumes the values $l = 0, 2, \dots, n-2, n$ if n is even, and $l = 1, 3, \dots, n-2, n$ if n is odd.

The expansion of the exponential function is given by

$$e^{i(\mathbf{r}_1 \cdot \mathbf{r}_2)} = 4\pi \sum_{l=0}^{\infty} i^l j_l(r_1 r_2) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.14)$$

where $j_l(x)$ is a spherical Bessel function.

An arbitrary function $f(\mathbf{r}_1 \cdot \mathbf{r}_2)$ which can be written as a power series

$$f(r_1 \cdot r_2) = \sum_{n=0}^{\infty} c_n (r_1 \cdot r_2)^n \quad (5.17.15)$$

may be expanded in terms of the spherical harmonics:

$$f(\mathbf{r}_1 \cdot \mathbf{r}_2) = \sum_{l=0}^{\infty} f_l(r_1 r_2) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.16)$$

The expansion coefficients are determined by

$$f_l(r_1 r_2) = 4\pi \sum_{n=l, l+2}^{\infty} c_n \frac{n! (r_1 r_2)^n}{(n-1)!!(n+l+1)!!} \quad (5.17.17)$$

5.17.4 Expansions of Some Functions Which Depend on $r = |\mathbf{r}_1 - \mathbf{r}_2|$

An expansion of the delta function $\delta(\mathbf{r}_1 - \mathbf{r}_2)$ is

$$\delta(\mathbf{r}_1 - \mathbf{r}_2) = \frac{\delta(r_1 - r_2)}{r_1^2} \sum_{l=0}^{\infty} (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.18)$$

This equation results from the completeness relation for the spherical harmonics (Eqs. (5.6.1)).

An expansion of the Green's function for the scalar Helmholtz equation may be written in the form

$$\frac{e^{ikr}}{r} = 4\pi i k \sum_{l=0}^{\infty} j_l(kr_1) h_l^{(1)}(kr_2) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)), \quad (r_1 < r_2) \quad (5.17.19)$$

where $j_l(x)$ and $h_l^{(1)}(x)$ are the spherical Bessel function and the Hankel function of the first kind; respectively. Equation (5.17.19) is valid at $r_1 < r_2$; for $r_1 > r_2$ one should make the replacement $r_1 \leftrightarrow r_2$. Equation (5.17.19) follows from a more general expansion which is given in Ref. [49].

$$z_0(kr) = 4\pi \sum_{l=0}^{\infty} j_l(kr_1) z_l(kr_2) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)), \quad (r_1 < r_2) \quad (5.17.20)$$

Here $z_l(x) = \sqrt{\pi/2x} Z_{l+\frac{1}{2}}(x)$, and $Z_{l+\frac{1}{2}}(x)$ is any of the cylinder functions.

When $k = 0$ Eq. (5.17.19) yields the important expansion of the Green's function for the scalar Laplace equation

$$\frac{1}{r} = \frac{4\pi}{r_2} \sum_{l=0}^{\infty} \frac{1}{(2l+1)} \left(\frac{r_1}{r_2}\right)^l (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)), \quad (r_1 < r_2) \quad (5.17.21)$$

This result as well as Eq. (5.17.19) are valid when $r_1 < r_2$, whereas if $r_1 > r_2$ it becomes valid after the interchange $r_1 \leftrightarrow r_2$. We may also write Eq. (5.17.21) in a symmetric form valid both at $r_1 < r_2$ and at $r_1 > r_2$:

$$\frac{1}{r} = \frac{4\pi}{\sqrt{r_1^2 + r_2^2}} \sum_{l=0}^{\infty} \left\{ \sum_{n=l, l+2}^{\infty} \frac{(2n-1)!!}{(n-l)!!(l+n+1)!!} \left(\frac{r_1 r_2}{r_1^2 + r_2^2} \right)^n \right\} (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.22)$$

Below we give some examples of expansions of r^n ($n = 1, -3, -5$)

$$r = 4\pi \sum_{l=0}^{\infty} \frac{1}{(2l+1)} \left(\frac{r_1^l}{r_2^{l+1}} \right) \left(\frac{r_1^2}{2l+3} - \frac{r_2^2}{2l-1} \right) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.23)$$

$$\frac{1}{r^3} = \frac{4\pi}{r_2^2 - r_1^2} \sum_{l=0}^{\infty} \left(\frac{r_1^l}{r_2^{l+1}} \right) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.24)$$

$$\frac{1}{r^5} = \frac{4\pi}{3(r_2^2 - r_1^2)^3} \sum_{l=0}^{\infty} (2l-1)(2l+3) \left(\frac{r_1^l}{r_2^{l+1}} \right) \left(\frac{r_2^2}{2l-1} - \frac{r_1^2}{2l+3} \right) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)) \quad (5.17.25)$$

These equations are valid when $r_1 < r_2$, whereas for $r_1 > r_2$ they are valid after interchanging r_1 and r_2 . Equations (5.17.23)-(5.17.25) follow from the expansions which will be considered below.

5.17.5 Expansions of $r^n \equiv |r_1 - r_2|^n$

Let n be an integer. Then the expansion under discussion is as follows

$$r^n = 4\pi \sum_{l=0}^{\infty} \frac{1}{2l+1} a_l^n(r_1, r_2) (\mathbf{Y}_l(\Omega_1) \cdot \mathbf{Y}_l(\Omega_2)), \quad (5.17.26)$$

where the expansion coefficients $a_l^n(r_1, r_2)$ are determined by (see Ref. [50])

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l}{\left(\frac{1}{2}\right)_l} r_2^n \left(\frac{r_1}{r_2}\right)^l F\left(l - \frac{n}{2}, -\frac{1}{2} - \frac{n}{2}; l + \frac{3}{2}; \frac{r_1^2}{r_2^2}\right), (r_1 < r_2), \quad (5.17.27)$$

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l}{\left(\frac{1}{2}\right)_l} \frac{r_1^l (r_2^2 - r_1^2)^{n+2}}{r_2^{l+n+4}} F\left(l + 2 + \frac{n}{2}, \frac{3}{2} + \frac{n}{2}; l + \frac{3}{2}; \frac{r_1^2}{r_2^2}\right), (r_1 < r_2), \quad (5.17.28)$$

$$a_l^n(r_1, r_2) = \frac{2^{n+1} \left(l + \frac{1}{2}\right) \left(-\frac{n}{2}\right)_l}{\left(1 + \frac{n}{2}\right)_{l+1} r_2^{n-l} F\left(l - \frac{n}{2}, -\frac{1}{2} - \frac{n}{2}; -1 - n; \frac{r_2^2 - r_1^2}{r_2^2}\right)} - \frac{2l+1}{2^{n+3}(n+2)} \cdot \frac{r_1^l (r_2^2 - r_1^2)^{n+2}}{r_2^{l+n+4}} F\left(l + \frac{n}{2} + 2, \frac{3}{2} + \frac{n}{2}; n + 3; \frac{r_2^2 - r_1^2}{r_2^2}\right), \quad (5.17.29)$$

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l (-1)^{\frac{n}{2}} \cos \frac{n\pi}{2} r_1^{n-l} r_2^l F\left(l - \frac{n}{2}, -\frac{1}{2} - \frac{n}{2}; l + \frac{3}{2}; \frac{r_2^2}{r_1^2}\right)}{\left(\frac{1}{2}\right)_l} + \frac{\Gamma\left(l + \frac{3}{2}\right) \sqrt{\pi} (-1)^{\frac{n+1}{2}} r_1^{n+l+1}}{\Gamma\left(-\frac{n}{2}\right) \Gamma\left(2 + l + \frac{n}{2}\right) r_2^{l+1}} F\left(-1 - l - \frac{n}{2}, -\frac{1}{2} - \frac{n}{2}; \frac{1}{2} - l; \frac{r_2^2}{r_1^2}\right), \quad (5.17.30)$$

If $r_1 > r_2$, one should interchange r_1 and r_2 in Eqs. (5.17.27)-(5.17.30). Let us present also symmetric expressions for $a_l^n(r_1, r_2)$ which are valid when $r_1 < r_2$ and $r_1 > r_2$:

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l}{\left(\frac{1}{2}\right)_l} \cdot \frac{(r_1 r_2)^l}{(r_1^2 + r_2^2)^{l - \frac{n}{2}}} F\left(\frac{l}{2} - \frac{n}{4}, \frac{l}{2} - \frac{n}{4} + \frac{1}{2}; l + \frac{3}{2}; \left(\frac{2r_1 r_2}{r_1^2 + r_2^2}\right)^2\right) \quad (5.17.31)$$

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l}{\left(\frac{1}{2}\right)_l} \cdot \frac{(r_1 r_2)^l}{(r_1 + r_2)^{2l-n}} F\left(l - \frac{n}{2}, 1 + l; 2l + 2; \frac{4r_1 r_2}{(r_1 + r_2)^2}\right) \quad (5.17.32)$$

A comparison of Eqs. (5.17.27) and (5.17.28) shows that the expansion coefficient for r^n and r^{-n-4} are related by

$$a_l^n(r_1, r_2) = \frac{\left(-\frac{n}{2}\right)_l}{\left(\frac{n}{2} + 2\right)_l} (r_2^2 - r_1^2)^{n+2} a_l^{-n-4}(r_1, r_2) \quad (5.17.33)$$

Note that all the expansions given in Secs. 5.17.4 and 5.17.5 remain valid after substituting $R = |\mathbf{r}_1 + \mathbf{r}_2|$ for $r = |r_1 - r_2|$ and multiplying the expansion coefficients by $(-1)^l$.

5.17.6 Expansions of Spherical Waves

The spherical waves have the form $z_L(kr)Y_{LM}(\vartheta, \varphi)$, where $\mathbf{r}(r, \vartheta, \varphi) = \mathbf{r}_1 - \mathbf{r}_2$, $z_L(x) = \sqrt{\pi/2x}Z_{L+\frac{1}{2}}(x)$ and $Z_{L+\frac{1}{2}}(x)$ is any of the cylinder functions. The spherical waves are expressed in terms of the bipolar spherical harmonics of rank L [49] (Sec. 5.16.1)

$$z_L(kr)Y_{LM}(\vartheta, \varphi) = \sqrt{\frac{4\pi}{2L+1}} \sum_{l_1, l_2=0}^{\infty} i^{l_1-l_2-L} \sqrt{(2l_1+1)(2l_2+1)} \\ \times C_{l_1 0 l_2 0}^{L 0} j_{l_1}(kr_1) z_{l_2}(kr_2) \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM}, (r_1 < r_2) \quad (5.17.34)$$

When $r_1 > r_2$, one should interchange r_1 and r_2 in Eq. (5.17.34). If $z_L(kr) = j_L(kr)$, $j_L(x)$ being a spherical Bessel function, one obtains as $k \rightarrow 0$ [22, 96]

$$r^L Y_{LM}(\vartheta, \varphi) = \sqrt{4\pi(2L+1)!} \sum_{\substack{l_1, l_2=0 \\ l_1+l_2=L}}^L (-1)^{l_2} \frac{r_1^{l_1} r_2^{l_2}}{\sqrt{(2l_1+1)!(2l_2+1)!}} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \quad (5.17.35)$$

Similarly, if $z_L(kr) = n_L(kr)$, $n_L(kr)$ being a spherical Neumann function, one obtains as $k \rightarrow 0$ [22, 96]

$$\frac{1}{r^{L+1}} Y_{LM}(\vartheta, \varphi) = \sqrt{\frac{4\pi}{(2L)!}} \sum_{\substack{l_1, l_2=0 \\ l_2-l_1=L}}^{\infty} (-1)^{l_2} \sqrt{\frac{(2l_2)!}{(2l_1+1)!}} \cdot \frac{r_1^{l_1}}{r_2^{l_2+1}} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM}, (r_1 < r_2) \quad (5.17.36)$$

For example, spherical components of r/r^3 read

$$\frac{r_\mu}{r^3} = 4\pi \sum_{l=1}^{\infty} (-1)^l \sqrt{\frac{l}{3}} \cdot \frac{r_1^{l-1}}{r_2^{l+1}} \{\mathbf{Y}_{l-1}(\Omega_1) \otimes \mathbf{Y}_l(\Omega_2)\}_{1\mu}, (r_1 < r_2) \quad (5.17.37)$$

5.17.7 Expansions of $r^N Y_{LM}(\vartheta, \varphi)$

The expansion of this function in terms of bipolar harmonics of rank L may be written as

$$r^N Y_{LM}(\vartheta, \varphi) = 4\pi \sum_{l_1, l_2=0}^{\infty} a_{l_1 l_2}^{NL}(r_1, r_2) \sqrt{\frac{(2l_1+1)(2l_2+1)}{4\pi(2L+1)}} C_{l_1 0 l_2 0}^{L 0} \{\mathbf{Y}_{l_1}(\Omega_1) \otimes \mathbf{Y}_{l_2}(\Omega_2)\}_{LM} \quad (5.17.38)$$

The sum is over all positive integer l_1 and l_2 allowed by the momentum addition rules and the parity selection rule ($l_1 + l_2 - L$ is even). The expansion coefficients $a_{l_1 l_2}^{NL}(r_1, r_2)$ are given by (see Ref. [51])

$$a_{l_1 l_2}^{NL}(r_1, r_2) = (-1)^{\frac{l_1-l_2-L}{2}} \frac{2^{l_1}}{(2l_1+1)!!} \frac{\Gamma(\frac{l_1+l_2-N}{2}) \Gamma(\frac{L+N+3}{2})}{\Gamma(\frac{L-N}{2}) \Gamma(\frac{-l_1+l_2+N+3}{2})} \\ \times r_2^N \left(\frac{r_1}{r_2}\right)^{l_1} F\left(\frac{l_1+l_2-N}{2}, \frac{l_1-l_2-N-1}{2}; l_1+\frac{3}{2}; \frac{r_1^2}{r_2^2}\right), (r_1 < r_2). \quad (5.17.39)$$

If $r_1 > r_2$, the expression for the expansion coefficients may be obtained from Eq. (5.17.39) by interchanging r_1 and r_2 , l_1 and l_2 and introducing the additional factor $(-1)^L$. The expression (5.17.39) was studied in detail in Ref. [52]. For different values of N and L the expansion coefficients are determined by the following relations. For $N = L, L+2, L+4, \dots$ we have

$$a_{l_1 l_2}^{NL}(r_1, r_2) = \begin{cases} (-1)^{l_2} (N+L+1)!! (N-L)!! \frac{r_1^{l_1+2e_1} r_2^{2e_2}}{(2e_1)!! (2e_1+2l_1+1)!! (2e_2)!! (2e_2+2l_2+1)!!} \\ \quad (e_1+e_2 = \frac{N-l_1-l_2}{5}) \\ \quad \text{if } l_1+l_2 \leq N, \\ 0 \\ \quad \text{if } l_1+l_2 > N. \end{cases} \quad (5.17.40)$$

For $N = L-2, L-4, L-6, \dots$ one gets

$$a_{l_1 l_2}^{NL}(r_1, r_2) = (-1)^{l_2+\frac{N-2}{2}} \frac{(N+L+1)!!}{2(L-N-2)!! (N+l_1+l_2+2)!} r_1^{l_1} r_2^{l_2} \\ \times \left(\frac{1}{r_1} \cdot \frac{\partial}{\partial r_1}\right)^{l_1} \left(\frac{1}{r_2} \cdot \frac{\partial}{\partial r_2}\right)^{l_2} \frac{1}{r_1 r_2} \left[(r_1-r_2)^{N+l_1+l_2+2} \ln|r_1-r_2| - (r_1+r_2)^{N+l_1+l_2+2} \ln(r_1+r_2) \right]. \quad (5.17.41)$$

If N is integer, $N - L$ is odd, and $r_1 < r_2$, the expansion coefficient is given by

$$a_{l_1 l_2}^{NL}(r_1, r_2) = \begin{cases} (-1)^{N+l_2+1} + l_1 & \frac{(N+L+1)!!}{(L-N-2)!!} r_1^{l_1} r_2^{N-l_1} \sum_{s=0}^{\frac{N-l_1+l_2+1}{2}} \frac{\left(\frac{r_1}{r_2}\right)^{2s}}{(2s)!!(N-l_1-l_3-2s)!!(N-l_1+l_2+1-2s)!!(2l_1+2s+1)!!} \\ 0 & \text{if } N - l_1 + l_2 + 1 \geq 0, \\ & \text{if } N - l_1 + l_2 + 1 < 0. \end{cases} \quad (5.17.42)$$

The expansion coefficient for $r_1 > r_2$ may be obtained from (5.17.42) by interchanging r_1 and r_2 , l_1 and l_2 and introducing the additional factor $(-1)^L$. It should be emphasised that if $N \leq -3$ the expansion coefficients contain additional terms involving the delta function $\delta(r_1 - r_2)$ and its derivatives [52]

$$a_{l_1 l_2}^{NL}(r_1, r_2)_\delta = (-1)^{N+l_1-1} \frac{(N+L+1)!!}{(L-N-2)!!} \sum_{e_1=0}^{l_1} \sum_{\substack{e_2=0 \\ e_1+e_2+N \leq -3}}^{l_2} (-1)^{e_2} \frac{(l_1+s_1)!(l_2+s_2)!}{(l_1-s_1)!(l_2-s_2)!} \\ \times \frac{1}{(2s_1)!!(2s_2)!!} \cdot \frac{1}{r_2^{\beta_2+1}} \left(\frac{\partial}{\partial r_2} \right)^{-N-s_1-e_2-3} \left[\frac{\delta(r_1 - r_2)}{r_1^2 r_2^{\beta_1-1}} \right]. \quad (5.17.43)$$

These additional terms give finite contributions into the integrals involving the expansion (5.17.38).

Finally, if N is not integer, the expansion coefficient is given by

$$a_{l_1 l_2}^{NL}(r_1, r_2) = (-1)^{l_1+L+1} \frac{2^{L-1} \Gamma\left(\frac{N+L+3}{2}\right) \Gamma(N-L+2)}{\Gamma\left(\frac{N-L+3}{2}\right) \Gamma(N+l_1+l_2+3)} r_1^{l_1} r_2^{l_2} \\ \times \left(\frac{1}{r_1} \frac{\partial}{\partial r_1} \right)^{l_1} \left(\frac{1}{r_2} \frac{\partial}{\partial r_2} \right)^{l_2} \frac{1}{r_1 r_2} \left\{ |r_1 + r_2|^{N+l_1+l_2+2} - |r_1 - r_2|^{N+l_1+l_2+2} \right\} \quad (5.17.44)$$

Note that all the expansions given in Secs. 5.17.6 and 5.17.7 remain valid after substituting $\mathbf{R}(R, \Theta, \Phi) = \mathbf{r}_1 + \mathbf{r}_2$ for $\mathbf{r}(r, \vartheta, \varphi) = \mathbf{r}_1 - \mathbf{r}_2$ and introducing the additional factor $(-1)^{l_2}$ into the expansion coefficients.

Chapter 6

Spin functions

6.1 Spin functions of particles with arbitrary spin

6.1.1 Definition

Spin functions describe polarization states of particles¹ of definite spin, i.e. definite intrinsic angular momentum.

The spin functions $\chi(\sigma)$ may be treated as functions of a discrete variable σ which is the spin projection on the z -axis. The variable σ takes $2S+1$ values, $\sigma = -S, -S+1, \dots, S-1, S$, where S is the particle spin (integer or half integer non-negative number). The quantity $|\chi(\sigma)|^2$ gives the probability that the spin projection on the z -axis in a given state equals σ .

The spin functions are commonly written as column matrices which contain $2S+1$ elements:

$$\chi = \begin{pmatrix} \chi(S) \\ \chi(S-1) \\ \vdots \\ \chi(-S+1) \\ \chi(-S) \end{pmatrix}. \quad (6.1.1)$$

The elements of this matrix give the values of the spin function $\chi(\sigma)$ for corresponding values of spin variable σ . The quantities $\chi(\sigma)$ are called the contravariant components of the spin function χ (see Sec. 6.1.4 below).

In such a representation operators acting on the spin variables take the form of square $(2S+1) \times (2S+1)$ matrices, and summation over the spin variable is replaced by matrix multiplication. The Hermitian conjugate function $\chi^\dagger$ has the form of a row matrix

$$\chi^\dagger = (\chi^*(S), \chi^*(S-1), \dots, \chi^*(-S+1), \chi^*(-S)). \quad (6.1.2)$$

The interpretation of $|\chi(\sigma)|^2$ as the probability for the spin projection on the z -axis to be equal to σ is possible only if $\chi(\sigma)$ satisfies the normalization condition

$$\sum_{\sigma=-S}^S |\chi(\sigma)|^2 = 1. \quad (6.1.3)$$

¹ This condition may be written also in a matrix form

$$\chi^\dagger \chi = 1. \quad (6.1.4)$$

The above representation of spin functions is called the spherical basis representation. When the spin value is integer, one may also use the cartesian basis representation. The latter representation is considered below (Sec. 6.3) for spin-1 particles.

6.1.2 Basis Spin Functions

According to definition, the basis spin functions describe the states with definite spin and spin projection on the z -axis. The basis spin functions are eigenfunctions of the operators $\hat{\mathbf{S}}^2$ and $\hat{S}_z$, where $\hat{\mathbf{S}}$ is the spin operator (see Sec. 2.3)

$$\hat{\mathbf{S}}^2 \chi_{Sm} = S(S+1) \chi_{Sm}, \hat{S}_z \chi_{Sm} = m \chi_{Sm}. \quad (6.1.5)$$

¹We use the name "particle" not only for an elementary particle (electron, nucleon, etc.) but also for any composite system (atom, molecule) which can be treated in phenomena under consideration as one object.

As follows from the definition, the dependence of the basis spin functions $\chi_{Sm}(\sigma)$ on the spin variable σ is given by

$$\chi_{Sm}(\sigma) = \delta_{m\sigma}. \quad (6.1.6)$$

In other words, contravariant components of the basis spin functions χ_{Sm} have the form

$$[\chi_{Sm}]^\sigma = \delta_{m\sigma}. \quad (6.1.7)$$

If the basis spin functions are written as column matrices, then

$$\chi_{SS} = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \chi_{SS-1} = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \dots, \quad \chi_{S-S} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}. \quad (6.1.8)$$

The collection of $2S + 1$ basis functions $\chi_{Sm} (m = S, S-1, \dots, -S)$ constitutes a complete orthonormal set of functions. The orthonormality condition for the basis functions is

$$\chi_{Sm}^\dagger \chi_{Sm'} = \delta_{mm'}. \quad (6.1.9)$$

The completeness condition for the basis spin functions may be written in matrix form

$$\sum_{m=-S}^S \chi_{Sm} \chi_{Sm}^\dagger = \hat{I}. \quad (6.1.10)$$

where $\hat{I}$ is the unit $(2S + 1) \times (2S + 1)$ matrix.

Matrix products $\chi_{Sm} \chi_{Sm'}^\dagger$ of the basis spin functions are the square $(2S + 1) \times (2S + 1)$ matrices. They may be expanded in terms of the polarization operators $\hat{T}_{LM}(S)$ (see 2.4),

$$\chi_{Sm} \chi_{Sm'}^\dagger = \sum_L \sqrt{\frac{2L+1}{2S+1}} C_{S m' L M}^S \hat{T}_{LM}(S). \quad (6.1.11)$$

Matrix elements of spherical components of the spin operator $\hat{S}_\mu (\mu = \pm 1, 0)$ between the basis spin functions are given by

$$\chi_{Sm'}^\dagger \hat{S}_\mu \chi_{Sm} = \sqrt{S(S+1)} C_{S m' 1 \mu}^S; \quad (6.1.12)$$

The only non-vanishing matrix elements are the following:

$$\begin{aligned} \chi_{Sm+1}^\dagger \hat{S}_{+1} \chi_{Sm} &= -\frac{1}{\sqrt{2}} \sqrt{(S-m)(S+m+1)}, \\ \chi_{Sm}^\dagger \hat{S}_0 \chi_{Sm} &= m, \\ \chi_{Sm-1}^\dagger \hat{S}_{-1} \chi_{Sm} &= \frac{1}{\sqrt{2}} \sqrt{(S+m)(S-m+1)}. \end{aligned} \quad (6.1.13)$$

For cartesian components of the spin operator $\hat{S}_i (i = x, y, z)$ the non-vanishing matrix elements are

$$\begin{aligned} \chi_{Sm\pm 1}^\dagger \hat{S}_x \chi_{Sm} &= \frac{1}{2} \sqrt{(S \mp m)(S \pm m + 1)}, \\ \chi_{Sm\pm 1}^\dagger \hat{S}_y \chi_{Sm} &= \mp \frac{i}{2} \sqrt{(S \mp m)(S \pm m + 1)}, \\ \chi_{Sm}^\dagger \hat{S}_z \chi_{Sm} &= m. \end{aligned} \quad (6.1.14)$$

The matrix elements of the polarization operators $\hat{T}_{LM}(S)$ between the basis spin functions may be written as

$$\chi_{Sm'}^\dagger \hat{T}_{LM}(S) \chi_{Sm} = \sqrt{\frac{2L+1}{2S+1}} C_{S m' L M}^S. \quad (6.1.15)$$

Under rotations of the coordinate system the basis spin functions are transformed either by the rotation operator $\hat{D}^S(\alpha, \beta, \gamma)$ (Eq. (2.4.17)), if rotations are defined by the Euler angles α, β, γ , or by the rotation operator $\hat{U}^S(\omega; \theta, \Phi)$ (Eq. (2.4.18)) if rotations are described by the rotation axis $\mathbf{n}(\theta, \Phi)$ and the rotation angle ω

$$\begin{aligned} \chi'_{Sm'} &\equiv \hat{D}^S(\alpha, \beta, \gamma) \chi_{Sm'} = \sum_m D_{mm'}^S(\alpha, \beta, \gamma) \chi_{Sm}, \\ \chi'_{Sm'} &\equiv \hat{U}^S(\omega; \Theta, \Phi) \chi_{Sm'} = \sum_m U_{mm'}^S(\omega; \Theta, \Phi) \chi_{Sm}, \end{aligned} \quad (6.1.16)$$

where $D_{mm'}^S$ are the Wigner D -functions (Chap. 4), and $U_{mm'}^S$ are the functions defined in Sec. 4.5. The functions $\chi'_{Sm'}$ describe quantum states with definite spin S and spin projection m' on the new z' -axis. They are eigenfunctions of the operators $\hat{\mathbf{S}}'^2$ and $\hat{S}'_z$,

$$\begin{aligned}\hat{\mathbf{S}}'^2 \chi'_{Sm'} &= S(S+1) \chi'_{Sm'}, \\ \hat{S}'_z \chi'_{Sm'} &= m' \chi'_{Sm'}.\end{aligned}\tag{6.1.17}$$

Here $\hat{\mathbf{S}}'$ is the spin operator in the rotated coordinate system

$$\hat{S}'_\mu = \sum_\nu D_{\nu\mu}^1(\alpha, \beta, \gamma) \hat{S}_\nu, \quad (\mu, \nu = \pm 1, 0).\tag{6.1.18}$$

6.1.3 Helicity Basis Functions

The spin projection on the linear momentum direction of a particle is called the helicity. For particles of spin S the helicity λ assumes $(2S+1)$ values, $\lambda = S, S-1, \dots, -S+1, S$.

The helicity basis functions $\chi_{S\lambda}(\vartheta, \varphi)$, by definition, are the eigenfunctions of the operators $\hat{\mathbf{S}}^2$ and $\hat{\mathbf{S}} \cdot \mathbf{n}$ where $\hat{\mathbf{S}}$ is the spin operator, $\mathbf{n} = \mathbf{p}/p$ is the momentum direction of the particles, ϑ and φ are the polar angles of the vector $\mathbf{n}$,

$$\begin{aligned}\hat{\mathbf{S}}^2 \chi_{S\lambda}(\vartheta, \varphi) &= S(S+1) \chi_{S\lambda}(\vartheta, \varphi), \\ \hat{\mathbf{S}} \cdot \mathbf{n} \chi_{S\lambda}(\vartheta, \varphi) &= \lambda \chi_{S\lambda}(\vartheta, \varphi).\end{aligned}\tag{6.1.19}$$

The function $\chi_{S\lambda}(\vartheta, \varphi)$ describes a quantum state with definite spin S and helicity λ .

The helicity basis functions $\chi_{S\lambda}(\vartheta, \varphi)$ may be expressed in terms of the basis functions χ_{Sm} by means of the rotation which turns the z -axis parallel to $\mathbf{n}(\vartheta, \varphi)$. Such a rotation is determined by the Euler angles $\alpha = \varphi$, $\beta = \vartheta$. The third Euler angle γ (the angle of rotation about the new z' -axis) is arbitrary. For simplicity, let us adopt $\gamma = 0$. In this case the rotated cartesian basis vectors $\mathbf{e}'_x, \mathbf{e}'_y, \mathbf{e}'_z$ will coincide with the polar basis vectors $\mathbf{e}_\theta, \mathbf{e}_\varphi, \mathbf{e}_\rho$, respectively (Sec. 1.1.2). Making use of the transformation properties of the basis spin functions χ_{Sm} under the rotation (6.1.16) we obtain

$$\begin{aligned}\chi_{S\lambda}(\vartheta, \varphi) &= \sum_m D_{m\lambda}^S(\varphi, \vartheta, 0) \chi_{Sm}, \\ \chi_{Sm} &= \sum_\lambda D_{-\lambda-m}^S(0, \vartheta, \varphi) \chi_{S\lambda}(\vartheta, \varphi).\end{aligned}\tag{6.1.20}$$

For Hermitian adjoint functions we have

$$\begin{aligned}\chi_{S\lambda}^\dagger(\vartheta, \varphi) &= \sum_m (-1)^{\lambda-m} D_{-m-\lambda}^S(\varphi, \vartheta, 0) \chi_{Sm}^\dagger, \\ \chi_{Sm}^\dagger &= \sum_\lambda (-1)^{\lambda-m} D_{\lambda m}^S(0, \vartheta, \varphi) \chi_{S\lambda}^\dagger(\vartheta, \varphi).\end{aligned}\tag{6.1.21}$$

It follows from (6.1.7), (6.1.20), (6.1.21) that the contravariant components of the helicity basis functions have the form

$$\begin{aligned}[\chi_{S\lambda}(\vartheta, \varphi)]^\sigma &= D_{\sigma\lambda}^S(\varphi, \vartheta, 0), \\ \left[\chi_{S\lambda}^\dagger(\vartheta, \varphi) \right]^\sigma &= (-1)^{\lambda-\sigma} D_{-\sigma-\lambda}^S(\varphi, \vartheta, 0). \quad (\sigma, \lambda = -S, -S+1, \dots, S-1, S).\end{aligned}\tag{6.1.22}$$

The collection of $2S+1$ helicity basis functions $\chi_{S\lambda}(\vartheta, \varphi)$ ($\lambda = S, S-1, \dots, -S$) as well as the set of the functions χ_{Sm} form complete orthonormal sets. The orthonormality condition for the helicity basis functions is

$$\chi_{S\lambda}^\dagger(\vartheta, \varphi) \chi_{S\lambda'}(\vartheta, \varphi) = \delta_{\lambda\lambda'}.\tag{6.1.23}$$

The completeness condition for these functions has the following matrix form

$$\sum_\lambda \chi_{S\lambda}(\vartheta, \varphi) \chi_{S\lambda}^\dagger(\vartheta, \varphi) = \hat{I}.\tag{6.1.24}$$

Matrix products $\chi_{S\lambda}(\vartheta, \varphi) \chi_{S\lambda'}^\dagger(\vartheta, \varphi)$ of the helicity basis functions are the square $(2S+1) \times (2S+1)$ matrices. They may be expanded in terms of the polarization operators $\hat{T}_{LM}(S)$ (see Sec. 2.4) as

$$\chi_{S\lambda}(\vartheta, \varphi) \chi_{S\lambda'}^\dagger(\vartheta, \varphi) = \sum_{LM} \sqrt{\frac{2L+1}{2S+1}} C_{S\lambda' L A}^S D_{MA}^L(\varphi, \vartheta, 0) \hat{T}_{LM}(S).\tag{6.1.25}$$

Matrix elements of the polarization operators between the helicity basis functions are given by

$$\chi_{S\lambda'}^\dagger(\vartheta, \varphi) \hat{T}_{LM}(S) \chi_{S\lambda}(\vartheta, \varphi) = \sqrt{\frac{2L+1}{2S+1}} (-1)^{M+\Lambda} C_{S\lambda' L A}^S D_{-M-A}^L(\varphi, \vartheta, 0). \quad (6.1.26)$$

In particular, the diagonal matrix elements are

$$\chi_{S\lambda}^\dagger(\vartheta, \varphi) \hat{T}_{LM}(S) \chi_{S\lambda}(\vartheta, \varphi) = \sqrt{\frac{4\pi}{2S+1}} C_{S\lambda L 0}^S Y_{LM}(\vartheta, \varphi). \quad (6.1.27)$$

6.1.4 General Spin Functions

Any spin function (6.1.1) of particles of spin S can be expanded in sum of the basis spin functions χ_{Sm} ,

$$\chi = \sum_{m=-S}^S a^m \chi_{Sm}, \quad (6.1.28)$$

where $a^m \equiv \chi(m)$ is a contravariant component of some irreducible tensor of rank S . Similarly, the expansion of the Hermitian adjoint function $\chi^\dagger$ has the form

$$\chi^\dagger = \sum_{m=-S}^S (a^m)^* \chi_{Sm}^\dagger, \quad (6.1.29)$$

In general, $(a^m)^* \neq a_m$, i.e., the corresponding irreducible tensor is not real. The normalization condition (6.1.4) imposes a restriction on a^m :

$$\sum_{m=-S}^S |a^m|^2 = 1$$

The product $\chi\chi^\dagger$ of two spin functions represents the square $(2S+1) \times (2S+1)$ matrix which may be expanded in terms of the polarization operators $\hat{T}_{LM}(S)$ (Sec. 2.4)

$$\chi\chi^\dagger = \sum_{L=0}^{2S} \sum_{M=-L}^L (-1)^M P_{L-M} \hat{T}_{LM}(S), \quad (6.1.30)$$

where the expansion coefficients may be expressed through a^m as

$$P_{LM} = \sqrt{\frac{2L+1}{2S+1}} \sum_{m,n=-S}^S C_{S m L M}^S (a^n)^* a^m. \quad (6.1.31)$$

In particular, $P_{00} = 1/\sqrt{2S+1}$.

The quantities $P_{LM}(M = -L, -L+1, \dots, L)$ are covariant components of some irreducible tensor of rank L . They represent the mean values of the polarization operators $\hat{T}_{LM}(S)$ in the states described by the spin functions χ :

$$\chi^\dagger \hat{T}_{LM}(S) \chi = P_{LM}. \quad (6.1.32)$$

Under complex conjugation P_{LM} transforms as

$$(P_{LM})^* = (-1)^M P_{L-M}. \quad (6.1.33)$$

The equality $\chi\chi^\dagger\chi\chi^\dagger = \chi\chi^\dagger$ implies the following conditions for the coefficients P_{LM} ,

$$P_{LM} = \sum_{L_1 M_1 L_2 M_2} (-1)^{2S+L_1+L_2} \sqrt{(2L_1+1)(2L_2+1)} \begin{Bmatrix} L_1 & L_2 & L \\ S & S & S \end{Bmatrix} C_{L_1 M_1 L_2 M_2}^{L M} P_{L_1 M_1} P_{L_2 M_2}. \quad (6.1.34)$$

In particular, Eq. (6.1.34) gives the normalization condition

$$\sum_{L=0}^{2S} \sum_{M=-L}^L |P_{LM}|^2 = 1. \quad (6.1.35)$$

Under rotations of the coordinate system specified by the Euler angles α, β, γ the spin function χ is transformed by the rotation operator $\hat{D}^S(\alpha, \beta, \gamma)$ (Eq. (2.4.17)) into

$$\chi' = \hat{D}^S(\alpha, \beta, \gamma) \chi. \quad (6.1.36)$$

The spin function χ' in the rotated coordinate system may be expanded in sum of basis spin functions χ_{Sm} or χ'_{Sm} in the original or rotated coordinate systems:

$$\chi' = \sum_{m=-S}^S a'^m \chi_{Sm} = \sum_{m=-S}^S a^m \chi'_{Sm}, \quad (6.1.37)$$

where a^m and a'^m are components of the spin function in the original and rotated systems, respectively. The relations between the basis functions χ_{Sm} and χ'_{Sm} are given by Eq. (6.1.16). The transformation properties of a^m under rotations are

$$\begin{aligned} a'^m &= \sum_{n=-S}^S D_{mn}^S(\alpha, \beta, \gamma) a^n, \\ a^n &= \sum_{m=-S}^S D_{mn}^{S*}(\alpha, \beta, \gamma) a'^m, \end{aligned} \quad (6.1.38)$$

in accordance with general transformation rule for contravariant components (Sec. 4.1.2).

6.1.5 Polarization Density Matrix

Spin functions enable one to describe only the totally polarized (pure) particle states.

For describing partially polarized states which are incoherent statistical mixtures of the pure states, the polarization density matrix is used.

The polarization density matrix $\hat{\rho}$ of the particles of spin S is the square $(2S+1) \times (2S+1)$ matrix defined by

$$\rho_{\sigma\sigma'} = \langle \chi(\sigma) \chi^*(\sigma') \rangle_{\xi}, \quad (6.1.39)$$

or in a matrix form by

$$\hat{\rho} = \langle \chi \chi^\dagger \rangle_{\xi}, \quad (6.1.40)$$

where $(\)_{\xi}$ denotes the statistical average. In particular, for pure states one gets

$$\hat{\rho} = \chi \chi^\dagger. \quad (6.1.41)$$

The density matrix is Hermitian,

$$\hat{\rho}^\dagger = \hat{\rho}, \quad \text{i.e., } \rho_{\sigma\sigma'}^* = \rho_{\sigma'\sigma}; \quad (6.1.42)$$

and normalized,

$$\text{Tr } \hat{\rho} = 1, \quad \text{i.e., } \sum_{\sigma} \rho_{\sigma\sigma} = 1. \quad (6.1.43)$$

The conditions (6.1.42), (6.1.43) reveal that the density matrix of particles of spin S is completely determined by $(2S+1)^2 - 1 = 4S(S+1)$ real parameters.

An expectation value of any polarization operator $\hat{T}$ in a state described by the density matrix $\hat{\rho}$ may be evaluated by

$$\langle \hat{T} \rangle = \text{Tr} \{ \hat{T} \hat{\rho} \} = \text{Tr} \{ \hat{\rho} \hat{T} \}. \quad (6.1.44)$$

Note also the following property of the density matrix

$$\text{Tr} \{ \hat{\rho}^2 \} \leq 1, \quad (6.1.45)$$

where the equality holds only for pure states. The density matrix of pure states satisfies the relation

$$\hat{\rho}^2 = \hat{\rho}. \quad (6.1.46)$$

The density matrix $\hat{\rho}$ may be expanded into a sum of the polarisation operators $\hat{T}_{LM}(S)$,

$$\hat{\rho} = \sum_{L=0}^{2S} \sum_{M=-L}^L (-1)^M t_{L-M} \hat{T}_{LM}(S). \quad (6.1.47)$$

The expansion coefficients t_{LM} are called the statistical tensors. They represent expectation values of the polarization operators $\hat{T}_{LM}(S)$ in a state described by the density matrix $\hat{\rho}$,

$$t_{LM} = \langle \hat{T}_{LM}(S) \rangle = \text{Tr} \{ \hat{\rho} \hat{T}_{LM}(S) \}. \quad (6.1.48)$$

The statistical tensors are related to the density matrix elements by

$$t_{LM} = \sqrt{\frac{2L+1}{2S+1}} \sum_{\sigma, \sigma'=-S}^S C_{S\sigma L M}^{S\sigma'} \rho_{\sigma\sigma'}, \quad (6.1.49)$$

$$\rho_{\sigma\sigma'} = \sum_{L,M} \sqrt{\frac{2L+1}{2S+1}} C_{S\sigma L M}^{S\sigma'} t_{LM}. \quad (6.1.50)$$

In particular,

$$t_{00} = \frac{1}{\sqrt{2S+1}}. \quad (6.1.51)$$

Under complex conjugation the statistical tensors transform as

$$t_{LM}^* = (-1)^M t_{L-M}.$$

The properties of the statistical tensors for pure (completely polarized) states have been considered above (Eqs. (6.1.31)-(6.1.35)). For unpolarised states all statistical tensors vanish except t_{00} , thus

$$\hat{\rho}_{\text{unpol}} = \frac{1}{2S+1} \hat{I}, \quad (6.1.52)$$

where $\hat{I}$ is the unit matrix.

The quantities t_{LM} are covariant components of some irreducible tensor of rank L . Under rotations they transform in accordance with Eq. (4.1.2).

6.1.6 Two Particles with Arbitrary Spins

Let us consider a system of two particles, 1 and 2, with spins S_1 and S_2 . Total spin S of the system takes the values $|S_1 - S_2|, |S_1 - S_2| + 1, \dots, S_1 + S_2$. The spin function $\chi_{Sm}(1, 2)$ which describes the system with total spin S and spin projection m may be written as

$$\chi_{Sm}(1, 2) = \sum_{m_1 m_2} C_{S_1 m_1 S_2 m_2}^{S m} \chi_{S_1 m_1}(1) \chi_{S_2 m_2}(2). \quad (6.1.53)$$

The inverse relation is

$$\chi_{S_1 m_1}(1) \chi_{S_2 m_2}(2) = \sum_S C_{S_1 m_1 S_2 m_2}^{S m} \chi_{Sm}(1, 2). \quad (6.1.54)$$

Spin functions (6.1.53) are orthonormalized

$$\chi_{Sm}^\dagger(1, 2) \chi_{S'm'}(1, 2) = \delta_{SS'} \delta_{mm'}, \quad (6.1.55)$$

and satisfy the completeness condition

$$\sum_{S,m} \chi_{Sm}(1, 2) \chi_{Sm}^\dagger(1, 2) = \hat{I}. \quad (6.1.56)$$

Let the polarization operator $\hat{T}_{LM}(i)$ ($i = 1, 2$) act on spin variables of a particle i only. Then we have

$$\begin{aligned} \hat{T}_{L_1 M_1}(1) \chi_{Sm}(1, 2) &= (-1)^{S_1+S_2+S+L_1} \sqrt{(2L_1+1)(2S+1)} \\ &\times \sum_{S'} \left\{ \begin{matrix} S_1 & S_2 & S \\ S' & L_1 & S_1 \end{matrix} \right\} C_{S' m' L_1 M_1}^{S' m'} \chi_{S'm'}(1, 2), \end{aligned} \quad (6.1.57)$$

$$\begin{aligned} \hat{T}_{L_2 M_2}(2) \chi_{Sm}(1, 2) &= \sqrt{(2L_2+1)(2S+1)} \\ &\times \sum_{S'} (-1)^{S_1+S_2+S'+L_2} \left\{ \begin{matrix} S_1 & S_2 & S \\ L_2 & S' & S_2 \end{matrix} \right\} C_{S' m' L_2 M_2}^{S' m'} \chi_{S'm'}(1, 2). \end{aligned} \quad (6.1.58)$$

One may also consider a set of scalar operators $\hat{Q}_L(1, 2)$ with $0 \leq L \leq \min\{2S_1, 2S_2\}$

$$\hat{Q}_L(1, 2) = \sum_M (-1)^M \hat{T}_{LM}(1) \hat{T}_{L-M}(2). \quad (6.1.59)$$

These operators leave the spin of a system and spin projection unchanged:

$$\hat{Q}_L(1, 2) \chi_{Sm}(1, 2) = (-1)^{S_1+S_2+S} (2L+1) \left\{ \begin{matrix} S_1 & S_2 & S \\ S_2 & S_1 & L \end{matrix} \right\} \chi_{Sm}(1, 2). \quad (6.1.60)$$

Among scalar operators the so-called projection operators $\hat{P}_S(1, 2)$ ($|S_1 - S_2| \leq S \leq S_1 + S_2$) are of special interest. When applied to arbitrary spin function $\chi(1, 2)$, the operator $\hat{P}_S(1, 2)$ gives the component of this function with definite total spin S . Thus, the projection operator $\hat{P}_S(1, 2)$ may be defined by the equations

$$\hat{P}_S(1, 2)\chi_{S'm'}(1, 2) = \delta_{SS'}\chi_{S'm'}(1, 2). \quad (6.1.61)$$

From Eqs. (6.1.61) one may obtain the following properties of the projection operators

$$\hat{P}_S(1, 2)\hat{P}_{S'}(1, 2) = \delta_{SS'}\hat{P}_S(1, 2), \quad (6.1.62)$$

$$\sum_S \hat{P}_S(1, 2) = \hat{I}. \quad (6.1.63)$$

The form of $\hat{P}_S(1, 2)$ may be written explicitly as

$$\hat{P}_S(1, 2) = \sum_m \chi_{Sm}(1, 2)\chi_{Sm}^\dagger(1, 2), \quad (6.1.64)$$

or, in terms of scalar operators $\hat{Q}_L(1, 2)$,

$$\hat{P}_S(1, 2) = (-1)^{S_1+S_2+S}(2S+1) \sum_L \left\{ \begin{matrix} S_1 & S_2 & S \\ S_2 & S_1 & L \end{matrix} \right\} \hat{Q}_L(1, 2). \quad (6.1.65)$$

6.2 Spin functions for $S = 1/2$

6.2.1 Basis Spin Functions

The basis functions $\chi_{\frac{1}{2}m}(m = \pm 1/2)$ are the eigenfunctions of the operators $\hat{S}^2$ and $\hat{S}_z$:

$$\begin{aligned} \hat{S}^2 \chi_{\frac{1}{2}m} &= \frac{3}{4} \chi_{\frac{1}{2}m}, \\ \hat{S}_z \chi_{\frac{1}{2}m} &= m \chi_{\frac{1}{2}m}. \end{aligned} \quad (6.2.1)$$

A function $\chi_{\frac{1}{2}m}$ describes the state of a spin- $\frac{1}{2}$ particle with definite spin projection, m , on the z -axis. The dependence of the basis functions $\chi_{\frac{1}{2}m}(\sigma)$ on the spin variable σ is given by

$$\chi_{\frac{1}{2}m}(\sigma) = \delta_{m\sigma}. \quad (6.2.2)$$

According to Eq. (6.1.1), the basis functions $\chi_{\frac{1}{2}m}$ may be written as

$$\chi_{\frac{1}{2}\frac{1}{2}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \chi_{\frac{1}{2}-\frac{1}{2}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (6.2.3)$$

The Hermitian conjugate functions $\chi_{\frac{1}{2}m}^\dagger$ have the form

$$\chi_{\frac{1}{2}\frac{1}{2}}^\dagger = (1, 0), \chi_{\frac{1}{2}-\frac{1}{2}}^\dagger = (0, 1). \quad (6.2.4)$$

The following compact notations for the basis functions are widely used:

$$\chi_{\frac{1}{2}\frac{1}{2}} \equiv \alpha, \chi_{\frac{1}{2}-\frac{1}{2}} \equiv \beta. \quad (6.2.5)$$

The basis functions $\chi_{\frac{1}{2}m}$ satisfy the orthonormality condition

$$\chi_{\frac{1}{2}m'}^\dagger \chi_{\frac{1}{2}m} = \delta_{m'm}. \quad (6.2.6)$$

The completeness condition for the basis functions is

$$\sum_m \chi_{\frac{1}{2}m} \chi_{\frac{1}{2}m}^\dagger = \hat{I}, \quad (6.2.7)$$

where $\hat{I}$ is the unit 2×2 matrix.

6.2.2 Expansions of Products of Basis Functions

The products $\chi_{\frac{1}{2}m}\chi_{\frac{1}{2}m'}^\dagger$ represent 2×2 square matrices whose elements are

$$\left[\chi_{\frac{1}{2}m}\chi_{\frac{1}{2}m'}^\dagger\right]_{\sigma\sigma'} \equiv \chi_{\frac{1}{2}\sigma}^\dagger \left(\chi_{\frac{1}{2}m}\chi_{\frac{1}{2}m'}^\dagger\right) \chi_{\frac{1}{2}\sigma'} = \delta_{\sigma m}\delta_{\sigma'm'}. \quad (6.2.8)$$

The expansions of these products in terms of the spin matrices are given by

$$\chi_{\frac{1}{2}m}\chi_{\frac{1}{2}m'}^\dagger = \frac{1}{2}\delta_{mm'}\hat{I} - \sqrt{3}C_{1\mu\frac{1}{2}m'}^{\frac{1}{2}m}\hat{S}_\mu, \quad (6.2.9)$$

where $S_\mu (\mu = \pm 1, 0)$ are spherical components of the spin operator (Sec. 2.5). In more detailed form Eq. (6.2.9) may be written as

$$\begin{aligned} \alpha\alpha^\dagger &\equiv \chi_{\frac{1}{2}\frac{1}{2}}\chi_{\frac{1}{2}\frac{1}{2}}^\dagger = \frac{1}{2}\hat{I} + \hat{S}_z = \frac{1}{2}\hat{I} + \hat{S}_0, \\ \alpha\beta^\dagger &\equiv \chi_{\frac{1}{2}\frac{1}{2}}\chi_{\frac{1}{2}-\frac{1}{2}}^\dagger = \hat{S}_x + i\hat{S}_y = -\sqrt{2}\hat{S}_{+1}, \\ \beta\alpha^\dagger &\equiv \chi_{\frac{1}{2}-\frac{1}{2}}\chi_{\frac{1}{2}\frac{1}{2}}^\dagger = \hat{S}_x - i\hat{S}_y = \sqrt{2}\hat{S}_{-1}, \\ \beta\beta^\dagger &\equiv \chi_{\frac{1}{2}-\frac{1}{2}}\chi_{\frac{1}{2}-\frac{1}{2}}^\dagger = \frac{1}{2}\hat{I} - \hat{S}_z = \frac{1}{2}\hat{I} - \hat{S}_0. \end{aligned} \quad (6.2.10)$$

6.2.3 Action of Spin Operators on Basis Functions

Cartesian components of a spin operator $\hat{S}_i$ (see Sec. 2.5) applied to the basis functions yield

$$\begin{aligned} \hat{S}_x\chi_{\frac{1}{2}\frac{1}{2}} &= \frac{1}{2}\chi_{\frac{1}{2}-\frac{1}{2}}, & \hat{S}_x\chi_{\frac{1}{2}-\frac{1}{2}} &= \frac{1}{2}\chi_{\frac{1}{2}\frac{1}{2}}, \\ \hat{S}_y\chi_{\frac{1}{2}\frac{1}{2}} &= \frac{i}{2}\chi_{\frac{1}{2}-\frac{1}{2}}, & \hat{S}_y\chi_{\frac{1}{2}-\frac{1}{2}} &= -\frac{i}{2}\chi_{\frac{1}{2}\frac{1}{2}}, \\ \hat{S}_z\chi_{\frac{1}{2}\frac{1}{2}} &= \frac{1}{2}\chi_{\frac{1}{2}\frac{1}{2}}, & \hat{S}_z\chi_{\frac{1}{2}-\frac{1}{2}} &= -\frac{1}{2}\chi_{\frac{1}{2}-\frac{1}{2}}. \end{aligned} \quad (6.2.11)$$

Spherical components of the spin operator $\hat{S}_\mu$ (see Sec. 2.5) act on the basis functions as follows:

$$\hat{S}_\mu\chi_{\frac{1}{2}m} = -\frac{\sqrt{3}}{2}C_{1\mu\frac{1}{2}m}^{\frac{1}{2}m'}\chi_{\frac{1}{2}m'}. \quad (6.2.12)$$

More explicitly Eq. (6.2.12) is

$$\begin{aligned} \hat{S}_{+1}\chi_{\frac{1}{2}\frac{1}{2}} &= 0, & \hat{S}_{+1}\chi_{\frac{1}{2}-\frac{1}{2}} &= -\frac{1}{\sqrt{2}}\chi_{\frac{1}{2}\frac{1}{2}}, \\ \hat{S}_0\chi_{\frac{1}{2}\frac{1}{2}} &= \frac{1}{2}\chi_{\frac{1}{2}\frac{1}{2}}, & \hat{S}_0\chi_{\frac{1}{2}-\frac{1}{2}} &= -\frac{1}{2}\chi_{\frac{1}{2}-\frac{1}{2}}, \\ \hat{S}_{-1}\chi_{\frac{1}{2}\frac{1}{2}} &= \frac{1}{\sqrt{2}}\chi_{\frac{1}{2}-\frac{1}{2}}, & \hat{S}_{-1}\chi_{\frac{1}{2}-\frac{1}{2}} &= 0. \end{aligned} \quad (6.2.13)$$

According to (6.2.12), matrix elements of the operator $\hat{S}_\mu$ for any vector $\mathbf{a}$ have the form

$$[\hat{\mathbf{S}} \cdot \mathbf{a}]_{\sigma\sigma'} = \chi_{\frac{1}{2}\sigma}^\dagger \hat{\mathbf{S}} \cdot \mathbf{a} \chi_{\frac{1}{2}\sigma'} = (-1)^{1-\mu} \frac{\sqrt{3}}{2} C_{1\mu\frac{1}{2}\sigma'}^{\frac{1}{2}\sigma} a_{-\mu}, \quad (6.2.14)$$

where $a_{-\mu} (\mu = \pm 1, 0)$ are covariant spherical components of $\mathbf{a}$ (Sec. 1.2).

6.2.4 Transformation of the Basis Functions Under Rotations of the Coordinate System

The basis functions $\chi_{\frac{1}{2}m}$ are covariant. Under rotations $S \rightarrow S'$ of the coordinate system these functions are transformed by the rotation operator $\hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ (Eq. (2.5.32)) if rotations are described by the Euler angles α, β, γ , or by the rotation operator $\hat{U}^{\frac{1}{2}}(\omega; \Theta, \Phi)$ (Eq. (2.5.36)), if rotations are defined by the rotation axis $n(\theta, \Phi)$ and the rotation angle ω :

$$\begin{aligned} \chi'_{\frac{1}{2}m'} &= \hat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)\chi_{\frac{1}{2}m} = \sum_m \chi_{\frac{1}{2}m} D_{mm'}^{\frac{1}{2}}(\alpha, \beta, \gamma), \\ \chi'_{\frac{1}{2}m'} &= \hat{U}^{\frac{1}{2}}(\omega; \theta, \Phi)\chi_{\frac{1}{2}m} = \sum_m \chi_{\frac{1}{2}m} U_{mm'}^{\frac{1}{2}}(\omega; \theta, \Phi), \end{aligned} \quad (6.2.15)$$

where $D_{mm'}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ are the Wigner D-functions (Chap. 4), and $U_{mm'}^{\frac{1}{2}}$ are defined in Sec. 4.5. A function $\chi'_{\frac{1}{2}m'}$ describes a state in which a particle of spin $\frac{1}{2}$ has the spin projection m' on the new z' -axis. The functions $\chi'_{\frac{1}{2}m'}$ are eigenfunctions of the operators $\hat{\mathbf{S}}'^2$ and $\hat{S}'_x, \hat{\mathbf{S}}'$ being the spin operator in the rotated coordinate system (Eqs.(2.5.40) and (2.5.41)):

$$\begin{aligned}\hat{\mathbf{S}}'^2 \chi'_{\frac{1}{2}m'} &= \frac{3}{4} \chi'_{\frac{1}{2}m'}, \\ \hat{S}'_z \chi'_{\frac{1}{2}m'} &= m' \chi'_{\frac{1}{2}m'}.\end{aligned}\tag{6.2.16}$$

The explicit form of the basis spin functions in the rotated coordinate system is

$$\chi'_{\frac{1}{2}\frac{1}{2}} = \begin{pmatrix} \cos \frac{\beta}{2} e^{-i\frac{\alpha+\gamma}{2}} \\ \sin \frac{\beta}{2} e^{i\frac{\alpha-\gamma}{2}} \end{pmatrix}, \quad \chi'_{\frac{1}{2}-\frac{1}{2}} = \begin{pmatrix} -\sin \frac{\beta}{2} e^{-i\frac{\alpha-\gamma}{2}} \\ \cos \frac{\beta}{2} e^{i\frac{\alpha+\gamma}{2}} \end{pmatrix},\tag{6.2.17}$$

if the rotation is specified by the Euler angles α, β, γ and

$$\chi'_{\frac{1}{2}\frac{1}{2}} = \begin{pmatrix} \cos \frac{\omega}{2} - i \sin \frac{\omega}{2} \cos \theta \\ -i \sin \frac{\omega}{2} \sin \theta e^{i\phi} \end{pmatrix}, \quad \chi'_{\frac{1}{2}-\frac{1}{2}} = \begin{pmatrix} -i \sin \frac{\omega}{2} \sin \theta e^{-i\phi} \\ \cos \frac{\omega}{2} + i \sin \frac{\omega}{2} \cos \theta \end{pmatrix}.\tag{6.2.18}$$

in terms of the angles ω, θ, Φ .

The functions $\chi'_{\frac{1}{2}m'}$ satisfy the orthonormality and completeness conditions (6.2.6), (6.2.7) as well as the relations (6.2.9)-(6.2.13) in which the spin operator $\hat{\mathbf{S}}$ should be replaced by $\hat{\mathbf{S}}'$.

6.2.5 Helicity Basis Functions

The helicity basis functions $\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi)$ ($\lambda = \pm \frac{1}{2}$) describe states in which the spin projection on the linear momentum direction $\mathbf{n}(\vartheta, \varphi) \equiv \mathbf{p}/p$ is equal to λ . These functions are eigenfunctions of the operators $\hat{\mathbf{S}}^2$ and $\hat{\mathbf{S}} \cdot \mathbf{n}$

$$\begin{aligned}\hat{\mathbf{S}}^2 \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) &= \frac{3}{4} \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi), \\ \hat{\mathbf{S}} \cdot \mathbf{n} \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) &= \lambda \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi).\end{aligned}\tag{6.2.19}$$

According to Eq. (6.2.15) the helicity functions $\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi)$ are related to the basis functions $\chi_{\frac{1}{2}m}$ by

$$\begin{aligned}\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) &= \sum_m D_{m\lambda}^{\frac{1}{2}}(\varphi, \vartheta, 0) \chi_{\frac{1}{2}m}, \\ \chi_{\frac{1}{2}m} &= \sum_{\lambda} D_{-\lambda-m}^{\frac{1}{2}}(0, \vartheta, \varphi) \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi).\end{aligned}\tag{6.2.20}$$

For Hermitian adjoint functions Eqs. (6.2.20) yield

$$\begin{aligned}\chi_{\frac{1}{2}\lambda}^{\dagger}(\vartheta, \varphi) &= \sum_m (-1)^{\lambda-m} D_{-m-\lambda}^{\frac{1}{2}}(\varphi, \vartheta, 0) \chi_{\frac{1}{2}m}^{\dagger}, \\ \chi_{\frac{1}{2}m}^{\dagger} &= \sum_{\lambda} (-1)^{\lambda-m} D_{\lambda m}^{\frac{1}{2}}(0, \vartheta, \varphi) \chi_{\frac{1}{2}\lambda}^{\dagger}(\vartheta, \varphi).\end{aligned}\tag{6.2.21}$$

The explicit form of the helicity functions is given by

$$\chi_{\frac{1}{2}\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} \cos \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \\ \sin \frac{\vartheta}{2} e^{i\frac{\varphi}{2}} \end{pmatrix}, \quad \chi_{\frac{1}{2}-\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} -\sin \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \\ \cos \frac{\vartheta}{2} e^{i\frac{\varphi}{2}} \end{pmatrix},\tag{6.2.22}$$

$$\chi_{\frac{1}{2}\frac{1}{2}}^{\dagger}(\vartheta, \varphi) = \begin{pmatrix} \cos \frac{\vartheta}{2} e^{i\frac{\varphi}{2}}, \sin \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \end{pmatrix}, \quad \chi_{\frac{1}{2}-\frac{1}{2}}^{\dagger}(\vartheta, \varphi) = \begin{pmatrix} -\sin \frac{\vartheta}{2} e^{i\frac{\varphi}{2}}, \cos \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \end{pmatrix}.\tag{6.2.23}$$

The helicity functions $\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi)$ as well as $\chi_{\frac{1}{2}m}$ constitute an orthonormalised basis. The orthonormality condition for the helicity functions may be written as

$$\chi_{\frac{1}{2}\lambda'}^{\dagger}(\vartheta, \varphi) \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) = \delta_{\lambda'\lambda}.\tag{6.2.24}$$

The completeness condition is

$$\sum_{\lambda} \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) \chi_{\frac{1}{2}\lambda}^{\dagger}(\vartheta, \varphi) = \hat{I}.\tag{6.2.25}$$

The expansion of products of the helicity functions has the form

$$\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) \chi_{\frac{1}{2}\lambda'}^\dagger(\vartheta, \varphi) = \frac{1}{2} \delta_{\lambda\lambda'} \hat{I} + \sqrt{3} \sum_{\mu, \nu} C_{\frac{1}{2}\lambda' 1\mu}^{\frac{1}{2}\lambda} D_{\nu\mu}^1(\varphi, \vartheta, 0) \hat{S}_\nu. \quad (6.2.26)$$

Explicitly, this gives

$$\begin{aligned} \chi_{\frac{1}{2}\frac{1}{2}}(\vartheta, \varphi) \chi_{\frac{1}{2}\frac{1}{2}}^\dagger(\vartheta, \varphi) &= \frac{1}{2} \begin{pmatrix} 1 + \cos \vartheta & \sin \vartheta e^{-i\varphi} \\ \sin \vartheta e^{i\varphi} & 1 - \cos \vartheta \end{pmatrix} = \frac{1}{2} \hat{I} + \mathbf{n} \cdot \hat{\mathbf{S}}, \\ \chi_{\frac{1}{2}\frac{1}{2}}(\vartheta, \varphi) \chi_{\frac{1}{2}-\frac{1}{2}}^\dagger(\vartheta, \varphi) &= \frac{1}{2} \begin{pmatrix} -\sin \vartheta & (\cos \vartheta + 1)e^{-i\varphi} \\ (\cos \vartheta - 1)e^{i\varphi} & \sin \vartheta \end{pmatrix} = -\sqrt{2} \sum_{\nu} D_{\nu 1}^1(\varphi, \vartheta, 0) \hat{S}_\nu, \\ \chi_{\frac{1}{2}-\frac{1}{2}}(\vartheta, \varphi) \chi_{\frac{1}{2}\frac{1}{2}}^\dagger(\vartheta, \varphi) &= \frac{1}{2} \begin{pmatrix} -\sin \vartheta & (\cos \vartheta - 1)e^{-i\varphi} \\ (\cos \vartheta + 1)e^{i\varphi} & \sin \vartheta \end{pmatrix} = \sqrt{2} \sum_{\nu} D_{\nu -1}^1(\varphi, \vartheta, 0) \hat{S}_\nu, \\ \chi_{\frac{1}{2}-\frac{1}{2}}(\vartheta, \varphi) \chi_{\frac{1}{2}-\frac{1}{2}}^\dagger(\vartheta, \varphi) &= \frac{1}{2} \begin{pmatrix} 1 - \cos \vartheta & -\sin \vartheta e^{-i\varphi} \\ -\sin \vartheta e^{i\varphi} & 1 + \cos \vartheta \end{pmatrix} = \frac{1}{2} \hat{I} - \mathbf{n} \cdot \hat{\mathbf{S}}. \end{aligned} \quad (6.2.27)$$

Matrix elements of the spherical components of the spin operator $\hat{S}_\mu (\mu = \pm 1, 0)$ between the helicity states are given by

$$\chi_{\frac{1}{2}\lambda'}^\dagger(\vartheta, \varphi) \hat{S}_\mu \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) = (-1)^{\mu+\nu} \frac{\sqrt{3}}{2} C_{\frac{1}{2}\lambda' 1\mu}^{\frac{1}{2}\lambda} D_{-\mu-\nu}^1(\varphi, \vartheta, 0). \quad (6.2.28)$$

or, in a detailed form

$$\begin{aligned} \langle + | \hat{S}_{+1} | + \rangle &= -\frac{\sin \vartheta}{2\sqrt{2}} e^{i\varphi}, & \langle + | \hat{S}_{+1} | - \rangle &= -\frac{1 + \cos \vartheta}{2\sqrt{2}} e^{i\varphi}, \\ \langle + | \hat{S}_0 | + \rangle &= \frac{1}{2} \cos \vartheta, & \langle + | \hat{S}_0 | - \rangle &= -\frac{1}{2} \sin \vartheta, \\ \langle + | \hat{S}_{-1} | + \rangle &= \frac{\sin \vartheta}{2\sqrt{2}} e^{-i\varphi}, & \langle + | \hat{S}_{-1} | - \rangle &= -\frac{1 - \cos \vartheta}{2\sqrt{2}} e^{-i\varphi}, \\ \langle - | \hat{S}_{+1} | + \rangle &= \frac{1 - \cos \vartheta}{2\sqrt{2}} e^{i\varphi}, & \langle - | \hat{S}_{+1} | - \rangle &= \frac{\sin \vartheta}{2\sqrt{2}} e^{i\varphi}, \\ \langle - | \hat{S}_0 | + \rangle &= -\frac{1}{2} \sin \vartheta, & \langle - | \hat{S}_0 | - \rangle &= -\frac{1}{2} \cos \vartheta, \\ \langle - | \hat{S}_{-1} | + \rangle &= \frac{1 + \cos \vartheta}{2\sqrt{2}} e^{-i\varphi}, & \langle - | \hat{S}_{-1} | - \rangle &= -\frac{\sin \vartheta}{2\sqrt{2}} e^{-i\varphi}. \end{aligned} \quad (6.2.29)$$

Here we use the abbreviations

$$\begin{aligned} \langle + | \hat{S}_\mu | + \rangle &\equiv \chi_{\frac{1}{2}\frac{1}{2}}^\dagger(\vartheta, \varphi) \hat{S}_\mu \chi_{\frac{1}{2}\frac{1}{2}}(\vartheta, \varphi) \\ \langle + | \hat{S}_\mu | - \rangle &\equiv \chi_{\frac{1}{2}\frac{1}{2}}^\dagger(\vartheta, \varphi) \hat{S}_\mu \chi_{\frac{1}{2}-\frac{1}{2}}(\vartheta, \varphi), \text{ etc.} \end{aligned}$$

Matrix elements of cartesian components of the spin matrices $\hat{S}_i (i = x, y, z)$ have the form

$$\begin{aligned} \langle + | \hat{S}_x | + \rangle &= \frac{1}{2} \sin \vartheta \cos \varphi, & \langle + | \hat{S}_x | - \rangle &= \frac{1}{2} (\cos \vartheta \cos \varphi + i \sin \varphi), \\ \langle + | \hat{S}_y | + \rangle &= \frac{1}{2} \sin \vartheta \sin \varphi, & \langle + | \hat{S}_y | - \rangle &= \frac{1}{2} (\cos \vartheta \sin \varphi - i \cos \varphi), \\ \langle + | \hat{S}_z | + \rangle &= \frac{1}{2} \cos \vartheta, & \langle + | \hat{S}_z | - \rangle &= -\frac{1}{2} \sin \vartheta, \\ \langle - | \hat{S}_x | + \rangle &= \frac{1}{2} (\cos \vartheta \cos \varphi - i \sin \varphi), & \langle - | \hat{S}_x | - \rangle &= -\frac{1}{2} \sin \vartheta \cos \varphi, \\ \langle - | \hat{S}_y | + \rangle &= \frac{1}{2} (\cos \vartheta \sin \varphi + i \cos \varphi), & \langle - | \hat{S}_y | - \rangle &= -\frac{1}{2} \sin \vartheta \sin \varphi, \\ \langle - | \hat{S}_z | + \rangle &= -\frac{1}{2} \sin \vartheta, & \langle - | \hat{S}_z | - \rangle &= -\frac{1}{2} \cos \vartheta \end{aligned} \quad (6.2.30)$$

Note that the diagonal matrix elements of the spin operator may be written as

$$\chi_{\frac{1}{2}\lambda}^\dagger(\vartheta, \varphi) \hat{\mathbf{S}} \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi) = \lambda \mathbf{n}(\vartheta, \varphi). \quad (6.2.31)$$

6.2.6 General Spin Functions for $S = \frac{1}{2}$

An arbitrary spin function $\chi_{\frac{1}{2}}$ of a particle of spin $\frac{1}{2}$ may be expanded in terms of the basis spin functions $\chi_{\frac{1}{2}m}$,

$$\chi_{\frac{1}{2}} = \sum_{m=-\frac{1}{2}}^{\frac{1}{2}} a^m \chi_{\frac{1}{2}m} = \begin{pmatrix} a^{\frac{1}{2}} \\ a^{-\frac{1}{2}} \end{pmatrix}. \quad (6.2.32)$$

An analogous expansion of the Hermitian conjugate function $\chi_{\frac{1}{2}}^\dagger$ is

$$\chi_{\frac{1}{2}}^\dagger = \sum_{m=-\frac{1}{2}}^{\frac{1}{2}} a^{m*} \chi_{\frac{1}{2}m}^\dagger = \left(a^{\frac{1}{2}*}, a^{-\frac{1}{2}*}\right). \quad (6.2.33)$$

Here a^m ($m = \pm\frac{1}{2}$) is the contravariant component of the spinor $\chi_{\frac{1}{2}}$ (see below). The normalization condition

$$\chi_{\frac{1}{2}}^\dagger \chi_{\frac{1}{2}} = 1. \quad (6.2.34)$$

imposes the following restriction

$$\left|a^{\frac{1}{2}}\right|^2 + \left|a^{-\frac{1}{2}}\right|^2 = 1.$$

The coefficient a^m may be interpreted as the probability amplitude that the spin projection on the z -axis for the particle in a given state $\chi_{\frac{1}{2}}$ is equal to m .

(a) *Special cases*

(b) *Spin is directed along the z -axis*

$$a^{\frac{1}{2}} = 1, a^{-\frac{1}{2}} = 0, \quad \chi_{\frac{1}{2}} = \chi_{\frac{1}{2}\frac{1}{2}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (6.2.35)$$

(c) *Spin is directed along the negative z -axis*

$$a^{\frac{1}{2}} = 0, a^{-\frac{1}{2}} = 1, \quad \chi_{\frac{1}{2}} = \chi_{\frac{1}{2}-\frac{1}{2}} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (6.2.36)$$

(d) *Spin is parallel to $\mathbf{n}(\vartheta, \varphi)$*

$$a^{\frac{1}{2}} = \cos \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}}, a^{-\frac{1}{2}} = \sin \frac{\vartheta}{2} e^{i\frac{\varphi}{2}}, \chi_{\frac{1}{2}} = \chi_{\frac{1}{2}\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} \cos \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \\ \sin \frac{\vartheta}{2} e^{i\frac{\varphi}{2}} \end{pmatrix}. \quad (6.2.37)$$

(e) *Spin is antiparallel to $\mathbf{n}(\vartheta, \varphi)$*

$$a^{\frac{1}{2}} = -\sin \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}}, a^{-\frac{1}{2}} = \cos \frac{\vartheta}{2} e^{i\frac{\varphi}{2}}, \chi_{\frac{1}{2}} = \chi_{\frac{1}{2}-\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} -\sin \frac{\vartheta}{2} e^{-i\frac{\varphi}{2}} \\ \cos \frac{\vartheta}{2} e^{i\frac{\varphi}{2}} \end{pmatrix}. \quad (6.2.38)$$

In general, the spin function (6.2.32) describes a spin- $\frac{1}{2}$ particle whose spin is directed along some unit vector $\mathbf{n}$ with cartesian components

$$n_x = 2 \operatorname{Re} \left(a^{\frac{1}{2}*} a^{-\frac{1}{2}} \right), n_y = 2 \operatorname{Im} \left(a^{\frac{1}{2}*} a^{-\frac{1}{2}} \right), n_z = \left| a^{\frac{1}{2}} \right|^2 - \left| a^{-\frac{1}{2}} \right|^2. \quad (6.2.39)$$

The spherical components of this unit vector $\mathbf{n}$ are given by

$$n_\mu = \sqrt{3} \sum_{m,m'} C_{\frac{1}{2}m1\mu}^{\frac{1}{2}m'} a^{m'*} a^m, \quad (6.2.40)$$

or more explicitly

$$n_{+1} = -\sqrt{2} a^{\frac{1}{2}*} a^{-\frac{1}{2}}, n_0 = \left| a^{\frac{1}{2}} \right|^2 - \left| a^{-\frac{1}{2}} \right|^2, n_{-1} = \sqrt{2} a^{-\frac{1}{2}*} a^{\frac{1}{2}}. \quad (6.2.41)$$

The product of the spin functions $\chi_{\frac{1}{2}} \chi_{\frac{1}{2}}^\dagger$ may be expressed in terms of the spin matrices

$$\chi_{\frac{1}{2}} \chi_{\frac{1}{2}}^\dagger = \begin{pmatrix} \left| a^{\frac{1}{2}} \right|^2 & a^{-\frac{1}{2}*} a^{\frac{1}{2}} \\ a^{\frac{1}{2}*} a^{-\frac{1}{2}} & \left| a^{-\frac{1}{2}} \right|^2 \end{pmatrix} = \frac{1}{2} \hat{I} + \mathbf{n} \cdot \hat{\mathbf{S}}, \quad (6.2.42)$$

where $\mathbf{n}$ specifies the spin direction (6.2.39)-(6.2.41).

Matrix elements of the spin operator $\hat{\mathbf{S}}$ may be written in terms of the vector $\mathbf{n}$ as

$$\chi_{\frac{1}{2}}^\dagger \hat{\mathbf{S}} \chi_{\frac{1}{2}} = \frac{1}{2} \mathbf{n}. \quad (6.2.43)$$

The transformation of spin functions under rotations of coordinate system is effected by the rotation operators $\widehat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)$ (Eq. (2.5.32)) or $\widehat{U}^{\frac{1}{2}}(\omega; \theta, \Phi)$ (Eq. (2.5.36)), depending on the choice of the parameters to describe rotations:

$$\chi'_{\frac{1}{2}} = \widehat{D}^{\frac{1}{2}}(\alpha, \beta, \gamma)\chi_{\frac{1}{2}} = \widehat{U}^{\frac{1}{2}}(\omega; \theta, \Phi)\chi_{\frac{1}{2}}. \quad (6.2.44)$$

The spin functions $\chi'_{\frac{1}{2}}$ in the rotated coordinate system may be expressed in terms of the basis spin functions, $\chi_{\frac{1}{2}m}$ or $\chi'_{\frac{1}{2}m}$, referred to the original or rotated coordinate systems, respectively.

$$\chi'_{\frac{1}{2}} = \begin{pmatrix} a'^{\frac{1}{2}} \\ a'^{-\frac{1}{2}} \end{pmatrix} = \sum_{\dot{m}} a'^m \chi_{\frac{1}{2}m} = \sum_m a^m \chi'_{\frac{1}{2}m}. \quad (6.2.45)$$

In this case a^m and a'^m are spinor components in the original and rotated coordinate systems. The relation between the basis functions $\chi'_{\frac{1}{2}m}$ and $\chi_{\frac{1}{2}m}$ is given by Eq. (6.2.15). The transformation properties of a^m are as follows:

$$\begin{aligned} a'^m &= \sum_n D_{mn}^{\frac{1}{2}}(\alpha, \beta, \gamma) a^n, \\ a^n &= \sum_m D_{mn}^{\frac{1}{2}*}(\alpha, \beta, \gamma) a'^m. \end{aligned} \quad \left(\dot{m}, n = \pm \frac{1}{2} \right). \quad (6.2.46)$$

Thus, a^m are contravariant spinor components (see Eq. (4.1.2)).

6.2.7 Polarization Density Matrix

The polarisation density matrix for particles of spin $\frac{1}{2}$ may be written in the form

$$\hat{\rho} = \frac{1}{2} \{ \hat{I} + 2\mathbf{P}\hat{\mathbf{S}} \}, \quad (6.2.47)$$

where the real vector $\mathbf{P}$ is called the polarisation vector. This vector gives the expectation value of the spin operator multiplied by 2,

$$\mathbf{P} = 2\langle \mathbf{S} \rangle = 2 \text{Tr} \{ \hat{\rho} \hat{\mathbf{S}} \}. \quad (6.2.48)$$

The absolute value of the vector $\mathbf{P}$ is called the polarization degree; it ranges from 0 (for unpolarised states) to 1 (for pure, i.e., totally polarized states):

$$0 \leq |P| \leq 1. \quad (6.2.49)$$

The spherical components of the polarization vector are related to the elements of the density matrix by

$$P_{\mu} = \sqrt{3} \sum_{\sigma\sigma'} C_{\frac{1}{2}\sigma\frac{1}{2}\mu}^{\frac{1}{2}\sigma'} \rho_{\sigma\sigma'}, \quad (6.2.50)$$

or in an expanded form

$$P_{+1} = -\sqrt{2}\rho_{-\frac{1}{2}\frac{1}{2}}, P_0 = \rho_{\frac{1}{2}\frac{1}{2}} - \rho_{-\frac{1}{2}-\frac{1}{2}}, P_{-1} = \sqrt{2}\rho_{\frac{1}{2}-\frac{1}{2}}. \quad (6.2.51)$$

Cartesian components of the polarization vector are given by

$$P_x = \rho_{\frac{1}{2}-\frac{1}{2}} + \rho_{-\frac{1}{2}\frac{1}{2}}, P_y = i \left(\rho_{\frac{1}{2}-\frac{1}{2}} - \rho_{-\frac{1}{2}\frac{1}{2}} \right), P_z = \rho_{\frac{1}{2}\frac{1}{2}} - \rho_{-\frac{1}{2}-\frac{1}{2}}. \quad (6.2.52)$$

For a pure state described by a spin function $\chi_{\frac{1}{2}}$ the density matrix $\hat{\rho}$ has the form

$$\hat{\rho}_{\text{pure}} = \chi_{\frac{1}{2}} \chi_{\frac{1}{2}}^{\dagger}, \quad (6.2.53)$$

and the polarization vector $\mathbf{P}$ coincides with the unit vector $\mathbf{n}$ (see Eqs. (6.2.39)-(6.2.41)). For an unpolarized state

$$\hat{\rho}_{\text{unpol}} = \frac{1}{2} \hat{I}. \quad (6.2.54)$$

6.3 Spin functions for $S = 1$

6.3.1 Basis Spin Functions

The basis functions χ_{1m} ($m = \pm 1, 0$) are eigenfunctions of the operators $\widehat{S}^2$ and $\widehat{S}_z$

$$\begin{aligned} \widehat{S}^2 \chi_{1m} &= 2\chi_{1m}, \\ \widehat{S}_z \chi_{1m} &= m\chi_{1m}. \end{aligned} \quad (6.3.1)$$

The function χ_{1m} describes the state of a spin-1 particle with definite spin projection m on the z -axis. The three basis functions $\chi_{1m}(m = \pm 1, 0)$ may be treated as spherical covariant basis vectors $\mathbf{e}_m$ (see Sec. 1.1) written in column form. Instead of χ_{1m} one may use the basis functions $\chi_i(i = x, y, z)$ which are the cartesian basis vectors $\mathbf{e}_i$:

$$\begin{aligned}\chi_{11} &= -\frac{1}{\sqrt{2}}(\chi_x + i\chi_y), & \chi_x &= \frac{1}{\sqrt{2}}(\chi_{1-1} - \chi_{11}) \\ \chi_{10} &= \chi_z, & \chi_y &= \frac{i}{\sqrt{2}}(\chi_{1-1} + \chi_{11}), \\ \chi_{1-1} &= \frac{1}{\sqrt{2}}(\chi_x - i\chi_y), & \chi_z &= \chi_{10}.\end{aligned}\quad (6.3.2)$$

The functions $\chi_{1m}(m = \pm 1, 0)$ as well as $\chi_i(i = x, y, z)$ constitute an orthonormal basis. The orthonormality conditions read

$$\chi_{1m'}^\dagger \chi_{1m} = \delta_{m'm}, \quad \chi_i^\dagger \chi_k = \delta_{ik}. \quad (6.3.3)$$

The completeness condition for basis functions may be written as

$$\sum_{m=\pm 1, 0} \chi_{1m} \chi_{1m}^\dagger = \hat{I}, \quad \sum_{i=x, y, z} \chi_i \chi_i^\dagger = \hat{I}, \quad (6.3.4)$$

where $\hat{I}$ is the unit 3×3 matrix.

To describe the states of particles with spin 1 one may use the spherical basis representation or the cartesian basis representation.

(a) Spherical basis representation

In the spherical basis representation the dependence of the functions $\chi_{1m}(\sigma)$ on the spin variable σ is given by

$$\chi_{1m}(\sigma) = \delta_{m\sigma}. \quad (6.3.5)$$

According to Eq. (6.1.1), the basis spin functions χ_{1m} may be written as

$$\chi_{11} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \chi_{10} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \chi_{1-1} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad (6.3.6)$$

and the Hermitian conjugate functions $\chi_{1m}^\dagger$ read

$$\chi_{11}^\dagger = (1, 0, 0), \quad \chi_{10}^\dagger = (0, 1, 0), \quad \chi_{1-1}^\dagger = (0, 0, 1). \quad (6.3.7)$$

In the spherical basis representation the functions $\chi_i(i = x, y, z)$ have the form

$$\chi_x = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}, \quad \chi_y = \frac{i}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \quad \chi_z = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad (6.3.8)$$

$$\chi_x^\dagger = \frac{1}{\sqrt{2}}(-1, 0, 1), \quad \chi_y^\dagger = -\frac{i}{\sqrt{2}}(1, 0, 1), \quad \chi_z^\dagger = (0, 1, 0). \quad (6.3.9)$$

It follows from (6.3.6), (6.3.8) that in the spherical basis representation the functions χ_{1m} are real,

$$\chi_{1m}^* = \chi_{1m}, \quad (m = \pm 1, 0), \quad (6.3.10)$$

and the functions χ_i satisfy the relations

$$\chi_x^* = \chi_x, \quad \chi_y^* = -\chi_y, \quad \chi_z^* = \chi_z. \quad (6.3.11)$$

Explicit forms of the spin matrices and the polarization operators in the spherical basis representation are given by Eqs. (2.6.9)-(2.6.22).

(b) Cartesian basis representation

In the cartesian basis representation the spin variable σ assumes three possible values, $\sigma = x, y, z$. The dependence of the functions $\chi_i(\sigma)$ on σ is given by

$$\chi_i(\sigma) = \delta_{i\sigma}. \quad (6.3.12)$$

Thus, the basis spin functions $\chi_i (i = x, y, z)$ in the cartesian basis representation may be written in the following form (see also Eq. (1.4.37))

$$\chi_x = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \chi_y = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \chi_z = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad (6.3.13)$$

$$\chi_x^\dagger = (1, 0, 0), \quad \chi_y^\dagger = (0, 1, 0), \quad \chi_z^\dagger = (0, 0, 1). \quad (6.3.14)$$

The basis functions $\chi_{1m} (m = \pm 1, 0)$ read

$$\chi_{11} = -\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \\ 0 \end{pmatrix}, \quad \chi_{10} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad \chi_{1-1} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \\ 0 \end{pmatrix}, \quad (6.3.15)$$

$$\chi_{11}^\dagger = -\frac{1}{\sqrt{2}}(1, -i, 0), \quad \chi_{10}^\dagger = (0, 0, 1), \quad \chi_{1-1}^\dagger = \frac{1}{\sqrt{2}}(1, i, 0). \quad (6.3.16)$$

It follows from Eqs. (6.3.13), (6.3.15) that in the cartesian basis representation the functions χ_i are real,

$$\chi_i^* = \chi_i \quad (i = x, y, z), \quad (6.3.17)$$

and the functions χ_{1m} satisfy the relations

$$\chi_{1m}^* = (-1)^m \chi_{1-m} (m = \pm 1, 0). \quad (6.3.18)$$

The spin matrices and polarization operators in the cartesian basis representation are given by Eqs. (2.6.23) and (2.6.35).

The direct and reverse transformations from a spherical basis to a cartesian one can be performed by the use of the unitary 3×3 matrix U ,

$$\begin{aligned} \chi(\text{ spherical basis }) &= U \chi(\text{ cartesian basis }), \\ \chi(\text{ cartesian basis }) &= U^{-1} \chi(\text{ spherical basis }). \end{aligned} \quad (6.3.19)$$

The explicit form of U is given by Eq. (2.6.37). The expressions below are independent of the representation used unless the contrary is indicated.

6.3.2 Expansions of Products of Spin Functions

Products of the basis functions $\chi_{1m} \chi_{1m'}^\dagger$ and $\chi_i \chi_{k1}^\dagger$ are square 3×3 matrices which may be expanded in terms of the polarization matrices $\hat{I}, \hat{S}_\mu (\mu = \pm 1, 0), \hat{T}_{2M} (M = \pm 2, \pm 1, 0)$ or, equivalently, of the matrices $\hat{I}, \hat{S}_i, \hat{Q}_{ik} (i, k = x, y, z)$ (Sec. 2.6). These expansions are given by

$$\chi_{1m} \chi_{1m'}^\dagger = \frac{1}{3} \delta_{mm'} \hat{I} + \frac{1}{\sqrt{2}} C_{1m'1\mu}^1 \hat{S}_\mu + \sqrt{\frac{5}{3}} C_{1m'2M}^1 \hat{T}_{2M}, \quad (6.3.20)$$

$$\chi_i \chi_k^\dagger = \frac{1}{3} \delta_{ik} \hat{I} + \frac{i}{2} \varepsilon_{ikl} \hat{S}_l - \hat{Q}_{ik}. \quad (6.3.21)$$

Written in component form these expansions yield

$$\begin{aligned}
\chi_{11}\chi_{11}^\dagger &= \frac{1}{3}\hat{I} + \frac{1}{2}\hat{S}_0 + \frac{1}{\sqrt{6}}\hat{T}_{20} = \frac{1}{3}\hat{I} + \frac{1}{2}\hat{S}_z + \frac{1}{2}\hat{Q}_{zz}, \\
\chi_{11}\chi_{10}^\dagger &= -\frac{1}{2}\hat{S}_{+1} - \frac{1}{\sqrt{2}}\hat{T}_{21} = \frac{1}{2\sqrt{2}}\left(\hat{S}_x + i\hat{S}_y + 2\hat{Q}_{xz} + 2i\hat{Q}_{yz}\right), \\
\chi_{11}\chi_{1-1}^\dagger &= \hat{T}_{22} = \frac{1}{2}\left(\hat{Q}_{xx} - \hat{Q}_{yy} + 2i\hat{Q}_{xy}\right), \\
\chi_{10}\chi_{11}^\dagger &= \frac{1}{2}\hat{S}_{-1} + \frac{1}{\sqrt{2}}\hat{T}_{2-1} = \frac{1}{2\sqrt{2}}\left(\hat{S}_x - i\hat{S}_y + 2\hat{Q}_{xz} - 2i\hat{Q}_{yz}\right), \\
\chi_{10}\chi_{10}^\dagger &= \frac{1}{3}\hat{I} - \sqrt{\frac{2}{3}}\hat{T}_{20} = \frac{1}{3}\hat{I} - \hat{Q}_{zz}, \\
\chi_{10}\chi_{1-1}^\dagger &= -\frac{1}{2}\hat{S}_{+1} + \frac{1}{\sqrt{2}}\hat{T}_{21} = \frac{1}{2\sqrt{2}}\left(\hat{S}_x + i\hat{S}_y - 2\hat{Q}_{xz} - 2i\hat{Q}_{yz}\right), \\
\chi_{1-1}\chi_{11}^\dagger &= \hat{T}_{2-2} = \frac{1}{2}\left(\hat{Q}_{xx} - \hat{Q}_{yy} - 2i\hat{Q}_{xy}\right), \\
\chi_{1-1}\chi_{10}^\dagger &= \frac{1}{2}\hat{S}_{-1} - \frac{1}{\sqrt{2}}\hat{T}_{2-1} = \frac{1}{2\sqrt{2}}\left(\hat{S}_x - i\hat{S}_y - 2\hat{Q}_{xz} + 2i\hat{Q}_{yz}\right), \\
\chi_{1-1}\chi_{1-1}^\dagger &= \frac{1}{3}\hat{I} - \frac{1}{2}\hat{S}_0 + \frac{1}{\sqrt{6}}\hat{T}_{20} = \frac{1}{3}\hat{I} - \frac{1}{2}\hat{S}_z + \frac{1}{2}\hat{Q}_{zz}.
\end{aligned} \tag{6.3.22}$$

$$\begin{aligned}
\chi_x\chi_x^\dagger &= \frac{1}{3}\hat{I} - \frac{1}{2}\hat{T}_{2-2} - \frac{1}{2}\hat{T}_{22} + \frac{1}{\sqrt{6}}\hat{T}_{20} = \frac{1}{3}\hat{I} - \hat{Q}_{xx}, \\
\chi_x\chi_y^\dagger &= \frac{i}{2}\left(\hat{S}_0 - \hat{T}_{2-2} + \hat{T}_{22}\right) = \frac{i}{2}\hat{S}_z - \hat{Q}_{xy}, \\
\chi_x\chi_z^\dagger &= \frac{1}{2\sqrt{2}}\left(\hat{S}_{-1} + \hat{S}_{+1}\right) - \frac{1}{2}\left(\hat{T}_{2-1} - \hat{T}_{21}\right) = -\frac{i}{2}\hat{S}_y - \hat{Q}_{xz}, \\
\chi_y\chi_x^\dagger &= -\frac{i}{2}\left(\hat{S}_0 + \hat{T}_{2-2} - \hat{T}_{22}\right) = -\frac{i}{2}\hat{S}_z - \hat{Q}_{yx}, \\
\chi_y\chi_y^\dagger &= \frac{1}{3}\hat{I} + \frac{1}{2}\hat{T}_{2-2} + \frac{1}{2}\hat{T}_{22} + \frac{1}{\sqrt{6}}\hat{T}_{20} = \frac{1}{3}\hat{I} - \hat{Q}_{yy}, \\
\chi_y\chi_z^\dagger &= \frac{i}{2\sqrt{2}}\left(\hat{S}_{-1} - \hat{S}_{+1}\right) - \frac{i}{2}\left(\hat{T}_{2-1} + \hat{T}_{21}\right) = \frac{i}{2}\hat{S}_x - \hat{Q}_{yz}, \\
\chi_z\chi_x^\dagger &= -\frac{1}{2\sqrt{2}}\left(\hat{S}_{-1} + \hat{S}_{+1}\right) - \frac{1}{2}\left(\hat{T}_{2-1} - \hat{T}_{21}\right) = \frac{i}{2}\hat{S}_y - \hat{Q}_{zx}, \\
\chi_z\chi_y^\dagger &= -\frac{i}{2\sqrt{2}}\left(\hat{S}_{-1} - \hat{S}_{+1}\right) - \frac{i}{2}\left(\hat{T}_{2-1} + \hat{T}_{21}\right) = -\frac{i}{2}\hat{S}_x - \hat{Q}_{zy}, \\
\chi_z\chi_z^\dagger &= \frac{1}{3}\hat{I} - \sqrt{\frac{2}{3}}\hat{T}_{20} = \frac{1}{3}\hat{I} - \hat{Q}_{zz}.
\end{aligned} \tag{6.3.23}$$

6.3.3 Action of Spin Operators on Basis Functions

Spherical components of the spin operator $\hat{S}_\mu(\mu = \pm 1, 0)$ and its cartesian components $\hat{S}_i(i = x, y, z)$ act on the basis functions as follows

$$\begin{aligned}
\hat{S}_\mu\chi_{1m} &= \sqrt{2}C_{1m1\mu}^{1m'}\chi_{1m'}, \\
\hat{S}_i\chi_k &= i\varepsilon_{ikl}\chi_l.
\end{aligned} \tag{6.3.24}$$

In a more detailed form

$$\begin{aligned}\hat{S}_{+1}\chi_{11} &= 0, & \hat{S}_{+1}\chi_{10} &= -\chi_{11}, & \hat{S}_{+1}\chi_{1-1} &= -\chi_{10}, \\ \hat{S}_0\chi_{11} &= \chi_{11}, & \hat{S}_0\chi_{10} &= 0, & \hat{S}_0\chi_{1-1} &= -\chi_{1-1}, \\ \hat{S}_{-1}\chi_{11} &= \chi_{10}, & \hat{S}_{-1}\chi_{10} &= \chi_{1-1}, & \hat{S}_{-1}\chi_{1-1} &= 0.\end{aligned}\tag{6.3.25}$$

$$\begin{aligned}\hat{S}_{+1}\chi_x &= -\frac{1}{\sqrt{2}}\chi_z, & \hat{S}_{+1}\chi_y &= -\frac{i}{\sqrt{2}}\chi_z, & \hat{S}_{+1}\chi_z &= \frac{1}{\sqrt{2}}(\chi_x + i\chi_y), \\ \hat{S}_0\chi_x &= i\chi_y, & \hat{S}_0\chi_y &= -i\chi_x, & \hat{S}_0\chi_z &= 0, \\ \hat{S}_{-1}\chi_x &= -\frac{1}{\sqrt{2}}\chi_z, & \hat{S}_{-1}\chi_y &= \frac{i}{\sqrt{2}}\chi_z, & \hat{S}_{-1}\chi_z &= \frac{1}{\sqrt{2}}(\chi_x - i\chi_y).\end{aligned}\tag{6.3.26}$$

$$\begin{aligned}\hat{S}_x\chi_{11} &= \frac{1}{\sqrt{2}}\chi_{10}, & \hat{S}_x\chi_{10} &= \frac{1}{\sqrt{2}}(\chi_{1-1} + \chi_{11}), & \hat{S}_x\chi_{1-1} &= \frac{1}{\sqrt{2}}\chi_{10}, \\ \hat{S}_y\chi_{11} &= \frac{i}{\sqrt{2}}\chi_{10}, & \hat{S}_y\chi_{10} &= \frac{i}{\sqrt{2}}(\chi_{1-1} - \chi_{11}), & \hat{S}_y\chi_{1-1} &= -\frac{i}{\sqrt{2}}\chi_{10}, \\ \hat{S}_z\chi_{11} &= \chi_{11}, & \hat{S}_z\chi_{10} &= 0, & \hat{S}_z\chi_{1-1} &= -\chi_{1-1}.\end{aligned}\tag{6.3.27}$$

$$\begin{aligned}\hat{S}_x\chi_x &= 0, & \hat{S}_x\chi_y &= i\chi_z, & \hat{S}_x\chi_z &= -i\chi_y, \\ \hat{S}_y\chi_x &= -i\chi_z, & \hat{S}_y\chi_y &= 0, & \hat{S}_y\chi_z &= i\chi_x, \\ \hat{S}_z\chi_x &= i\chi_y, & \hat{S}_z\chi_y &= -i\chi_x, & \hat{S}_z\chi_z &= 0.\end{aligned}\tag{6.3.28}$$

6.3.4 Action of Quadrupole Operators on Basis Functions

The operators $\hat{T}_{2M}(M = \pm 2, \pm 1, 0)$ and $\hat{Q}_{ik}(i, k = x, y, z)$ act on basis functions in accordance with

$$\begin{aligned}\hat{T}_{2M}\chi_{1m} &= \sqrt{\frac{5}{3}}C_{1m2M}^{1m'}\chi_{1m'}, \\ \hat{Q}_{ik}\chi_l &= \frac{1}{2}\left(\frac{2}{3}\delta_{ik}\chi_l - \delta_{il}\chi_k - \delta_{kl}\chi_i\right).\end{aligned}\tag{6.3.29}$$

In detailed form we have

$$\begin{aligned}\hat{T}_{22}\chi_{11} &= 0, & \hat{T}_{22}\chi_{10} &= 0, & \hat{T}_{22}\chi_{1-1} &= \chi_{11}, \\ \hat{T}_{21}\chi_{11} &= 0, & \hat{T}_{21}\chi_{10} &= -\frac{1}{\sqrt{2}}\chi_{11}, & \hat{T}_{21}\chi_{1-1} &= \frac{1}{\sqrt{2}}\chi_{10}, \\ \hat{T}_{20}\chi_{11} &= \frac{1}{\sqrt{6}}\chi_{11}, & \hat{T}_{20}\chi_{10} &= -\sqrt{\frac{2}{3}}\chi_{10}, & \hat{T}_{20}\chi_{1-1} &= \frac{1}{\sqrt{6}}\chi_{1-1}, \\ \hat{T}_{2-1}\chi_{11} &= \frac{1}{\sqrt{2}}\chi_{10}, & \hat{T}_{2-1}\chi_{10} &= -\frac{1}{\sqrt{2}}\chi_{1-1}, & \hat{T}_{2-1}\chi_{1-1} &= 0, \\ \hat{T}_{2-2}\chi_{11} &= \chi_{1-1}, & \hat{T}_{2-2}\chi_{10} &= 0, & \hat{T}_{2-2}\chi_{1-1} &= 0.\end{aligned}\tag{6.3.30}$$

$$\begin{aligned}\hat{T}_{22}\chi_x &= -\frac{1}{2}(\chi_x + i\chi_y), & \hat{T}_{22}\chi_y &= \frac{1}{2}(\chi_y - i\chi_x), & \hat{T}_{22}\chi_z &= 0, \\ \hat{T}_{21}\chi_x &= \frac{1}{2}\chi_z, & \hat{T}_{21}\chi_y &= \frac{i}{2}\chi_z, & \hat{T}_{21}\chi_z &= \frac{1}{2}(\chi_x + i\chi_y), \\ \hat{T}_{20}\chi_x &= \frac{1}{\sqrt{6}}\chi_x, & \hat{T}_{20}\chi_y &= \frac{1}{\sqrt{6}}\chi_y, & \hat{T}_{20}\chi_z &= -\sqrt{\frac{2}{3}}\chi_z, \\ \hat{T}_{2-1}\chi_x &= -\frac{1}{2}\chi_z, & \hat{T}_{2-1}\chi_y &= \frac{i}{2}\chi_z, & \hat{T}_{2-1}\chi_z &= -\frac{1}{2}(\chi_x - i\chi_y), \\ \hat{T}_{2-2}\chi_x &= -\frac{1}{2}(\chi_x - i\chi_y), & \hat{T}_{2-2}\chi_y &= \frac{1}{2}(\chi_y + i\chi_x), & \hat{T}_{2-2}\chi_z &= 0.\end{aligned}\tag{6.3.31}$$

$$\begin{aligned}\hat{Q}_{xx}\chi_{11} &= \frac{1}{2}\chi_{1-1} - \frac{1}{6}\chi_{11}, & \hat{Q}_{xx}\chi_{10} &= \frac{1}{3}\chi_{10}, & \hat{Q}_{xx}\chi_{1-1} &= \frac{1}{2}\chi_{11} - \frac{1}{6}\chi_{1-1}, \\ \hat{Q}_{yy}\chi_{11} &= -\frac{1}{2}\chi_{1-1} - \frac{1}{6}\chi_{11}, & \hat{Q}_{yy}\chi_{10} &= \frac{1}{3}\chi_{10}, & \hat{Q}_{yy}\chi_{1-1} &= -\frac{1}{2}\chi_{11} - \frac{1}{6}\chi_{1-1}, \\ \hat{Q}_{zz}\chi_{11} &= \frac{1}{3}\chi_{11}, & \hat{Q}_{zz}\chi_{10} &= -\frac{2}{3}\chi_{10}, & \hat{Q}_{zz}\chi_{1-1} &= \frac{1}{3}\chi_{1-1}, \\ \hat{Q}_{xy}\chi_{11} &= \frac{i}{2}\chi_{1-1}, & \hat{Q}_{xy}\chi_{10} &= 0, & \hat{Q}_{xy}\chi_{1-1} &= -\frac{i}{2}\chi_{11}, \\ \hat{Q}_{xz}\chi_{11} &= \frac{1}{2\sqrt{2}}\chi_{10}, & \hat{Q}_{xz}\chi_{10} &= \frac{1}{2\sqrt{2}}(\chi_{11} - \chi_{1-1}), & \hat{Q}_{xz}\chi_{1-1} &= -\frac{1}{2\sqrt{2}}\chi_{10}, \\ \hat{Q}_{yz}\chi_{11} &= \frac{i}{2\sqrt{2}}\chi_{10}, & \hat{Q}_{yz}\chi_{10} &= -\frac{i}{2\sqrt{2}}(\chi_{11} + \chi_{1-1}), & \hat{Q}_{yz}\chi_{1-1} &= \frac{i}{2\sqrt{2}}\chi_{10}.\end{aligned}\tag{6.3.32}$$

$$\begin{aligned}\hat{Q}_{xx}\chi_x &= -\frac{2}{3}\chi_x, & \hat{Q}_{xx}\chi_y &= \frac{1}{3}\chi_y, & \hat{Q}_{xx}\chi_z &= \frac{1}{3}\chi_z, \\ \hat{Q}_{yy}\chi_x &= \frac{1}{3}\chi_x, & \hat{Q}_{yy}\chi_y &= -\frac{2}{3}\chi_y, & \hat{Q}_{yy}\chi_z &= \frac{1}{3}\chi_z, \\ \hat{Q}_{zz}\chi_x &= \frac{1}{3}\chi_x, & \hat{Q}_{zz}\chi_y &= \frac{1}{3}\chi_y, & \hat{Q}_{zz}\chi_z &= -\frac{2}{3}\chi_z, \\ \hat{Q}_{xy}\chi_x &= -\frac{1}{2}\chi_y, & \hat{Q}_{xy}\chi_y &= -\frac{1}{2}\chi_x, & \hat{Q}_{xy}\chi_z &= 0, \\ \hat{Q}_{xz}\chi_x &= -\frac{1}{2}\chi_z, & \hat{Q}_{xz}\chi_y &= 0, & \hat{Q}_{xz}\chi_z &= -\frac{1}{2}\chi_x, \\ \hat{Q}_{yz}\chi_x &= 0, & \hat{Q}_{yz}\chi_y &= -\frac{1}{2}\chi_z, & \hat{Q}_{yz}\chi_z &= -\frac{1}{2}\chi_y.\end{aligned}\tag{6.3.33}$$

6.3.5 Transformation of Basis Functions Under Rotations of the Coordinate Systems

The spin functions $\chi_{1m}(m = \pm 1, 0)$ and $\chi_i(i = x, y, z)$ constitute the covariant spherical basis and cartesian basis, respectively. Under rotations they transform through the rotation operator $\hat{D}^1(\alpha, \beta, \gamma)$ (Eqs. (2.6.75), (2.6.78)) if rotations are specified by the Euler angles α, β, γ or through the rotation operator $\hat{U}^1(\omega; \Theta, \Phi)$ (Eqs. (2.6.79)-(2.6.83)) if rotations are described by the rotation angle ω and the rotation axis $n(\Theta, \Phi)$.

$$\begin{aligned}\chi'_{1m'} &= \hat{D}^1(\alpha, \beta, \gamma)\chi_{1m} = \sum_{m=\pm 1, 0} D^1_{mm'}(\alpha, \beta, \gamma)\chi_{1m}, \\ \chi'_i &= \hat{D}^1(\alpha, \beta, \gamma)\chi_i = \sum_{k=x, y, z} a_{ki}\chi_k.\end{aligned}\quad (6.3.34)$$

Here $D^1_{mm'}(\alpha, \beta, \gamma)$ are the Wigner D -functions (Chap. 4) and a_{ik} are elements of the rotation matrix (Sec. 1.4.6).

The functions $\chi'_{1m'}$ describe quantum states in which a particle of spin 1 has the spin projection m' on the new z' -axis. These functions are eigenfunctions of the operators $\hat{\mathbf{S}}'^2$ and $\hat{S}'_z$ where $\hat{\mathbf{S}}'$ is the spin operator in the rotated coordinate system (Eqs. (2.6.85) and (2.6.87))

$$\begin{aligned}\hat{\mathbf{S}}'^2\chi'_{1m'} &= 2\chi'_{1m'}, \\ \hat{S}'_z\chi'_{1m'} &= m'\chi'_{1m'}.\end{aligned}\quad (6.3.35)$$

In the spherical basis representation the functions $\chi'_{1m'}$ have the form

$$\chi'_{11} = \begin{pmatrix} \frac{1+\cos\beta}{2}e^{-i(\alpha+\gamma)} \\ \frac{\sin\beta}{\sqrt{2}}e^{-i\gamma} \\ \frac{1-\cos\beta}{2}e^{i(\alpha-\gamma)} \end{pmatrix}, \quad \chi'_{10} = \begin{pmatrix} -\frac{\sin\beta}{\sqrt{2}}e^{-i\alpha} \\ \cos\beta \\ \frac{\sin\beta}{\sqrt{2}}e^{i\alpha} \end{pmatrix}, \quad \chi'_{1-1} = \begin{pmatrix} \frac{1-\cos\beta}{2}e^{-i(\alpha-\gamma)} \\ -\frac{\sin\beta}{\sqrt{2}}e^{i\gamma} \\ \frac{1+\cos\beta}{2}e^{i(\alpha+\gamma)} \end{pmatrix}, \quad (6.3.36)$$

and the functions χ'_i are given by

$$\chi'_x = \begin{pmatrix} -\frac{\cos\beta\cos\gamma-i\sin\gamma}{\sqrt{2}}e^{-i\alpha} \\ -\sin\beta\cos\gamma \\ \frac{\cos\beta\cos\gamma+i\sin\gamma}{\sqrt{2}}e^{i\alpha} \end{pmatrix}, \quad \chi'_y = \begin{pmatrix} \frac{\cos\beta\sin\gamma+i\cos\gamma}{\sqrt{2}}e^{-i\alpha} \\ \sin\beta\sin\gamma \\ -\frac{\cos\beta\sin\gamma-i\cos\gamma}{\sqrt{2}}e^{i\alpha} \end{pmatrix}, \quad \chi'_z = \begin{pmatrix} -\frac{\sin\beta}{\sqrt{2}}e^{-i\alpha} \\ \cos\beta \\ \frac{\sin\beta}{\sqrt{2}}e^{i\alpha} \end{pmatrix}. \quad (6.3.37)$$

In the cartesian basis representation these functions read

$$\begin{aligned}\chi'_{11} &= \begin{pmatrix} -\frac{\cos\beta\cos\alpha-i\sin\alpha}{\sqrt{2}}e^{-i\gamma} \\ -\frac{\cos\beta\sin\alpha+i\cos\alpha}{\sqrt{2}}e^{-i\gamma} \\ \frac{\sin\beta}{\sqrt{2}}e^{-i\gamma} \end{pmatrix}, \quad \chi'_{10} = \begin{pmatrix} \sin\beta\cos\alpha \\ \sin\beta\sin\alpha \\ \cos\beta \end{pmatrix}, \quad \chi'_{1-1} = \begin{pmatrix} \frac{\cos\beta\cos\alpha+i\sin\alpha}{\sqrt{2}}e^{i\gamma} \\ \frac{\cos\beta\sin\alpha-i\cos\alpha}{\sqrt{2}}e^{i\gamma} \\ -\frac{\sin\beta}{\sqrt{2}}e^{i\gamma} \end{pmatrix}, \\ \chi'_x &= \begin{pmatrix} \cos\beta\cos\alpha\cos\gamma - \sin\alpha\sin\gamma \\ \cos\beta\sin\alpha\cos\gamma + \cos\alpha\sin\gamma \\ -\sin\beta\cos\gamma \end{pmatrix}, \quad \chi'_y = \begin{pmatrix} -\cos\beta\cos\alpha\sin\gamma - \sin\alpha\cos\gamma \\ -\cos\beta\sin\alpha\sin\gamma + \cos\alpha\cos\gamma \\ \sin\beta\sin\gamma \end{pmatrix}, \quad \chi'_z = \begin{pmatrix} \sin\beta\cos\alpha \\ \sin\beta\sin\alpha \\ \cos\beta \end{pmatrix}.\end{aligned}\quad (6.3.38)$$

The expansions for the basis functions in the rotated coordinate system in terms of the angles ω, Θ, Φ may be obtained by the use of the rotation operator $\hat{U}^1(\omega; \Theta, \Phi)$ (Eqs. (2.6.80)-(2.6.81)).

6.3.6 Helicity Basis Functions for $S = 1$

The helicity basis functions $\chi_{1\lambda}(\vartheta, \varphi)(\lambda = \pm 1, 0)$ describe the states in which the spin projection on the linear momentum direction $\mathbf{n}(\vartheta, \varphi) \equiv \mathbf{p}/|\mathbf{p}|$ is equal to λ . The functions $\chi_{1\lambda}(\vartheta, \varphi)$ are eigenfunctions of the operators $\hat{\mathbf{S}}^2$ and $\hat{\mathbf{S}} \cdot \mathbf{n}$

$$\begin{aligned}\hat{\mathbf{S}}^2\chi_{1\lambda}(\vartheta, \varphi) &= 2\chi_{1\lambda}(\vartheta, \varphi), \\ \hat{\mathbf{S}} \cdot \mathbf{n}\chi_{1\lambda}(\vartheta, \varphi) &= \lambda\chi_{1\lambda}(\vartheta, \varphi).\end{aligned}\quad (6.3.40)$$

According to Eq. (6.1.20), the helicity basis functions may be derived from the functions χ_{1m} by a coordinate rotation

$$\begin{aligned}\chi_{1\lambda}(\vartheta, \varphi) &= \sum_m D^1_{m\lambda}(\varphi, \vartheta, 0)\chi_{1m}, \\ \chi_{1m} &= \sum_{\lambda} D^1_{-\lambda-m}(0, \vartheta, \varphi)\chi_{1\lambda}(\vartheta, \varphi).\end{aligned}\quad (6.3.41)$$

For Hermitian adjoint functions Eqs. (6.3.41) assume the form

$$\begin{aligned}\chi_{1\lambda}^\dagger(\vartheta, \varphi) &= \sum_m (-1)^{\lambda-m} D_{-m-\lambda}^1(\varphi, \vartheta, 0) \chi_{1m}^\dagger, \\ \chi_{1m}^\dagger &= \sum_\lambda (-1)^{\lambda-m} D_{\lambda m}^1(0, \vartheta, \varphi) \chi_{1\lambda}^\dagger(\vartheta, \varphi).\end{aligned}\quad (6.3.42)$$

The helicity functions $\chi_{1\lambda}(\vartheta, \varphi)$ may be treated as the covariant helicity basis vectors $\mathbf{e}'_\lambda$ (see Sec. 1.1.4) written as column matrices

$$\chi_{1\lambda}(\vartheta, \varphi) = \mathbf{e}'_\lambda \quad (\lambda = \pm 1, 0). \quad (6.3.43)$$

One can construct the following linear combinations $\chi_i(\vartheta, \varphi)$ ($i = x, y, z$) of the helicity functions $\chi_{1\lambda}(\vartheta, \varphi)$:

$$\begin{aligned}\chi_x(\vartheta, \varphi) &= \frac{1}{\sqrt{2}} \{ \chi_{1-1}(\vartheta, \varphi) - \chi_{11}(\vartheta, \varphi) \}, \\ \chi_y(\vartheta, \varphi) &= \frac{i}{\sqrt{2}} \{ \chi_{1-1}(\vartheta, \varphi) + \chi_{11}(\vartheta, \varphi) \}, \\ \chi_z(\vartheta, \varphi) &= \chi_{10}(\vartheta, \varphi).\end{aligned}\quad (6.3.44)$$

The functions $\chi_i(\vartheta, \varphi)$ ($i = x, y, z$) coincide with the polar basis vectors (see Sec. 1.1.3):

$$\chi_x(\vartheta, \varphi) \equiv \mathbf{e}_\vartheta, \quad \chi_y(\vartheta, \varphi) \equiv \mathbf{e}_\varphi, \quad \chi_z(\vartheta, \varphi) \equiv \mathbf{e}_p. \quad (6.3.45)$$

It should be emphasized that the functions $\chi_i(\vartheta, \varphi)$ describe the states with no definite helicity. These functions are related to the basis functions χ_k ($k = x, y, z$) (Sec. 6.3.1) by

$$\begin{aligned}\chi_i(\vartheta, \varphi) &= \sum_k a_{ki}(\varphi, \vartheta, 0) \chi_k, \\ \chi_k &= \sum_i a_{ik}(\varphi, \vartheta, 0) \chi_i(\vartheta, \varphi),\end{aligned}\quad (6.3.46)$$

where $a_{ik}(\alpha, \beta, \gamma)$ are elements of the rotation matrix given by Eqs. (1.4.54). The relations between the Hermitian conjugate functions $\chi_i^\dagger(\vartheta, \varphi)$ and $\chi_k^\dagger$ are also given by Eq. (6.3.46) because the elements a_{ik} are real.

Three helicity functions $\chi_{1\lambda}(\vartheta, \varphi)$ with $\lambda = \pm 1, 0$ or, equivalently, three functions $\chi_i(\vartheta, \varphi)$ with $i = x, y, z$ constitute an orthonormalized basis. The orthogonality and normalization conditions have the form

$$\begin{aligned}\chi_{1\lambda}^\dagger(\vartheta, \varphi) \chi_{1\lambda'}(\vartheta, \varphi) &= \delta_{\lambda\lambda'}, \\ \chi_i^\dagger(\vartheta, \varphi) \chi_k(\vartheta, \varphi) &= \delta_{ik}.\end{aligned}\quad (6.3.47)$$

The completeness conditions are

$$\begin{aligned}\sum_{\lambda=\pm 1, 0} \chi_{1\lambda}(\vartheta, \varphi) \chi_{1\lambda}^\dagger(\vartheta, \varphi) &= \widehat{I}, \\ \sum_{i=x, y, z} \chi_i(\vartheta, \varphi) \chi_i^\dagger(\vartheta, \varphi) &= \widehat{I}.\end{aligned}\quad (6.3.48)$$

Explicit form of the helicity functions $\chi_{1\lambda}(\vartheta, \varphi)$ and the functions $\chi_i(\vartheta, \varphi)$

(a) *Spherical basis representation*

$$\chi_{11}(\vartheta, \varphi) = \begin{pmatrix} \frac{1+\cos\vartheta}{2} e^{-i\varphi} \\ \frac{\sin\vartheta}{\sqrt{2}} \\ \frac{1-\cos\vartheta}{2} e^{i\varphi} \end{pmatrix}, \quad \chi_{10}(\vartheta, \varphi) = \begin{pmatrix} -\frac{\sin\vartheta}{\sqrt{2}} e^{-i\varphi} \\ \cos\vartheta \\ \frac{\sin\vartheta}{\sqrt{2}} e^{i\varphi} \end{pmatrix}, \quad \chi_{1-1}(\vartheta, \varphi) = \begin{pmatrix} \frac{1-\cos\vartheta}{2} e^{-i\varphi} \\ -\frac{\sin\vartheta}{\sqrt{2}} \\ \frac{1+\cos\vartheta}{2} e^{i\varphi} \end{pmatrix}, \quad (6.3.49)$$

$$\chi_x(\vartheta, \varphi) = \begin{pmatrix} -\frac{\cos\vartheta}{\sqrt{2}} e^{-i\varphi} \\ -\sin\vartheta \\ \frac{\cos\vartheta}{\sqrt{2}} e^{i\varphi} \end{pmatrix}, \quad \chi_y(\vartheta, \varphi) = \begin{pmatrix} \frac{i}{\sqrt{2}} e^{-i\varphi} \\ 0 \\ \frac{i}{\sqrt{2}} e^{i\varphi} \end{pmatrix}, \quad \chi_z(\vartheta, \varphi) = \begin{pmatrix} -\frac{\sin\vartheta}{\sqrt{2}} e^{-i\varphi} \\ \cos\vartheta \\ \frac{\sin\vartheta}{\sqrt{2}} e^{i\varphi} \end{pmatrix}. \quad (6.3.50)$$

(b) Cartesian basis representation

$$\begin{aligned}\chi_{11}(\vartheta, \varphi) &= -\frac{1}{\sqrt{2}} \begin{pmatrix} \cos \vartheta \cos \varphi - i \sin \varphi \\ \cos \vartheta \sin \varphi + i \cos \varphi \\ -\sin \vartheta \end{pmatrix}, \quad \chi_{10}(\vartheta, \varphi) = \begin{pmatrix} \sin \vartheta \cos \varphi \\ \sin \vartheta \sin \varphi \\ \cos \vartheta \end{pmatrix}, \\ \chi_{1-1}(\vartheta, \varphi) &= \frac{1}{\sqrt{2}} \begin{pmatrix} \cos \vartheta \cos \varphi + i \sin \varphi \\ \cos \vartheta \sin \varphi - i \cos \varphi \\ -\sin \vartheta \end{pmatrix},\end{aligned}\quad (6.3.51)$$

$$\chi_x(\vartheta, \varphi) = \begin{pmatrix} \cos \vartheta \cos \varphi \\ \cos \vartheta \sin \varphi \\ -\sin \vartheta \end{pmatrix}, \quad \chi_y(\vartheta, \varphi) = \begin{pmatrix} -\sin \varphi \\ \cos \varphi \\ 0 \end{pmatrix}, \quad \chi_z(\vartheta, \varphi) = \begin{pmatrix} \sin \vartheta \cos \varphi \\ \sin \vartheta \sin \varphi \\ \cos \vartheta \end{pmatrix}. \quad (6.3.52)$$

In the cartesian basis representation these functions satisfy the relations

$$\begin{aligned}\chi_{1\lambda}^*(\vartheta, \varphi) &= (-1)^\lambda \chi_{1-\lambda}(\vartheta, \varphi), \quad (\lambda = \pm 1, 0), \\ \chi_i^*(\vartheta, \varphi) &= \chi_i(\vartheta, \varphi), \quad (i = x, y, z).\end{aligned}\quad (6.3.53)$$

The expansions of products of the helicity basis functions in series of spin matrices and quadrupole operators (Sec. 2.6) are given by

$$\begin{aligned}\chi_{1\lambda}(\vartheta, \varphi) \chi_{1\lambda'}^\dagger(\vartheta, \varphi) &= \frac{1}{3} \delta_{\lambda\lambda'} \hat{I} + \frac{1}{\sqrt{2}} C_{1\lambda' 1 \nu}^{1\lambda} \sum_{\mu=-1}^1 D_{\mu\nu}^1(\varphi, \vartheta, 0) \hat{S}_\mu, \\ &+ \sqrt{\frac{5}{3}} C_{1\lambda' 2 N}^{1\lambda} \sum_{M=-2}^2 D_{MN}^2(\varphi, \vartheta, 0) \hat{T}_{2M}.\end{aligned}\quad (6.3.54)$$

Analogous expansions for $\chi_i(\vartheta, \varphi)$ can be written as

$$\chi_i(\vartheta, \varphi) \chi_k^\dagger(\vartheta, \varphi) = \frac{1}{3} \delta_{ik} \hat{I} + \frac{i}{2} \varepsilon_{ikl} \sum_{m=x,y,z} a_{ml}(\varphi, \vartheta, 0) \hat{S}_m - \sum_{m,n=x,y,z} a_{mi}(\varphi, \vartheta, 0) a_{nk}(\varphi, \vartheta, 0) \hat{Q}_{mn}, \quad (6.3.55)$$

where $a_{ml}(\alpha, \beta, \gamma)$ are elements of the rotation matrix (Eq. (1.4.54)).

Matrix elements of the spin operator between the helicity states $\chi_{1\lambda}(\vartheta, \varphi)$ or between the states $\chi_i(\vartheta, \varphi)$ are

$$\chi_{1\lambda}^\dagger(\vartheta, \varphi) \hat{S}_\mu \chi_{1\lambda'}(\vartheta, \varphi) = (-1)^{\mu+\nu} \sqrt{2} C_{1\lambda' 1 \nu}^{1\lambda} D_{-\mu-\nu}^1(\varphi, \vartheta, 0), \quad (6.3.56)$$

$$\chi_i^\dagger(\vartheta, \varphi) \hat{S}_l \chi_k(\vartheta, \varphi) = -i \sum_m \varepsilon_{ikm} a_{lm}(\varphi, \vartheta, 0). \quad (6.3.57)$$

In particular, the expectation value of the spin operator $\hat{\mathbf{S}}$ in a state with the helicity λ is given by

$$\chi_{1\lambda}^\dagger(\vartheta, \varphi) \hat{\mathbf{S}} \chi_{1\lambda}(\vartheta, \varphi) = \lambda \mathbf{n}(\vartheta, \varphi), \quad (\lambda = \pm 1, 0), \quad (6.3.58)$$

The expectation value of the operator $\hat{\mathbf{S}}$ in the states $\chi_i(\vartheta, \varphi)$ is zero.

$$\chi_i^\dagger(\vartheta, \varphi) \hat{\mathbf{S}} \chi_i(\vartheta, \varphi) = 0, \quad (i = x, y, z). \quad (6.3.59)$$

The evaluation of matrix elements of the quadrupole operators $\hat{T}_{2M}$ ($M = \pm 2, \pm 1, 0$) and $\hat{Q}_{ik}$ ($i, k = x, y, z$) gives

$$\chi_{1\lambda}^\dagger(\vartheta, \varphi) \hat{T}_{2M} \chi_{1\lambda'}(\vartheta, \varphi) = (-1)^{M+N} \sqrt{\frac{5}{3}} C_{1\lambda' 2 N}^{1\lambda} D_{-M-N}^2(\varphi, \vartheta, 0), \quad (6.3.60)$$

$$\chi_i^\dagger(\vartheta, \varphi) \hat{Q}_{lm} \chi_k(\vartheta, \varphi) = -\frac{1}{2} \left\{ a_{li}(\varphi, \vartheta, 0) a_{mk}(\varphi, \vartheta, 0) + a_{lk}(\varphi, \vartheta, 0) a_{mi}(\varphi, \vartheta, 0) - \frac{2}{3} \delta_{ik} \delta_{lm} \right\}. \quad (6.3.61)$$

In particular, the expectation values of the quadrupole operators $\hat{T}_{2M}$ in states with helicity λ are equal to

$$\begin{aligned}\chi_{1\lambda}^\dagger(\vartheta, \varphi) \hat{T}_{2M} \chi_{1\lambda}(\vartheta, \varphi) &= \frac{(-1)^{1+\lambda}}{1+|\lambda|} \sqrt{\frac{8\pi}{15}} Y_{2M}(\vartheta, \varphi), \\ (\lambda = \pm 1, 0; M = \pm 2, \pm 1, 0).\end{aligned}\quad (6.3.62)$$

6.3.7 General Spin Functions for $S = 1$

An arbitrary spin function χ_1 of a particle of spin 1 may be expanded in terms of basis spin functions, i.e., written in the following form:

$$\chi_1 = \sum_{m=\pm 1,0} a^m \chi_{1m} = \sum_{m=\pm 1,0} (-1)^m a_{-m} \chi_{1m} = \sum_{i=x,y,z} a_i \chi_i. \quad (6.3.63)$$

The expansion of the Hermitian conjugate function $\chi_1^\dagger$ is given by

$$\chi_1^\dagger = \sum_{m=\pm 1,0} (a^m)^* \chi_{1m}^\dagger = \sum_{m=\pm 1,0} (-1)^m (a_{-m})^* \chi_{1m}^\dagger = \sum_{i=x,y,z} a_i^* \chi_i^\dagger. \quad (6.3.64)$$

In a spherical basis the function χ_1 has the form

$$\chi_1 = \begin{pmatrix} a^{+1} \\ a^0 \\ a^{-1} \end{pmatrix} = \begin{pmatrix} -a_{-1} \\ a_0 \\ -a_{+1} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\sqrt{2}}(a_x - ia_y) \\ a_z \\ \frac{1}{\sqrt{2}}(a_x + ia_y) \end{pmatrix}, \quad (6.3.65)$$

and in a cartesian basis this function reads

$$\chi_1 = \begin{pmatrix} \frac{1}{\sqrt{2}}(a^{-1} - a^{+1}) \\ -\frac{i}{\sqrt{2}}(a^{-1} + a^{+1}) \\ a^0 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}}(a_{-1} - a_{+1}) \\ \frac{i}{\sqrt{2}}(a_{-1} + a_{+1}) \\ a_0 \end{pmatrix} = \begin{pmatrix} a_x \\ a_y \\ a_z \end{pmatrix}. \quad (6.3.66)$$

The expansion coefficients a^m are contravariant spherical components of some generally complex vector $\mathbf{a}$, a_m are covariant spherical components of $\mathbf{a}$ and a_i are cartesian components. Sometimes $\mathbf{a}$ is called the polarization vector of particles of spin 1.

Under complex conjugation components of this vector transform as

$$(a^m)^* = (a^*)_m, (a_m)^* = (a^*)^m, (a_i)^* = (a^*)_i. \quad (6.3.67)$$

The normalization condition

$$\chi_1^\dagger \chi_1 = 1 \quad (6.3.68)$$

imposes a restriction to the components of $\mathbf{a}$

$$|a^{+1}|^2 + |a^0|^2 + |a^{-1}|^2 = |a_{+1}|^2 + |a_0|^2 + |a_{-1}|^2 = |a_x|^2 + |a_y|^2 + |a_z|^2 = 1, \quad (6.3.69)$$

or, in compact form

$$|\mathbf{a}|^2 = \mathbf{a}^* \cdot \mathbf{a} = 1. \quad (6.3.70)$$

The spin function product $\chi_1 \chi_1^\dagger$ may be expanded in series of the spin matrices $\hat{\mathbf{S}}$ and quadrupole momentum operators, $\hat{T}_{2M}$ or $\hat{Q}_{ik}$ (see Sec. 2.6),

$$\begin{aligned} \chi_1 \chi_1^\dagger &= \frac{1}{3} \hat{I} + \frac{1}{2} \sum_{\mu=-1}^1 (-1)^\mu P_{-\mu} \hat{S}_\mu + \sum_{M=-2}^2 (-1)^M P_{2-M} \hat{T}_{2M}, \\ \chi_1 \chi_1^\dagger &= \frac{1}{3} \hat{I} + \frac{1}{2} \sum_{i=x,y,z} P_i \hat{S}_i + \sum_{i,k=x,y,z} P_{ik} \hat{Q}_{ik}, \end{aligned} \quad (6.3.71)$$

or in more compact form,

$$\chi_1 \chi_1^\dagger = \frac{1}{3} \hat{\mathbf{I}} + \frac{1}{2} \mathbf{P} \cdot \hat{\mathbf{S}} + (\mathbf{P}_2 \cdot \hat{\mathbf{T}}_2), \quad (6.3.72)$$

where $(\mathbf{P}_2 \cdot \hat{\mathbf{T}}_2)$ is scalar product to two irreducible tensors of rank 2 (see Eq. (3.1.31)).

The real vector $\mathbf{P}$ can be expressed in terms of the polarization vector $\mathbf{a}$ as

$$\mathbf{P} = i [\mathbf{a} \times \mathbf{a}^*], \quad (6.3.73)$$

or, in components,

$$\begin{aligned} P_\mu &= \sqrt{2} \sum_{m,n} C_{1m1\mu}^1 (a^n)^* a^m, \quad (m, n, \mu = \pm 1, 0), \\ P_i &= i \sum_{k,l} \varepsilon_{ikl} a_k a_l^*, \quad (i, k, l = x, y, z). \end{aligned} \quad (6.3.74)$$

Spherical and cartesian components of a real irreducible tensor $\mathbf{P}_2$ of rank 2 may be expressed in terms of components of $\mathbf{a}$ by

$$\begin{aligned} P_{2M} &= \sqrt{\frac{5}{3}} \sum_{m,n} C_{1m2M}^1 (a^n)^* a^m, \quad (m, n = \pm 1, 0; M = \pm 2, \pm 1, 0), \\ P_{ik} &= -\frac{1}{2} \left\{ a_i a_k^* + a_k a_i^* - \frac{2}{3} \delta_{ik} \right\}, \quad (i, k = x, y, z), \\ P_{ik} &= P_{ki}, \quad \sum_i P_{ii} = 0. \end{aligned} \quad (6.3.75)$$

The vector $\mathbf{P}$ and tensor $\mathbf{P}_2$ give the following expectation values of the spin operator and the quadrupole momentum operator, respectively:

$$\begin{aligned} \langle \mathbf{S} \rangle &\equiv \chi_1^\dagger \hat{\mathbf{S}} \chi_1 = \mathbf{P}, \\ \langle \hat{T}_{2M} \rangle &\equiv \chi_1^\dagger \hat{T}_{2M} \chi_1 = P_{2M}, \\ \langle \hat{Q}_{ik} \rangle &\equiv \chi_1^\dagger \hat{Q}_{ik} \chi_1 = P_{ik}. \end{aligned} \quad (6.3.76)$$

The transformation of the spin function χ_1 under rotations of coordinate systems is performed by the rotation operators $\hat{D}^1(\alpha, \beta, \gamma)$ (Eqs. (2.6.76)-(2.6.77)) or $\hat{U}^1(\omega; \Theta, \Phi)$ (Eqs. (2.6.80)-(2.6.81)) depending on the choice of the parameters to describe rotations

$$\chi'_1 = \hat{D}^1(\alpha, \beta, \gamma) \chi_1 = \hat{U}^1(\omega; \Theta, \Phi) \chi_1. \quad (6.3.77)$$

Spin functions in a rotated coordinate system may be expanded into sums of basis spin functions χ_{1m}, χ_i in the original coordinate system as well as into sums of functions χ'_{1m}, χ'_i in the rotated coordinate system:

$$\begin{aligned} \chi'_1 &= \sum_{m=\pm 1,0} a'^m \chi_{1m} = \sum_{m=\pm 1,0} a^m \chi'_{1m}, \\ \chi'_1 &= \sum_{m=\pm 1,0} (-1)^m a'_{-m} \chi_{1m} = \sum_{m=\pm 1,0} (-1)^m a_{-m} \chi'_{1m}, \\ \chi'_1 &= \sum_{i=x,y,z} a'_i \chi_i = \sum_{i=x,y,z} a_i \chi'_i. \end{aligned} \quad (6.3.78)$$

The relations between basis functions in the rotated and original coordinate systems are given by Eqs. (6.3.34). The components of $\mathbf{a}$ in the above systems are related by

$$\begin{aligned} a'^m &= \sum_n D_{mn}^1(\alpha, \beta, \gamma) a^n, & a^n &= \sum_m D_{mn}^{1*}(\alpha, \beta, \gamma) a'^m, \\ a'_m &= \sum_n D_{nm}^1(\alpha, \beta, \gamma) a_n, & a_n &= \sum_m D_{nm}^{1*}(\alpha, \beta, \gamma) a'_m, \\ a'_i &= \sum_k a_{ik} a_k, & a_k &= \sum_i a_{ik} a'_i \end{aligned} \quad (6.3.79)$$

$(m, n = \pm 1, 0; i, k = x, y, z)$

Chapter 7

Tensor spherical harmonics

7.1 General properties of tensor spherical harmonics

7.1.1 Definition

The tensor spherical harmonics $Y_{JM}^{LS}(\vartheta, \varphi)$, by definition, are eigenfunctions of the operators $\hat{\mathbf{J}}^2$, $\hat{J}_z$, $\hat{\mathbf{L}}^2$ and $\hat{\mathbf{S}}^2$ where $\hat{\mathbf{L}}$ is the operator of orbital angular momentum (Sec. 2.2), $\hat{\mathbf{S}}$ is the spin operator (Sec. 2.3), and $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ is the operator of total angular momentum (Sec. 2.1):

$$\begin{aligned}\hat{\mathbf{J}}^2 Y_{JM}^{LS}(\vartheta, \varphi) &= J(J+1) Y_{JM}^{LS}(\vartheta, \varphi), \\ \hat{J}_z Y_{JM}^{LS}(\vartheta, \varphi) &= M Y_{JM}^{LS}(\vartheta, \varphi), \\ \hat{\mathbf{L}}^2 Y_{JM}^{LS}(\vartheta, \varphi) &= L(L+1) Y_{JM}^{LS}(\vartheta, \varphi), \\ \hat{\mathbf{S}}^2 Y_{JM}^{LS}(\vartheta, \varphi) &= S(S+1) Y_{JM}^{LS}(\vartheta, \varphi).\end{aligned}\tag{7.1.1}$$

A tensor spherical harmonic describes the angular distribution and polarization of spin- S particles in a state with definite total angular momentum J , projection M , and orbital angular momentum L . The spin value S is sometimes called the rank of the tensor spherical harmonic. Accordingly, one often uses such names as spinor spherical harmonics ($S = \frac{1}{2}$), vector spherical harmonics ($S = 1$), etc. However, strictly speaking, these terms are not entirely adequate because, in fact, the transformation properties of the tensor spherical harmonics under rotation of coordinate system are determined by J , but not by S (see Sec. 7.1.4).

The tensor spherical harmonics may be constructed from the spherical harmonics $Y_{LM}(\vartheta, \varphi)$ (eigenfunctions of $\hat{\mathbf{L}}^2$ and $\hat{L}_z$) and the spin functions $\chi_{S\sigma}$ (eigenfunctions of $\hat{\mathbf{S}}^2$ and $\hat{S}_z$) in accordance with the coupling scheme of two angular momenta

$$Y_{JM}^{LS}(\vartheta, \varphi) = \sum_{m, \sigma} C_{LmS\sigma}^J Y_{Lm}(\vartheta, \varphi) \chi_{S\sigma}.\tag{7.1.2}$$

Thus, the tensor spherical harmonics are irreducible tensor products (of rank J) of scalar spherical harmonics and spin functions:

$$Y_{JM}^{LS}(\vartheta, \varphi) = \{Y_L \otimes \chi_S\}_{JM}.\tag{7.1.3}$$

The indices J and S are integer or half-integer nonnegative numbers, and L is always an integer nonnegative number. For given J and S , the momentum L takes the values $L = |J - S|, |J - S| + 1, \dots, J + S$. The possible values of M are $M = -J, -J + 1, \dots, J - 1, J$.

The tensor spherical harmonics with fixed indices are functions of three variables: two polar angles ϑ, φ ($0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$) and spin variable ξ ($\xi = -S, -S + 1, \dots, S - 1, S$) which is the argument of spin function $\chi_{S\sigma}$. The spin variable ξ is usually not mentioned as an argument of tensor spherical harmonics. Instead, the tensor spherical harmonics are written in matrix form by analogy with spin functions. More precisely, $Y_{JM}^{LS}(\vartheta, \varphi)$ is represented by a column matrix with $2S + 1$ elements, and $Y_{JM}^{LS+}(\vartheta, \varphi)$ is represented by a row matrix. Summation over spin variable is replaced by matrix multiplication.

The tensor spherical harmonics $Y_{JM}^{LS}(\vartheta, \varphi)$ constitute a complete orthonormal set for series expansion of rank S tensor functions within the domain of arguments $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$ (see Sec. 7.1.8).

Note also the following orthogonality relations for $Y_{JM}^{LS}(\vartheta, \varphi)$ which have the same J, M and S but different L :

$$\sum_M Y_{JM}^{L'S+}(\vartheta, \varphi) \cdot Y_{JM}^{LS}(\vartheta, \varphi) = 0, \quad \text{if } L' \neq L.\tag{7.1.4}$$

7.1.2 Components of Tensor Spherical Harmonics

- (a) Spherical contravariant components of tensor spherical harmonics may be written, in accordance with (7.1.2), as

$$[Y_{JM}^{LS}(\vartheta, \varphi)]^\mu = C_{LM-\mu S \mu}^{JM} Y_{LM-\mu}(\vartheta, \varphi). \quad (7.1.5)$$

$$(\mu = -S, \dots, S-1, S).$$

Covariant spherical components may be defined by the relation

$$[Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = (-1)^{S-\mu} [Y_{JM}^{LS}(\vartheta, \varphi)]^{-\mu}, \quad (7.1.6)$$

from which one obtains

$$[Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = (-1)^{S-\mu} C_{LM+\mu S -\mu}^{JM} Y_{LM+\mu}(\vartheta, \varphi). \quad (7.1.7)$$

If S is integer, an alternative definition of the covariant spherical components that is widely used is (see Sec. 7.3.2)

$$[Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = (-1)^\mu [Y_{JM}^{LS}(\vartheta, \varphi)]^{-\mu}, \quad (7.1.8)$$

so that

$$[Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = (-1)^\mu C_{LM+\mu S -\mu}^{JM} Y_{LM+\mu}(\vartheta, \varphi). \quad (7.1.9)$$

Note that the equations of this chapter are independent of definition of covariant components, unless the contrary is indicated.

- (b) The expansion of tensor spherical harmonics in terms of helicity spin functions may be written as

$$Y_{JM}^{LS}(\vartheta, \varphi) = \sum_{\lambda=-S}^S [Y_{JM}^{LS}(\vartheta, \varphi)]'^\lambda \chi_{s\lambda}(\vartheta, \varphi), \quad (7.1.10)$$

where contravariant helicity components are given by

$$[Y_{JM}^{LS}(\vartheta, \varphi)]'^\lambda = \sqrt{\frac{2L+1}{4\pi}} C_{L0S\lambda}^{J\lambda} D_{-\lambda-M}^J(0, \vartheta, \varphi). \quad (7.1.11)$$

If covariant helicity components are defined by

$$[Y_{JM}^{LS}(\vartheta, \varphi)]'_\lambda = (-1)^{S-\lambda} [Y_{JM}^{LS}(\vartheta, \varphi)]'^{-\lambda}, \quad (7.1.12)$$

then

$$[Y_{JM}^{LS}(\vartheta, \varphi)]'_\lambda = (-1)^{J+L-\lambda} \sqrt{\frac{2L+1}{4\pi}} C_{L0S\lambda}^{J\lambda} D_{\lambda-M}^J(0, \vartheta, \varphi). \quad (7.1.13)$$

An alternative definition of covariant components (for integer S)

$$[Y_{JM}^{LS}(\vartheta, \varphi)]'_\lambda = (-1)^\lambda [Y_{JM}^{LS}(\vartheta, \varphi)]'^{-\lambda}, \quad (7.1.14)$$

yields

$$[Y_{JM}^{LS}(\vartheta, \varphi)]'_\lambda = (-1)^{J+L-S+\lambda} \sqrt{\frac{2L+1}{4\pi}} C_{L0S\lambda}^{J\lambda} D_{\lambda-M}^J(0, \vartheta, \varphi). \quad (7.1.15)$$

7.1.3 Complex Conjugation

Spherical components of a tensor spherical harmonic transform under complex conjugation according to

$$[Y_{JM}^{LS}(\vartheta, \varphi)]^{\mu*} = (-1)^{J+L+S-M-\mu} [Y_{J-M}^{LS}(\vartheta, \varphi)]^{-\mu},$$

$$[Y_{JM}^{LS}(\vartheta, \varphi)]_\mu^* = (-1)^{J+L+S-M-\mu} [Y_{J-M}^{LS}(\vartheta, \varphi)]_{-\mu}. \quad (7.1.16)$$

These transformation rules are valid also for the corresponding helicity components.

7.1.4 Transformations of Coordinate Systems

(a) Coordinate inversion

The operator for coordinate inversion $\hat{P}_r$, when applied to a tensor spherical harmonic, gives

$$\hat{P}_r Y_{JM}^{LS}(\vartheta, \varphi) = \eta_p Y_{JM}^{LS}(\pi - \vartheta, \pi + \varphi) = \eta_p (-1)^L Y_{JM}^{LS}(\vartheta, \varphi), \quad (7.1.17)$$

where η_p is a phase factor which describes the intrinsic parity of the tensor.

(b) Rotations of coordinate systems

Under rotations of the coordinate system specified by the Euler angles α, β, γ the tensor spherical harmonics transform according to

$$Y_{JM'}^{LS}(\vartheta', \varphi') \equiv \hat{D}(\alpha, \beta, \gamma) Y_{JM}^{LS}(\vartheta, \varphi) = \sum_{M=-J}^J D_{MM'}^J(\alpha, \beta, \gamma) Y_{JM}^{LS}(\vartheta, \varphi), \quad (7.1.18)$$

where M is the projection of total angular momentum on the initial z -axis, M' is its projection on the final z' axis; ϑ, φ and ϑ', φ' are polar angles of the unit vector $\mathbf{n}$ in the initial and final coordinate systems, respectively. The relations between ϑ', φ' and ϑ, φ are given by Eqs. (1.4.2), (1.4.3). The coefficients $D_{MM'}^J(\alpha, \beta, \gamma)$ are matrix elements of the rotation operator, i.e., the Wigner D -functions (Chap. 4). Note that the transformation properties of tensor spherical harmonics are determined by the total angular momentum $\mathbf{J}$, but not by the spin $\mathbf{S}$. Thus, it is J that determines true rank of a tensor spherical harmonic.

7.1.5 Differential Equations

(a) Tensor spherical harmonics are eigenfunctions of the operator $\hat{\mathbf{L}}^2$; hence, they satisfy the equation

$$\{\Delta_\Omega + L(L+1)\} Y_{JM}^{LS}(\vartheta, \varphi) = 0, \quad (7.1.19)$$

where Δ_Ω is the angular part of the Laplace operator (see Sec. 1.3.2). The expanded form of (7.1.19) is

$$\frac{1}{\sin \vartheta} \cdot \frac{\partial}{\partial \vartheta} \left\{ \sin \vartheta \frac{\partial}{\partial \vartheta} Y_{JM}^{LS}(\vartheta, \varphi) \right\} + \frac{1}{\sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2} Y_{JM}^{LS}(\vartheta, \varphi) + L(L+1) Y_{JM}^{LS}(\vartheta, \varphi) = 0. \quad (7.1.20)$$

(b) Components of the tensors $r^L Y_{JM}^{LS}(\vartheta, \varphi)$ are harmonic polynomials of degree L and satisfy the Laplace equation

$$\Delta \{r^L Y_{JM}^{LS}(\vartheta, \varphi)\} = 0. \quad (7.1.21)$$

(c) The quantities $z_L(kr) Y_{JM}^{LS}(\vartheta, \varphi)$ obey the Helmholtz equation

$$(\Delta + k^2) \{z_L(kr) Y_{JM}^{LS}(\vartheta, \varphi)\} = 0, \quad (7.1.22)$$

where

$$z_L(kr) = \sqrt{\frac{\pi}{2kr}} Z_{L+\frac{1}{2}}(kr). \quad (7.1.23)$$

$Z_{L+\frac{1}{2}}$ is any of the cylinder functions of half-integer order.

7.1.6 Action of Operators $\nabla, \mathbf{n}$ and Angular Momentum Operators

In the equations given below $\mathbf{r} = \{r, \vartheta, \varphi\}$, $\mathbf{n} = \mathbf{r}/r$, $\Phi(r)$ is an arbitrary function of $r = |\mathbf{r}|$, $\hat{\mathbf{L}}$ is the operator of orbital angular momentum, $\hat{\mathbf{S}}$ is the spin operator, $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ is the operator of total angular momentum, and $n_\mu, \nabla_\mu, \hat{L}_\mu$, etc. are spherical components of the corresponding operators ($\mu = \pm 1, 0$).

(a)

$$\begin{aligned} \nabla_\mu \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} &= (-1)^{J+L+S} \sqrt{(2J+1)(L+1)} \left\{ \frac{d\Phi(r)}{dr} - \frac{L}{r} \Phi(r) \right\}, \\ &\times \sum_{J'} \left\{ \begin{matrix} J & J' & 1 \\ L+1 & L & S \end{matrix} \right\} C_{JM+1\mu}^{J' M+\mu} Y_{J'M+\mu}^{L+1S}(\vartheta, \varphi) + (-1)^{J+L+S+1} \sqrt{(2J+1)L} \left\{ \frac{d\Phi(r)}{dr} + \frac{L+1}{r} \Phi(r) \right\}, \\ &\times \sum_{J'} \left\{ \begin{matrix} J & J' & 1 \\ L-1 & L & S \end{matrix} \right\} C_{JM-1\mu}^{J' M+\mu} Y_{J'M+\mu}^{L-1S}(\vartheta, \varphi). \end{aligned} \quad (7.1.24)$$

In particular,

$$r\nabla_\mu Y_{JM}^{LS}(\vartheta, \varphi) = (-1)^{J+L+S+1} \sqrt{2J+1} \sum_{J'L'} A_{L'}(L) \left\{ \begin{matrix} J & J' & 1 \\ L' & L & S \end{matrix} \right\} C_{JM1\mu}^{J'M+\mu} Y_{J'M+\mu}^{L'S}(\vartheta, \varphi), \quad (7.1.25)$$

where

$$A_{L'}(L) = \begin{cases} L\sqrt{L+1}, & \text{if } L' = L+1 \\ (L+1)\sqrt{L}, & \text{if } L' = L-1 \\ 0 & \text{if } L' \neq L \pm 1 \end{cases}$$

$$n_\mu \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = (-1)^{J+S+1} \Phi(r) \sqrt{(2J+1)(2L+1)} \sum_{J'L'} (-1)^{L'} \left\{ \begin{matrix} J & J' & 1 \\ L' & L & S \end{matrix} \right\} C_{L'010}^{L'0} C_{JM1\mu}^{J'M'} Y_{J'M'}^{L'S}(\vartheta, \varphi), \quad (7.1.26)$$

$$\begin{aligned} \widehat{L}_\mu \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} &= \Phi(r) \widehat{L}_\mu \{ Y_{JM}^{LS}(\vartheta, \varphi) \} \\ &= (-1)^{J+L+S+1} \Phi(r) \sqrt{(2J+1)L(L+1)(2L+1)} \sum_{J'} \left\{ \begin{matrix} J & J' & 1 \\ L & L & S \end{matrix} \right\} C_{JM1\mu}^{J'M+\mu} Y_{J'M+\mu}^{LS}(\vartheta, \varphi), \end{aligned} \quad (7.1.27)$$

$$\begin{aligned} \widehat{S}_\mu \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} &= \Phi(r) \widehat{S}_\mu \{ Y_{JM}^{LS}(\vartheta, \varphi) \} \\ &= (-1)^{L+1} \Phi(r) \sqrt{(2J+1)S(S+1)(2S+1)} \sum_{J'} (-1)^{J'+S} \left\{ \begin{matrix} J & J' & 1 \\ S & S & L \end{matrix} \right\} C_{JM1\mu}^{J'M+\mu} Y_{J'M+\mu}^{LS}(\vartheta, \varphi) \end{aligned} \quad (7.1.28)$$

$$\widehat{J}_\mu \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = \Phi(r) \widehat{J}_\mu \{ Y_{JM}^{LS}(\vartheta, \varphi) \} = \Phi(r) \sqrt{J(J+1)} C_{JM1\mu}^{J'M+\mu} Y_{J'M+\mu}^{LS}(\vartheta, \varphi). \quad (7.1.29)$$

(b)

$$(\mathbf{r} \cdot \nabla) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = r \frac{d\Phi(r)}{dr} Y_{JM}^{LS}(\vartheta, \varphi), \quad (7.1.30)$$

$$(\mathbf{r} \cdot \widehat{\mathbf{L}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = 0, \quad (7.1.31)$$

$$(\mathbf{r} \cdot \widehat{\mathbf{S}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = r \Phi(r) (-1)^{J+L+S} \sqrt{S(S+1)(2S+1)(2L+1)} \sum_{L'} \left\{ \begin{matrix} L & L' & 1 \\ S & S & L \end{matrix} \right\} C_{L'010}^{L'0} Y_{JM}^{L'S}(\vartheta, \varphi). \quad (7.1.32)$$

An expanded form of (7.1.32) is

$$\begin{aligned} (\widehat{\mathbf{S}} \cdot \mathbf{n}) Y_{JM}^{LS}(\vartheta, \varphi) &= -\frac{1}{2} \left\{ \sqrt{\frac{(J+L+S+2)(J+L-S+1)(J-L+S)(-J+L+S+1)}{(2L+1)(2L+3)}} Y_{JM}^{L+1S}(\vartheta, \varphi) \right. \\ &\quad \left. + \sqrt{\frac{(J+L+S+1)(J+L-S)(J-L+S+1)(-J+L+S)}{(2L-1)(2L+1)}} Y_{JM}^{L-1S}(\vartheta, \varphi) \right\}, \end{aligned} \quad (7.1.33)$$

$$(\mathbf{r} \cdot \widehat{\mathbf{J}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = (\mathbf{r} \cdot \widehat{\mathbf{S}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \}. \quad (7.1.34)$$

(c)

$$\begin{aligned} &(\widehat{\mathbf{S}} \cdot \nabla) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = \\ &= -\frac{1}{2} \left\{ \frac{d\Phi(r)}{dr} - \frac{L}{r} \Phi(r) \right\} \sqrt{\frac{(J+L+S+2)(J+L-S+1)(J-L+S)(-J+L+S+1)}{(2L+1)(2L+3)}} Y_{JM}^{L+1S}(\vartheta, \varphi) \\ &\quad - \frac{1}{2} \left\{ \frac{d\Phi(r)}{dr} + \frac{L+1}{r} \Phi(r) \right\} \sqrt{\frac{(J+L+S+1)(J+L-S)(J-L+S+1)(-J+L+S)}{(2L-1)(2L+1)}} Y_{JM}^{L-1S}(\vartheta, \varphi). \end{aligned} \quad (7.1.35)$$

$$(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = \frac{1}{2} \{ J(J+1) - L(L+1) - S(S+1) \} \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi). \quad (7.1.36)$$

$$(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{J}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = \frac{1}{2} \{ J(J+1) - L(L+1) + S(S+1) \} \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi). \quad (7.1.37)$$

$$(\widehat{\mathbf{L}} \cdot \widehat{\mathbf{J}}) \{ \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi) \} = \frac{1}{2} \{ J(J+1) + L(L+1) - S(S+1) \} \Phi(r) Y_{JM}^{LS}(\vartheta, \varphi). \quad (7.1.38)$$

7.1.7 Sums of Tensor Spherical Harmonics

In what follows ((7.1.39)-(7.1.42)) S is integer, and covariant components of tensor spherical harmonics are defined in accordance with (7.1.8), (7.1.9):

$$\sum_{\mu=-S}^S Y_{S\mu}(\vartheta, \varphi) [Y_{JM}^{LS}(\vartheta, \varphi)]^\mu = \sqrt{\frac{(2S+1)(2L+1)}{4\pi(2J+1)}} C_{L0S0}^{J0} Y_{JM}(\vartheta, \varphi). \quad (7.1.39)$$

$$\sum_{\mu=-S}^S Y_{S\mu}^*(\vartheta, \varphi) [Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = \sqrt{\frac{(2S+1)(2L+1)}{4\pi(2J+1)}} C_{L0S0}^{J0} Y_{JM}^*(\vartheta, \varphi), \quad (7.1.40)$$

$$\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) [Y_{JM}^{LS}(\vartheta, \varphi)]^\mu = (-1)^\mu \sqrt{\frac{(2J+1)(2L+1)}{4\pi(2S+1)}} C_{S0L0}^{J0} Y_{S-\mu}(\vartheta, \varphi), \quad (7.1.41)$$

$$\sum_{M=-J}^J Y_{JM}(\vartheta, \varphi) [Y_{JM}^{LS}(\vartheta, \varphi)]_\mu = \sqrt{\frac{(2J+1)(2L+1)}{4\pi(2S+1)}} C_{S0L0}^{J0} Y_{S\mu}(\vartheta, \varphi). \quad (7.1.42)$$

7.1.8 Orthogonality, Normalization and Completeness

A collection of tensor spherical harmonics with the same S and all possible J, L, M forms a complete orthonormal set of tensor functions in the domain of the arguments $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$.

The orthonormality condition for tensor spherical harmonics has the form

$$\int_0^\pi \int_0^{2\pi} Y_{J'M'}^{L'} S + (\vartheta, \varphi) Y_{JM}^{LS}(\vartheta, \varphi) \sin \vartheta d\vartheta d\varphi = \delta_{JJ'} \delta_{MM'} \delta_{LL'}. \quad (7.1.43)$$

The completeness of these functions is given by

$$\sum_J \sum_{L=|J-S|}^{J+S} \sum_{M=-J}^J [Y_{JM}^{LS}(\vartheta, \varphi)]^\mu [Y_{JM}^{LS}(\vartheta', \varphi')]^{\nu*} = \delta_{\mu\nu} \delta(\cos \vartheta - \cos \vartheta') \delta(\varphi - \varphi'), \quad (7.1.44)$$

$$(\mu; \nu = -S, \dots, S-1, S).$$

7.1.9 Expansion in a Series of Tensor Spherical Harmonics

Any tensor $F_S(\vartheta, \varphi)$ of rank S which depends on polar angles ϑ, φ and possesses components which satisfy the conditions

$$\int |F_{S\mu}(\vartheta, \varphi)|^2 d\Omega < \infty, \quad (\mu = -S, \dots, S-1, S), \quad (7.1.45)$$

may be expanded in a series of tensor spherical harmonics $Y_{JM}^{LS}(\vartheta, \varphi)$, i.e., written in the form

$$F_S(\vartheta, \varphi) = \sum_{JLM} A_{JLM} Y_{JM}^{LS}(\vartheta, \varphi). \quad (7.1.46)$$

The expansion coefficients are given by

$$A_{JLM} = \int d\Omega Y_{JM}^{LS} + (\vartheta, \varphi) F_S(\vartheta, \varphi). \quad (7.1.47)$$

These coefficients satisfy the relation

$$\sum_{JLM} |A_{JLM}|^2 = \int d\Omega |F_S(\vartheta, \varphi)|^2. \quad (7.1.48)$$

7.2 Spinor spherical harmonics

7.2.1 Definition

The spinor spherical harmonics $\Omega_{JM}^L(\vartheta, \varphi)$ are defined as tensor spherical harmonics with $S = \frac{1}{2}$ (Sec. 7.1.1)

$$\Omega_{JM}^L(\vartheta, \varphi) \equiv Y_{JM}^{L\frac{1}{2}}(\vartheta, \varphi). \quad (7.2.1)$$

They are eigenfunctions of the operators $\hat{\mathbf{J}}^2, \hat{J}_z, \hat{\mathbf{L}}^2, \hat{\mathbf{S}}^2$ where $\hat{\mathbf{L}}$ is the operator of orbital angular momentum (Sec. 2.2), $\hat{\mathbf{S}}$ is the spin operator for $S = \frac{1}{2}$ (Sec. 2.5) and $\hat{\mathbf{J}}$ is the operator of total angular momentum (Sec. 2.1)

$$\begin{aligned}\hat{J}^2 \Omega_{JM}^L(\vartheta, \varphi) &= J(J+1) \Omega_{JM}^L(\vartheta, \varphi), \\ \hat{J}_z \Omega_{JM}^L(\vartheta, \varphi) &= M \Omega_{JM}^L(\vartheta, \varphi), \\ \hat{\mathbf{L}}^2 \Omega_{JM}^L(\vartheta, \varphi) &= L(L+1) \Omega_{JM}^L(\vartheta, \varphi), \\ \hat{\mathbf{S}}^2 \Omega_{JM}^L(\vartheta, \varphi) &= \frac{3}{4} \Omega_{JM}^L(\vartheta, \varphi).\end{aligned}\tag{7.2.2}$$

The spinor spherical harmonics may be expressed in terms of scalar spherical harmonics $Y_{LM}(\vartheta, \varphi)$ (Chap. 5) and spin functions $\chi_{\frac{1}{2}\sigma}$ (Sec. 6.2) as

$$\Omega_{JM}^L(\vartheta, \varphi) = \sum_{m\sigma} C_{Lm\frac{1}{2}\sigma}^{JM} Y_{Lm}(\vartheta, \varphi) \chi_{\frac{1}{2}\sigma}.\tag{7.2.3}$$

For the spinor spherical harmonics J is a half-integer nonnegative number because L is always integer. For given J only two values of L are possible, $L = J \pm \frac{1}{2}$, while M assumes $2J+1$ values: $M = -J, -J+1, \dots, J-1, J$. The spinor spherical harmonics are orthonormalized in the domain of angles $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$ (Sec. 7.2.7).

7.2.2 Components of Spinor Spherical Harmonics

(a) The spinor spherical harmonics may be expanded in terms of basis spin functions $\chi_{\frac{1}{2}\mu}$ as

$$\Omega_{JM}^L(\vartheta, \varphi) = \sum_{\mu=-\frac{1}{2}}^{\frac{1}{2}} [\Omega_{JM}^L(\vartheta, \varphi)]^\mu \chi_{\frac{1}{2}\mu},\tag{7.2.4}$$

where contravariant components $[\Omega_{JM}^L]^\mu$ are given by

$$[\Omega_{JM}^L(\vartheta, \varphi)]^\mu = C_{LM-\mu\frac{1}{2}\mu}^{JM} Y_{LM-\mu}(\vartheta, \varphi).\tag{7.2.5}$$

In more detailed form (7.2.5) reads

$$\begin{aligned}[\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi)]^{\frac{1}{2}} &= -\sqrt{\frac{J-M+1}{2(J+1)}} Y_{J+\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi), \\ [\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi)]^{-\frac{1}{2}} &= \sqrt{\frac{J+M+1}{2(J+1)}} Y_{J+\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi),\end{aligned}\tag{7.2.6}$$

$$\begin{aligned}[\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi)]^{\frac{1}{2}} &= \sqrt{\frac{J+M}{2J}} Y_{J-\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi), \\ [\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi)]^{-\frac{1}{2}} &= \sqrt{\frac{J-M}{2J}} Y_{J-\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi).\end{aligned}\tag{7.2.7}$$

Covariant components of spinor spherical harmonics are defined by

$$[\Omega_{JM}^L(\vartheta, \varphi)]_\mu = (-1)^{\frac{1}{2}-\mu} [\Omega_{JM}^L(\vartheta, \varphi)]^{-\mu},\tag{7.2.8}$$

and have the form

$$[\Omega_{JM}^L(\vartheta, \varphi)]_\mu = (-1)^{\frac{1}{2}-\mu} C_{LM+\mu\frac{1}{2}-\mu}^{JM} Y_{LM+\mu}(\vartheta, \varphi).\tag{7.2.9}$$

A more detailed form of (7.2.9) is

$$\begin{aligned}[\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi)]_{\frac{1}{2}} &= \sqrt{\frac{J+M+1}{2(J+1)}} Y_{J+\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi), \\ [\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi)]_{-\frac{1}{2}} &= \sqrt{\frac{J-M+1}{2(J+1)}} Y_{J+\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi),\end{aligned}\tag{7.2.10}$$

$$\begin{aligned}[\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi)]_{\frac{1}{2}} &= \sqrt{\frac{J+M}{2J}} Y_{J-\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi), \\ [\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi)]_{-\frac{1}{2}} &= -\sqrt{\frac{J-M}{2J}} Y_{J-\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi).\end{aligned}\tag{7.2.11}$$

- (b) If spinors $\chi_{\frac{1}{2}\sigma}$ are written as column matrices, the spinor spherical harmonics assume the form

$$\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} -\sqrt{\frac{J-M+1}{2(J+1)}} Y_{J+\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi) \\ \sqrt{\frac{J+M+1}{2(J+1)}} Y_{J+\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi) \end{pmatrix}, \quad (7.2.12)$$

$$\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi) = \begin{pmatrix} \sqrt{\frac{J+M}{2J}} Y_{J-\frac{1}{2}M-\frac{1}{2}}(\vartheta, \varphi) \\ \sqrt{\frac{J-M}{2J}} Y_{J-\frac{1}{2}M+\frac{1}{2}}(\vartheta, \varphi) \end{pmatrix}. \quad (7.2.13)$$

- (c) An expansion of $\Omega_{JM}^L(\vartheta, \varphi)$ in terms of helicity basis spin functions $\chi_{\frac{1}{2}\lambda}(\vartheta, \varphi)$ (Sec. 6.2.5) may be written as

$$\Omega_{JM}^L(\vartheta, \varphi) = \sqrt{\frac{2L+1}{4\pi}} \sum_{\lambda} C_{L0\frac{1}{2}\lambda}^{J\lambda} D_{-\lambda-M}^J(0, \vartheta, \varphi) \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi), \quad (7.2.14)$$

or, in a detailed form

$$\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi) = \sqrt{\frac{2J+1}{8\pi}} \sum_{\lambda} (-1)^{\frac{1}{2}+\lambda} D_{-\lambda-M}^J(0, \vartheta, \varphi) \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi), \quad (7.2.15)$$

$$\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi) = \sqrt{\frac{2J+1}{8\pi}} \sum_{\lambda} D_{-\lambda-M}^J(0, \vartheta, \varphi) \chi_{\frac{1}{2}\lambda}(\vartheta, \varphi).$$

7.2.3 Complex Conjugation. Time Reversal

- (a) Here we use the representation in which the basis spin functions are real, $\chi_{\frac{1}{2}\sigma}^* = \chi_{\frac{1}{2}\sigma}$. For this reason, the transformation properties of spinor spherical harmonics with respect to complex conjugation may be written as

$$\Omega_{JM}^{L*}(\vartheta, \varphi) = (-1)^{J+L-M} i \hat{\sigma}_y \Omega_{J-M}^L(\vartheta, \varphi), \quad (7.2.16)$$

where 2×2 matrix $i \hat{\sigma}_y$ has the form

$$i \hat{\sigma}_y = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (7.2.17)$$

Components of spinor spherical harmonics transform under complex conjugation according to

$$[\Omega_{JM}^L(\vartheta, \varphi)]_{\mu}^* = [\Omega_{JM}^{L*}(\vartheta, \varphi)]_{\mu} = (-1)^{J+L+M} [\Omega_{J-M}^L(\vartheta, \varphi)]^{\mu} = (-1)^{J+L-M-\mu+\frac{1}{2}} [\Omega_{J-M}^L(\vartheta, \varphi)]_{-\mu}, \quad (7.2.18)$$

$$[\Omega_{JM}^L(\vartheta, \varphi)]^{\mu*} = [\Omega_{JM}^{L*}(\vartheta, \varphi)]^{\mu} = (-1)^{J+L-M} [\Omega_{J-M}^L(\vartheta, \varphi)]_{\mu} = (-1)^{J+L-M-\mu+\frac{1}{2}} [\Omega_{J-M}^L(\vartheta, \varphi)]^{-\mu}. \quad (7.2.19)$$

- (b) The time-reversal operator $\hat{P}_t$ (as defined by Wigner) acts on a spinor spherical harmonic in the following manner

$$\hat{P}_t \Omega_{JM}^L(\vartheta, \varphi) = \hat{\sigma}_y \Omega_{J-M}^{L*}(\vartheta, \varphi) = (-1)^{J+L-M+\frac{1}{2}} \Omega_{J-M}^L(\vartheta, \varphi). \quad (7.2.20)$$

7.2.4 Transformation of Coordinate Systems.

- (a) *Coordinate inversion*

$$\hat{P}_r \Omega_{JM}^L(\vartheta, \varphi) = \Omega_{JM}^L(\pi - \vartheta, \pi + \varphi) = (-1)^L \Omega_{JM}^L(\vartheta, \varphi). \quad (7.2.21)$$

- (b) *Rotations of coordinate system*

Under rotations specified by the Euler angles α, β, γ the spinor spherical harmonics transform according to

$$\Omega_{JM'}^L(\vartheta', \varphi') = \hat{D}(\alpha, \beta, \gamma) \Omega_{JM'}^L(\vartheta, \varphi) = \sum_{M=-J}^J D_{MM'}^J(\alpha, \beta, \gamma) \Omega_{JM}^L(\vartheta, \varphi), \quad (7.2.22)$$

where $D_{MM'}^J$ are the Wigner D -functions (Chap. 4), M' is the projection of total angular momentum on the rotated z' -axis; ϑ, φ and ϑ', φ' are polar angles of the position vector $\mathbf{r}$ in the initial and final coordinate systems. The relations between ϑ', φ' and ϑ, φ are given by Eqs. (1.4.2)-(1.4.3).

7.2.5 Action of ∇ and Angular Momentum Operators

The spinor spherical harmonics satisfy Eqs. (7.1.24)-(7.1.38) for $S = \frac{1}{2}$. Note also the following relations in which $\hat{\mathbf{S}}$ is the spin operator (Sec. 2.5), $\hat{\mathbf{L}}$ is the operator of orbital angular momentum (Sec. 2.2), $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ is the operator of total angular momentum, and $\mathbf{n}(\vartheta, \varphi) = \mathbf{r}/r$.

(a)

$$(\hat{\mathbf{S}} \cdot \mathbf{n})\Omega_{JM}^L(\vartheta, \varphi) = -\frac{1}{2}\Omega_{JM}^{L'}(\vartheta, \varphi), \quad (7.2.23)$$

where

$$L' = 2J - L = \begin{cases} J - \frac{1}{2}, & \text{if } L = J + \frac{1}{2} \\ J + \frac{1}{2}, & \text{if } L = J - \frac{1}{2} \end{cases}. \quad (7.2.24)$$

An expanded form of (7.2.23) is

$$\begin{aligned} (\hat{\mathbf{S}} \cdot \mathbf{n})\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi) &= -\frac{1}{2}\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi), \\ (\hat{\mathbf{S}} \cdot \mathbf{n})\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi) &= -\frac{1}{2}\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi). \end{aligned} \quad (7.2.25)$$

(b)

$$r(\hat{\mathbf{S}} \cdot \nabla)\Omega_{JM}^L(\vartheta, \varphi) = -\frac{1+\kappa}{2}\Omega_{JM}^{L'}(\vartheta, \varphi), \quad (7.2.26)$$

where $L' = 2J - L$,

$$\kappa = \begin{cases} J + \frac{1}{2} = L, & \text{if } L = J + \frac{1}{2} \\ -(J + \frac{1}{2}) = -(L + 1), & \text{if } L = J - \frac{1}{2} \end{cases}. \quad (7.2.27)$$

In a more detailed form (7.2.26) gives

$$\begin{aligned} r(\hat{\mathbf{S}} \cdot \nabla)\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi) &= -\frac{2J+3}{4}\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi), \\ r(\hat{\mathbf{S}} \cdot \nabla)\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi) &= \frac{2J-1}{4}\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi). \end{aligned} \quad (7.2.28)$$

(c)

$$(\hat{\mathbf{L}} \cdot \hat{\mathbf{S}})\Omega_{JM}^L(\vartheta, \varphi) = \frac{1}{2} \left\{ J(J+1) - L(L+1) - \frac{3}{4} \right\} \Omega_{JM}^L(\vartheta, \varphi) = -\frac{1+\kappa}{2}\Omega_{JM}^L(\vartheta, \varphi), \quad (7.2.29)$$

or, in detailed form

$$\begin{aligned} (\hat{\mathbf{L}} \cdot \hat{\mathbf{S}})\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi) &= -\frac{2J+3}{4}\Omega_{JM}^{J+\frac{1}{2}}(\vartheta, \varphi), \\ (\hat{\mathbf{L}} \cdot \hat{\mathbf{S}})\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi) &= \frac{2J-1}{4}\Omega_{JM}^{J-\frac{1}{2}}(\vartheta, \varphi). \end{aligned} \quad (7.2.30)$$

7.2.6 Recursion Relations

$$\begin{aligned} \cos \vartheta \Omega_{JM}^L(\vartheta, \varphi) &= \frac{\sqrt{(J-M+1)(J+M+1)}}{2(J+1)} \Omega_{J+1M}^{L+1}(\vartheta, \varphi) - \frac{M}{2J(J+1)} \Omega_{JM}^{L'}(\vartheta, \varphi) \\ &\quad + \frac{\sqrt{(J-M)(J+M)}}{2J} \Omega_{J-1M}^{L-1}(\vartheta, \varphi) \end{aligned} \quad (7.2.31)$$

$$\begin{aligned} \sin \vartheta e^{i\varphi} \Omega_{JM}^L(\vartheta, \varphi) &= -\frac{\sqrt{(J+M+1)(J+M+2)}}{2(J+1)} \Omega_{J+1M+1}^{L+1}(\vartheta, \varphi) \\ &\quad - \frac{\sqrt{(J+M+1)(J-M)}}{2J(J+1)} \Omega_{JM+1}^{L'}(\vartheta, \varphi) - \frac{\sqrt{(J-M)(J-M-1)}}{2J} \Omega_{J-1M+1}^{L-1}(\vartheta, \varphi), \end{aligned} \quad (7.2.32)$$

$$\begin{aligned} \sin \vartheta e^{-i\varphi} \Omega_{JM}^L(\vartheta, \varphi) &= \frac{\sqrt{(J-M+1)(J-M+2)}}{2(J+1)} \Omega_{J+1M-1}^{L+1}(\vartheta, \varphi) \\ &\quad - \frac{\sqrt{(J+M)(J-M+1)}}{2J(J+1)} \Omega_{JM-1}^{L'}(\vartheta, \varphi) - \frac{\sqrt{(J+M)(J+M-1)}}{2J} \Omega_{J-1M-1}^{L-1}(\vartheta, \varphi), \end{aligned} \quad (7.2.33)$$

where L' is defined by (7.2.24). Note that Eqs. (7.2.31)-(7.2.33) may be written in a compact form as

$$n_\mu \Omega_{JM}^L(\vartheta, \varphi) = (-1)^{J+L+\frac{1}{2}} \sqrt{(2J+1)(2L+1)} \sum_{J'L'} \left\{ \begin{matrix} J' & J & 1 \\ L & L' & \frac{1}{2} \end{matrix} \right\} C_{L'010}^{L'0} C_{JM1\mu}^{J'M+\mu} \Omega_{J'M+\mu}^{L'}(\vartheta, \varphi) \quad (7.2.34)$$

where $n_\mu (\mu = \pm 1, 0)$ are spherical components of the unit vector $n(\vartheta, \varphi)$.

7.2.7 Orthonormality and Completeness. Expansion in a Series of Spinor Spherical Harmonics

The collection of spinor spherical harmonics $\Omega_{JM}^L(\vartheta, \varphi)$ with half-integer nonnegative J ($\frac{1}{2} \leq J < \infty$), $L = J \pm \frac{1}{2}$ and $M = -J, -J+1, \dots, J$ forms complete set of spinors for $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$.

The orthonormality condition has the form

$$\int_0^\pi d\vartheta \sin \vartheta \int_0^{2\pi} d\varphi \Omega_{J'M'}^{L'}(\vartheta, \varphi) \Omega_{JM}^L(\vartheta, \varphi) = \delta_{J'J} \delta_{L'L} \delta_{M'M}. \quad (7.2.35)$$

The completeness condition may be written as

$$\sum_{J=\frac{1}{2}}^\infty \sum_{L=J-\frac{1}{2}}^{J+\frac{1}{2}} \sum_{M=-J}^J \Omega_{JM}^L(\vartheta, \varphi) \Omega_{JM}^{L+}(\vartheta', \varphi') = \hat{I} \delta(\cos \vartheta - \cos \vartheta') \delta(\varphi - \varphi'), \quad (7.2.36)$$

where $\hat{I}$ is the unit 2×2 matrix.

Any spinor $\chi(\vartheta, \varphi)$ which satisfies

$$\int \chi^+(\vartheta, \varphi) \chi(\vartheta, \varphi) d\Omega < \infty, \quad (7.2.37)$$

may be expanded in a series of spinor spherical harmonics

$$\chi(\vartheta, \varphi) = \sum_{JLM} A_{JM}^L \Omega_{JM}^L(\vartheta, \varphi), \quad (7.2.38)$$

where the expansion coefficients A_{JM}^L are given by

$$A_{JM}^L = \int_0^\pi d\vartheta \sin \vartheta \int_0^{2\pi} d\varphi \Omega_{JM}^{L+}(\vartheta, \varphi) \chi(\vartheta, \varphi). \quad (7.2.39)$$

The expansion (7.2.38) is valid for the range $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$.

7.2.8 Clebsch-Gordan Series

$$\begin{aligned} \Omega_{J_1 M_1}^{L_1+}(\vartheta, \varphi) \Omega_{J_2 M_2}^{L_2}(\vartheta, \varphi) &= \sum_L (-1)^{J_1+M_1+J_2+L+\frac{1}{2}} \left\{ \begin{matrix} L_1 & L_2 & L \\ J_2 & J_1 & \frac{1}{2} \end{matrix} \right\} \\ &\times \sqrt{\frac{(2J_1+1)(2J_2+1)(2L_1+1)(2L_2+1)}{4\pi(2L+1)}} C_{L_1 0 L_2 0}^{L 0} C_{J_1 -M_1 J_2 M_2}^{L M} Y_{LM}(\vartheta, \varphi), \end{aligned} \quad (7.2.40)$$

$$\begin{aligned} \Omega_{J_1 M_1}^{L_1+}(\vartheta, \varphi) \hat{S} \Omega_{J_2 M_2}^{L_2}(\vartheta, \varphi) &= (-1)^{J_2+L_1+M_1} \sqrt{\frac{3(2J_1+1)(2J_2+1)(2L_1+1)(2L_2+1)}{8\pi}} \\ &\times \sum_{JL} (-1)^J C_{L_1 0 L_2 0}^{L 0} \left\{ \begin{matrix} L_1 & J_1 & \frac{1}{2} \\ L_2 & J_2 & \frac{1}{2} \\ L & J & 1 \end{matrix} \right\} C_{J_1 -M_1 J_2 M_2}^{J M} \mathbf{Y}_{JM}^L(\vartheta, \varphi), \end{aligned} \quad (7.2.41)$$

where $\hat{S}$ is the spin operator, $\mathbf{Y}_{JM}^L$ is a vector spherical harmonic (see Sec. 7.3).

7.2.9 Addition Theorems

Let $\mathbf{n}_1$ and $\mathbf{n}_2$ be unit vectors with polar angles ϑ_1, φ_1 and ϑ_2, φ_2 , $\cos \omega_{12} \equiv \mathbf{n}_1 \cdot \mathbf{n}_2 = \cos \vartheta_1 \cos \vartheta_2 + \sin \vartheta_1 \sin \vartheta_2 \cos(\varphi_1 - \varphi_2)$. The addition theorem for the spinor spherical harmonics may be written in the form

$$4\pi \sum_{M=-J}^J \Omega_{JM}^{L+}(\vartheta_1, \varphi_1) \Omega_{JM}^{L'}(\vartheta_2, \varphi_2) = \delta_{LL'} (2J+1) P_L(\cos \omega_{12}), \quad (7.2.42)$$

$$4\pi \sum_{M=-J}^J \Omega_{JM}^{L+}(\vartheta_1, \varphi_1) \hat{S} \Omega_{JM}^{L'}(\vartheta_2, \varphi_2) = \begin{cases} i |\mathbf{n}_1 \times \mathbf{n}_2| P_L'(\cos \omega_{12}) & \text{if } L' = L \\ (L' - L) \{ \mathbf{n}_1 P_{L'}'(\cos \omega_{12}) - \mathbf{n}_2 P_{L'}'(\cos \omega_{12}) \}, & \text{if } L' = 2J - L \end{cases} \quad (7.2.43)$$

where $P_L(x)$ is a Legendre polynomial, $P'_L(x) = dP_L(x)/dx$, $\hat{\mathbf{S}}$ is the spin operator (Sec. 2.5). Another form of the addition theorem may be obtained for matrix products of the type

$$\begin{aligned} & 4\pi \sum_{M=-J}^J \Omega_{JM}^L(\vartheta_1, \varphi_1) \Omega_{JM}^{L+}(\vartheta_2, \varphi_2) \\ &= \left(J + \frac{1}{2}\right) \hat{I} P_L(\cos \omega_{12}) - i4(J-L) \hat{\mathbf{S}} \cdot [\mathbf{n}_1 \times \mathbf{n}_2] P'_L(\cos \omega_{12}), \end{aligned} \quad (7.2.44)$$

$$\begin{aligned} & 4\pi \sum_{M=-J}^J \Omega_{JM}^L(\vartheta_1, \varphi_1) \Omega_{JM}^{L'+}(\vartheta_2, \varphi_2) \\ &= 2(L' - L) \left\{ (\hat{\mathbf{S}} \cdot \mathbf{n}_1) P'_L(\cos \omega_{12}) - (\hat{\mathbf{S}} \cdot \mathbf{n}_2) P'_{L'}(\cos \omega_{12}) \right\}, \quad (L' = 2J - L). \end{aligned} \quad (7.2.45)$$

7.2.10 Quadratic Forms of Spinor Spherical Harmonics

The quadratic forms $\Omega_{JM}^{L+}(\vartheta, \varphi) \Omega_{JM}^L(\vartheta, \varphi)$ describe angular distributions of spin- $\frac{1}{2}$ particles in states with definite total angular momentum J , angular momentum projection M and orbital angular momentum L . Note that in fact these forms are independent of L and angle φ .

Let us denote

$$W_{JM}(\vartheta) \equiv \Omega_{JM}^{L+}(\vartheta, \varphi) \Omega_{JM}^L(\vartheta, \varphi). \quad (7.2.46)$$

Then one obtains

$$\begin{aligned} W_{JM}(\vartheta) &= \frac{1}{2(J+1)} \left\{ (J+M+1) \left| Y_{J+\frac{1}{2}, M+\frac{1}{2}}(\vartheta, \varphi) \right|^2 + (J-M+1) \left| Y_{J+\frac{1}{2}, M-\frac{1}{2}}(\vartheta, \varphi) \right|^2 \right\} \\ &= \frac{1}{2J} \left\{ (J+M) \left| Y_{J-\frac{1}{2}, M-\frac{1}{2}}(\vartheta, \varphi) \right|^2 + (J-M) \left| Y_{J-\frac{1}{2}, M+\frac{1}{2}}(\vartheta, \varphi) \right|^2 \right\}. \end{aligned} \quad (7.2.47)$$

An expansion of $W_{JM}(\vartheta)$ in terms of the Legendre polynomials may be written as

$$W_{JM}(\vartheta) = \sum_{n=0}^{J-\frac{1}{2}} a_n(J, M) P_{2n}(\cos \vartheta), \quad (7.2.48)$$

where

$$\begin{aligned} a_n(J, M) &= -\frac{\sqrt{2J(2J+1)}}{4\pi} (4n+1) \left\{ \begin{matrix} J & J & 2n \\ J-\frac{1}{2} & J-\frac{1}{2} & \frac{1}{2} \end{matrix} \right\} C_{J-\frac{1}{2}, 0, 2n, 0}^{J-\frac{1}{2}, 0} C_{J, M, 2n, 0}^{J, M} \\ &= (-1)^n \frac{(4n+1)(2J+2n+1)}{4\pi\sqrt{2J+1}} \cdot \frac{(J-\frac{1}{2}+n)! (2n)!}{(J-\frac{1}{2}-n)! (n!)^2} \sqrt{\frac{(2J-2n)!}{(2J+2n+1)!}} C_{J, M, 2n, 0}^{J, M}. \end{aligned} \quad (7.2.49)$$

In particular,

$$\begin{aligned} a_0(J, M) &= \frac{1}{4\pi}, \quad a_1(J, M) = \frac{5}{16\pi} \cdot \frac{J(J+1) - 3M^2}{J(J+1)}, \\ a_2(J, M) &= \frac{27}{256\pi} \cdot \frac{3(J^2 + 2J - 5M^2)(J^2 - 5M^2 - 1) - 10M^2(4M^2 - 1)}{(J-1)J(J+1)(J+2)}. \end{aligned} \quad (7.2.50)$$

The functions $W_{JM}(\vartheta)$ are normalized according to

$$\begin{aligned} & \int W_{JM}(\vartheta) d\Omega = 1, \\ & \sum_{M=-J}^J W_{JM}(\vartheta) = \frac{2J+1}{4\pi}. \end{aligned} \quad (7.2.51)$$

The symmetries of $W_{JM}(\vartheta)$ are

$$W_{JM}(\vartheta) = W_{J-M}(\vartheta) = W_{JM}(\pi - \vartheta) = W_{J-M}(\pi - \vartheta). \quad (7.2.52)$$

For $\vartheta = 0$ and $\vartheta = \pi$ we have

$$W_{JM}(0) = W_{JM}(\pi) = \begin{cases} \frac{2J+1}{8\pi}, & \text{if } M = \pm \frac{1}{2} \\ 0, & \text{otherwise.} \end{cases} \quad (7.2.53)$$

For special M values one obtains

$$W_{J\pm\frac{1}{2}}(\vartheta) = \frac{1}{2\pi(2J+1)} \left\{ \sin^2 \vartheta \left[P'_{J-\frac{1}{2}}(\cos \vartheta) \right]^2 + \left(J + \frac{1}{2} \right)^2 \left[P_{J-\frac{1}{2}}(\cos \vartheta) \right]^2 \right\}, \quad (7.2.54)$$

$$W_{J\pm(J-1)}(\vartheta) = \frac{(2J-1)!(\sin \vartheta)^{2J-3}}{\pi 2^{2J+1} \left[\left(J - \frac{1}{2} \right)! \right]^2} \left\{ 1 + 4J(J-1) \cos^2 \vartheta \right\}, \quad (7.2.55)$$

$$W_{J\pm J}(\vartheta) = \frac{(2J)!}{\pi 2^{2J+1} \left[\left(J - \frac{1}{2} \right)! \right]^2} \sin^{2J-1} \vartheta. \quad (7.2.56)$$

The explicit forms of $W_{JM}(\vartheta)$ for $J = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}, \frac{9}{2}, \frac{11}{2}$ ($\frac{1}{2} \leq M \leq J$) are given in Table 7.1. For negative M one may use the relation $W_{JM}(\vartheta) = W_{J-M}(\vartheta)$.

7.3 Vector spherical harmonics

7.3.1 Definition

The vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are defined as the tensor spherical harmonics at $S = 1$ (Sec. 7.1.1).

$$\mathbf{Y}_{JM}^L(\vartheta, \varphi) \equiv Y_{JM}^{L1}(\vartheta, \varphi). \quad (7.3.1)$$

They are eigenfunctions of the operators $\hat{\mathbf{J}}^2, \hat{J}_z, \hat{\mathbf{L}}^2, \hat{\mathbf{S}}^2$ where $\hat{\mathbf{L}}$ is the orbital angular momentum operator (Sec. 2.2), $\hat{\mathbf{S}}$ is the spin operator for $S = 1$ (Sec. 2.6), and $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ is the total angular momentum operator (Sec. 2.1)

$$\begin{aligned} \hat{\mathbf{J}}^2 \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= J(J+1) \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\ \hat{J}_z \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= M \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\ \hat{\mathbf{L}}^2 \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= L(L+1) \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\ \hat{\mathbf{S}}^2 \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= 2 \mathbf{Y}_{JM}^L(\vartheta, \varphi). \end{aligned} \quad (7.3.2)$$

According to (7.1.2), the vector spherical harmonics $\mathbf{Y}_{JM}^L$ may be written as

$$\mathbf{Y}_{JM}^L(\vartheta, \varphi) = \sum_{m, \sigma} C_{Lm1\sigma}^{JM} Y_{Lm}(\vartheta, \varphi) \mathbf{e}_\sigma, \quad (7.3.3)$$

where $C_{Lm1\sigma}^{JM}$ is the Clebsch-Gordan coefficient (Chap. 8), Y_{LM} is a scalar spherical harmonic (Chap. 5), and $\mathbf{e}_\sigma$ is a covariant spherical basis vector (spin function for an $S = 1$ particle, Sec. 1.1.3); J and L are nonnegative integers. For a given J , three values of L are possible: $L = J, J \pm 1$ (with the only exception that $L = 1$ for $J = 0$), and M takes the values $M = -J, -J+1, \dots, J-1, J$.

The vectors $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ which have the same J, M but different L satisfy the orthogonality relations

$$\sum_M \mathbf{Y}_{JM}^{L'*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^L(\vartheta, \varphi) = 0, \quad \text{if } L \neq L'. \quad (7.3.4)$$

Along with $\mathbf{Y}_{JM}^L$, other vector spherical harmonics $\mathbf{Y}_{JM}^{(\lambda)}(\lambda = 0, \pm 1)$ are widely used. The $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$'s (unlike $\mathbf{Y}_{JM}^L$) are not eigenfunctions of the operator $\hat{\mathbf{L}}^2$ but are suitably oriented with respect to the vector $\mathbf{n} \equiv \mathbf{r}/r$ specified by polar angles ϑ, φ . The vector spherical harmonics $\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)$ and $\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)$ are transverse, and $\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)$ is longitudinal with respect to $\mathbf{n}(\vartheta, \varphi)$.

$$\begin{aligned} \mathbf{n} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= \mathbf{n} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) = 0, \\ \mathbf{n} \times \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= 0. \end{aligned} \quad (7.3.5)$$

Sometimes $\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)$ and $\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)$ are called the electric and magnetic multipoles, respectively. They have the form

$$\begin{aligned} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= \frac{1}{\sqrt{J(J+1)}} \nabla_\Omega Y_{JM}(\vartheta, \varphi), \\ \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= \frac{-i}{\sqrt{J(J+1)}} (\mathbf{n} \times \nabla_\Omega) Y_{JM}(\vartheta, \varphi) = \frac{\hat{\mathbf{L}}}{\sqrt{J(J+1)}} Y_{JM}(\vartheta, \varphi). \end{aligned} \quad (7.3.6)$$

Table 7.1. Explicit forms of $W_{JM}(\vartheta)$.

J	M	$W_{JM}(\vartheta) = \Omega_{JM}^{L+}(\vartheta, \varphi) \Omega_{JM}^L(\vartheta, \varphi)$
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{4\pi}$
$\frac{3}{2}$	$\frac{1}{2}$	$\frac{1}{8\pi} \{3 \cos^2 \vartheta + 1\}$
$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{8\pi} \sin^2 \vartheta$
$\frac{5}{2}$	$\frac{1}{2}$	$\frac{3}{16\pi} \{5 \cos^4 \vartheta - 2 \cos^2 \vartheta + 1\}$
$\frac{5}{2}$	$\frac{3}{2}$	$\frac{3}{32\pi} \sin^2 \vartheta (15 \cos^2 \vartheta + 1)$
$\frac{5}{2}$	$\frac{5}{2}$	$\frac{15}{32\pi} \sin^4 \vartheta$
$\frac{7}{2}$	$\frac{1}{2}$	$\frac{1}{64\pi} \{175 \cos^6 \vartheta - 165 \cos^4 \vartheta + 45 \cos^2 \vartheta + 9\}$
$\frac{7}{2}$	$\frac{3}{2}$	$\frac{15}{64\pi} \sin^2 \vartheta \{21 \cos^4 \vartheta - 6 \cos^2 \vartheta + 1\}$
$\frac{7}{2}$	$\frac{5}{2}$	$\frac{5}{64\pi} \sin^4 \vartheta \{35 \cos^2 \vartheta + 1\}$
$\frac{7}{2}$	$\frac{7}{2}$	$\frac{35}{64\pi} \sin^6 \vartheta$
$\frac{9}{2}$	$\frac{1}{2}$	$\frac{5}{256\pi} \{441 \cos^8 \vartheta - 644 \cos^6 \vartheta + 294 \cos^4 \vartheta - 36 \cos^2 \vartheta + 9\}$
$\frac{9}{2}$	$\frac{3}{2}$	$\frac{15}{128\pi} \sin^2 \vartheta (147 \cos^6 \vartheta - 105 \cos^4 \vartheta + 21 \cos^2 \vartheta + 1)$
$\frac{9}{2}$	$\frac{5}{2}$	$\frac{35}{128\pi} \sin^4 \vartheta (45 \cos^4 \vartheta - 10 \cos^2 \vartheta + 1)$
$\frac{9}{2}$	$\frac{7}{2}$	$\frac{35}{512\pi} \sin^6 \vartheta \{63 \cos^2 \vartheta + 1\}$
$\frac{9}{2}$	$\frac{9}{2}$	$\frac{315}{512\pi} \sin^8 \vartheta$
$\frac{11}{2}$	$\frac{1}{2}$	$\frac{3}{512\pi} \{4851 \cos^{10} \vartheta - 9555 \cos^8 \vartheta + 6510 \cos^6 \vartheta - 1750 \cos^4 \vartheta + 175 \cos^2 \vartheta + 25\}$
$\frac{11}{2}$	$\frac{3}{2}$	$\frac{105}{512\pi} \sin^2 \vartheta \{297 \cos^8 \vartheta - 348 \cos^6 \vartheta + 126 \cos^4 \vartheta - 12 \cos^2 \vartheta + 1\}$
$\frac{11}{2}$	$\frac{5}{2}$	$\frac{105}{1024\pi} \sin^4 \vartheta \{495 \cos^6 \vartheta - 285 \cos^4 \vartheta + 45 \cos^2 \vartheta + 1\}$
$\frac{11}{2}$	$\frac{7}{2}$	$\frac{315}{1024\pi} \sin^6 \vartheta \{77 \cos^4 \vartheta - 14 \cos^2 \vartheta + 1\}$
$\frac{11}{2}$	$\frac{9}{2}$	$\frac{63}{1024\pi} \sin^8 \vartheta (99 \cos^2 \vartheta + 1)$
$\frac{11}{2}$	$\frac{11}{2}$	$\frac{693}{1024\pi} \sin^{10} \vartheta$

The longitudinal vector $\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)$ is given by

$$\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) = n Y_{JM}(\vartheta, \varphi). \quad (7.3.7)$$

In (7.3.6) ∇_Ω denotes the angular part of the ∇ operator (see Eqs. (1.3.9)-(1.3.10)), and $\hat{\mathbf{L}}$ is the orbital angular momentum operator. Three vectors $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ with different λ are mutually orthogonal:

$$\mathbf{Y}_{JM}^{(\lambda')}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) = 0, \quad \text{if } \lambda' \neq \lambda. \quad (7.3.8)$$

The vector spherical harmonics $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ are linear superpositions of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ with different L . Thus

$$\begin{aligned} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) + \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi), \\ \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= \mathbf{Y}_{JM}^J(\vartheta, \varphi), \\ \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi). \end{aligned} \quad (7.3.9)$$

The inverse relations are written as

$$\begin{aligned}\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi), \\ \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\ \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) + \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi).\end{aligned}\tag{7.3.10}$$

The vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$, as well as $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ constitute a complete orthonormal vector set for the range $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$ (see Sec. 7.3.13 below).

7.3.2 Components of Vector Spherical Harmonics

- (a) The vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ may be expanded in terms of the spherical covariant basis vectors $\mathbf{e}_\mu (\mu = \pm 1, 0)$ as

$$\mathbf{Y}_{JM}^L(\vartheta, \varphi) = \sum_{\mu} [\mathbf{Y}_{JM}^L(\vartheta, \varphi)]^{\mu} \mathbf{e}_{\mu} = \sum_{\mu} (-1)^{\mu} [\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_{-\mu} \mathbf{e}_{\mu},\tag{7.3.11}$$

where $[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]^{\mu}$ and $[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_{\mu}$ are contravariant and covariant components, respectively.

The contravariant spherical components of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are given, according to (7.1.5), by

$$[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]^{\mu} = C_{LM-\mu 1}^{JM} Y_{LM-\mu}(\vartheta, \varphi).\tag{7.3.12}$$

The covariant spherical components of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are related to the contravariant ones by

$$[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_{\mu} = (-1)^{\mu} [\mathbf{Y}_{JM}^L(\vartheta, \varphi)]^{-\mu},\tag{7.3.13}$$

and are given by

$$[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_{\mu} = (-1)^{\mu} C_{LM+\mu 1}^{JM} Y_{LM+\mu}(\vartheta, \varphi).\tag{7.3.14}$$

In more detailed form Eqs. (7.3.12), (7.3.14) may be written as

$$\begin{aligned}[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]^{+1} &= -[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]_{-1} = \left[\frac{(J-M+1)(J-M+2)}{2(J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1M-1}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]^0 &= [\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]_0 = - \left[\frac{(J-M+1)(J+M+1)}{(J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1M}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]^{-1} &= -[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]_{+1} = \left[\frac{(J+M+1)(J+M+2)}{2(J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1M+1}(\vartheta, \varphi),\end{aligned}\tag{7.3.15}$$

$$\begin{aligned}[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]^{+1} &= -[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]_{-1} = - \left[\frac{(J+M)(J-M+1)}{2J(J+1)} \right]^{\frac{1}{2}} Y_{JM-1}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^J(\vartheta, \varphi)]^0 &= [\mathbf{Y}_{JM}^J(\vartheta, \varphi)]_0 = \frac{M}{\sqrt{J(J+1)}} Y_{JM}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^J(\vartheta, \varphi)]^{-1} &= -[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]_{+1} = \left[\frac{(J-M)(J+M+1)}{2J(J+1)} \right]^{\frac{1}{2}} Y_{JM+1}(\vartheta, \varphi).\end{aligned}\tag{7.3.16}$$

$$\begin{aligned}[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]^{+1} &= -[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]_{-1} = \left[\frac{(J+M)(J+M-1)}{2J(2J-1)} \right]^{\frac{1}{2}} Y_{J-1M-1}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]^0 &= [\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]_0 = \left[\frac{(J-M)(J+M)}{J(2J-1)} \right]^{\frac{1}{2}} Y_{J-1M}(\vartheta, \varphi), \\ [\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]^{-1} &= -[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]_{+1} = \left[\frac{(J-M)(J-M-1)}{2J(2J-1)} \right]^{\frac{1}{2}} Y_{J-1M+1}(\vartheta, \varphi).\end{aligned}\tag{7.3.17}$$

(b) An expansion of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ in terms of the spherical covariant basis vectors $\mathbf{e}_\mu$ ($\mu = \pm 1, 0$) has the form

$$\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) = \sum_{\mu} \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]^{\mu} \mathbf{e}_{\mu} = \sum_{\mu} (-1)^{\mu} \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]_{-\mu} \mathbf{e}_{\mu}. \quad (7.3.18)$$

The contravariant spherical components of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ may be written as

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right]^{\mu} &= \sqrt{\frac{J+1}{2J+1}} C_{J-1 M-\mu 1}^{JM} Y_{J-1 M-\mu}(\vartheta, \varphi) + \sqrt{\frac{J}{2J+1}} C_{J+1 M-\mu 1}^{JM} Y_{J+1 M-\mu}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]^{\mu} &= C_{J M-\mu 1}^{JM} Y_{JM-\mu}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]^{\mu} &= \sqrt{\frac{J}{2J+1}} C_{J-1 M-\mu 1}^{JM} Y_{J-1 M-\mu}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} C_{J+1 M-\mu 1}^{JM} Y_{J+1 M-\mu}(\vartheta, \varphi). \end{aligned} \quad (7.3.19)$$

The covariant spherical components of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ are related to contravariant ones by

$$\left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]_{\mu} = (-1)^{\mu} \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]^{-\mu}, \quad (7.3.20)$$

and may be written as

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]^{+1} &= - \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]_{-1} = \left[\frac{(J+M)(J+M-1)(J+1)}{2J(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M-1}(\vartheta, \varphi), \\ &\quad + \left[\frac{(J-M+1)(J-M+2)J}{2(J+1)(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M-1}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]^0 &= \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]_0 = \left[\frac{(J-M)(J+M)(J+1)}{J(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M}(\vartheta, \varphi), \\ &\quad - \left[\frac{(J-M+1)(J+M+1)J}{(J+1)(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]^{-1} &= - \left[\mathbf{Y}_{JM}^{(+1)}(\vartheta, \varphi) \right]_{+1} = \left[\frac{(J-M)(J-M-1)(J+1)}{2J(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M+1}(\vartheta, \varphi), \\ &\quad + \left[\frac{(J+M+1)(J+M+2)J}{2(J+1)(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M+1}(\vartheta, \varphi). \end{aligned} \quad (7.3.21)$$

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]^{+1} &= - \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_{-1} = - \left[\frac{(J+M)(J-M+1)}{2J(J+1)} \right]^{\frac{1}{2}} Y_{JM-1}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]^0 &= \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_0 = \frac{M}{\sqrt{J(J+1)}} Y_{JM}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]^{-1} &= - \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_{+1} = \left[\frac{(J-M)(J+M+1)}{2J(J+1)} \right]^{\frac{1}{2}} Y_{JM+1}(\vartheta, \varphi), \end{aligned} \quad (7.3.22)$$

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]^{+1} &= - \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_{-1} = \left[\frac{(J+M)(J+M-1)}{2(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M-1}(\vartheta, \varphi), \\ &\quad - \left[\frac{(J-M+1)(J-M+2)}{2(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M-1}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]^0 &= \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_0 = \left[\frac{(J-M)(J+M)}{(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M}(\vartheta, \varphi), \\ &\quad + \left[\frac{(J-M+1)(J+M+1)}{(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]^{-1} &= - \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_{+1} = \left[\frac{(J-M)(J-M-1)}{2(2J-1)(2J+1)} \right]^{\frac{1}{2}} Y_{J-1 M+1}(\vartheta, \varphi), \\ &\quad - \left[\frac{(J+M+1)(J+M+2)}{2(2J+1)(2J+3)} \right]^{\frac{1}{2}} Y_{J+1 M+1}(\vartheta, \varphi). \end{aligned} \quad (7.3.23)$$

(c) An expansion of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ in terms of the polar basis vectors $\mathbf{e}_r$, $\mathbf{e}_\vartheta$ and $\mathbf{e}_\varphi$ (Sec. 1.1.2) is given by

$$\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) = \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]_r \mathbf{e}_r + \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]_\vartheta \mathbf{e}_\vartheta + \left[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \right]_\varphi \mathbf{e}_\varphi, \quad (7.3.24)$$

where the polar components of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ have the form

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right]_\sigma &= 0, \\ \left[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right]_\vartheta &= \frac{1}{\sqrt{J(J+1)}} \cdot \frac{\partial}{\partial \vartheta} Y_{JM}(\vartheta, \varphi), \\ &= \frac{1}{2} \sqrt{\frac{(J-M)(J+M+1)}{J(J+1)}} e^{-i\varphi} Y_{JM+1}(\vartheta, \varphi) - \frac{1}{2} \sqrt{\frac{(J+M)(J-M+1)}{J(J+1)}} e^{i\varphi} Y_{JM-1}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right]_\varphi &= \frac{1}{\sqrt{J(J+1)}} \cdot \frac{1}{\sin \vartheta} \cdot \frac{\partial}{\partial \varphi} Y_{JM}(\vartheta, \varphi) = \frac{iM}{\sqrt{J(J+1)}} \cdot \frac{1}{\sin \vartheta} Y_{JM}(\vartheta, \varphi). \end{aligned} \quad (7.3.25)$$

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_r &= 0, \\ \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_\vartheta &= \frac{i}{\sqrt{J(J+1)}} \cdot \frac{1}{\sin \vartheta} \cdot \frac{\partial}{\partial \varphi} Y_{JM}(\vartheta, \varphi) = \frac{-M}{\sqrt{J(J+1)}} \cdot \frac{1}{\sin \vartheta} Y_{JM}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right]_\varphi &= -\frac{i}{\sqrt{J(J+1)}} \cdot \frac{\partial}{\partial \vartheta} Y_{JM}(\vartheta, \varphi), \\ &= -\frac{i}{2} \sqrt{\frac{(J-M)(J+M+1)}{J(J+1)}} e^{-i\varphi} Y_{JM+1}(\vartheta, \varphi) + \frac{i}{2} \sqrt{\frac{(J+M)(J-M+1)}{J(J+1)}} e^{i\varphi} Y_{JM-1}(\vartheta, \varphi). \end{aligned} \quad (7.3.26)$$

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_r &= Y_{JM}(\vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_\vartheta &= 0, \\ \left[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right]_\varphi &= 0. \end{aligned} \quad (7.3.27)$$

(d) An expansion of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ in terms of helicity basis vectors $\mathbf{e}'_\nu(\vartheta, \varphi)$ (Sec. 1.1.4) may be written as

$$\mathbf{Y}_{JM}^L(\vartheta, \varphi) = \sum_\nu \left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]^\nu \mathbf{e}'_\nu(\vartheta, \varphi) = \sum_\nu (-1)^\nu \left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]'_{-\nu} \mathbf{e}'_\nu(\vartheta, \varphi), \quad (7.3.28)$$

where $\left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]^\nu$ and $\left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]'_\nu$ are contravariant and covariant helicity components, respectively.

$$\left| \mathbf{Y}_{JM}^L(\vartheta, \varphi) \right|'^\nu = \sqrt{\frac{2L+1}{4\pi}} C_{L01\nu}^J D_{-\nu-M}^J(0, \vartheta, \varphi). \quad (7.3.29)$$

The covariant helicity components of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are related to contravariant ones by

$$\left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]'_\nu = (-1)^\nu \left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]'^{-\nu}, \quad (7.3.30)$$

and may be written as

$$\left[\mathbf{Y}_{JM}^L(\vartheta, \varphi) \right]'_\nu = (-1)^{J+L+\nu+1} \sqrt{\frac{2L+1}{4\pi}} C_{L01\nu}^J D_{\nu-M}^J(0, \vartheta, \varphi). \quad (7.3.31)$$

Explicitly we may write Eqs. (7.3.29)-(7.3.31) as follows:

$$\begin{aligned} \left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'^{+1} &= - \left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'_{-1} = \sqrt{\frac{J}{8\pi}} D_{-1-M}^J(0, \vartheta, \varphi), \\ \left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'^0 &= \left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'_0 = -\sqrt{\frac{J+1}{4\pi}} D_{0-M}^J(0, \vartheta, \varphi), \end{aligned} \quad (7.3.32)$$

$$\left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'^{-1} = - \left[\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \right]'_{+1} = \sqrt{\frac{J}{8\pi}} D_{1-M}^J(0, \vartheta, \varphi),$$

$$\begin{aligned}
[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'^{+1} &= -[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'_{-1} = -\sqrt{\frac{2J+1}{8\pi}} D_{-1-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'^0 &= [\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'_0 = 0, \\
[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'^{-1} &= -[\mathbf{Y}_{JM}^J(\vartheta, \varphi)]'_{+1} = \sqrt{\frac{2J+1}{8\pi}} D_{1-M}^J(0, \vartheta, \varphi).
\end{aligned} \tag{7.3.33}$$

$$\begin{aligned}
[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'^{+1} &= -[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'_{-1} = \sqrt{\frac{J+1}{8\pi}} D_{-1-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'^0 &= [\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'_0 = \sqrt{\frac{J}{4\pi}} D_{0-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'^{-1} &= -[\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]'_{+1} = \sqrt{\frac{J+1}{8\pi}} D_{1-M}^J(0, \vartheta, \varphi).
\end{aligned} \tag{7.3.34}$$

(e) An expansion of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ in terms of the helicity basis vectors $\mathbf{e}'_\nu(\vartheta, \varphi)$ ($\nu = \pm 1, 0$) has the form

$$\begin{aligned}
\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= \sqrt{\frac{2J+1}{8\pi}} \{D_{-1-M}^J(0, \vartheta, \varphi) \mathbf{e}'_1(\vartheta, \varphi) + D_{1-M}^J(0, \vartheta, \varphi) \mathbf{e}'_{-1}(\vartheta, \varphi)\}, \\
\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= \sqrt{\frac{2J+1}{8\pi}} \{-D_{-1-M}^J(0, \vartheta, \varphi) \mathbf{e}'_1(\vartheta, \varphi) + D_{1-M}^J(0, \vartheta, \varphi) \mathbf{e}'_{-1}(\vartheta, \varphi)\}, \\
\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= \sqrt{\frac{2J+1}{4\pi}} D_{0-M}^J(0, \vartheta, \varphi) \mathbf{e}'_0(\vartheta, \varphi).
\end{aligned} \tag{7.3.35}$$

The contravariant and covariant helicity components, $[\mathbf{Y}_{JM}^{(\lambda)}]^\nu$ and $[\mathbf{Y}_{JM}^{(\lambda)}]_\nu'$, are given by

$$\begin{aligned}
[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'^{+1} &= -[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'_{-1} = \sqrt{\frac{2J+1}{8\pi}} D_{-1-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'^0 &= [\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'_0 = 0, \\
[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'^{-1} &= -[\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]'_{+1} = \sqrt{\frac{2J+1}{8\pi}} D_{1-M}^J(0, \vartheta, \varphi),
\end{aligned} \tag{7.3.36}$$

$$\begin{aligned}
[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'^{+1} &= -[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'_{-1} = -\sqrt{\frac{2J+1}{8\pi}} D_{-1-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'^0 &= [\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'_0 = 0, \\
[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'^{-1} &= -[\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]'_{+1} = \sqrt{\frac{2J+1}{8\pi}} D_{1-M}^J(0, \vartheta, \varphi),
\end{aligned} \tag{7.3.37}$$

$$\begin{aligned}
[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'^{+1} &= -[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'_{-1} = 0, \\
[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'^0 &= [\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'_0 = \sqrt{\frac{2J+1}{4\pi}} D_{0-M}^J(0, \vartheta, \varphi), \\
[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'^{-1} &= -[\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]'_{+1} = 0.
\end{aligned} \tag{7.3.38}$$

7.3.3 Complex Conjugation

The vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ and $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ are complex vectors. Under complex conjugation they transform as follows

$$\begin{aligned}
\mathbf{Y}_{JM}^{L*}(\vartheta, \varphi) &= (-1)^{J+L+M+1} \mathbf{Y}_{J-M}^L(\vartheta, \varphi), \\
\mathbf{Y}_{JM}^{(\lambda)*}(\vartheta, \varphi) &= (-1)^{M+\lambda+1} \mathbf{Y}_{J-M}^{(\lambda)}(\vartheta, \varphi).
\end{aligned} \tag{7.3.39}$$

The complex conjugates of the spherical components of $\mathbf{Y}_{JM}^L$ and $\mathbf{Y}_{JM}^{(\lambda)}$ are given by

$$[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_\mu^* = [\mathbf{Y}_{JM}^{L*}(\vartheta, \varphi)]_\mu = (-1)^{J+L+M+\mu+1} [\mathbf{Y}_{J-M}^L(\vartheta, \varphi)]_{-\mu} = (-1)^{J+L+M+1} [\mathbf{Y}_{J-M}^L(\vartheta, \varphi)]_\mu, \quad (7.3.40)$$

$$[\mathbf{Y}_{JM}^L(\vartheta, \varphi)]^{\mu*} = [\mathbf{Y}_{JM}^{L*}(\vartheta, \varphi)]_\mu = (-1)^{J+L+M+\mu+1} [\mathbf{Y}_{J-M}^L(\vartheta, \varphi)]^{-\mu} = (-1)^{J+L+M+1} [\mathbf{Y}_{J-M}^L(\vartheta, \varphi)]_\mu, \quad (7.3.41)$$

$$[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)]_\mu^* = [\mathbf{Y}_{JM}^{(\lambda)*}(\vartheta, \varphi)]_\mu = (-1)^{M+\mu+\lambda+1} [\mathbf{Y}_{J-M}^{(\lambda)}(\vartheta, \varphi)]_{-\mu} = (-1)^{M+\lambda+1} [\mathbf{Y}_{J-M}^{(\lambda)}(\vartheta, \varphi)]_\mu, \quad (7.3.42)$$

$$[\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)]^{\mu*} = [\mathbf{Y}_{JM}^{(\lambda)*}(\vartheta, \varphi)]_\mu = (-1)^{M+\mu+\lambda+1} [\mathbf{Y}_{J-M}^{(\lambda)}(\vartheta, \varphi)]^{-\mu} = (-1)^{M+\lambda+1} [\mathbf{Y}_{J-M}^{(\lambda)}(\vartheta, \varphi)]_\mu. \quad (7.3.43)$$

7.3.4 Transformations of Coordinate Systems

(a) *Coordinate inversion*

$$\begin{aligned} \hat{P}_r \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^L(\pi - \vartheta, \pi + \varphi) = (-1)^{L+1} \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\ \hat{P}_r \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(\lambda)}(\pi - \vartheta, \pi + \varphi) = (-1)^{J+\lambda+1} \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi). \end{aligned} \quad (7.3.44)$$

(b) *Rotations of the coordinate system*

Under rotations specified by the Euler angles α, β, γ (Sec. 1.4) the vector spherical harmonics transform according to

$$\begin{aligned} \mathbf{Y}_{JM'}^L(\vartheta', \varphi') &= \hat{D}(\alpha, \beta, \gamma) \mathbf{Y}_{JM'}^L(\vartheta, \varphi) = \sum_M D_{MM'}^J(\alpha, \beta, \gamma) \mathbf{Y}_{JM}^L(\vartheta, \varphi) \\ \mathbf{Y}_{JM'}^{(\lambda)}(\vartheta', \varphi') &= \hat{D}(\alpha, \beta, \gamma) \mathbf{Y}_{JM'}^{(\lambda)}(\vartheta, \varphi) = \sum_M D_{MM'}^J(\alpha, \beta, \gamma) \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi). \end{aligned} \quad (7.3.45)$$

where $D_{MM'}^J(\alpha, \beta, \gamma)$ are the Wigner D -functions (Chap. 4), M' is the projection of total angular momentum on the new z' -axis; ϑ, φ and ϑ', φ' are polar angles of r in the initial and final coordinate systems, respectively. The relations between ϑ', φ' and ϑ, φ are given by eqs. (1.4.2), (1.4.3).

7.3.5 Differential Equations

(a) The vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are eigenfunctions of the operator $\hat{\mathbf{L}}^2$; hence, they satisfy the equation

$$\frac{1}{\sin \vartheta} \cdot \frac{\partial}{\partial \vartheta} \left\{ \sin \vartheta \frac{\partial}{\partial \vartheta} \mathbf{Y}_{JM}^L(\vartheta, \varphi) \right\} + \frac{1}{\sin^2 \vartheta} \frac{\partial^2}{\partial \varphi^2} \mathbf{Y}_{JM}^L(\vartheta, \varphi) + L(L+1) \mathbf{Y}_{JM}^L(\vartheta, \varphi) = 0. \quad (7.3.46)$$

This equation may be rewritten in a compact form as

$$\{\Delta_\Omega + L(L+1)\} \mathbf{Y}_{JM}^L(\vartheta, \varphi) = 0, \quad (7.3.47)$$

where $\Delta_\Omega \equiv \nabla_\Omega^2$ is the angular part of the Laplace operator (see Eq. (1.3.15)).

(b) The vector functions $r^L \mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are solutions of the Laplace equation

$$\Delta \{r^L \mathbf{Y}_{JM}^L(\vartheta, \varphi)\} = 0. \quad (7.3.48)$$

It follows from (7.3.48) that the components of $r^L \mathbf{Y}_{JM}^L(\vartheta, \varphi)$ are harmonic polynomials of degree L .

(c) The vectors $z_L(kr) \mathbf{Y}_{JM}^L(\vartheta, \varphi)$ satisfy the Helmholtz equation

$$(\Delta + k^2) \{z_L(kr) \mathbf{Y}_{JM}^L(\vartheta, \varphi)\} = 0, \quad (7.3.49)$$

where

$$z_L(kr) = \sqrt{\frac{\pi}{2kr}} Z_{L+\frac{1}{2}}(kr), \quad (7.3.50)$$

and $Z_{L+\frac{1}{2}}$ is any of the cylinder functions of order $L + \frac{1}{2}$.

7.3.6 Differential Operations

Below we present the results of the action of the ∇ operator on scalar and vector spherical harmonics; here $f(r)$ denotes an arbitrary function of $r \equiv |r|$.

(a)

$$\begin{aligned} \nabla \{f(r)Y_{JM}(\vartheta, \varphi)\} &\equiv \text{grad} \{f(r)Y_{JM}(\vartheta, \varphi)\} \\ &= \sqrt{\frac{J}{2J+1}} \left\{ \frac{d}{dr} + \frac{J+1}{r} \right\} f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} \left\{ \frac{d}{dr} - \frac{J}{r} \right\} f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi), \end{aligned} \quad (7.3.51)$$

$$\begin{aligned} \nabla \{f(r)Y_{JM}(\vartheta, \varphi)\} &\equiv \text{grad} \{f(r)Y_{JM}(\vartheta, \varphi)\} = \frac{df(r)}{dr} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) + \sqrt{J(J+1)} \frac{1}{r} f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi), \\ (\mathbf{r} \cdot \nabla) \{f(r) \mathbf{Y}_{JM}^L(\vartheta, \varphi)\} &= r \frac{d}{dr} f(r) \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\ (\mathbf{r} \cdot \nabla) \{f(r) \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)\} &= r \frac{d}{dr} f(r) \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi). \end{aligned} \quad (7.3.52)$$

(b)

$$\begin{aligned} \nabla \cdot [f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)] &\equiv \text{div} [f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)] = -\sqrt{\frac{J+1}{2J+1}} \left(\frac{d}{dr} + \frac{J+2}{r} \right) f(r) Y_{JM}(\vartheta, \varphi), \\ \nabla \cdot [f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi)] &\equiv \text{div} [f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi)] = 0, \end{aligned} \quad (7.3.53)$$

$$\nabla \cdot [f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)] \equiv \text{div} [f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)] = \sqrt{\frac{J}{2J+1}} \left(\frac{d}{dr} - \frac{J-1}{r} \right) f(r) Y_{JM}(\vartheta, \varphi),$$

$$\begin{aligned} \nabla \cdot [f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)] &\equiv \text{div} [f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)] = -\sqrt{J(J+1)} \frac{1}{r} f(r) Y_{JM}(\vartheta, \varphi), \\ \nabla \cdot [f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)] &\equiv \text{div} [f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)] = 0, \end{aligned} \quad (7.3.54)$$

$$\nabla \cdot [f(r) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)] \equiv \text{div} [f(r) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)] = \left(\frac{d}{dr} + \frac{2}{r} \right) f(r) Y_{JM}(\vartheta, \varphi).$$

(c)

$$\begin{aligned} \nabla \times [f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)] &\equiv \text{curl} [f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)] = i \sqrt{\frac{J}{2J+1}} \left(\frac{d}{dr} + \frac{J+2}{r} \right) f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi), \\ \nabla \times [f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi)] &\equiv \text{curl} [f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi)], \\ &= i \sqrt{\frac{J}{2J+1}} \left(\frac{d}{dr} - \frac{J}{r} \right) f(r) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) + i \sqrt{\frac{J+1}{2J+1}} \left(\frac{d}{dr} + \frac{J+1}{r} \right) f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi), \end{aligned} \quad (7.3.55)$$

$$\nabla \times [f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)] \equiv \text{curl} [f(r) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)] = i \sqrt{\frac{J+1}{2J+1}} \left(\frac{d}{dr} - \frac{J-1}{r} \right) f(r) \mathbf{Y}_{JM}^J(\vartheta, \varphi),$$

$$\begin{aligned} \nabla \times [f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)] &\equiv \text{curl} [f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)] = i \left(\frac{d}{dr} + \frac{1}{r} \right) f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\ \nabla \times [f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)] &\equiv \text{curl} [f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)], \\ &= i \left(\frac{d}{dr} + \frac{1}{r} \right) f(r) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) + i \sqrt{J(J+1)} \frac{1}{r} f(r) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi), \\ \nabla \times [f(r) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)] &\equiv \text{curl} [f(r) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)] = -i \sqrt{J(J+1)} \frac{1}{r} f(r) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi). \end{aligned} \quad (7.3.56)$$

(d) Some applications of spherical harmonics to physical problems involve the functions

$$\begin{aligned} F_{JM}(\mathbf{r}) &\equiv z_J(kr) Y_{JM}(\vartheta, \varphi), \\ \mathbf{F}_{JM}^L(\mathbf{r}) &\equiv z_L(kr) \mathbf{Y}_{JM}^L(\vartheta, \varphi). \end{aligned} \quad (7.3.57)$$

where $z_L(kr) \equiv \sqrt{\pi/(2kr)}Z_{L+\frac{1}{2}}(kr)$, and $Z_{L+\frac{1}{2}}$ is any cylinder functions; k is an arbitrary parameter. For these special functions Eqs. (7.3.51)-(7.3.56) are reduced to the following

$$\begin{aligned} \frac{1}{k} \text{grad} \{F_{JM}(\mathbf{r})\} &= \sqrt{\frac{J}{2J+1}} z_{J-1}(kr) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) + \sqrt{\frac{J+1}{2J+1}} z_{J+1}(kr) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) \\ &= \sqrt{\frac{J}{2J+1}} \mathbf{F}_{JM}^{J-1}(\mathbf{r}) + \sqrt{\frac{J+1}{2J+1}} \mathbf{F}_{JM}^{J+1}(\mathbf{r}), \end{aligned} \quad (7.3.58)$$

$$\begin{aligned} \frac{1}{k} \text{div} \{\mathbf{F}_{JM}^{J+1}(\mathbf{r})\} &= -\sqrt{\frac{J+1}{2J+1}} z_J(kr) Y_{JM}(\vartheta, \varphi) = -\sqrt{\frac{J+1}{2J+1}} F_{JM}(\mathbf{r}), \\ \frac{1}{k} \text{div} \{\mathbf{F}_{JM}^J(\mathbf{r})\} &= 0, \\ \frac{1}{k} \text{div} \{\mathbf{F}_{JM}^{J-1}(\mathbf{r})\} &= -\sqrt{\frac{J}{2J+1}} z_J(kr) Y_{JM}(\vartheta, \varphi) = -\sqrt{\frac{J}{2J+1}} F_{JM}(\mathbf{r}), \end{aligned} \quad (7.3.59)$$

$$\begin{aligned} \frac{1}{k} \text{curl} \{\mathbf{F}_{JM}^{J+1}(\mathbf{r})\} &= i\sqrt{\frac{J}{2J+1}} z_J(kr) \mathbf{Y}_{JM}^J(\vartheta, \varphi) = i\sqrt{\frac{J}{2J+1}} \mathbf{F}_{JM}^J(\mathbf{r}), \\ \frac{1}{k} \text{curl} \{\mathbf{F}_{JM}^J(\mathbf{r})\} &= -i\sqrt{\frac{J}{2J+1}} z_{J+1}(kr) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) + i\sqrt{\frac{J+1}{2J+1}} z_{J-1}(kr) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) \\ &= -i\sqrt{\frac{J}{2J+1}} \mathbf{F}_{JM}^{J+1}(\mathbf{r}) + i\sqrt{\frac{J+1}{2J+1}} \mathbf{F}_{JM}^{J-1}(\mathbf{r}), \\ \frac{1}{k} \text{curl} \{\mathbf{F}_{JM}^{J-1}(\mathbf{r})\} &= -i\sqrt{\frac{J+1}{2J+1}} z_J(kr) \mathbf{Y}_{JM}^J(\vartheta, \varphi) = -i\sqrt{\frac{J+1}{2J+1}} \mathbf{F}_{JM}^J(\mathbf{r}). \end{aligned} \quad (7.3.60)$$

7.3.7 Action of Angular Momentum Operators

The action of the angular momentum operators on vector spherical harmonics is given by Eqs. (7.1.27) and (7.1.29) for $S = 1$. In addition, we note the following relations in which $\hat{\mathbf{S}}, \hat{\mathbf{L}}$ and $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ are the spin, orbital and total angular momentum operators, respectively; $\mathbf{n}$ is the unit vector defined by the polar angles ϑ, φ .

(a)

$$\begin{aligned} (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= -\frac{1}{2} \sqrt{\frac{(J+L+3)(J+L)(-J+L+2)(J-L+1)}{(2L+1)(2L+3)}} \mathbf{Y}_{JM}^{L+1}(\vartheta, \varphi), \\ &\quad -\frac{1}{2} \sqrt{\frac{(J+L+2)(J+L-1)(-J+L+1)(J-L+2)}{(2L-1)(2L+1)}} \mathbf{Y}_{JM}^{L-1}(\vartheta, \varphi). \end{aligned} \quad (7.3.61)$$

In particular,

$$\begin{aligned} (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= -\sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi), \\ (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= -\sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi), \\ (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= -\sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi), \end{aligned} \quad (7.3.62)$$

$$\begin{aligned} (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\ (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi), \\ (\hat{\mathbf{S}} \cdot \mathbf{n}) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= 0. \end{aligned} \quad (7.3.63)$$

(b)

$$\begin{aligned} r(\hat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^L(\vartheta, \varphi) &= \frac{L}{2} \sqrt{\frac{(J+L+3)(J+L)(-J+L+2)(J-L+1)}{(2L+1)(2L+3)}} \mathbf{Y}_{JM}^{L+1}(\vartheta, \varphi), \\ &\quad -\frac{L+1}{2} \sqrt{\frac{(J+L+2)(J+L-1)(-J+L+1)(J-L+2)}{(2L-1)(2L+1)}} \mathbf{Y}_{JM}^{L-1}(\vartheta, \varphi). \end{aligned} \quad (7.3.64)$$

In particular,

$$\begin{aligned}
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= -(J+2) \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi), \\
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= J \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) - (J+1) \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi), \\
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= (J-1) \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi),
\end{aligned} \tag{7.3.65}$$

$$\begin{aligned}
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) - \sqrt{J(J+1)} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi), \\
r(\widehat{\mathbf{S}} \cdot \nabla) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= \sqrt{J(J+1)} \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi).
\end{aligned} \tag{7.3.66}$$

(c)

$$(\widehat{\mathbf{S}} \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^L(\vartheta, \varphi) = \frac{1}{2} \{J(J+1) - L(L+1) - 2\} \mathbf{Y}_{JM}^L(\vartheta, \varphi). \tag{7.3.67}$$

In particular,

$$\begin{aligned}
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= -(J+2) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi), \\
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^J(\vartheta, \varphi), \\
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= (J-1) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi),
\end{aligned} \tag{7.3.68}$$

$$\begin{aligned}
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) + \sqrt{J(J+1)} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi), \\
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= -\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\
(\widehat{\mathbf{S}} \cdot \widehat{\mathbf{L}}) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= \sqrt{J(J+1)} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) - 2\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi).
\end{aligned} \tag{7.3.69}$$

7.3.8 Algebraic Relations

In the equations given below $\mathbf{n}$ is the unit vector specified by the polar angles ϑ, φ .

(a)

$$\begin{aligned}
\mathbf{n} Y_{JM}(\vartheta, \varphi) &= \sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) - \sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi), \\
\mathbf{n} Y_{JM}(\vartheta, \varphi) &= \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi).
\end{aligned} \tag{7.3.70}$$

(b)

$$\begin{aligned}
\mathbf{n} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= -\sqrt{\frac{J+1}{2J+1}} Y_{JM}(\vartheta, \varphi), \\
\mathbf{n} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= 0, \\
\mathbf{n} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= \sqrt{\frac{J}{2J+1}} Y_{JM}(\vartheta, \varphi).
\end{aligned} \tag{7.3.71}$$

$$\begin{aligned}
\mathbf{n} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= 0, \\
\mathbf{n} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= 0, \\
\mathbf{n} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= Y_{JM}(\vartheta, \varphi).
\end{aligned} \tag{7.3.72}$$

(c)

$$\begin{aligned}
\mathbf{n} \times \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= i\sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi), \\
\mathbf{n} \times \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= i\sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) + i\sqrt{\frac{J}{2J+1}} \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi), \\
\mathbf{n} \times \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= i\sqrt{\frac{J+1}{2J+1}} \mathbf{Y}_{JM}^J(\vartheta, \varphi),
\end{aligned} \tag{7.3.73}$$

$$\begin{aligned}
\mathbf{n} \times \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) &= i\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi), \\
\mathbf{n} \times \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) &= i\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi), \\
\mathbf{n} \times \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) &= 0.
\end{aligned} \tag{7.3.74}$$

7.3.9 Sums of Vector Spherical Harmonics

(a)

$$\begin{aligned}
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_{\mu} &= \sqrt{\frac{3}{4\pi}} \mathbf{n} \cdot \mathbf{Y}_{JM}^L(\vartheta, \varphi), \\
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)]_{\mu} &= \sqrt{\frac{3}{4\pi}} \mathbf{n} \cdot \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi).
\end{aligned} \tag{7.3.75}$$

From (7.3.75) one obtains

$$\begin{aligned}
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)]_{\mu} &= -\sqrt{\frac{3(J+1)}{4\pi(2J+1)}} Y_{JM}(\vartheta, \varphi), \\
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^J(\vartheta, \varphi)]_{\mu} &= 0, \\
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)]_{\mu} &= \sqrt{\frac{3J}{4\pi(2J+1)}} Y_{JM}(\vartheta, \varphi).
\end{aligned} \tag{7.3.76}$$

$$\begin{aligned}
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)]_{\mu} &= 0, \\
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)]_{\mu} &= 0, \\
\sum_{\mu=-1}^1 Y_{1\mu}^*(\vartheta, \varphi) [\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)]_{\mu} &= \sqrt{\frac{3}{4\pi}} Y_{JM}(\vartheta, \varphi).
\end{aligned} \tag{7.3.77}$$

(b)

$$\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^L(\vartheta, \varphi) = \frac{\sqrt{(2J+1)(2L+1)}}{4\pi} C_{10L0}^{J0} \mathbf{n}. \tag{7.3.78}$$

In more detailed form (7.3.78) reads

$$\begin{aligned}
\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi) &= -\frac{\sqrt{(J+1)(2J+1)}}{4\pi} \mathbf{n}, \\
\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^J(\vartheta, \varphi) &= 0, \\
\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi) &= \frac{\sqrt{J(2J+1)}}{4\pi} \mathbf{n}.
\end{aligned} \tag{7.3.79}$$

Analogous relations for $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ have the form

$$\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) = \sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) = 0, \quad (7.3.80)$$

$$\sum_{M=-J}^J Y_{JM}^*(\vartheta, \varphi) \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) = \frac{2J+1}{4\pi} \mathbf{n}.$$

(c)

$$\sum_{M=-J}^J \mathbf{Y}_{JM}^{L'*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^L(\vartheta, \varphi) = \frac{2J+1}{4\pi} \delta_{LL'}, \quad (7.3.81)$$

$$\sum_{M=-J}^J \mathbf{Y}_{JM}^{(\lambda')*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) = \frac{2J+1}{4\pi} \delta_{\lambda\lambda'}.$$

(d) Below $\mathbf{a}$ is an arbitrary complex vector, $\mathbf{n}$ is the unit vector specified by the polar angles ϑ, φ .

$$\sum_{M=-J}^J |\mathbf{a} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)|^2 = \frac{1}{8\pi} \{J|\mathbf{a}|^2 + (J+2)|\mathbf{n} \cdot \mathbf{a}|^2\}, \quad (7.3.82)$$

$$\sum_{M=-J}^J |\mathbf{a} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi)|^2 = \frac{2J+1}{8\pi} \{|\mathbf{a}|^2 - |\mathbf{n} \cdot \mathbf{a}|^2\}, \quad (7.3.83)$$

$$\sum_{M=-J}^J |\mathbf{a} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)|^2 = \frac{1}{8\pi} \{(J+1)|\mathbf{a}|^2 + (J-1)|\mathbf{n} \cdot \mathbf{a}|^2\}, \quad (7.3.84)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi)) = -i \frac{\sqrt{J(2J+1)}}{8\pi} \mathbf{n} \cdot [\mathbf{a}^* \times \mathbf{a}], \quad (7.3.85)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)) = \frac{\sqrt{J(J+1)}}{8\pi} \{|\mathbf{a}|^2 - 3|\mathbf{n} \cdot \mathbf{a}|^2\}, \quad (7.3.86)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)) = -i \frac{\sqrt{J(2J+1)}}{8\pi} \mathbf{n} \cdot [\mathbf{a}^* \times \mathbf{a}], \quad (7.3.87)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi)) = -i \frac{\sqrt{(J+1)(2J+1)}}{8\pi} \mathbf{n} \cdot [\mathbf{a}^* \times \mathbf{a}]. \quad (7.3.88)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta, \varphi)) = \frac{\sqrt{J(J+1)}}{8\pi} \{|\mathbf{a}|^2 - 3|\mathbf{n} \cdot \mathbf{a}|^2\}, \quad (7.3.89)$$

$$\sum_{M=-J}^J (\mathbf{a} \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta, \varphi))^* (\mathbf{a} \cdot \mathbf{Y}_{JM}^J(\vartheta, \varphi)) = -i \frac{\sqrt{(J+1)(2J+1)}}{8\pi} \mathbf{n} \cdot [\mathbf{a}^* \times \mathbf{a}]. \quad (7.3.90)$$

(e) Analogous relations for $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ have the form:

$$\sum_{M=-J}^J \left| \mathbf{a} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right|^2 = \frac{2J+1}{8\pi} \{ |\mathbf{a}|^2 - |\mathbf{n} \cdot \mathbf{a}|^2 \}, \quad (7.3.91)$$

$$\sum_{M=-J}^J \left| \mathbf{a} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right|^2 = \frac{2J+1}{8\pi} \{ |\mathbf{a}|^2 - |\mathbf{n} \cdot \mathbf{a}|^2 \}, \quad (7.3.92)$$

$$\sum_{M=-J}^J \left| \mathbf{a} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right|^2 = \frac{2J+1}{4\pi} |\mathbf{n} \cdot \mathbf{a}|^2, \quad (7.3.93)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right) = -i \frac{2J+1}{8\pi} \mathbf{n} \cdot [\mathbf{a}^* \times \mathbf{a}], \quad (7.3.94)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right) = 0, \quad (7.3.95)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right) = -i \frac{2J+1}{8\pi} \mathbf{n} [\mathbf{a}^* \times \mathbf{a}], \quad (7.3.96)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right) = 0, \quad (7.3.97)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) \right) = 0, \quad (7.3.98)$$

$$\sum_{M=-J}^J \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) \right)^* \left(\mathbf{a} \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) \right) = 0. \quad (7.3.99)$$

7.3.10 Clebsch-Gordan Series

$$\begin{aligned} & \mathbf{Y}_{J_1 M_1}^{L_1}(\vartheta, \varphi) \cdot \mathbf{Y}_{J_2 M_2}^{L_2}(\vartheta, \varphi) \\ &= \sum_L (-1)^{J_2+L_1+L} \sqrt{\frac{(2J_1+1)(2J_2+1)(2L_1+1)(2L_2+1)}{4\pi(2L+1)}} \left\{ \begin{matrix} L_1 & L_2 & L \\ J_2 & J_1 & 1 \end{matrix} \right\} C_{L_1 0 L_2 0}^{L 0} C_{J_1 M_1 J_2 M_2}^{L M} Y_{LM}(\vartheta, \varphi), \end{aligned} \quad (7.3.100)$$

$$\begin{aligned} & \mathbf{Y}_{J_1 M_1}^{L_1}(\vartheta, \varphi) \times \mathbf{Y}_{J_2 M_2}^{L_2}(\vartheta, \varphi) = i \sqrt{\frac{3}{2\pi}} (2J_1+1)(2J_2+1)(2L_1+1)(2L_2+1) \\ & \times \sum_{JL} \left\{ \begin{matrix} J_1 & L_1 & 1 \\ J_2 & L_2 & 1 \\ J & L & 1 \end{matrix} \right\} C_{L_1 0 L_2 0}^{L 0} C_{J_1 M_1 J_2 M_2}^{J M} \mathbf{Y}_{JM}^L(\vartheta, \varphi). \end{aligned} \quad (7.3.101)$$

7.3.11 Addition Theorems for Vector Spherical Harmonics

Let $\mathbf{n}_1$ and $\mathbf{n}_2$ be the unit vectors determined by the polar angles ϑ_1, φ_1 and ϑ_2, φ_2 , respectively; $\cos \omega_{12} \equiv \mathbf{n}_1 \cdot \mathbf{n}_2 = \cos \vartheta_1 \cos \vartheta_2 + \sin \vartheta_1 \sin \vartheta_2 \cos(\varphi_1 - \varphi_2)$. Then we have the following addition theorems for the vector spherical harmonics $\mathbf{Y}_{JM}^L$.

(a)

$$4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{L'*}(\vartheta_1, \varphi_1) \cdot \mathbf{Y}_{JM}^L(\vartheta_2, \varphi_2) = \delta_{L'L}(2J+1)P_L(\cos \omega_{12}). \quad (7.3.102)$$

(b)

$$\begin{aligned}
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J+1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2) &= \frac{2J+1}{J+1} [\mathbf{n}_1 \times \mathbf{n}_2] P'_{J+1}(\cos \omega_{12}), \\
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J+1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2) &= -i \frac{\sqrt{J(2J+1)}}{J+1} \{ \mathbf{n}_1 P'_{J+1}(\cos \omega_{12}) - \mathbf{n}_2 P'_J(\cos \omega_{12}) \}, \\
\end{aligned} \tag{7.3.103}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J+1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2) &= 0, \\
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2) &= i \frac{\sqrt{J(2J+1)}}{J+1} \{ \mathbf{n}_1 P'_J(\cos \omega_{12}) - \mathbf{n}_2 P'_{J+1}(\cos \omega_{12}) \}, \\
\end{aligned}$$

$$4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2) = \frac{(2J+1)}{J(J+1)} [\mathbf{n}_1 \times \mathbf{n}_2] P'_J(\cos \omega_{12}), \tag{7.3.104}$$

$$4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2) = -i \frac{\sqrt{(J+1)(2J+1)}}{J} \{ \mathbf{n}_1 P'_J(\cos \omega_{12}) - \mathbf{n}_2 P'_{J-1}(\cos \omega_{12}) \},$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J-1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2) &= 0, \\
4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J-1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2) &= i \frac{\sqrt{(J+1)(2J+1)}}{J} \{ \mathbf{n}_1 P'_{J-1}(\cos \omega_{12}) - \mathbf{n}_2 P'_J(\cos \omega_{12}) \}, \\
\end{aligned} \tag{7.3.105}$$

$$4\pi \sum_{M=-J}^J \mathbf{Y}_{JM}^{J-1*}(\vartheta_1, \varphi_1) \times \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2) = -\frac{2J+1}{J} [\mathbf{n}_1 \times \mathbf{n}_2] P'_{J-1}(\cos \omega_{12}).$$

In these equations $P_L(x)$ is a Legendre polynomial, and $P'_L(x) \equiv dP_L(x)/dx$. Analogous equations for $\mathbf{Y}_{JM}^{(\lambda)}$ may be obtained from Eqs. (7.3.9)-(7.3.10).

- (c) The most general form of the addition theorems for $\mathbf{Y}_{JM}^L$ may be written by introducing arbitrary complex vectors $\mathbf{a}_1$ and $\mathbf{a}_2$. For brevity, we shall omit the arguments $(\cos \omega_{12})$ of derivatives of the Legendre polynomials, writing P'_L instead of $P'_L(\cos \omega_{12})$, etc.

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) &= \frac{1}{J+1} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_1) (\mathbf{a}_2 \cdot \mathbf{n}_2) [P''_J + (2J+1)P'_{J+1}] \\
&+ (\mathbf{a}_1^* \cdot \mathbf{n}_2) (\mathbf{a}_2 \cdot \mathbf{n}_1) P''_J - [(\mathbf{a}_1^* \cdot \mathbf{n}_1) (\mathbf{a}_2 \cdot \mathbf{n}_1) + (\mathbf{a}_1^* \cdot \mathbf{n}_2) (\mathbf{a}_2 \cdot \mathbf{n}_2)] P''_{J+1} + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P'_J \}, \\
\end{aligned} \tag{7.3.106a}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) &= \frac{i}{2(J+1)} \sqrt{\frac{2J+1}{J}}, \\
&\times \{ -[\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 J P'_{J+1} + [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 J P'_J - [(\mathbf{a}_1^* \cdot \mathbf{n}_1) (\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) (\mathbf{a}_2 \cdot \mathbf{n}_1)] P''_{J+1} \\
&+ [(\mathbf{a}_1^* \cdot \mathbf{n}_2) (\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) (\mathbf{a}_2 \cdot \mathbf{n}_2)] P''_J \}, \\
\end{aligned} \tag{7.3.106b}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) &= \frac{1}{\sqrt{J(J+1)}}, \\
&\times \{ [(\mathbf{a}_1^* \cdot \mathbf{n}_1) (\mathbf{a}_2 \cdot \mathbf{n}_2) + (\mathbf{a}_1^* \cdot \mathbf{n}_2) (\mathbf{a}_2 \cdot \mathbf{n}_1)] P''_J - (\mathbf{a}_1^* \cdot \mathbf{n}_1) (\mathbf{a}_2 \cdot \mathbf{n}_1) P''_{J+1} \\
&- (\mathbf{a}_1^* \cdot \mathbf{n}_2) (\mathbf{a}_2 \cdot \mathbf{n}_2) P''_{J-1} + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P'_J \}. \\
\end{aligned} \tag{7.3.106c}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) &= \frac{i}{2(J+1)} \sqrt{\frac{2J+1}{J}} \\
&\times \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 J P_J' - [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 J P_{J+1}' + [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_1)] P_J'' \\
&- [(\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_2)] P_{J+1}'' \} \quad (7.3.107a)
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) &= \frac{2J+1}{J(J+1)} \{ -(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_2) [P_{J-1}'' + (J-1)P_J'] \\
&- (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) [P_{J-1}'' + J P_J'] + [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_1) + (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_2)] P_J'' + (\mathbf{a}_1^* \cdot \mathbf{a}_2) [J^2 P_J - P_{J-1}'] \} . \quad (7.3.107b)
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) &= \frac{i}{2J} \sqrt{\frac{2J+1}{J+1}} \{ -[\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 (J+1) P_J' \\
&+ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 (J+1) P_{J-1}' + [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_1)] P_J'' \\
&- [(\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_2)] P_{J-1}'' \} . \quad (7.3.107c)
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) &= \frac{1}{\sqrt{J(J+1)}} \{ [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_2) + (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1)] P_J'' \\
&- (\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_1) P_{J-1}'' - (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_2) P_{J+1}'' + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P_J' \} , \quad (7.3.108a)
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) &= \frac{i}{2J} \sqrt{\frac{2J+1}{J+1}} \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 (J+1) P_{J-1}' \\
&- [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 (J+1) P_J' - [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_1)] P_{J-1}'' \\
&+ [(\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot [\mathbf{n}_1 \times \mathbf{n}_2]) + (\mathbf{a}_1^* \cdot [\mathbf{n}_1 \times \mathbf{n}_2])(\mathbf{a}_2 \cdot \mathbf{n}_2)] P_J'' \} , \quad (7.3.108b)
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) &= \frac{1}{J} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_2) [P_J'' - (2J-1)P_{J-1}'] \\
&+ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P_J'' - [(\mathbf{a}_1^* \cdot \mathbf{n}_1)(\mathbf{a}_2 \cdot \mathbf{n}_1) + (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_2)] P_{J-1}'' + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P_J' \} . \quad (7.3.108c)
\end{aligned}$$

- (d) The case when the vectors $\mathbf{a}_1$ and $\mathbf{a}_2$ are transverse (i.e., $\mathbf{a}_1 \cdot \mathbf{n}_1 = \mathbf{a}_2 \cdot \mathbf{n}_2 = 0$) is of special interest (e.g., to describe the multipole electromagnetic fields). In this case Eqs. (7.3.106)-(7.3.108c) are considerably simplified to

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) &= \frac{1}{J+1} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P_J'' + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P_J' \} , \\
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) &= \frac{i}{J+1} \sqrt{\frac{2J+1}{J}} \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 P_J'' + [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 [(J+1)P_J' - P_{J+1}''] \} , \quad (7.3.109) \\
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) &= \frac{1}{\sqrt{J(J+1)}} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P_J'' + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P_J' \} ,
\end{aligned}$$

$$\begin{aligned}
& 4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) \\
&= \frac{i}{J+1} \sqrt{\frac{2J+1}{J}} \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 [(J+1)P'_J - P''_{J+1}] + [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 P''_J \}, \\
& 4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) \\
&= \frac{2J+1}{J(J+1)} \{ -(\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) [P''_{J-1} + JP'_J] + (\mathbf{a}_1^* \cdot \mathbf{a}_2) [J^2 P_J - P'_{J-1}] \}, \tag{7.3.110} \\
& 4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^J(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) \\
&= \frac{i}{J} \sqrt{\frac{2J+1}{J+1}} \{ -[\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 [P''_{J-1} + JP'_J] + [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 P''_J \},
\end{aligned}$$

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J+1}(\vartheta_2, \varphi_2)) &= \frac{1}{\sqrt{J(J+1)}} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P''_J + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P'_J \} \\
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^J(\vartheta_2, \varphi_2)) & \tag{7.3.111} \\
&= \frac{i}{J} \sqrt{\frac{2J+1}{J+1}} \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 P''_J - [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 [P''_{J-1} + JP'_J] \},
\end{aligned}$$

$$4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{J-1}(\vartheta_2, \varphi_2)) = \frac{1}{J} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P''_J + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P'_J \}$$

Note that $P''_{J+1} - (J+1)P'_J = P''_{J-1} + JP'_J$.

(e) Analogous relations for $\mathbf{Y}_{JM}^{(\lambda)}$, when $\mathbf{a}_1 \cdot \mathbf{n}_1 = \mathbf{a}_2 \cdot \mathbf{n}_2 = 0$, acquire the form

$$\begin{aligned}
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta_2, \varphi_2)) &= \frac{2J+1}{J(J+1)} \{ (\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) P''_J + (\mathbf{a}_1^* \cdot \mathbf{a}_2) P'_J \} \\
4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta_2, \varphi_2)) & \tag{7.3.112}
\end{aligned}$$

$$\begin{aligned}
&= \frac{i}{J(J+1)} \{ [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 (2J+1)P''_J - [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 [(J+1)P''_{J-1} + JP''_{J+1}] \} \\
& 4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{(1)}(\vartheta_2, \varphi_2)) \\
&= \frac{i}{J(J+1)} \{ -[\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_1 [(J+1)P''_{J-1} + JP''_{J+1}] + [\mathbf{a}_1^* \times \mathbf{a}_2] \cdot \mathbf{n}_2 (2J+1)P''_J \} \\
& 4\pi \sum_{M=-J}^J (\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta_1, \varphi_1))^* (\mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{(0)}(\vartheta_2, \varphi_2)) \tag{7.3.113} \\
&= \frac{2J+1}{J(J+1)} \{ -(\mathbf{a}_1^* \cdot \mathbf{n}_2)(\mathbf{a}_2 \cdot \mathbf{n}_1) [P''_{J-1} + JP'_J] + (\mathbf{a}_1^* \cdot \mathbf{a}_2) [J^2 P_J - P'_{J-1}] \}.
\end{aligned}$$

The sums involving the longitudinal vectors $\mathbf{Y}_{JM}^{(-1)}$ are equal to zero, because

$$\mathbf{a}_1 \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta_1, \varphi_1) = \mathbf{a}_2 \cdot \mathbf{Y}_{JM}^{(-1)}(\vartheta_2, \varphi_2) = 0.$$

7.3.12 Integrals Involving Vector Spherical Harmonics

Let us use the notation $\int f(\vartheta, \varphi) d\Omega \equiv \int_0^\pi d\vartheta \sin \vartheta \int_0^{2\pi} d\varphi f(\vartheta, \varphi)$.

$$\int e^{ikr} \mathbf{Y}_{JM}^L(\vartheta, \varphi) d\Omega = i^L 4\pi j_L(kr) \mathbf{Y}_{JM}^L(\vartheta_k, \varphi_k), \tag{7.3.114}$$

where

$$\mathbf{k} = (k, \vartheta_k, \varphi_k), \mathbf{r} = (r, \vartheta, \varphi), j_L(x) = \sqrt{\frac{\pi}{2x}} J_{L+\frac{1}{2}}(x).$$

$$\begin{aligned} \int e^{ikr} \mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi) d\Omega &= 4\pi i^{J+1} \left\{ \sqrt{\frac{J}{2J+1}} j_{J+1}(kr) \mathbf{Y}_{JM}^{J+1}(\vartheta_k, \varphi_k) - \sqrt{\frac{J+1}{2J+1}} j_{J-1}(kr) \mathbf{Y}_{JM}^{J-1}(\vartheta_k, \varphi_k) \right\} \\ &= \frac{4\pi i^{J+1}}{2J+1} \left\{ [J \cdot j_{J+1}(kr) - (J+1)j_{J-1}(kr)] \mathbf{Y}_{JM}^{(1)}(\vartheta_k, \varphi_k) - \sqrt{J(J+1)} [j_{J+1}(kr) + j_{J-1}(kr)] \mathbf{Y}_{JM}^{(-1)}(\vartheta_k, \varphi_k) \right\}, \end{aligned} \quad (7.3.115)$$

$$\int e^{ikr} \mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi) d\Omega = 4\pi i^J j_J(kr) \mathbf{Y}_{JM}^J(\vartheta_k, \varphi_k) = 4\pi i^J j_J(kr) \mathbf{Y}_{JM}^{(0)}(\vartheta_k, \varphi_k), \quad (7.3.116)$$

$$\begin{aligned} \int e^{ikr} \mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi) d\Omega &= 4\pi i^{J-1} \left\{ \sqrt{\frac{J+1}{2J+1}} j_{J+1}(kr) \mathbf{Y}_{JM}^{J+1}(\vartheta_k, \varphi_k) + \sqrt{\frac{J}{2J+1}} j_{J-1}(kr) \mathbf{Y}_{JM}^{J-1}(\vartheta_k, \varphi_k) \right\} \\ &= \frac{4\pi i^{J-1}}{2J+1} \left\{ \sqrt{J(J+1)} [j_{J+1}(kr) + j_{J-1}(kr)] \mathbf{Y}_{JM}^{(1)}(\vartheta_k, \varphi_k) - [(J+1)j_{J+1}(kr) - Jj_{J-1}(kr)] \mathbf{Y}_{JM}^{(-1)}(\vartheta_k, \varphi_k) \right\}, \end{aligned} \quad (7.3.117)$$

$$\int \mathbf{Y}_{JM}^L(\vartheta, \varphi) d\Omega = \sqrt{4\pi} \delta_{J1} \delta_{L0} \mathbf{e}_M,$$

$$\int \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) d\Omega = \sqrt{\frac{4\pi}{3}} \delta_{J1} \left(\sqrt{2} \delta_{\lambda 1} + \delta_{\lambda -1} \right) \mathbf{e}_M; \quad (7.3.118)$$

$$\int \mathbf{Y}_{J'M'}^{L'*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^L(\vartheta, \varphi) d\Omega = \delta_{J'J} \delta_{L'L} \delta_{M'M}, \quad (7.3.119)$$

$$\int \mathbf{Y}_{J'M'}^{(\lambda')*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) d\Omega = \delta_{J'J} \delta_{\lambda'\lambda} \delta_{M'M};$$

$$\int \mathbf{Y}_{J'M'}^{L'*}(\vartheta, \varphi) \times \mathbf{Y}_{JM}^L(\vartheta, \varphi) d\Omega = i(-1)^{J+L} \delta_{LL'} \sqrt{6(2J'+1)} \begin{Bmatrix} 1 & 1 & 1 \\ J' & J & L \end{Bmatrix} C_{J'M'1m}^{JM} \mathbf{e}_m. \quad (7.3.120)$$

7.3.13 Orthogonality, Normalization and Completeness

The collection of vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ with integer nonnegative $J(0 \leq J < \infty)$, $L = J, J \pm 1$ and integer $M(|M| \leq J)$ constitutes a complete orthonormal set of vector functions in the domain of arguments $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$. The orthonormality condition for $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ has the form

$$\int_0^\pi \int_0^{2\pi} \mathbf{Y}_{J'M'}^{L'*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^L(\vartheta, \varphi) \sin \vartheta d\vartheta d\varphi = \delta_{J'J} \delta_{L'L} \delta_{M'M}. \quad (7.3.121)$$

The completeness condition for spherical components of $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ may be written as

$$\sum_{J=0}^{\infty} \sum_{L=J-1}^{J+1} \sum_{M=-J}^J [\mathbf{Y}_{JM}^L(\vartheta, \varphi)]_\mu [\mathbf{Y}_{JM}^L(\vartheta', \varphi')]_\nu^* = \delta_{\mu\nu} \delta(\cos \vartheta - \cos \vartheta') \delta(\varphi - \varphi'). \quad (7.3.122)$$

Similarly, the set of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ constitutes a complete orthonormal set. The orthonormality property for $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ is the following

$$\int_0^\pi \int_0^{2\pi} \mathbf{Y}_{J'M'}^{(\lambda')*}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi) \sin \vartheta d\vartheta d\varphi = \delta_{J'J} \delta_{\lambda'\lambda} \delta_{M'M}, \quad (7.3.123)$$

and the completeness relation for spherical components of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ reads

$$\sum_{J=0}^{\infty} \sum_{\lambda=-1}^1 \sum_{M=-J}^J [\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)]_\mu [\mathbf{Y}_{JM}^{(\lambda)}(\vartheta', \varphi')]_\nu^* = \delta_{\mu\nu} \delta(\cos \vartheta - \cos \vartheta') \delta(\varphi - \varphi'). \quad (7.3.124)$$

7.3.14 Expansion in Series of Vector Spherical Harmonics

Any vector $\mathbf{F}(\vartheta, \varphi)$ which depends on polar angles ϑ, φ and satisfies the condition

$$\int |\mathbf{F}(\vartheta, \varphi)|^2 d\Omega < \infty, \quad (7.3.125)$$

with $\int d\Omega \equiv \int_0^\pi d\vartheta \sin \vartheta \int_0^{2\pi} d\varphi$, may be expanded in series of vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ or $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$. In other words, for $0 \leq \vartheta \leq \pi, 0 \leq \varphi < 2\pi$ $\mathbf{F}(\vartheta, \varphi)$ may be written as

$$\mathbf{F}(\vartheta, \varphi) = \sum_{JLM} A_{JM}^L \mathbf{Y}_{JM}^L(\vartheta, \varphi), \quad (7.3.126)$$

or

$$\mathbf{F}(\vartheta, \varphi) = \sum_{J\lambda M} A_{JM}^{(\lambda)} \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi). \quad (7.3.127)$$

The expansion coefficients are given by

$$\begin{aligned} A_{JM}^L &= \int d\Omega \mathbf{F}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{L*}(\vartheta, \varphi), \\ A_{JM}^{(\lambda)} &= \int d\Omega \mathbf{F}(\vartheta, \varphi) \cdot \mathbf{Y}_{JM}^{(\lambda)*}(\vartheta, \varphi). \end{aligned} \quad (7.3.128)$$

Note also the following useful rule. If a scalar function $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ of vector arguments $\mathbf{r}_1(r_1, \vartheta_1, \varphi_1)$ and $\mathbf{r}_2(r_2, \vartheta_2, \varphi_2)$ may be expanded in a series of scalar spherical harmonics as

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{Lm} A_L(r_1, r_2) Y_{Lm}^*(\vartheta_2, \varphi_2) Y_{Lm}(\vartheta_1, \varphi_1), \quad (7.3.129)$$

then the expansion of the vector function $\mathbf{F}(\mathbf{r}_2) \Phi(\mathbf{r}_1, \mathbf{r}_2)$ in a series of $\mathbf{Y}_{JM}^L(\vartheta_1, \varphi_1)$ has the form

$$\mathbf{F}(\mathbf{r}_2) \Phi(\mathbf{r}_1, \mathbf{r}_2) = \sum_{JLM} A_L(r_1, r_2) [\mathbf{F}(\mathbf{r}_2) \cdot \mathbf{Y}_{JM}^{L*}(\vartheta_2, \varphi_2)] \mathbf{Y}_{JM}^L(\vartheta_1, \varphi_1). \quad (7.3.130)$$

Some examples of such expansions are given below.

(a) *Expansion of a plane wave*

$$\boldsymbol{\varepsilon}(\mathbf{k}) e^{i\mathbf{k}\mathbf{r}} = 4\pi \sum_{JLM} i^L j_L(kr) \{ \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{L*}(\vartheta_k, \varphi_k) \} \mathbf{Y}_{JM}^L(\vartheta, \varphi), \quad (7.3.131)$$

where $j_L(x) = \sqrt{\pi/(2x)} J_{L+\frac{1}{2}}(x)$ is a spherical Bessel function, $\mathbf{k} = (k, \vartheta_k, \varphi_k)$, $\mathbf{r} = (r, \vartheta, \varphi)$. The expansion in a series of $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ has the form

$$\boldsymbol{\varepsilon}(\mathbf{k}) e^{i\mathbf{k}\mathbf{r}} = \sum_{J\lambda M} A_{JM}^{(\lambda)} \mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi), \quad (7.3.132)$$

where the expansion coefficients are given by

$$\begin{aligned} A_{JM}^{(1)} &= 4\pi i^{J+1} \left\{ \sqrt{\frac{J}{2J+1}} j_{J+1}(kr) \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{J+1*}(\vartheta_k, \varphi_k) - \sqrt{\frac{J+1}{2J+1}} j_{J-1}(kr) \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{J-1*}(\vartheta_k, \varphi_k) \right\} \\ &= \frac{4\pi i^{J+1}}{2J+1} \left\{ [J \cdot j_{J+1}(kr) - (J+1)j_{J-1}(kr)] \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(1)*}(\vartheta_k, \varphi_k) \right. \\ &\quad \left. - \sqrt{J(J+1)} [j_{J+1}(kr) + j_{J-1}(kr)] \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(-1)*}(\vartheta_k, \varphi_k) \right\}, \end{aligned} \quad (7.3.133)$$

$$A_{JM}^{(0)} = 4\pi i^J j_J(kr) \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^J(\vartheta_k, \varphi_k) = 4\pi i^J j_J(kr) \cdot \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(0)*}(\vartheta_k, \varphi_k), \quad (7.3.134)$$

$$\begin{aligned} A_{JM}^{(-1)} &= 4\pi i^{J-1} \left\{ \sqrt{\frac{J+1}{2J+1}} j_{J+1}(kr) \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{J+1*}(\vartheta_k, \varphi_k) + \sqrt{\frac{J}{2J+1}} j_{J-1}(kr) \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{J-1*}(\vartheta_k, \varphi_k) \right\} \\ &= \frac{4\pi i^{J-1}}{2J+1} \left\{ \sqrt{J(J+1)} [j_{J+1}(kr) + j_{J-1}(kr)] \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(1)*}(\vartheta_k, \varphi_k) \right. \\ &\quad \left. - [(J+1)j_{J+1}(kr) - J \cdot j_{J-1}(kr)] \boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(-1)*}(\vartheta_k, \varphi_k) \right\}. \end{aligned} \quad (7.3.135)$$

If the plane wave is transverse, i.e., $\mathbf{k} \cdot \boldsymbol{\varepsilon}(\mathbf{k}) = 0$, then $\boldsymbol{\varepsilon}(\mathbf{k}) \cdot \mathbf{Y}_{JM}^{(-1)*}(\vartheta_k, \varphi_k) = 0$.

(b) *Green's function for Laplace equation*

$$\mathbf{J}(\mathbf{R}) \frac{1}{|\mathbf{R} - \mathbf{r}|} = 4\pi \sum_{JLM} A_L(R, r) \frac{1}{(2L+1)} \{ \mathbf{J}(\mathbf{R}) \cdot \mathbf{Y}_{JM}^{L*}(\Theta, \Phi) \} \mathbf{Y}_{JM}^L(\vartheta, \varphi), \quad (7.3.136)$$

where $\mathbf{r} = (r, \vartheta, \varphi)$, $\mathbf{R} = (R, \Theta, \Phi)$,

$$A_L(R, r) = \begin{cases} \frac{r^L}{R^{L+1}}, & \text{if } r < R, \\ \frac{R^L}{r^{L+1}}, & \text{if } r > R. \end{cases} \quad (7.3.137)$$

(c) *Green's function for Helmholtz equation*

$$\mathbf{J}(\mathbf{R}) \frac{e^{ik|\mathbf{R}-\mathbf{r}|}}{|\mathbf{R} - \mathbf{r}|} = 4\pi ik \sum_{JLM} A_L(R, r) \{ \mathbf{J}(\mathbf{R}) \mathbf{Y}_{JM}^{L*}(\Theta, \Phi) \} \mathbf{Y}_{JM}^L(\vartheta, \varphi), \quad (7.3.138)$$

where

$$A_L(R, r) = \begin{cases} j_L(kr)h_L^{(1)}(kR), & \text{if } r < R, \\ h_L^{(1)}(kr)j_L(kR), & \text{if } r > R. \end{cases} \quad (7.3.139)$$

$h_L^{(1)}(x) = \sqrt{\pi/(2x)} H_{L+\frac{1}{2}}^{(1)}(x)$ is the spherical Hankel function.

7.3.15 Vector Spherical Harmonics for $\vartheta = 0$ or $\vartheta = \pi$

If $\vartheta = 0$ or $\vartheta = \pi$, the vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ and $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ are equal to zero unless $M = 0, \pm 1$. The expressions for these harmonics in terms of spherical basis vectors $\mathbf{e}_\mu$ ($\mu = 0, \pm 1$) have the form

$$\mathbf{Y}_{JM}^{J+1}(0, \varphi) = (-1)^{J-1} \mathbf{Y}_{JM}^{J+1}(\pi, \varphi) = \begin{cases} \sqrt{\frac{J}{8\pi}} \mathbf{e}_M, & \text{if } M = \pm 1; \\ -\sqrt{\frac{J+1}{4\pi}} \mathbf{e}_0, & \text{if } M = 0 \end{cases}, \quad (7.3.140)$$

$$\mathbf{Y}_{JM}^J(0, \varphi) = (-1)^J \mathbf{Y}_{JM}^J(\pi, \varphi) = \begin{cases} -M \sqrt{\frac{2J+1}{8\pi}} \mathbf{e}_M, & \text{if } M = \pm 1; \\ 0, & \text{otherwise} \end{cases}, \quad (7.3.141)$$

$$\mathbf{Y}_{JM}^{J-1}(0, \varphi) = (-1)^{J-1} \mathbf{Y}_{JM}^{J-1}(\pi, \varphi) = \begin{cases} \sqrt{\frac{J+1}{8\pi}} \mathbf{e}_M, & \text{if } M = \pm 1; \\ \sqrt{\frac{J}{4\pi}} \mathbf{e}_0, & \text{if } M = 0 \end{cases}, \quad (7.3.142)$$

$$\mathbf{Y}_{JM}^{(1)}(0, \varphi) = (-1)^{J-1} \mathbf{Y}_{JM}^{(1)}(\pi, \varphi) = \begin{cases} \sqrt{\frac{2J+1}{8\pi}} \mathbf{e}_M, & \text{if } M = \pm 1; \\ 0, & \text{otherwise} \end{cases}, \quad (7.3.143)$$

$$\mathbf{Y}_{JM}^{(0)}(0, \varphi) = (-1)^J \mathbf{Y}_{JM}^{(0)}(\pi, \varphi) = \begin{cases} -M \sqrt{\frac{2J+1}{8\pi}} \mathbf{e}_M, & \text{if } M = \pm 1; \\ 0, & \text{otherwise} \end{cases}, \quad (7.3.144)$$

$$\mathbf{Y}_{JM}^{(-1)}(0, \varphi) = (-1)^{J-1} \mathbf{Y}_{JM}^{(-1)}(\pi, \varphi) = \begin{cases} \sqrt{\frac{2J+1}{4\pi}} \mathbf{e}_0, & \text{if } M = 0; \\ 0, & \text{otherwise} \end{cases}. \quad (7.3.145)$$

7.3.16 Vector Spherical Harmonics at $J = 0, 1$

If $J = 0$ or $J = 1$, the vector spherical harmonics $\mathbf{Y}_{JM}^L(\vartheta, \varphi)$ may be expressed in terms of spherical basis vectors $\mathbf{e}_\mu$ ($\mu = 0, \pm 1$) and the unit vector $\mathbf{n}(\vartheta, \varphi)$ as

$$\mathbf{Y}_{00}^1(\vartheta, \varphi) = -\frac{1}{\sqrt{4\pi}} \mathbf{n}, \quad (7.3.146)$$

$$\mathbf{Y}_{1M}^0(\vartheta, \varphi) = \frac{1}{\sqrt{4\pi}} \mathbf{e}_M, \quad (7.3.147)$$

$$\mathbf{Y}_{1M}^1(\vartheta, \varphi) = i \sqrt{\frac{3}{8\pi}} [\mathbf{e}_M \times \mathbf{n}], \quad (7.3.148)$$

$$\mathbf{Y}_{1M}^2(\vartheta, \varphi) = \frac{1}{\sqrt{8\pi}} \{ \mathbf{e}_M - 3\mathbf{n}(\mathbf{e}_M \cdot \mathbf{n}) \}. \quad (7.3.149)$$

Analogous expressions for $\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)$ are

$$\mathbf{Y}_{00}^{(-1)}(\vartheta, \varphi) = \frac{1}{\sqrt{4\pi}} \mathbf{n}, \quad (7.3.150)$$

$$\mathbf{Y}_{1M}^{(1)}(\vartheta, \varphi) = \sqrt{\frac{3}{8\pi}} \{ \mathbf{e}_M - \mathbf{n} (\mathbf{e}_M \cdot \mathbf{n}) \}, \quad (7.3.151)$$

$$\mathbf{Y}_{1M}^{(0)}(\vartheta, \varphi) = i \sqrt{\frac{3}{8\pi}} [\mathbf{e}_M \times \mathbf{n}], \quad (7.3.152)$$

$$\mathbf{Y}_{1M}^{(-1)}(\vartheta, \varphi) = \sqrt{\frac{3}{4\pi}} \mathbf{n} (\mathbf{e}_M \cdot \mathbf{n}). \quad (7.3.153)$$

7.3.17 Quadratic Forms of the Vector Spherical Harmonics

The functions $|\mathbf{Y}_{JM}^{(\lambda)}(\vartheta, \varphi)|^2$ are important for physical applications. In particular, they describe the angular distribution of spin-1 particles in quantum states with total angular momentum J . These functions are independent of φ and are the same at $\lambda = 1$ and $\lambda = 0$. Let us use the notation

$$W_{JM}^{\perp}(\vartheta) \equiv |\mathbf{Y}_{JM}^{(1)}(\vartheta, \varphi)|^2 = |\mathbf{Y}_{JM}^{(0)}(\vartheta, \varphi)|^2 \quad (7.3.154)$$

$$W_{JM}^{\parallel}(\vartheta) \equiv |\mathbf{Y}_{JM}^{(-1)}(\vartheta, \varphi)|^2.$$

Then one has

$$W_{JM}^{\perp}(\vartheta) = \frac{1}{2J(J+1)} \left\{ (J-M)(J+M+1) |Y_{JM+1}(\vartheta, \varphi)|^2 + 2M^2 |Y_{JM}(\vartheta, \varphi)|^2 \right. \\ \left. + (J+M)(J-M+1) |Y_{JM-1}(\vartheta, \varphi)|^2 \right\}, \quad (7.3.155)$$

$$W_{JM}^{\parallel}(\vartheta) = |Y_{JM}(\vartheta, \varphi)|^2.$$

The expansions of $W_{JM}^{\perp, \parallel}(\vartheta)$ in terms of the Legendre polynomials have the form

$$W_{JM}^{\perp}(\vartheta) = \sum_{n=0}^J a_n(J, M) P_{2n}(\cos \vartheta), \\ W_{JM}^{\parallel}(\vartheta) = \sum_{n=0}^J b_n(J, M) P_{2n}(\cos \vartheta), \quad (7.3.156)$$

where

$$a_n(J, M) = -\frac{(2J+1)(4n+1)}{4\pi} \left\{ \begin{matrix} J & J & 2n \\ J & J & 1 \end{matrix} \right\} C_{J02n0}^{J0} C_{JM2n0}^{JM} \\ = (-1)^n \frac{4n+1}{4\pi} \frac{J(J+1) - n(2n+1)}{J(J+1)} \frac{(J+n)!(2n)!}{(J-n)!(n!)^2} \sqrt{\frac{(2J+1)(2J-2n)!}{(2J+2n+1)!}} C_{JM2n0}^{JM}, \quad (7.3.157)$$

$$b_n(J, M) = \frac{4n+1}{4\pi} C_{J02n0}^{J0} C_{JM2n0}^{JM} = (-1)^n \frac{4n+1}{4\pi} \frac{(J+n)!(2n)!}{(J-n)!(n!)^2} \sqrt{\frac{(2J+1)(2J-2n)!}{(2J+2n+1)!}} C_{JM2n0}^{JM}. \quad (7.3.158)$$

In particular,

$$a_0(J, M) = b_0(J, M) = \frac{1}{4\pi}, \quad (7.3.159)$$

$$a_1(J, M) = \frac{5}{4\pi} \frac{[J(J+1) - 3][J(J+1) - 3M^2]}{J(J+1)(2J-1)(2J+3)}, \quad (7.3.160)$$

$$b_1(J, M) = \frac{5}{4\pi} \frac{J(J+1) - 3M^2}{(2J-1)(2J+3)}, \quad (7.3.161)$$

$$a_2(J, M) = \frac{27}{16\pi} \frac{[J(J+1) - 10]}{J(J+1)} \cdot \frac{[3(J^2 + 2J - 5M^2)(J^2 - 5M^2 - 1) - 10M^2(4M^2 - 1)]}{(2J-3)(2J-1)(2J+3)(2J+5)}, \quad (7.3.162)$$

$$b_2(J, M) = \frac{27}{16\pi} \frac{[3(J^2 + 2J - 5M^2)(J^2 - 5M^2 - 1) - 10M^2(4M^2 - 1)]}{(2J-3)(2J-1)(2J+3)(2J+5)}. \quad (7.3.163)$$

The functions $W_{JM}^{\perp\parallel}(\vartheta)$ satisfy the normalization conditions

$$\begin{aligned} \int W_{JM}^{\perp}(\vartheta) d\Omega &= \int W_{JM}^{\parallel}(\vartheta) d\Omega = 1 \\ \sum_{M=-J}^J W_{JM}^{\perp}(\vartheta) &= \sum_{M=-J}^J W_{JM}^{\parallel}(\vartheta) = \frac{2J+1}{4\pi}. \end{aligned} \quad (7.3.164)$$

Note the following symmetry properties of $W_{JM}^{\perp\parallel}(\vartheta)$

$$\begin{aligned} W_{JM}^{\perp}(\vartheta) &= W_{J-M}^{\perp}(\vartheta) = W_{JM}^{\perp}(\pi - \vartheta) = W_{J-M}^{\perp}(\pi - \vartheta) \\ W_{JM}^{\parallel}(\vartheta) &= W_{J-M}^{\parallel}(\vartheta) = W_{JM}^{\parallel}(\pi - \vartheta) = W_{J-M}^{\parallel}(\pi - \vartheta). \end{aligned} \quad (7.3.165)$$

For $\vartheta = 0$ and $\vartheta = \pi$ we have

$$W_{JM}^{\perp}(0) = W_{JM}^{\perp}(\pi) = \begin{cases} \frac{2J+1}{8\pi}, & \text{if } M = \pm 1 \\ 0, & \text{otherwise} \end{cases}, \quad (7.3.166)$$

$$W_{JM}^{\parallel}(0) = W_{JM}^{\parallel}(\pi) = \begin{cases} \frac{2J+1}{4\pi}, & \text{if } M = 0 \\ 0, & \text{otherwise} \end{cases}. \quad (7.3.167)$$

The explicit forms of $W_{JM}^{\perp\parallel}(\vartheta)$ for $J = 0, 1, 2, 3, 4, 5$ ($0 \leq M \leq J$) are given in Table 7.2. For negative M one may use the relation $W_{J-M} = W_{JM}$. Note also the explicit form of $W_{JM}^{\perp\parallel}(\vartheta)$ for special values of M :

$$\begin{aligned} W_{J0}^{\perp}(\vartheta) &= \frac{2J+1}{4\pi(J+1)J} \sin^2 \vartheta [P_J'(\cos \vartheta)]^2, \\ W_{J0}^{\parallel}(\vartheta) &= \frac{2J+1}{4\pi} [P_J(\cos \vartheta)]^2, \\ W_{JJ}^{\perp}(\vartheta) &= W_{J-J}^{\perp}(\vartheta) = \frac{1}{4\pi} \frac{(2J+1)!(\sin \vartheta)^{2J-2}}{2^{2J}(J+1)!(J-1)!} (1 + \cos^2 \vartheta), \\ W_{JJ}^{\parallel}(\vartheta) &= W_{J-J}^{\parallel}(\vartheta) = \frac{1}{4\pi} \frac{(2J+1)!}{2^{2J}(J!)^2} (\sin \vartheta)^{2J}. \end{aligned} \quad (7.3.168) \quad (7.3.169)$$

7.4 Other notations for tensor spherical harmonics

The spherical harmonics defined by other authors are presented on the left-hand sides of the equalities given below. On the right-hand sides of these equalities we exhibit the corresponding harmonics in our notation.

(a) *Tensor spherical harmonics*

Berestetskii, Dolginov and Ter-Martirosian [53]

$$Y_{lm}^{L\lambda} = \eta Y_{lm}^{l+\lambda L}, \quad \text{where } \eta = \begin{cases} (-1)^L & \text{for integer } L \\ 1 & \text{for half-integer } L. \end{cases}$$

Newton [5]

$$\mathcal{U}_{jLS}^M = i^{l+S} Y_{jM}^{lS}$$

(b) *Spinor spherical harmonics*

Akhiezer and Berestetskii [26]: $\Omega_{jlm} = \Omega_{jm}^l$;

Berestetskii, Dolginov and Ter-Martirosian [53]: $Y_{j\mu}^{(\lambda)} = \Omega_{j\mu}^{j+\lambda}$;

Berestetskii, Lifshitz and Pitaevsky [17]: $\Omega_{jlm} = i^l \Omega_{jm}^l$.

Table 7.2. Explicit forms of $W_{JM}^\perp(\vartheta)$ and $W_{JM}^\parallel(\vartheta)$.

J	M	$W_{JM}^\perp(\vartheta)$	$W_{JM}^\parallel(\vartheta)$
0	0	$1/4\pi$	$1/4\pi$
1	0	$\frac{3}{8\pi} \sin^2 \vartheta$	$\frac{3}{4\pi} \cos^2 \vartheta$
1	1	$\frac{3}{16\pi} (1 + \cos^2 \vartheta)$	$\frac{3}{8\pi} \sin^2 \vartheta$
2	0	$\frac{15}{8\pi} \sin^2 \vartheta \cos^2 \vartheta$	$\frac{5}{16\pi} (1 - 3 \cos^2 \vartheta)^2$
2	1	$\frac{5}{16\pi} (1 - 3 \cos^2 \vartheta + 4 \cos^4 \vartheta)$	$\frac{15}{8\pi} \sin^2 \vartheta \cos^2 \vartheta$
2	2	$\frac{5}{16\pi} (1 - \cos^4 \vartheta)$	$\frac{15}{32\pi} \sin^4 \vartheta$
3	0	$\frac{21}{64\pi} \sin^2 \vartheta (1 - 5 \cos^2 \vartheta)^2$	$\frac{7}{16\pi} \cos^2 \vartheta (3 - 5 \cos^2 \vartheta)^2$
3	1	$\frac{7}{256\pi} (1 + 111 \cos^2 \vartheta - 305 \cos^4 \vartheta + 225 \cos^6 \vartheta)$	$\frac{21}{64\pi} \sin^2 \vartheta (1 - 5 \cos^2 \vartheta)^2$
3	2	$\frac{35}{128\pi} \sin^2 \vartheta (1 - 2 \cos^2 \vartheta + 9 \cos^4 \vartheta)$	$\frac{105}{32\pi} \sin^4 \vartheta \cos^2 \vartheta$
3	3	$\frac{105}{256\pi} \sin^4 \vartheta (1 + \cos^2 \vartheta)$	$\frac{35}{64\pi} \sin^6 \vartheta$
4	0	$\frac{45}{64\pi} \sin^2 \vartheta \cos^2 \vartheta (3 - 7 \cos^2 \vartheta)^2$	$\frac{9}{256\pi} (3 - 30 \cos^2 \vartheta + 35 \cos^4 \vartheta)^2$
4	1	$\frac{9}{256\pi} (9 - 153 \cos^2 \vartheta + 855 \cos^4 \vartheta - 1463 \cos^6 \vartheta + 784 \cos^8 \vartheta)$	$\frac{45}{64\pi} \sin^2 \vartheta \cos^2 \vartheta (3 - 7 \cos^2 \vartheta)^2$
4	2	$\frac{9}{128\pi} \sin^2 \vartheta (1 + 50 \cos^2 \vartheta - 175 \cos^4 \vartheta + 196 \cos^6 \vartheta)$	$\frac{45}{128\pi} \sin^4 \vartheta (1 - 7 \cos^2 \vartheta)^2$
4	3	$\frac{63}{256\pi} \sin^4 \vartheta (1 + \cos^2 \vartheta + 16 \cos^4 \vartheta)$	$\frac{315}{64\pi} \sin^6 \vartheta \cos^2 \vartheta$
4	4	$\frac{63}{128\pi} \sin^6 \vartheta (1 + \cos^2 \vartheta)$	$\frac{315}{512\pi} \sin^8 \vartheta$
5	0	$\frac{165}{512\pi} \sin^2 \vartheta (1 - 14 \cos^2 \vartheta + 21 \cos^4 \vartheta)^2$	$\frac{11}{256\pi} \cos^2 \vartheta (15 - 70 \cos^2 \vartheta + 63 \cos^4 \vartheta)^2$
5	1	$\frac{11}{1024\pi} (1 + 813 \cos^2 \vartheta - 7070 \cos^4 \vartheta + 21378 \cos^6 \vartheta - 26019 \cos^8 \vartheta + 11025 \cos^{10} \vartheta)$	$\frac{165}{512\pi} \sin^2 \vartheta (1 - 14 \cos^2 \vartheta + 21 \cos^4 \vartheta)^2$
5	2	$\frac{77}{256\pi} \sin^2 \vartheta (1 - 20 \cos^2 \vartheta + 150 \cos^4 \vartheta - 324 \cos^6 \vartheta + 225 \cos^8 \vartheta)$	$\frac{1155}{128\pi} \sin^4 \vartheta \cos^2 \vartheta (1 - 3 \cos^2 \vartheta)^2$
5	3	$\frac{231}{2048\pi} \sin^4 \vartheta (1 + 31 \cos^2 \vartheta - 129 \cos^4 \vartheta + 225 \cos^6 \vartheta)$	$\frac{385}{1024\pi} \sin^6 \vartheta (1 - 9 \cos^2 \vartheta)^2$
5	4	$\frac{231}{1024\pi} \sin^6 \vartheta (1 + 6 \cos^2 \vartheta + 25 \cos^4 \vartheta)$	$\frac{3465}{512\pi} \sin^8 \vartheta \cos^2 \vartheta$
5	5	$\frac{1155}{2048\pi} \sin^8 \vartheta (1 + \cos^2 \vartheta)$	$\frac{693}{1024\pi} \sin^{10} \vartheta$

(c) Vector spherical harmonics

Akhiezer and Berestetskii [26]: $\mathbf{Y}_{JM}^{(\lambda)} = \mathbf{Y}_{JM}^{(\lambda)}, \mathbf{Y}_{JLM} = \mathbf{Y}_{JM}^L$;

Blatt and Weisskopf [27]; Rose [4]: $\mathbf{T}_{L\lambda}^M = (-1)^{\lambda+1-L} \mathbf{Y}_{LM}^{(\lambda)}$;

Berestetskii, Dolginov and Ter-Martirosian [53]: $\mathbf{Y}_{lm}^{(\lambda)} = -\mathbf{Y}_{lm}^{l+\lambda}$.

Jackson [28]: $\mathbf{X}_{lm} = \mathbf{Y}_{lm}^{(0)}$.

Newton [5]: $\mathbf{Y}_{JM}^{(e)} = i^J \mathbf{Y}_{JM}^{(1)}, \mathbf{Y}_{JM}^{(m)} = i^{J+1} \mathbf{Y}_{JM}^{(0)}, \mathbf{Y}_{JM}^{(0)} = i^J \mathbf{Y}_{JM}^{(-1)}, \mathbf{Y}_{Jl}^M = i^{l+1} \mathbf{Y}_{JM}^l$.

Berestetskii, Lifshitz and Pitaevskii [17]: $\mathbf{Y}_{JM}^{(e)} = i^J \mathbf{Y}_{JM}^{(1)}, \mathbf{Y}_{JM}^{(M)} = i^{J+1} \mathbf{Y}_{JM}^{(0)}, \mathbf{Y}_{JM}^{(l)} = i^J \mathbf{Y}_{JM}^{(-1)}$.

Chapter 8

Clebsch-Gordan coefficients and $3jm$ symbols

8.1 Definition of vector addition coefficients

The Clebsch-Gordan coefficients are vector addition coefficients. They play an important role in the decomposition of reducible representations of rotation group into irreducible representations. All recoupling coefficients or $3nj$ symbols can be determined as the sums of products of the Clebsch-Gordan coefficients. The aforesaid explains the extensive application of these coefficients in the quantum theory of angular momentum.

8.1.1 The Clebsch-Gordan Coefficients

Let $\mathbf{j}_1$ and $\mathbf{j}_2$ be two angular momenta with projections m_1 and m_2 on the quantization axis. A Clebsch-Gordan coefficient represents the probability amplitude that $\mathbf{j}_1$ and $\mathbf{j}_2$ are coupled into a resultant angular momentum $\mathbf{j}$ with projection m . In accordance with the vector addition rules $\mathbf{j}_1 + \mathbf{j}_2 = \mathbf{j}$, the Clebsch-Gordan coefficient vanishes unless the triangular conditions (triangular inequalities) are fulfilled, i.e.,

$$|j_1 - j_2| \leq j \leq j_1 + j_2, \quad (8.1.1)$$

and the requirement

$$m_1 + m_2 = m, \quad (8.1.2)$$

is satisfied. It will also be assumed that arguments of the Clebsch-Gordan coefficients satisfy the following conditions:

- (a) j_1, j_2, j are integer or half-integer non-negative numbers; ¹
- (b) m_1, m_2, m are integer or half-integer (positive or negative) numbers;
- (c) $|m_1| \leq j_1, |m_2| \leq j_2, |m| \leq j$;
- (d) $j_1 + m_1, j_2 + m_2, j + m, j_1 + j_2 + j$ are integer non-negative numbers.

The absolute value of a Clebsch-Gordan coefficient is given by

$$\left[C_{j_1 m_1 j_2 m_2}^{j m} \right]^2 = \frac{2j+1}{8\pi} \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{m_1 m_1}^{j_1}(\alpha, \beta, \gamma) D_{m_2 m_2}^{j_2}(\alpha, \beta, \gamma) D_{m m}^{j*}(\alpha, \beta, \gamma). \quad (8.1.4)$$

The phase of the Clebsch-Gordan coefficients may be chosen in different ways. The phase convention proposed by Condon and Shortley [22] is universally accepted. In accordance with this convention the Clebsch-Gordan coefficients are real and their phases are fixed by the additional relation

$$C_{j_1 m_1 j_2 m_2}^{j m} C_{j_1 j_1 j_2 -j_2}^{j j_1 - j_2} = \frac{2j+1}{8\pi} \int_0^{2\pi} d\alpha \int_0^\pi d\beta \sin \beta \int_0^{2\pi} d\gamma D_{m_1 j_1}^{j_1}(\alpha, \beta, \gamma) D_{m_2 -j_2}^{j_2}(\alpha, \beta, \gamma) D_{m j_1 - j_2}^{j*}(\alpha, \beta, \gamma), \quad (8.1.5)$$

where

$$C_{j_1 j_1 j_2 -j_2}^{j j_1 - j_2} > 0.$$

¹The extension of the Clebsch-Gordan coefficients to negative values of momenta will be considered in Sec. 8.4.5.

Under such a phase definition the Clebsch-Gordan coefficients satisfy the relations

$$\begin{aligned} C_{j\ 0\ 0}^{j\ m} = 1, \quad C_{j_1\ j_1\ j_2\ j_2}^{j_1+j_2\ j_1+j_2} = 1, \quad C_{j_1\ m_1\ j_2\ m_2}^{j_1+j_2\ m_1+m_2} > 0, \quad C_{j_1\ m_1\ j_2\ -j_2}^{j\ m} > 0, \quad (-1)^{j_1+j_2-j} C_{j_1\ m_1\ j_2\ j_2}^{j\ m} > 0, \\ (-1)^{j_1-m_1} C_{j_1\ m_1\ j_2\ m_2}^{j\ j} > 0, \quad (-1)^{j_2+m_2} C_{j_1\ m_1\ j_2\ m_2}^{j_1-j_2\ m} > 0. \end{aligned} \quad (8.1.6)$$

The Clebsch-Gordan coefficients are elements of the unitary matrix which performs direct and inverse transformations between state vectors $|j_1 m_1 j_2 m_2\rangle$ and $|j_1 j_2 j m\rangle$

$$C_{j_1\ m_1\ j_2\ m_2}^{j\ m} = \langle j_1 m_1 j_2 m_2 | j_1 j_2 j m \rangle = \langle j_1 j_2 j m | j_1 m_1 j_2 m_2 \rangle. \quad (8.1.7)$$

The unitarity relation is

$$\begin{aligned} \sum_{m_1 m_2} C_{j_1\ m_1\ j_2\ m_2}^{j\ m} C_{j_1\ m_1\ j_2\ m_2}^{j'\ m'} &= \delta_{jj'} \delta_{mm'}. \\ \sum_{j(m)} C_{j_1\ m_1\ j_2\ m_2}^{j\ m} C_{j_1\ m_1' j_2\ m_2'}^{j\ m} &= \delta_{m_1 m_1'} \delta_{m_2 m_2'}. \end{aligned} \quad (8.1.8)$$

The direct product of two irreducible tensors $\mathfrak{M}_{j_1 m_1}$ and $\mathfrak{N}_{j_2 m_2}$ may be decomposed into irreducible tensors. The coefficients of this decomposition are just the Clebsch-Gordan coefficients:

$$\mathfrak{M}_{j_1 m_1} \mathfrak{N}_{j_2 m_2} = \sum_{j(m)} C_{j_1\ m_1\ j_2\ m_2}^{j\ m} \{\mathfrak{N}_{j_1} \otimes \mathfrak{N}_{j_2}\}_{jm}. \quad (8.1.9)$$

The inverse relation is

$$\{\mathfrak{M}_{j_1} \otimes \mathfrak{N}_{j_2}\}_{jm} = \sum_{m_1 m_2} C_{j_1\ m_1\ j_2\ m_2}^{j\ m} \mathfrak{M}_{j_1 m_1} \mathfrak{N}_{j_2 m_2}. \quad (8.1.10)$$

8.1.2 The Wigner $3jm$ Symbols

Not infrequently the Wigner $3j$ symbols [54] are used instead of the Clebsch-Gordan coefficients. These symbols possess simpler symmetry properties. The $3j$ symbols are related to the Clebsch-Gordan coefficients by

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = (-1)^{j_3+m_3+2j_1} \frac{1}{\sqrt{2j_3+1}} C_{j_1\ -m_1\ j_2\ -m_2}^{j_3\ m_3}. \quad (8.1.11)$$

The inverse relation is

$$C_{j_1\ m_1\ j_2\ m_2}^{j_3\ m_3} = (-1)^{j_1-j_2+m_3} \sqrt{2j_3+1} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & -m_3 \end{pmatrix}. \quad (8.1.12)$$

The $3j$ symbol represents the probability amplitude that three angular momenta j_1, j_2 and j_3 with projections m_1, m_2 and m_3 are coupled to yield zero angular momentum:

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \eta \sum_{j' m'} C_{j_1\ m_1\ j_2\ m_2}^{j'\ m'} C_{j'\ m'\ j_3\ m_3}^{0\ 0} \quad (8.1.13)$$

The phase factor $\eta = (-1)^{j_1-j_2+j_3}$ is chosen in such a way that any cyclic permutation of columns leaves the $3jm$ symbol unchanged. Below we shall use, along with the letters j, m other Latin letters (a, b, c , etc.) to denote angular momenta, and Greek letters (α, β, γ , etc.) to denote momentum projections in the arguments of the $3jm$ symbols and Clebsch-Gordan coefficients.

8.1.3 Regge R-Symbols

The Wigner $3jm$ symbol may be represented by a square 3×3 array $\|R_{ik}\|$ ($i, k = 1, 2, 3$) which is called the Regge R -symbol [55]

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| \quad (8.1.14)$$

where

$$\begin{aligned} R_{11} &= -a + b + c, & R_{12} &= a - b + c, & R_{13} &= a + b - c, \\ R_{21} &= a + \alpha, & R_{22} &= b + \beta, & R_{23} &= c + \gamma, \\ R_{31} &= a - \alpha, & R_{32} &= b - \beta, & R_{33} &= c - \gamma. \end{aligned} \quad (8.1.15)$$

The inverse relations are

$$\begin{aligned} 2a &= R_{21} + R_{31} = R_{12} + R_{13}, & 2\alpha &= R_{21} - R_{31} = R_{32} - R_{22} + R_{33} - R_{23}, \\ 2b &= R_{22} + R_{32} = R_{11} + R_{13}, & 2\beta &= R_{22} - R_{32} = R_{31} - R_{21} + R_{33} - R_{23}, \\ 2c &= R_{23} + R_{33} = R_{11} + R_{12}, & 2\gamma &= R_{23} - R_{33} = R_{31} - R_{21} + R_{32} - R_{22}. \end{aligned} \quad (8.1.16)$$

All nine elements R_{ik} are non-negative integers which satisfy the relations

$$\sum_i R_{ik} = J, \quad \sum_k R_{ik} = J, \quad (8.1.17)$$

where

$$J = a + b + c. \quad (8.1.18)$$

8.2 EXplicit forms of the Clebsch-Gordan coefficients and their relations to other functions

Below we shall assume that the arguments of the Clebsch-Gordan coefficients and $3jm$ symbols satisfy Eqs. 8.1.1-8.1.3. In addition we introduce the Δ -symbol defined by

$$\Delta(abc) = \left[\frac{(a+b-c)!(a-b+c)!(-a+b+c)!}{(a+b+c+1)!} \right]^{\frac{1}{2}}. \quad (8.2.1)$$

The Δ -symbol is invariant under permutations of a, b, c . Numerical values of the Δ -symbol are given in Table 8.12 for $\frac{1}{2} \leq a, b, c \leq 5$. If one of the momenta a, b, c equals zero, the Δ -symbol is reduced to

$$\Delta(aa0) = \frac{1}{\sqrt{2a+1}}. \quad (8.2.2)$$

8.2.1 Representations of the Clebsch-Gordan Coefficients in the Form of Algebraic Sums

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} \Delta(abc) [(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!(2c+1)]^{\frac{1}{2}}, \\ &\times \sum_z \frac{(-1)^z}{z!(a+b-c-z)!(a-\alpha-z)!(b+\beta-z)!(c-b+\alpha+z)!(c-a-\beta+z)!} \end{aligned} \quad (8.2.3)$$

(Van der Waerden [29], Racah [56])

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} \frac{\Delta(abc)}{(c+a-b)!(c-a+b)!} \left[\frac{(a-\alpha)!(b-\beta)!(c+\gamma)!(c-\gamma)!(2c+1)}{(a+\alpha)!(b+\beta)!} \right]^{\frac{1}{2}} \\ &\times \sum_z \frac{(-1)^{a-\alpha+z} (a+\alpha+z)!(c+b-\alpha-z)!}{z!(a-\alpha-z)!(c-\gamma-z)!(b-c+\alpha+z)!} \end{aligned} \quad (8.2.4)$$

(Racah [56], Fock [57])

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} \Delta(abc) \left[\frac{(c+\gamma)!(c-\gamma)!(2c+1)}{(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!} \right]^{\frac{1}{2}} \\ &\sum_z \frac{(-1)^{b+\beta+z} (c+b+\alpha-z)!(a-\alpha+z)!}{(c-a+b-z)!(c+\gamma-z)!(a-b-\gamma+z)!} \end{aligned} \quad (8.2.5)$$

(Wigner [15])

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} \frac{\Delta(abc)}{(a-b+c)!(a+b-c)!} \left[\frac{(a+\alpha)!(a-\alpha)!(b-\beta)!(c+\gamma)!(2c+1)}{(b+\beta)!(c-\gamma)!} \right]^{\frac{1}{2}} \\ &\times \sum_z \frac{(-1)^{b+\beta+x} (2c-z)!(a+b-c+z)!}{z!(c-a+b-z)!(c+\gamma-z)!(a-c-\beta+z)!}. \end{aligned} \quad (8.2.6)$$

(Majumdar [58])

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} \frac{\Delta(abc)(a+b+c+1)!}{(a-b+c)!} \left[\frac{(a-\alpha)!(c+\gamma)!(2c+1)}{(a+\alpha)!(b+\beta)!(b-\beta)!(c-\gamma)!} \right]^{\frac{1}{2}} \times \sum_z \frac{(-1)^{b+\beta+z}(2c-z)!(c+b+\alpha-z)!}{z!(c-a+b-z)!(c+\gamma-z)!(a+b+c+1-z)!}. \quad (8.2.7)$$

(Bandzaitis and Yutsis [59])

$$C_{a\alpha b\beta}^{c\gamma} = \frac{\delta_{\gamma, \alpha+\beta}}{\Delta(abc)} \left[\frac{(a+\alpha)!(a-\alpha)!(c+\gamma)!(c-\gamma)!(2c+1)}{(b+\beta)!(b-\beta)!} \right]^{\frac{1}{2}} \times \sum_z \frac{(-1)^{a-\alpha+z}(a+b-\gamma-z)!(b+c-\alpha-z)!}{z!(a-\alpha-z)!(c-\gamma-z)!(a+b+c+1-z)!}. \quad (8.2.8)$$

(Bandzaitis and Yutsis [59]).

In Eqs. (8.2.3)-(8.2.8) the summation index z assumes integer values for which all the factorial arguments are nonnegative.

8.2.2 Quasi-Binomial Representation of the Clebsch-Gordan Coefficients

The Clebsch-Gordan coefficients can be presented in the form of quasi-binomials $(u \pm v)^{(k)}$ [60]

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} \frac{\Delta(abc)}{(a+b-c)!} \left[\frac{(c-\gamma)!(c+\gamma)!(2c+1)}{(a-\alpha)!(a+\alpha)!(b-\beta)!(b+\beta)!} \right]^{\frac{1}{2}} \left[(a+\alpha)(b-\beta) - (a-\alpha)(b+\beta) \right]^{(a+b-c)}. \quad (8.2.9)$$

For calculating the quasi-binomial one should use the binomial formula with all powers replaced by quasipowers.

$$(u \pm v)^{(k)} = \sum_z (\pm 1)^z \binom{k}{z} u^{(k-z)} v^{(z)}. \quad (8.2.10)$$

Quasi-powers (or generalized powers) are defined by

$$u^{(z)} \equiv u^{(1)(z)} = \frac{\Gamma(u+1)}{\Gamma(u-z+1)},$$

$$u^{(-1)(z)} = (u+z)^{(z)} = \frac{\Gamma(u+z+1)}{\Gamma(u+1)}. \quad (8.2.11)$$

If z is integer and positive, we get

$$u^{(z)} = u(u-1)\dots(u-z+1),$$

$$u^{(-1)(z)} = (u+1)(u+2)\dots(u+z). \quad (8.2.12)$$

8.2.3 Clebsch-Gordan Coefficients and Finite Differences

The Clebsch-Gordan coefficient are related to finite differences [19]

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} (-1)^{a-c+\beta} \left[\frac{(c+\gamma)!(a+b-c)!(2c+1)}{(c-\gamma)!(c+a-b)!(c-a+b)!(a+b+c+1)!} \right]^{\frac{1}{2}} \times \left[\frac{(a-\alpha)^{(a-b-\gamma)}}{(a+\alpha)^{(a-b+\gamma)}} \right]^{\frac{1}{2}} \Delta_a^{c-\gamma} \left[\frac{(a+\alpha)^{(c+a-b)}}{(a-\alpha)^{(a-b-c)}} \right]. \quad (8.2.13)$$

where the difference of order k with respect to the argument a is given by

$$\Delta_a^k f(a) = \sum_{n=0}^k (-1)^{k-n} \frac{k!}{n!(k-n)!} f(a+n). \quad (8.2.14)$$

In particular,

$$\Delta_a f(a) = f(a+1) - f(a),$$

$$\Delta_a a^{(n)} = n a^{(n-1)}. \quad (8.2.15)$$

Finite differences are analogous to differentials, while quasi-powers are analogous to ordinary powers. From this it can be concluded that the Clebsch-Gordan coefficients are analogous to the Wigner D -functions [18, 61]. To illustrate this similarity let us introduce the notations

$$j = b, \quad \mu = c - a, \quad \nu = \beta, \quad P = \frac{1}{2}(a + b + c), \quad Q = a + \alpha - \frac{1}{2}(a + b + c). \quad (8.2.16)$$

Now, Eq. (8.2.3) may be rewritten in terms of quasi-powers

$$\begin{aligned} \sqrt{\frac{a+b+c+1}{2c+1}} C_{a\alpha b\beta}^{c\gamma} &= \left[\frac{(j+\mu)!(j-\mu)!(j+\nu)!(j-\nu)!}{(2P)^{(2j)}(P-Q)^{(j+\mu)}(P-Q)^{(j+\nu)}(P+Q)^{(-\mu-\nu)}} \right]^{\frac{1}{2}} \\ &\times \sum_x (-1)^x \frac{(P-Q)^{(j+\mu+z)}(P+Q)^{(j-\mu-z)}}{z!(j-\mu-z)!(j+\nu-z)!(\mu-\nu+z)!}. \end{aligned} \quad (8.2.17)$$

Replacing quasi-powers in (8.2.17) by ordinary powers and introducing the angle ϑ according to

$$\cos \theta = \frac{Q}{P}, \quad \cos \frac{\theta}{2} = \sqrt{\frac{P+Q}{2P}}, \quad \sin \frac{\theta}{2} = \sqrt{\frac{P-Q}{2P}}, \quad (8.2.18)$$

we see that (8.2.17) reduces to the following relation for the Wigner D -function

$$\begin{aligned} d_{\nu\mu}^j(\theta) &= \left[\frac{(j+\mu)!(j-\mu)!(j+\nu)!(j-\nu)!}{(2P)^{2j}(P-Q)^{j+\mu}(P-Q)^{j+\nu}(P+Q)^{-\mu-\nu}} \right]^{\frac{1}{2}} \\ &\times \sum_x (-1)^x \frac{(P-Q)^{j+\mu+x}(P+Q)^{j-\mu-x}}{z!(j-\mu-z)!(j+\nu-z)!(\mu-\nu+z)!}. \end{aligned} \quad (8.2.19)$$

(8.2.19) is similar to Eq. (4.3.4). The analogy between (8.2.17) and (8.2.19) permits us to connect the recursion relations, symmetry properties, etc. of the Wigner D -functions and the Clebsch-Gordan coefficients [61]. Since for $u \gg z, u \gg 1$ quasi-powers become asymptotically equal to ordinary powers, a Clebsch-Gordan coefficient turns into a D -function in the large-momentum limit, see Eq. (8.9.1).

8.2.4 Expressions for the Clebsch-Gordan Coefficients in Terms of Binomial Coefficients [116]

We can express the Clebsch-Gordan coefficients in terms of a sum over binomials as

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} \left[\frac{\binom{2a}{J-2c} \binom{2b}{J-2c}}{\binom{J+1}{J-2c} \binom{2a}{a-\alpha} \binom{2b}{b-\beta} \binom{2c}{c-\gamma}} \right]^{\frac{1}{2}} \sum_x (-1)^x \binom{J-2c}{z} \binom{J-2b}{a-\alpha-z} \binom{J-2a}{b+\beta-z}, \quad (8.2.20)$$

where $J = a + b + c$.

8.2.5 Representations of the Clebsch-Gordan Coefficients in Terms of the Hypergeometric Functions

The Clebsch-Gordan coefficients may be expressed as generalized hypergeometric functions ${}_3F_2$ of unit argument,

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} \frac{\Delta(abc)}{(a+b-c)!(-b+c+\alpha)!(-a+c-\beta)!} \left[\frac{(a+\alpha)!(b-\beta)!(c+\gamma)!(c-\gamma)!(2c+1)}{(a-\alpha)!(b+\beta)!} \right]^{\frac{1}{2}} \\ &\times {}_3F_2 \left[\begin{matrix} -a-b+c, -a+\alpha, -b-\beta \\ -a+c-\beta+1, -b+c+\alpha+1 \end{matrix} \middle| 1 \right], \end{aligned} \quad (8.2.21)$$

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} (-1)^{a-\alpha} \frac{\Delta(abc)(b+c-\alpha)!}{(a-b+c)!(-a+b+c)!(b-c+\alpha)!} \left[\frac{(a+\alpha)!(b-\beta)!(c+\gamma)!(2c+1)}{(a-\alpha)!(b+\beta)!(c-\gamma)!} \right]^{\frac{1}{2}} \\ &\times {}_3F_2 \left[\begin{matrix} a+\alpha+1, -a+\alpha, -c+\gamma \\ -b-c+\alpha, b-c+\alpha+1 \end{matrix} \middle| 1 \right], \end{aligned} \quad (8.2.22)$$

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \delta_{\gamma, \alpha+\beta} (-1)^{b+\beta} \frac{\Delta(abc)(b+c+\alpha)!}{(-a+b+c)!(a-b-\gamma)!} \left[\frac{(a-\alpha)!(c-\gamma)!(2c+1)}{(a+\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!} \right]^{\frac{1}{2}} \\ &\times {}_3F_2 \left[\begin{matrix} a-b-c, a-\alpha+1, -c-\gamma \\ a-b-\gamma+1, -b-c-\alpha \end{matrix} \middle| 1 \right] \end{aligned} \quad (8.2.23)$$

¹² This follows from the fact that the Clebsch-Gordan coefficients may be expressed in terms of orthogonal polynomials of a discrete variable (Hahn polynomials); see Refs. [134, 135].

Rose [4]

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} (-1)^{b+\beta} \frac{\Delta(abc)(2c)!}{(a-b+c)!(-a+b+c)!(a-c-\beta)!} \left[\frac{(a+\alpha)!(a-\alpha)!(b-\beta)!(2c+1)}{(b+\beta)!(c+\gamma)!(c-\gamma)!} \right]^{\frac{1}{2}} \quad (8.2.24)$$

$$\times {}_3F_2 \left[\begin{matrix} a-b-c, a+b-c+1, -c-\gamma \\ -2c, a-c-\beta+1 \end{matrix} \middle| 1 \right]$$

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} (-1)^{b+\beta} \frac{\Delta(abc)(2c)!(b+c+\alpha)!}{(a-b+c)!(-a+b+c)!} \left[\frac{(a+\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!}{(a-\alpha)} \right]^{\frac{1}{2}} \quad (8.2.25)$$

$$\times {}_3F_2 \left[\begin{matrix} a-b-c, -a-b-c-1, -c-\gamma \\ -2c, -b-c-\alpha \end{matrix} \middle| 1 \right]$$

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} (-1)^{a-\alpha} \frac{(a+b-\gamma)!(b+c-\alpha)!}{\Delta(abc)(a+b+c+1)!} \left[\frac{(a+\alpha)!(c+\gamma)!(2c+1)}{(a-\alpha)!(b+\beta)!(b-\beta)!(c-\gamma)!} \right]^{\frac{1}{2}} \quad (8.2.26)$$

$$\times {}_3F_2 \left[\begin{matrix} -a-b-c-1, -a+\alpha, -c+\gamma \\ -a-b+\gamma, -b-c+\alpha \end{matrix} \middle| 1 \right].$$

Equations (8.2.21)-(8.2.26) may be associated with Eqs. (8.2.3)-(8.2.8), respectively.

According to Ref. [30], all ${}_3F_2$ -functions of unit argument, which are given by finite sums, may be expressed in terms of the Clebsch-Gordan coefficients. Any Clebsch-Gordan coefficient may also be represented in the form of derivatives of the hypergeometric functions [62]:

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma, \alpha+\beta} (-1)^{a+b-c} \frac{\Delta(abc)}{(a+b-c)!(-a+b+c)!(a-b+\gamma)!} \left[\frac{(a+\alpha)!(a-\alpha)!(b+\beta)!(c+\gamma)!(2c+1)}{(b-\beta)!(c-\gamma)!} \right]^{\frac{1}{2}} \quad (8.2.27)$$

$$\times \left\{ \frac{d^{b-\beta}}{(dt)^{b-\beta}} [(1-t)^{a+b-c} F(a-b-c, -c+\gamma; a-b+\gamma+1; t)] \right\}_{t=0}.$$

8.2.6 Representations of the $3j$ m Symbols in the Form of Algebraic Sums

The explicit form of the Wigner $3jm$ symbols may be obtained from Eqs. (8.2.3)-(8.2.8) for the Clebsch-Gordan coefficients. Rewriting these equations in terms of the R -symbol elements, we get the following relations for the $3jm$ symbols [20]

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{32}} \left[\frac{\prod_{i,j=1}^3 (R_{ij})!}{(J+1)!} \right]^{\frac{1}{2}} \quad (8.2.28)$$

$$\times \sum_x (-1)^x \frac{1}{z! (R_{13}-z)! (R_{22}-z)! (R_{31}-z)! (R_{33}-R_{22}+z)! (R_{11}-R_{22}+z)!},$$

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{31}+R_{32}} \left[\frac{R_{13}!R_{23}!R_{31}!R_{32}!R_{33}!}{(J+1)!R_{11}!R_{12}!R_{21}!R_{22}!} \right]^{\frac{1}{2}} \quad (8.2.29)$$

$$\times \sum_x (-1)^x \frac{(R_{21}+z)! (R_{11}+R_{31}-z)!}{z! (R_{31}-z)! (R_{23}-z)! (R_{13}-R_{31}+z)!},$$

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{22}+R_{32}} \left[\frac{R_{11}!R_{12}!R_{13}!R_{23}!R_{33}!}{(J+1)!R_{21}!R_{22}!R_{31}!R_{32}!} \right]^{\frac{1}{2}} \sum_z (-1)^z \frac{(R_{31}+z)! (R_{32}+R_{33}-z)!}{z! (R_{11}-z)! (R_{33}-z)! (R_{12}-R_{33}+z)!}, \quad (8.2.30)$$

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{22}+R_{32}} \left[\frac{R_{11}!R_{21}!R_{31}!R_{32}!R_{33}!}{(J+1)!R_{12}!R_{13}!R_{22}!R_{33}!} \right]^{\frac{1}{2}} \sum_z (-1)^z \frac{(R_{13}+z)! (R_{23}+R_{33}-z)!}{z! (R_{11}-z)! (R_{33}-z)! (R_{21}-R_{33}+z)!}, \quad (8.2.31)$$

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{22}+R_{32}} \left[\frac{(J+1)!R_{11}!R_{13}!R_{31}!R_{39}!}{R_{12}!R_{21}!R_{22}!R_{23}!R_{32}!} \right]^{\frac{1}{2}} \sum_z (-1)^z \frac{(R_{11}+R_{21}-z)! (R_{23}+R_{33}-z)!}{z! (R_{11}-z)! (R_{33}-z)! (J+1-z)!}, \quad (8.2.32)$$

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{31}+R_{32}} \left[\frac{(J+1)!R_{21}!R_{23}!R_{31}!R_{39}!}{R_{11}!R_{12}!R_{13}!R_{22}!R_{32}!} \right]^{\frac{1}{2}} \sum_z (-1)^z \frac{(R_{11}+R_{31}-z)! (R_{13}+R_{23}-z)!}{z! (R_{23}-z)! (R_{31}-z)! (J+1-z)!}, \quad (8.2.33)$$

where $J = a + b + c$.

8.2.7 Quasi-binomial Representations of the $3jm$ Symbols

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{R_{21}+R_{32}} \left[\frac{R_{21}^{(-1)(R_{31}-R_{13})} R_{31}^{(-1)(R_{32}-R_{13})} R_{32}^{(-1)(R_{22}-R_{13})} R_{22}^{(-1)(R_{21}-R_{13})}}{(J+1)!R_{13}!} \right]^{\frac{1}{4}} (u-v)^{(R_{13})}. \quad (8.2.34)$$

The quasi-binomial should be evaluated according to Eqs. (8.2.10) and (8.2.11). The variables u and v may be chosen in different ways [20]:

$$\begin{aligned} (a) \quad u &= R_{21}^{(1)} R_{32}^{(1)}, & v &= R_{22}^{(1)} R_{31}^{(1)}, \\ (b) \quad u &= R_{21}^{(1)} R_{11}^{(-1)}, & v &= R_{22}^{(1)} R_{12}^{(-1)}, \\ (c) \quad u &= R_{11}^{(-1)} R_{23}^{(-1)}, & v &= R_{32}^{(1)} (J+1)^{(1)}. \end{aligned} \quad (8.2.35)$$

8.3 Integral representations

8.3.1 Integrals Involving Algebraic Functions

The Clebsch-Gordan coefficients may be represented by the following integrals [19] ($J = a + b + c$)

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \frac{(-1)^{a-c+\beta}}{2^{J+1}} \left[\frac{(c+\gamma)!(J-2c)!(J+1)!(2c+1)}{(a-\alpha)!(a+\alpha)!(b-\beta)!(b+\beta)!(c-\gamma)!(J-2a)!(J-2b)!} \right]^{\frac{1}{2}} \\ &\quad \times \int_{-1}^1 (1-x)^{a-\alpha} (1+x)^{b-\beta} \frac{d^{c-\gamma}}{(dx)^{c-\gamma}} [(1-x)^{J-2a} (1+x)^{J-2b}] dx, \end{aligned} \quad (8.3.1)$$

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \frac{(-1)^{a-c+\beta}}{2^{J+1}} \left[\frac{(J-2b)!(J-2c)!(J+1)!(2c+1)}{(a-\alpha)!(a+\alpha)!(b-\beta)!(b+\beta)!(c-\gamma)!(c+\gamma)!(J-2a)!} \right]^{\frac{1}{2}} \\ &\quad \times \int_{-1}^1 (1-x)^{b+\beta} (1+x)^{b-\beta} \frac{d^{J-2a}}{(dx)^{J-2a}} [(1-x)^{c-\gamma} (1+x)^{c+\gamma}] dx. \end{aligned} \quad (8.3.2)$$

In particular, if $\alpha = \beta = \gamma = 0$, then

$$C_{a0b0}^{c0} = \frac{(-1)^{a-c}}{2^{J+1}a!b!c!} \left[\frac{(J-2b)!(J-2c)!(J+1)!(2c+1)}{(J-2a)!} \right]^{\frac{1}{2}} \int_{-1}^1 (1-x^2)^b \frac{d^{J-2a}}{(dx)^{J-2a}} [(1-x^2)^c] dx. \quad (8.3.3)$$

8.3.2 Integrals Involving the Wigner D-Functions

The Clebsch-Gordan coefficients can be written as the integral over three Wigner D functions,

$$C_{a\alpha b\beta}^{c\gamma} = \frac{1}{8\pi^2} \left[\frac{(a+b+c+1)!(a+b-c)!(2c+1)}{(2a)!(2b)!} \right]^{\frac{1}{2}} \int dR D_{\alpha a}^a(R) D_{\beta -b}^b(R) D_{\gamma a-b}^{c*}(R), \quad (8.3.4)$$

where R means the set of the Euler angles (see Sec. 4.12). We also have a more compact relation involving one d function (Akim and Levin, [62])

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \frac{(-1)^{a+c-\alpha-\gamma}}{2^{a+b+1}} \left[\frac{(J-2c)!(J+1)!(2c+1)}{(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!} \right]^{\frac{1}{2}} \\ &\quad \times \int_{-1}^1 (\sin \theta)^{a+b} \left(\frac{1-\cos \theta}{1+\cos \theta} \right)^{\frac{\alpha-\beta}{2}} d_{b-a\gamma}^c(\theta) d \cos \theta. \end{aligned} \quad (8.3.5)$$

8.3.3 Integral Involving the Spherical Harmonics

If a, b, c are integers and $J = a + b + c = 2g$ is even (i.e., g is integer), then

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{g-c} \left[\frac{4\pi}{(2a+1)(2b+1)} \right]^{\frac{1}{2}} \frac{(g-a)!(g-b)!(g-c)!}{\Delta(abc)g!} \int_0^{2\pi} d\varphi \int_0^\pi d\vartheta \sin \vartheta Y_{a\alpha}(\vartheta, \varphi) Y_{b\beta}(\vartheta, \varphi) Y_{c\gamma}^*(\vartheta, \varphi). \quad (8.3.6)$$

8.3.4 Integral Representations for Products of the Clebsch-Gordan Coefficients

$$C_{a\alpha b\beta}^{c\gamma} C_{a\alpha' b\beta'}^{c\gamma'} = \frac{2c+1}{8\pi^2} \int dR D_{\alpha\alpha'}^a(R) D_{\beta\beta'}^b(R) D_{\gamma\gamma'}^{c*}(R). \quad (8.3.7)$$

In particular,

$$[C_{a0b0}^{c0}]^2 = \frac{2c+1}{2} \int_0^\pi d\theta \sin\theta P_a(\cos\theta) P_b(\cos\theta) P_c(\cos\theta). \quad (8.3.8)$$

8.4 Symmetry properties

The simplest way to formulate the symmetry properties of the Clebsch-Gordan coefficients and $3jm$ symbols is to use the representations of these coefficients in the form of the Regge R -symbols [55] (see Sec. 8.1.3).

8.4.1 Symmetry Properties of the R -Symbols

The R -symbol possesses the following symmetry properties [55].

(a) *Permutations of columns*

$$\left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| = \varepsilon \left\| \begin{array}{ccc} R_{1i} & R_{1k} & R_{1l} \\ R_{2i} & R_{2k} & R_{2l} \\ R_{3i} & R_{3k} & R_{3l} \end{array} \right\|, \quad (8.4.1)$$

where

$$c = \begin{cases} +1 & \text{for cyclic permutation (even number of permutations)} \\ (-1)^J & \text{for non-cyclic permutation (odd number of permutations)} \end{cases}, \quad (8.4.2)$$

and

$$J \equiv \sum_k R_{ik} = \sum_i R_{ik};$$

(b) *Permutations of rows*

$$\left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| = \left\| \begin{array}{ccc} R_{i1} & R_{i2} & R_{i3} \\ R_{k1} & R_{k2} & R_{k3} \\ R_{l1} & R_{l2} & R_{l3} \end{array} \right\|, \quad (8.4.3)$$

(c) *Transposition*

$$\left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| = \left\| \begin{array}{ccc} R_{11} & R_{21} & R_{31} \\ R_{12} & R_{22} & R_{32} \\ R_{13} & R_{23} & R_{33} \end{array} \right\|. \quad (8.4.4)$$

These symmetry transformations relate $6 \times 6 \times 2 = 72$ generally different coefficients.

8.4.2 Symmetry Properties of the $3jm$ Symbols

The above relations are equivalent to the following symmetry properties of the $3jm$ symbol.

(a) *Classical symmetries* [54]

(i) Permutations of columns (corresponding to permutations of columns of the R -symbol):

$$\begin{aligned} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= \begin{pmatrix} b & c & a \\ \beta & \gamma & \alpha \end{pmatrix} = \begin{pmatrix} c & a & b \\ \gamma & \alpha & \beta \end{pmatrix} \\ &= (-1)^{a+b+c} \begin{pmatrix} a & c & b \\ \alpha & \gamma & \beta \end{pmatrix} = (-1)^{a+b+c} \begin{pmatrix} b & a & c \\ \beta & \alpha & \gamma \end{pmatrix} = (-1)^{\alpha+b+c} \begin{pmatrix} c & b & a \\ \gamma & \beta & \alpha \end{pmatrix}. \end{aligned} \quad (8.4.5)$$

(ii) Change of signs of momentum projections (corresponding to the permutation of the second and third rows of the R -symbol)

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{a+b+c} \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} \quad (8.4.6)$$

(b) *Regge symmetries* [55]

(iii) Replacement of arguments:

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{a+b+c} \begin{pmatrix} \frac{b+c+\alpha}{2} & \frac{a+c+\beta}{2} & \frac{a+b+\gamma}{2} \\ a - \frac{b+c-\alpha}{2} & b - \frac{a+c-\beta}{2} & c - \frac{a+b-\gamma}{2} \end{pmatrix} \quad (8.4.7)$$

(corresponding to the permutation of the first and third rows of the R -symbol) and

(iv)

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = \begin{pmatrix} a & \frac{b+c-\alpha}{2} & \frac{b+c+\alpha}{2} \\ -b+c & \frac{b-c-\alpha}{2} - \gamma & \frac{b-c+\alpha}{2} + \gamma \end{pmatrix} \quad (8.4.8)$$

This corresponds to the transposition of the R -symbol. According to the above symmetry properties, 72 formally different $3jm$ symbols have the same absolute values. One can decompose these symbols into six groups (12 coefficients in each group). The $3jm$ symbols which belong to different groups correspond to triangles of different forms but the same perimeter $J = a + b + c$. On the other hand, all $3jm$ symbols which belong to one group are related by the classical symmetry properties (8.4.5) and (8.4.6). In (8.4.9) we list one $3jm$ symbol from each group.

$$\begin{aligned} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= \begin{pmatrix} a & \frac{b+c-\alpha}{2} & \frac{b+c+\alpha}{2} \\ c-b & \frac{b+\beta-c-\gamma}{2} & \frac{b-\beta-c+\gamma}{2} \end{pmatrix} = \begin{pmatrix} b & \frac{a+c-\beta}{2} & \frac{a+c+\beta}{2} \\ a-c & \frac{-a-\alpha+c+\gamma}{2} & \frac{-a+\alpha+c-\gamma}{2} \end{pmatrix} \\ &= \begin{pmatrix} c & \frac{a+b-\gamma}{2} & \frac{a+b+\gamma}{2} \\ b-a & \frac{a+\alpha-b-\beta}{2} & \frac{a-\alpha-b+\beta}{2} \end{pmatrix} = \begin{pmatrix} \frac{b+c-\alpha}{2} & \frac{a+c-\beta}{2} & \frac{a+b+\gamma}{2} \\ a-\alpha & b-\beta & c-\gamma \end{pmatrix} \\ &= \begin{pmatrix} \frac{b+c+\alpha}{2} & \frac{a+c+\beta}{2} & \frac{a+b+\gamma}{2} \\ \frac{b+c+\alpha}{2} - a - \alpha & \frac{a+c+\beta}{2} - b - \beta & \frac{a+b+\gamma}{2} - c - \gamma \end{pmatrix}. \end{aligned} \quad (8.4.9)$$

8.4.3 Symmetry Properties of the Clebsch-Gordan Coefficients

(i)

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= (-1)^{a+b-c} C_{b\beta a\alpha}^{c\gamma} = (-1)^{a-\alpha} \sqrt{\frac{2c+1}{2b+1}} C_{a\alpha c-\gamma}^{b-\beta} = (-1)^{a-\alpha} \sqrt{\frac{2c+1}{2b+1}} C_{c\gamma a-\alpha}^{b\beta} \\ &= (-1)^{b+\beta} \sqrt{\frac{2c+1}{2a+1}} C_{c-\gamma b\beta}^{a-\alpha} = (-1)^{b+\beta} \sqrt{\frac{2c+1}{2a+1}} C_{b-\beta c\gamma}^{a\alpha}. \end{aligned} \quad (8.4.10)$$

(ii)

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{a+b-c} C_{a-\alpha b-\beta}^{c-\gamma}. \quad (8.4.11)$$

(iii)

$$C_{a\alpha b\beta}^{c\gamma} = C_{a'\alpha' b'\beta'}^{c'\gamma'}, \quad (8.4.12)$$

where

$$\begin{aligned} 2a' &= (a + \alpha) + (b + \beta), & 2a &= (a' + a') + (b' + \beta'), \\ 2\alpha' &= (a + \alpha) - (b + \beta), & 2\alpha &= (a' + \alpha') - (b' + \beta'), \\ 2b' &= (a - \alpha) + (b - \beta), & 2b &= (a' - \alpha') + (b' - \beta'), \\ 2\beta' &= (a - \alpha) - (b - \beta), & 2\beta &= (a' - \alpha') - (b' - \beta'), \end{aligned} \quad (8.4.13)$$

$$\begin{aligned} c' &= c, & c &= c' \\ \gamma' &= a - b, & \gamma &= a' - b' \end{aligned}$$

(iv)

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{b+\beta} \sqrt{\frac{2c+1}{2c'+1}} C_{a'\alpha' b'\beta'}^{c'\gamma'}, \quad (8.4.14)$$

where

$$\begin{aligned}
2a' &= (b - \beta) + (c + \gamma), & 2a &= (b' - \beta') + (c' + \gamma'), \\
2\alpha' &= 2(a + \alpha) - (b - \beta) - (c + \gamma), & 2\alpha &= 2(a' + \alpha') - (b' - \beta') - (c' + \gamma'), \\
2b' &= (a - \alpha) + (c + \gamma), & 2b &= (a' - \alpha') + (c' + \gamma'), \\
2\beta' &= 2(b + \beta) - (a - \alpha) - (c + \gamma), & 2\beta &= 2(b' + \beta') - (a' - \alpha') - (c' + \gamma'), \\
2c' &= (a - \alpha) + (b - \beta), & 2c &= (a' - \alpha') + (b' - \beta'), \\
2\gamma' &= (a - \alpha) + (b - \beta) - 2(c - \gamma), & 2\gamma &= (a' - \alpha') + (b' - \beta') - 2(c' - \gamma').
\end{aligned} \tag{8.4.15}$$

8.4.4 "Mirror" Symmetry

The original relations which define the $3jm$ symbols and the Clebsch-Gordan coefficients may be extended to the domain of negative integer or half-integer arguments a, b, c . In this case the following symmetry relations are valid [20] (here we denote $\bar{c} \equiv -c - 1, \gamma \equiv -\gamma$, etc.).

(a) $3jm$ symbols

The relations for the $3jm$ symbols are

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = (-1)^{b-c-\alpha} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = i(-1)^{a+\alpha-\beta} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} = i(-1)^{a+b+c} \begin{pmatrix} \bar{a} & b & \bar{c} \\ \alpha & \beta & \gamma \end{pmatrix}. \tag{8.4.16}$$

The $3j$ m symbols with negative momenta obey the symmetry relations (8.4.5) and (8.4.6) provided $\bar{a}, \bar{b}, \bar{c}$ are replaced by $-a, -b, -c$, respectively, in the phase factors of these relations.

(b) Clebsch-Gordan symbols

The relations for the Clebsch-Gordan coefficients are

$$\begin{aligned}
C_{a\alpha b\beta}^{c\gamma} &= (-1)^{a+b-c} C_{a\alpha\delta\beta}^{c\gamma} = (-1)^{b-c-\alpha} C_{a\alpha b\beta}^{c\gamma} = (-1)^{b+\beta} C_{a\alpha b\beta}^{c\gamma} = (-1)^{a-\alpha} C_{a\alpha\delta\beta}^{c\gamma} = C_{a\alpha b\beta}^{c\gamma} \\
&= \sqrt{\frac{2c+1}{2b+1}} C_{a\alpha c\gamma}^{b\beta} = \sqrt{\frac{2c+1}{2a+1}} C_{c\gamma b\beta}^{a\alpha} = \sqrt{\frac{2c+1}{2a+1}} C_{c\gamma b\beta}^{a\alpha}.
\end{aligned} \tag{8.4.17}$$

From (8.4.17) one gets

$$C_{a\bar{a}b\bar{\beta}}^{c\bar{\gamma}} = C_{\bar{a}\alpha\bar{b}\bar{\beta}}^{\bar{c}\gamma}, \tag{8.4.18}$$

$$C_{\bar{a}\bar{a}b\beta}^{c\gamma} = C_{a\alpha\bar{b}\bar{\beta}}^{\bar{c}\bar{\gamma}}, \tag{8.4.19}$$

$$C_{a\alpha\bar{b}\bar{\beta}}^{c\gamma} = C_{\bar{a}\bar{a}b\beta}^{\bar{c}\bar{\gamma}}. \tag{8.4.20}$$

When tabulating formulas for the Clebsch-Gordan coefficients, it is convenient to use the "mirror" symmetry properties in the form

$$C_{a\alpha b\beta}^{a+k\gamma} = (-1)^{b+\beta} C_{a\alpha b\beta}^{a-k\gamma}. \tag{8.4.21}$$

8.4.5 Properties of the Vector-Addition Coefficients under Transformations of the Coordinate System and Time Reversal

The vector-addition coefficients are not invariant under transformations in question, although formally they are independent of coordinates and time.

(a) Rotation of the coordinate system.

The projections of angular momenta vary under rotations of the coordinate system. The $3jm$ symbols and the Clebsch-Gordan coefficients in a new coordinate system may be expressed as superpositions of the corresponding quantities in the initial coordinate system. Thus

$$\begin{aligned}
\hat{R} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &\equiv \begin{pmatrix} a & b & c \\ \alpha' & \beta' & \gamma' \end{pmatrix} = \sum_{\alpha\beta\gamma} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} D_{\alpha\alpha'}^a(R) D_{\beta\beta'}^b(R) D_{\gamma\gamma'}^c(R), \\
\hat{R} C_{a\alpha b\beta}^{c\gamma} &\equiv C_{a\alpha' b\beta'}^{c\gamma'} = \sum_{\alpha\beta\gamma} C_{a\alpha b\beta}^{c\gamma} D_{\alpha\alpha'}^a(R) D_{\beta\beta'}^b(R) D_{\gamma\gamma'}^{c*}(R).
\end{aligned} \tag{8.4.22}$$

(b) *Inversion of the coordinate system.*

The $3jm$ symbols and the Clebsch-Gordan coefficients are invariant under inversion of the coordinate system, because angular momentum is a pseudovector.

$$\begin{aligned}\hat{P}_r \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix}, \\ \hat{P}_r C_{a\alpha b\beta}^{c\gamma} &= C_{a\alpha b\beta}^{c\gamma}.\end{aligned}\tag{8.4.23}$$

(c) *(Time reversal.*

The time reversal changes signs of projections of all angular momenta (reversal of the rotation direction). Hence

$$\begin{aligned}\hat{P}_t \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} &= \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} = (-1)^{a+b+c} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix}, \\ \hat{P}_t C_{a\alpha b\beta}^{c\gamma} &= C_{a-\alpha b-\beta}^{c-\gamma} = (-1)^{a+b-c} C_{a\alpha b\beta}^{c\gamma}.\end{aligned}\tag{8.4.24}$$

8.5 Explicit forms of the Clebsch-Gordan coefficients for special values of the arguments

8.5.1 Special Values of Momenta a, b, c

(a) $c = 0$ or $b = 0$.

$$C_{a\alpha b\beta}^{00} = (-1)^{a-\alpha} \frac{\delta_{\alpha b} \delta_{\alpha, -\beta}}{\sqrt{2a+1}},\tag{8.5.1}$$

$$C_{a\alpha 00}^{\alpha\gamma} = \delta_{ac} \delta_{\alpha\gamma}.\tag{8.5.2}$$

(b) $c = a + b$

$$C_{a\alpha b\beta}^{a+b\alpha+\beta} = \left[\frac{(2a)!(2b)!(a+b+\alpha+\beta)!(a+b-\alpha-\beta)!}{(2a+2b)!(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!} \right]^{\frac{1}{2}}.\tag{8.5.3}$$

The following two formulas are equivalent to (8.5.3) (i and n are integer)

$$C_{a\alpha-n+i\ b\ b-i+1}^{a+b\ a+b-n+1} = \left[\frac{\binom{2a}{n-i} \binom{2b}{i-1}}{\binom{2a+2b}{n-1}} \right]^{\frac{1}{2}},\tag{8.5.4}$$

$$C_{a\alpha+n-i\ b\ -b+i-1}^{a+b\ a-b+n-1} = \left[\frac{\binom{2a}{i-n} \binom{2b}{i-1}}{\binom{2a+2b}{2a+n-1}} \right]^{\frac{1}{2}}.\tag{8.5.5}$$

In particular,

$$C_{a\alpha b\beta}^{a+b\ a+b} = 1,\tag{8.5.6}$$

$$C_{a\alpha b\beta}^{a+b\ a-b} = \left[\frac{(2a)!(2b)!}{(2a+2b)!} \right]^{\frac{1}{2}}.\tag{8.5.7}$$

(c) $c = a + b - 1$

$$C_{a\alpha b\beta}^{a+b-1\ \alpha+\beta} = 2(b\alpha - a\beta) \left[\frac{(2a+2b-1)(2a-1)!(2b-1)!(a+b+\alpha+\beta-1)!(a+b-\alpha-\beta-1)!}{(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(2a+2b)!} \right]^{\frac{1}{2}}.\tag{8.5.8}$$

In particular,

$$C_{a\alpha b\beta}^{a+b-1\alpha+\beta} = 0, \quad \text{if} \quad \frac{\alpha}{\beta} = \frac{a}{b}. \quad (8.5.9)$$

$$C_{a\alpha-1b\beta}^{a+b-1a+b-1} = -\sqrt{\frac{b}{a+b}}. \quad (8.5.10)$$

$$C_{a\alpha b\beta-1}^{a+b-1a+b-1} = \sqrt{\frac{a}{a+b}}. \quad (8.5.11)$$

$$C_{a\alpha b-b}^{a+b-1a-b} = \left[\frac{(2a)!(2b)!(2a+2b-1)!}{(2a+2b)!} \right]^{\frac{1}{2}}, \quad (8.5.12)$$

(d) $c = a - b (a \geq b)$

$$C_{a\alpha b\beta}^{a-b\alpha+\beta} = (-1)^{b+\beta} \left[\frac{(a+\alpha)!(a-\alpha)!(2b)!(2a-2b+1)!}{(2a+1)!(b+\beta)!(b-\beta)!(a-b+\alpha+\beta)!(a-b-\alpha-\beta)!} \right]^{\frac{1}{2}}. \quad (8.5.13)$$

The following two formulas are equivalent to (8.5.13) (i and n are integer)

$$C_{a\alpha-n+ib\beta-i+1}^{a-b\alpha+b-n+1} = (-1)^{2b-i+1} \left[\frac{2a-2b+1}{2a+1} \frac{\binom{2b}{i-1} \binom{2a-2b}{2a-n+1}}{\binom{2a}{n-i}} \right]^{\frac{1}{2}}, \quad (8.5.14)$$

$$C_{a\alpha-n+ib\beta-i+1}^{a-b\alpha+b-n+1} = (-1)^{i+1} \left[\frac{2a-2b+1}{2a+1} \frac{\binom{2b}{-i+1} \binom{2a-2b}{n-1}}{\binom{2a}{n-i}} \right]^{\frac{1}{2}}. \quad (8.5.15)$$

In particular

$$C_{a\alpha b-b}^{a-b\alpha-b} = \sqrt{\frac{2a-2b+1}{2a+1}}. \quad (8.5.16)$$

(e) $c = a - b + 1 (a + 1 \geq b)$

$$C_{a\alpha b\beta}^{a-b+1\alpha+\beta} = (-1)^{b+\beta+1} 2(a\beta + b\alpha + \beta) \left[\frac{(2a-2b+3)(2b-1)!(2a-2b+1)!(a+\alpha)!(a-\alpha)!}{(2a+2)!(b+\beta)!(b-\beta)!(a-b+\alpha+\beta+1)!(a-b-\alpha-\beta+1)!} \right]^{\frac{1}{2}}. \quad (8.5.17)$$

In particular,

$$C_{a\alpha b-b}^{a-b+1a-b} = \left[\frac{(2a-2b+3)2b}{(2a+2)(2a+1)} \right]^{\frac{1}{2}}. \quad (8.5.18)$$

(f) $c = a + b - 2$

$$C_{a\alpha b\beta}^{a+b-2\alpha+\beta} = \left[\frac{2a(2a-1)2b(2b-1)}{2(2a+2b-2)(2a+2b-1)} \right]^{\frac{1}{2}} \left[\binom{2a}{a-\alpha} \binom{2b}{b-\beta} \binom{2a+2b-4}{a+b-\alpha-\beta-2} \right]^{-\frac{1}{2}} \\ \times \left\{ \binom{2a-2}{a-\alpha} \binom{2b-2}{b+\beta} - 2 \binom{2a-2}{a-\alpha-1} \binom{2b-2}{b+\beta-1} + \binom{2a-2}{a-\alpha-2} \binom{2b-2}{b+\beta-2} \right\}. \quad (8.5.19)$$

In particular,

$$C_{a\alpha b-b}^{a+b-2a-b} = \left[\frac{(2a)!(2b)!(2a+2b-3)!}{2(2a+2b-1)!} \right]^{\frac{1}{2}}. \quad (8.5.20)$$

(g) $c = a - b + 2 (a + 2 \geq b)$

$$C_{a\alpha b\beta}^{a-b+2\alpha+\beta} = \left[\frac{(2b-1)2b(2a-2b+5)(2a-2b+4)(2a-2b+3)}{2(2a+1)(2a+2)(2a+3)} \right]^{\frac{1}{2}} \\ \times \left[\binom{2a}{a-\alpha} \binom{2b}{b-\beta} \binom{2a-2b+4}{a-b-\alpha-\beta+2} \right]^{-\frac{1}{2}} (-1)^{b+\beta} \left\{ \binom{2b-2}{b+\beta} \binom{2a-2b+2}{a-b-\alpha-\beta} \right. \\ \left. - 2 \binom{2b-2}{b+\beta-1} \binom{2a-2b+2}{a-b-\alpha-\beta+1} + \binom{2b-2}{b+\beta-2} \binom{2a-2b+2}{a-b-\alpha-\beta+2} \right\}. \quad (8.5.21)$$

In particular,

$$C_{a \ a \ b-b}^{a-b+2 \ a-b} = \left[\frac{(2a-2b+5)2b(2b-1)}{(2a+1)(2a+2)(2a+3)} \right]^{\frac{1}{2}}. \quad (8.5.22)$$

(h) $a = b, \alpha = \beta$

$$C_{a \ \alpha \ a \ \alpha}^{c \ \gamma} = \begin{cases} 0, & \text{if } 2a + c = 2g + 1, \\ \delta_{\gamma, 2\alpha} \frac{(-1)^{g-c} \sqrt{2c+1} g!}{\left(\frac{c+1}{2}\right)! \left(\frac{c-1}{2}\right)! (g-c)!} \left[\frac{(c+\gamma)!(c-\gamma)!(2g-2c)!}{(2g+1)!} \right]^{\frac{1}{2}}, & \text{if } 2a + c = 2g, \end{cases} \quad (8.5.23)$$

g is integer and positive.

(i) $b = a + \frac{1}{2}, \beta = \alpha \pm \frac{1}{2}, 2a + c = 2g$

$$\begin{aligned} C_{a \ \alpha \ a + \frac{1}{2} \ \alpha + \frac{1}{2}}^{c + \frac{1}{2} \ 2\alpha + \frac{1}{2}} &= \left[\frac{c + 2\alpha + 1}{a + \alpha + 1} \right]^{\frac{1}{2}} C, \\ C_{a \ \alpha \ a + \frac{1}{2} \ \alpha - \frac{1}{2}}^{c + \frac{1}{2} \ 2\alpha - \frac{1}{2}} &= \left[\frac{c - 2\alpha + 1}{a - \alpha + 1} \right]^{\frac{1}{2}} C, \end{aligned} \quad (8.5.24)$$

where

$$C = \frac{(-1)^{c/2-a} \left(a + \frac{c}{2}\right)!}{\left(a - \frac{c}{2}\right)! \left(\frac{c}{2} + \alpha\right)! \left(\frac{c}{2} - \alpha\right)!} \left[\frac{(2a + c + 2)(2a - c)!(c + 2\alpha)!(c - 2\alpha)!}{2(2a + c + 1)!} \right]^{\frac{1}{2}}, \quad (8.5.25)$$

(j) $b = a + \frac{1}{2}, \beta = \alpha \pm \frac{1}{2}, 2a + c = 2g + 1$

$$\begin{aligned} C_{a \ \alpha \ a + \frac{1}{2} \ \alpha + \frac{1}{2}}^{c + \frac{1}{2} \ 2\alpha + \frac{1}{2}} &= \left[\frac{c - 2\alpha + 1}{a + \alpha + 1} \right]^{\frac{1}{2}} D, \\ C_{a \ \alpha \ a + \frac{1}{2} \ \alpha - \frac{1}{2}}^{c + \frac{1}{2} \ 2\alpha - \frac{1}{2}} &= - \left[\frac{c + 2\alpha + 1}{a - \alpha + 1} \right]^{\frac{1}{2}} D, \end{aligned} \quad (8.5.26)$$

where

$$D = \frac{(-1)^{\frac{a-1}{2}-a} \left(a + \frac{c+1}{2}\right)!}{\left(a - \frac{c+1}{2}\right)! \left(\frac{c+1}{2} + \alpha\right)! \left(\frac{c+1}{2} - \alpha\right)!} \left[\frac{(2a - c)!(c + 2\alpha + 1)!(c - 2\alpha + 1)!}{2(2a + c + 2)!} \right]^{\frac{1}{2}}. \quad (8.5.27)$$

(k) $b = a + 1; \beta = \alpha, \alpha \pm 1; 2a + c = 2g$

$$\begin{aligned} C_{a \ \alpha \ a+1 \ a+1}^{c \ 2a+1} &= - \left[\frac{(c + 2\alpha + 1)(c - 2\alpha)}{(a + \alpha + 2)(a + \alpha + 1)} \right]^{\frac{1}{2}} E, \\ C_{a \ \alpha \ a+1 \ \alpha}^{c \ 2\alpha} &= \frac{2\alpha}{[(a + \alpha + 1)(a - \alpha + 1)]^{\frac{1}{2}}} E, \\ C_{a \ \alpha \ a+1 \ \alpha-1}^{c \ 2\alpha-1} &= \left[\frac{(c - 2\alpha + 1)(c + 2\alpha)}{(a - \alpha + 2)(a - \alpha + 1)} \right]^{\frac{1}{2}} E, \end{aligned} \quad (8.5.28)$$

where

$$E = \frac{(-1)^{\frac{c}{2}-a} \left(a + \frac{c}{2}\right)!}{2 \left(a - \frac{c}{2}\right)! \left(\frac{c}{2} + \alpha\right)! \left(\frac{c}{2} - \alpha\right)!} \left[\frac{(2c + 1)(2a + c + 2)(2a - c + 1)!(c + 2\alpha)!(c - 2\alpha)!}{c(c + 1)(2a + c + 1)!} \right]^{\frac{1}{2}}. \quad (8.5.29)$$

(l) $b = a + 1; \beta = \alpha, \alpha \pm 1; 2a + c = 2g + 1$

$$\begin{aligned} C_{a \ \alpha \ a+1 \ \alpha+1}^{c \ 2\alpha+1} &= \frac{c(c + 1) + (2\alpha + 1)(2a + 2)}{2 \left(\frac{c+1}{2} + \alpha\right)! \left(\frac{c+1}{2} - \alpha\right)!} \left[\frac{(c + 2\alpha + 1)!(c - 2\alpha - 1)!}{(a + \alpha + 2)(a + \alpha + 1)} \right]^{\frac{1}{2}} F, \\ C_{a \ \alpha \ a+1 \ \alpha}^{c \ 2\alpha} &= \frac{2a + 2}{\left(\frac{c-1}{2} + \alpha\right)! \left(\frac{c-1}{2} - \alpha\right)!} \left[\frac{(c + 2\alpha)!(c - 2\alpha)!}{(a + \alpha + 1)(a - \alpha + 1)} \right]^{\frac{1}{2}} F, \\ C_{a \ \alpha \ a+1 \ \alpha-1}^{c \ 2\alpha-1} &= \frac{c(c + 1) - (2\alpha - 1)(2a + 2)}{2 \left(\frac{c+1}{2} - \alpha\right)! \left(\frac{c-1}{2} + \alpha\right)!} \left[\frac{(c + 2\alpha - 1)!(c - 2\alpha + 1)!}{(a - \alpha + 2)(a - \alpha + 1)} \right]^{\frac{1}{2}} F, \end{aligned} \quad (8.5.30)$$

where

$$F = \frac{(-1)^{\frac{c-1}{2}-a} (a + \frac{c+1}{2})!}{(a - \frac{c-1}{2})!} \left[\frac{(2c+1)(2a-c+1)!}{c(c+1)(2a+c+2)!} \right]^{\frac{1}{2}}. \quad (8.5.31)$$

Equations (8.5.24)-(8.5.31) were obtained by Stone [63].

8.5.2 Special Values of Momentum Projections

(a) $\alpha = \beta = \gamma = 0$

$$C_{a0b0}^{c0} = \begin{cases} 0, & \text{if } a+b+c = 2g+1 \\ \frac{(-1)^{g-c} \sqrt{2c+1} g!}{(g-a)!(g-b)!(g-c)!} \left[\frac{(2g-2a)!(2g-2b)!(2g-2c)!}{(2g+1)!} \right]^{\frac{1}{2}}, & \text{if } a+b+c = 2g \end{cases}, \quad (8.5.32)$$

g is integer and positive. In particular,

$$C_{a0b0}^{a+b0} = \frac{(a+b)!}{a!b!} \left[\frac{(2a)!(2b)!}{(2a+2b)!} \right]^{\frac{1}{2}}, \quad (8.5.33)$$

$$C_{a0b0}^{a-b0} = (-1)^b \frac{a!}{b!(a-b)!} \left[\frac{(2b)!(2a-2b+1)!}{(2a+1)!} \right]^{\frac{1}{2}}. \quad (8.5.34)$$

(b) $\gamma = c$ or $\alpha = a$:

$$C_{a\alpha b\beta}^{cc} = \delta_{\alpha+\beta,c} (-1)^{a-\alpha} \left[\frac{(2c+1)!(a+b-c)!(a+\alpha)!(b+\beta)!}{(a+b+c+1)!(a-b+c)!(-a+b+c)!(a-\alpha)!(b-\beta)!} \right]^{\frac{1}{2}}, \quad (8.5.35)$$

$$C_{a\alpha b\beta}^{c\gamma} = \delta_{\gamma-\beta,\alpha} \left[\frac{(2c+1)(2a)!(-a+b+c)!(b-\beta)!(c+\gamma)!}{(a+b+c+1)!(a-b+c)!(a+b-c)!(b+\beta)!(c-\gamma)!} \right]^{\frac{1}{2}}. \quad (8.5.36)$$

In particular,

$$C_{aabb}^{cc} = \delta_{a+b,c}, \quad (8.5.37)$$

$$C_{aabb-b}^{cc} = \delta_{a-b,c} \left[\frac{2c+1}{2a+1} \right]^{\frac{1}{2}}, \quad (8.5.38)$$

$$C_{aabb-c-a}^{cc} = \left[\frac{(2a)!(2c+1)!}{(a+b+c+1)!(a-b+c)!} \right]^{\frac{1}{2}}, \quad (8.5.39)$$

$$C_{a-1bca+1}^{cc} = - \left[\frac{(2a-1)!(2c+1)!(a+b-c)(-a+b+c+1)}{(a+b+c+1)!(a-b+c)!} \right]^{\frac{1}{2}}, \quad (8.5.40)$$

$$C_{aabb-c-a-1}^{cc-1} = \left[\frac{(2a)!(2c+1)!(a+b-c+1)(-a+b+c)}{(a+b+c+1)!(a-b+c)!2c} \right]^{\frac{1}{2}}, \quad (8.5.41)$$

$$C_{ccb0}^{cc} = (2c)! \left[\frac{2c+1}{(2c-b)!(2c+b+1)!} \right]^{\frac{1}{2}}, \quad (8.5.42)$$

$$C_{cc-bbb}^{cc} = (-1)^b \left[\frac{(2c+1)!(2b)!}{(2c+b+1)!b!} \right]^{\frac{1}{2}}, \quad (8.5.43)$$

(c) $\gamma = c-1$ or $\alpha = a-1$:

$$C_{a\alpha b\beta}^{cc-1} = \delta_{\alpha+\beta,c-1} (-1)^{a-\alpha} \{ (b-\beta)(b+\beta+1) - (a-\alpha)(a+\alpha+1) \} \\ \times \left[\frac{(2c+1)(2c-1)!(a+b-c)!(a+\alpha)!(b+\beta)!}{(a+b+c+1)!(a-b+c)!(-a+b+c)!(a-\alpha)!(b-\beta)!} \right]^{\frac{1}{2}}, \quad (8.5.44)$$

$$C_{a-1b\beta}^{c\gamma} = \delta_{\gamma-\beta,a-1} \{ (c-\gamma)(c+\gamma+1) - (b+\beta)(b-\beta+1) \} \\ \times \left[\frac{(2c+1)(2a-1)!(-a+b+c)!(b-\beta)!(c+\gamma)!}{(a+b+c+1)!(a-b+c)!(a+b-c)!(b+\beta)!(c-\gamma)!} \right]^{\frac{1}{2}}, \quad (8.5.45)$$

In particular,

$$C_{a\alpha a\alpha}^{c\ c-1} = 0, \quad (8.5.46)$$

$$C_{a\ a-1\ c-\gamma}^{c\ \gamma} = 0, \quad (8.5.47)$$

$$C_{a\ a-1\ b\ \beta}^{c\ c-1} = \delta_{\beta, c-a} \{a(a+1) - b(b+1) + c(c+1) - 2ac\} \left[\frac{(2c+1)(2c-1)!(2a-1)!}{(a+b+c+1)!(a-b+c)!} \right]^{\frac{1}{2}}. \quad (8.5.48)$$

8.6 Recursion relations for the Clebsch-Gordan coefficients

8.6.1 General Recursion Relations

$$\begin{aligned} C_{a\alpha b\beta}^{c\ \gamma} &= \left[\frac{(b+\beta-2k)!(c+\gamma)!(a+b-c)!(-a+b+c)!(a+b+c+1)!(2c+1)}{(b+\beta)!(c-\gamma)!(a-b+c)!} \right]^{\frac{1}{2}} \\ &\times \sum_{c'=c-k}^{c+k} (-1)^{c'+k-c} C_{a\alpha b-k\beta-k}^{c'\ \gamma-k} \frac{(c'-k+c)!(2k)!}{(c+k-c')!(c+c'+k+1)!(c'+k-c)!} \\ &\times \left[\frac{(c'+k-\gamma)!(a-b+c'+k)!(2c'+1)}{(c'-k+\gamma)!(-a+b+c'-k)!(a+b-c'-k)!(a+b+c'-k+1)!} \right]^{\frac{1}{2}}. \end{aligned} \quad (8.6.1)$$

(Yutsis and Bandzaitis [20]) where k is integer or half-integer, $0 \leq k \leq (b+\beta)/2$. In accordance with the choice of k (8.6.1) yields different recursion relations.

Note also the relation obtained by Stone [63]

$$\begin{aligned} C_{a\alpha b\beta}^{c\ \gamma} &= \left[\frac{(a+b+c+1)!(a-b+c)!(a+b-c)!(b+\beta)!(b-\beta)!(a-b+\alpha-\beta)!(a-b-\alpha+\beta)!}{(-a+b+c)!(a+\alpha)!(a-\alpha)!} \right]^{\frac{1}{2}} \\ &\times \sum_{b'} (-1)^{b-b'/2} C_{a-b\alpha-\beta\ b'\ 2\beta}^{c\ \gamma} \frac{(2b'+1) \left(b + \frac{b'}{2}\right)!}{(2b+b'+1)!(b - \frac{b'}{2})! \left(\frac{b'}{2} + \beta\right)! \left(\frac{b'}{2} - \beta\right)!} \\ &\times \left[\frac{(-a+b+c+b')!(b'+2\beta)!(b'-2\beta)!}{(a-b+c+b'+1)!(a-b+c-b')!(a-b-c+b')!} \right]^{\frac{1}{2}}. \end{aligned} \quad (8.6.2)$$

Here $a-b \geq |\alpha-\beta| \geq 0$, b' is integer, $b'+2b$ is even, and $a-b+c \geq b' \geq |-a+b+c|$, $2b \geq b' \geq |2\beta|$. Equation (8.6.1) may be rewritten by the use of quasi-powers in the form [20]

$$\begin{aligned} C_{a\alpha b\beta}^{c\ \gamma} &= \left[\frac{2c+1}{(b+\beta)(2k)} \right]^{\frac{1}{2}} \sum_{k'=-k}^k (-1)^{k+k'} C_{a\alpha b-k\beta-k}^{c+k'\ \gamma-k} \left| (c+\gamma)^{(k-k')} (c-\gamma+k+k')^{(k+k')} (a+b-c)^{(k+k')} \right. \\ &\times (-a+b+c)^{(k-k')} (a+b+c+1)^{(k-k')} (a-b+c+k+k')^{(k+k')} (2c+2k'+1) \left. \right]^{\frac{1}{2}} \\ &\times \frac{(2k)^{(k+k')}}{(2c+k+k'+1)^{(2k+1)} (k+k')^{(k+k')}}, \end{aligned} \quad (8.6.3)$$

where

$$a^{(n)} = \frac{a!}{(a-n)!}, \quad \frac{1}{2} \leq k \leq \frac{b-\kappa}{2}, \quad \kappa = \begin{cases} 0 & \text{if } b \text{ is integer,} \\ \frac{1}{2} & \text{if } b \text{ is half-integer.} \end{cases}$$

8.6.2 Arguments α, β, γ Change by 1

$$[(c \pm \gamma)(c \mp \gamma + 1)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\ \gamma \mp 1} = [(a \mp \alpha)(a \pm \alpha + 1)]^{\frac{1}{2}} C_{a\alpha \pm 1\ b\ \beta}^{c\ \gamma} + [(b \mp \beta)(b \pm \beta + 1)]^{\frac{1}{2}} C_{a\alpha b\beta \pm 1}^{c\ \gamma}. \quad (8.6.4)$$

In particular, for $|\gamma| = c$ one has

$$C_{a\alpha \mp 1\ b\ \beta}^{c\ \pm c} = -C_{a\alpha b\beta \mp 1}^{c\ \pm c} \left[\frac{(b \pm \beta)(b \mp \beta + 1)}{(a \pm \alpha)(a \mp \alpha + 1)} \right]^{\frac{1}{2}}, \quad (8.6.5)$$

when $|\alpha| = |\beta| = \frac{1}{2}$

$$C_{a\frac{1}{2}\ b\frac{1}{2}}^{c\ 1} = C_{a\frac{1}{2}\ b-\frac{1}{2}}^{c\ 0} \frac{(2b+1) + (-1)^{a+b-c}(2a+1)}{2[c(c+1)]^{\frac{1}{2}}}, \quad (8.6.6)$$

and for $|\alpha| = |\beta| = 1, a + b + c$ even, one gets

$$C_{a1b-1}^{c0} = C_{a0b0}^{c0} \frac{c(c+1) - a(a+1) - b(b+1)}{2[a(a+1)b(b+1)]^{\frac{1}{2}}}, \quad (8.6.7)$$

$$C_{a1b1}^{c2} = C_{a0b0}^{c0} \frac{a(a+1)[c(c+1) - a(a+1) + b(b+1)] + b(b+1)[c(c+1) + a(a+1) - b(b+1)]}{2[a(a+1)b(b+1)(c-1)c(c+1)(c+2)]^{\frac{1}{2}}}. \quad (8.6.8)$$

8.6.3 Arguments Change by $1/2$

$$(2a+1)[b \pm \beta]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = \mp [(a \mp \alpha)(a+b-c)(a+b+c+1)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha \pm \frac{1}{2}b - \frac{1}{2}\beta \mp \frac{1}{2}}^{c\gamma} \\ + [(a \pm \alpha + 1)(-a+b+c)(a-b+c+1)]^{\frac{1}{2}} C_{a+\frac{1}{2}\alpha \pm \frac{1}{2}b - \frac{1}{2}\beta \mp \frac{1}{2}}^{c\gamma} \quad (8.6.9)$$

$$(2a+1)[b \mp \beta + 1]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = [(a \mp \alpha)(a-b+c)(-a+b+c+1)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha \pm \frac{1}{2}b + \frac{1}{2}\beta \mp \frac{1}{2}}^{c\gamma} \\ \pm [(a \pm \alpha + 1)(a+b-c+1)(a+b+c+2)]^{\frac{1}{2}} C_{a+\frac{1}{2}\alpha \pm \frac{1}{2}b + \frac{1}{2}\beta \mp \frac{1}{2}}^{c\gamma} \quad (8.6.10)$$

$$[(-a+b+c)(a-b+c+1)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = [(a-\alpha+1)(b-\beta)]^{\frac{1}{2}} C_{a+\frac{1}{2}\alpha - \frac{1}{2}b - \frac{1}{2}\beta + \frac{1}{2}}^{c\gamma} \\ + [(a+\alpha+1)(b+\beta)]^{\frac{1}{2}} C_{a+\frac{1}{2}\alpha + \frac{1}{2}b - \frac{1}{2}\beta - \frac{1}{2}}^{c\gamma} \quad (8.6.11)$$

$$[(a-b+c)(-a+b+c+1)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = [(a-\alpha)(b-\beta+1)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha + \frac{1}{2}b + \frac{1}{2}\beta - \frac{1}{2}}^{c\gamma} \\ + [(a+\alpha)(b+\beta+1)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha - \frac{1}{2}b + \frac{1}{2}\beta + \frac{1}{2}}^{c\gamma} \quad (8.6.12)$$

$$\left[\frac{2c(-a+b+c)(a+b+c+1)}{2c+1} \right]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = [(b-\beta)(c-\gamma)]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta + \frac{1}{2}}^{c-\frac{1}{2}\gamma + \frac{1}{2}} + [(b+\beta)(c+\gamma)]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta - \frac{1}{2}}^{c-\frac{1}{2}\gamma - \frac{1}{2}}, \quad (8.6.13)$$

$$\left[\frac{2c(c \mp \gamma)(a+b+c+1)}{2c+1} \right]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = [(a \mp \alpha)(a-b+c)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha \pm \frac{1}{2}b\beta}^{c-\frac{1}{2}\gamma \pm \frac{1}{2}} + [(b \mp \beta)(-a+b+c)]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta \pm \frac{1}{2}}^{c-\frac{1}{2}\gamma \pm \frac{1}{2}}, \quad (8.6.14)$$

$$\left[\frac{2c(c \mp \gamma)(a+b-c+1)}{2c+1} \right]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = \pm [(a \pm \alpha + 1)(-a+b+c)]^{\frac{1}{2}} C_{a+\frac{1}{2}\alpha \pm \frac{1}{2}b\beta}^{c-\frac{1}{2}\gamma \pm \frac{1}{2}} \\ \mp [(b \pm \beta + 1)(a-b+c)]^{\frac{1}{2}} C_{a\alpha b + \frac{1}{2}\beta \pm \frac{1}{2}}^{c-\frac{1}{2}\gamma \pm \frac{1}{2}}, \quad (8.6.15)$$

$$\left[\frac{2(c+1)(c \pm \gamma + 1)(a+b-c)}{2c+1} \right]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = \mp [(a \mp \alpha)(-a+b+c+1)]^{\frac{1}{2}} C_{a-\frac{1}{2}\alpha \pm \frac{1}{2}b\beta}^{c+\frac{1}{2}\gamma \pm \frac{1}{2}} \\ \pm [(b \mp \beta)(a-b+c+1)]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta \pm \frac{1}{2}}^{c+\frac{1}{2}\gamma \pm \frac{1}{2}}, \quad (8.6.16)$$

$$[(2c+1)(b \mp \beta)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} = \left[\frac{(c \mp \gamma)(-a+b+c)(a+b+c+1)}{2c} \right]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta \pm \frac{1}{2}}^{c-\frac{1}{2}\gamma \pm \frac{1}{2}} \\ \pm \left[\frac{(c \pm \gamma + 1)(a-b+c+1)(a+b-c)}{2(c+1)} \right]^{\frac{1}{2}} C_{a\alpha b - \frac{1}{2}\beta \pm \frac{1}{2}}^{c+\frac{1}{2}\gamma \pm \frac{1}{2}}. \quad (8.6.17)$$

In particular, Eqs. (8.6.9) and (8.6.10) yield

$$C_{a \pm \frac{1}{2}b \mp \frac{1}{2}}^{c0} = \pm C_{a-\frac{1}{2}0b-\frac{1}{2}0}^{c0} \left[\frac{(a+b-c)(a+b+c+1)}{(2a+1)(2b+1)} \right]^{\frac{1}{2}} = \mp C_{a+\frac{1}{2}0b+\frac{1}{2}0}^{c0} \left[\frac{(a+b-c+1)(a+b+c+2)}{(2a+1)(2b+1)} \right]^{\frac{1}{2}}, \quad (8.6.18)$$

if $a + b + c$ is odd, and

$$C_{a \pm \frac{1}{2} b \mp \frac{1}{2}}^{c0} = C_{a + \frac{1}{2} 0 b - \frac{1}{2} 0}^{c0} \left[\frac{(-a + b + c)(a - b + c + 1)}{(2a + 1)(2b + 1)} \right]^{\frac{1}{2}} = C_{a - \frac{1}{2} 0 b + \frac{1}{2} 0}^{c0} \left[\frac{(a - b + c)(-a + b + c + 1)}{(2a + 1)(2b + 1)} \right]^{\frac{1}{2}}, \quad (8.6.19)$$

if $a + b + c$ is even.

8.6.4 The Case When $\alpha = \beta = \gamma = 0$

Let p be integer and $2g = a + b + c$. Then

$$C_{a+p 0 b-p 0}^{c0} = C_{a0 b0}^{c0} \frac{(g-a)!(g-b)!}{(g-a-p)!(g-b+p)!} \left[\frac{(2g-2a-2p)!(2g-2b+2p)!}{(2g-2a)!(2g-2b)!} \right]^{\frac{1}{2}}. \quad (8.6.20)$$

In particular, (8.6.20) yields

$$C_{a+1 0 b-1 0}^{c0} = C_{a0 b0}^{c0} \left[\frac{(-a+b+c)(a-b+c+1)}{(-a+b+c-1)(a-b+c+2)} \right]^{\frac{1}{2}}, \quad (8.6.21)$$

$$\begin{aligned} C_{a0 b+2p 0}^{c0} &= C_{a0 b0}^{c0} (-1)^p \frac{(g+p)!(g-a)!(g-b)!(g-c)!}{g!(g-a+p)!(g-b-p)!(g-c+p)!}, \\ &\times \left[\frac{(a+b-c+2p)!(a-b+c-2p)!(-a+b+c+2p)!(a+b+c+1)!}{(a+b-c)!(a-b+c)!(-a+b+c)!(a+b+c+2p+1)!} \right]^{\frac{1}{2}}. \end{aligned} \quad (8.6.22)$$

Moreover, from (8.6.22) one gets

$$C_{a0 b+2 0}^{c0} = -C_{a0 b0}^{c0} \left[\frac{(a+b+c+2)(a+b-c+1)(a-b+c)(-a+b+c+1)}{(a+b+c+3)(a+b-c+2)(a-b+c-1)(-a+b+c+2)} \right]^{\frac{1}{2}}. \quad (8.6.23)$$

8.6.5 Arguments a, b, c Change by 1

$$\begin{aligned} 2[b^2 - \beta^2]^{\frac{1}{2}} C_{a\alpha b-1\beta}^{c\gamma} &= \frac{1}{a(2a+1)} [(a^2 - \alpha^2)(-a+b+c)(-a+b+c+1)(a-b+c)(a-b+c+1)]^{\frac{1}{2}} \\ &\times C_{a-1\alpha b\beta}^{c\gamma} + \frac{\alpha}{a(a+1)} [(-a+b+c)(a-b+c+1)(a+b-c)(a+b+c+1)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} \\ &- \frac{1}{(a+1)(2a+1)} \{ [(a+1)^2 - \alpha^2](a+b-c)(a+b-c+1)(a+b+c+1)(a+b+c+2) \}^{\frac{1}{2}} C_{a+1\alpha b\beta}^{c\gamma}, \end{aligned} \quad (8.6.24)$$

$$\begin{aligned} 2\beta C_{a\alpha b\beta}^{c\gamma} &= -\frac{1}{a(2a+1)} [(a^2 - \alpha^2)(-a+b+c+1)(a-b+c)(a+b-c)(a+b+c+1)]^{\frac{1}{2}} C_{a-1\alpha b\beta}^{c\gamma} \\ &- \frac{\alpha}{a(a+1)} \{ a(a+1) + b(b+1) - c(c+1) \} C_{a\alpha b\beta}^{c\gamma} \\ &- \frac{1}{(a+1)(2a+1)} \{ [(a+1)^2 - \alpha^2](-a+b+c)(a-b+c+1)(a+b-c+1)(a+b+c+2) \}^{\frac{1}{2}} C_{a+1\alpha b\beta}^{c\gamma}, \end{aligned} \quad (8.6.25)$$

$$\begin{aligned} 2[(b+1)^2 - \beta^2]^{\frac{1}{2}} C_{a\alpha b+1\beta}^{c\gamma} &= -\frac{1}{a(2a+1)} [(a^2 - \alpha^2)(a+b-c)(a+b-c+1)(a+b+c+1)(a+b+c+2)]^{\frac{1}{2}} \\ &\times C_{a-1\alpha b\beta}^{c\gamma} + \frac{\alpha}{a(a+1)} [(-a+b+c+1)(a-b+c)(a+b-c+1)(a+b+c+2)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} \\ &+ \frac{1}{(a+1)(2a+1)} \{ [(a+1)^2 - \alpha^2](-a+b+c)(-a+b+c+1)(a-b+c)(a-b+c+1) \}^{\frac{1}{2}} C_{a+1\alpha b\beta}^{c\gamma}, \end{aligned} \quad (8.6.26)$$

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &= \left[\frac{4c^2(2c+1)(2c-1)}{(c+\gamma)(c-\gamma)(-a+b+c)(a-b+c)(a+b-c+1)(a+b+c+1)} \right]^{\frac{1}{2}} \\ &\times \left\{ \frac{(\alpha-\beta)c(c-1) - \gamma a(a+1) + \gamma b(b+1)}{2c(c-1)} C_{a\alpha b\beta}^{c-1\gamma} \right. \\ &\left. - \left[\frac{(c-\gamma-1)(c+\gamma-1)(-a+b+c-1)(a-b+c-1)(a+b-c+2)(a+b+c)}{4(c-1)^2(2c-3)(2c-1)} \right]^{\frac{1}{2}} C_{a\alpha b\beta}^{c-2\gamma} \right\}. \end{aligned} \quad (8.6.27)$$

8.6.6 Arguments a, b, α, β Change by 1

$$\begin{aligned}
& [(b \pm \beta)(b \pm \beta + 1)]^{\frac{1}{2}} C_{a \alpha b - 1 \beta}^{c \gamma} \\
= & \frac{1}{2a(2a+1)} [(a \pm \alpha)(a \pm \alpha - 1)(-a + b + c)(-a + b + c + 1)(a - b + c)(a - b + c + 1)]^{\frac{1}{2}} C_{a-1 \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& \mp \frac{1}{2a(a+1)} [(a \pm \alpha)(a \mp \alpha + 1)(-a + b + c)(a - b + c + 1)(a + b - c)(a + b + c + 1)]^{\frac{1}{2}} C_{a \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& + \frac{1}{2(a+1)(2a+1)} [(a \mp \alpha + 1)(a \mp \alpha + 2)(a + b - c)(a + b - c + 1)(a + b + c + 1)(a + b + c + 2)]^{\frac{1}{2}} \\
& \times C_{a+1 \alpha \mp 1 b \beta \pm 1}^{c \gamma}. \tag{8.6.28}
\end{aligned}$$

$$\begin{aligned}
& [(b \mp \beta)(b \pm \beta + 1)]^{\frac{1}{2}} C_{a \alpha b \beta}^{c \gamma} \\
= & \pm \frac{1}{2a(2a+1)} [(a \pm \alpha)(a \pm \alpha - 1)(-a + b + c + 1)(a - b + c)(a + b - c)(a + b + c + 1)]^{\frac{1}{2}} C_{a-1 \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& - \frac{1}{2a(a+1)} \{a(a+1) + b(b+1) - c(c+1)\} [(a \pm \alpha)(a \mp \alpha + 1)]^{\frac{1}{2}} C_{a \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& \mp \frac{1}{2(a+1)(2a+1)} [(a \mp \alpha + 1)(a \mp \alpha + 2)(-a + b + c)(a - b + c + 1)(a + b - c + 1)(a + b + c + 2)]^{\frac{1}{2}} \\
& \times C_{a+1 \alpha \mp 1 b \beta \pm 1}^{c \gamma}. \tag{8.6.29}
\end{aligned}$$

$$\begin{aligned}
& [(b \mp \beta)(b \mp \beta + 1)]^{\frac{1}{2}} C_{a \alpha b + 1 \beta}^{c \gamma} \\
= & \frac{1}{2a(2a+1)} [(a \pm \alpha)(a \pm \alpha - 1)(a + b - c)(a + b - c + 1)(a + b + c + 1)(a + b + c + 2)]^{\frac{1}{2}} C_{a-1 \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& \pm \frac{1}{2a(a+1)} [(a \pm \alpha)(a \mp \alpha + 1)(-a + b + c + 1)(a - b + c)(a + b - c + 1)(a + b + c + 2)]^{\frac{1}{2}} C_{a \alpha \mp 1 b \beta \pm 1}^{c \gamma} \\
& + \frac{1}{2(a+1)(2a+1)} [(a \mp \alpha + 1)(a \mp \alpha + 2)(-a + b + c)(-a + b + c + 1)(a - b + c)(a - b + c + 1)]^{\frac{1}{2}} \\
& \times C_{a+1 \alpha \mp 1 b \beta \pm 1}^{c \gamma}. \tag{8.6.30}
\end{aligned}$$

8.6.7 Arguments c, b, γ, β Change by 1

$$\begin{aligned}
& \left[\frac{(-a + b + c)(a - b + c)(a + b - c + 1)(a + b + c + 1)(2c - 1)}{(2c + 1)} \right]^{\frac{1}{2}} C_{a \alpha b \beta}^{c \gamma} \\
= & [(b + \beta)(b - \beta + 1)(c + \gamma)(c + \gamma - 1)]^{\frac{1}{2}} C_{a \alpha b \beta - 1}^{c-1 \gamma-1} - 2\beta [c^2 - \gamma^2]^{\frac{1}{2}} C_{a \alpha b \beta}^{c-1 \gamma} \\
& - [(b - \beta)(b + \beta + 1)(c - \gamma)(c - \gamma - 1)]^{\frac{1}{2}} C_{a \alpha b \beta + 1}^{c-1 \gamma+1}. \tag{8.6.31}
\end{aligned}$$

$$\begin{aligned}
& [(b \mp \beta)(b \mp \beta + 1)]^{\frac{1}{2}} C_{a \alpha b - 1 \beta}^{c \gamma} \\
= & \frac{1}{2c} \left[\frac{(c \pm \gamma)(c \pm \gamma - 1)(a - b + c)(a - b + c + 1)(a + b - c)(a + b - c + 1)}{(2c - 1)(2c + 1)} \right]^{\frac{1}{2}} C_{a \alpha b \beta \mp 1}^{c-1 \gamma \mp 1} \\
& \pm \frac{1}{2c(c+1)} [(c \pm \gamma)(c \mp \gamma + 1)(-a + b + c)(a - b + c + 1)(a + b - c)(a + b + c + 1)]^{\frac{1}{2}} C_{a \alpha b \beta \mp 1}^{c \gamma \mp 1} \\
& + \frac{1}{2(c+1)} \left[\frac{(c \mp \gamma + 1)(c \mp \gamma + 2)(-a + b + c)(-a + b + c + 1)(a + b + c + 1)(a + b + c + 2)}{(2c + 1)(2c + 3)} \right]^{\frac{1}{2}} C_{a \alpha b \beta \mp 1}^{c+1 \gamma \mp 1} \tag{8.6.32}
\end{aligned}$$

$$\begin{aligned}
& [(b \pm \beta)(b \mp \beta + 1)]^{\frac{1}{2}} C_{a\alpha b\beta}^{c\gamma} \\
&= \pm \frac{1}{2c} \left[\frac{(c \pm \gamma)(c \pm \gamma - 1)(-a + b + c)(a - b + c)(a + b - c + 1)(a + b + c + 1)}{(2c - 1)(2c + 1)} \right]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c-1\gamma \mp 1} \\
&\quad + \frac{1}{2c(c+1)} \{-a(a+1) + b(b+1) + c(c+1)\} [(c \pm \gamma)(c \mp \gamma + 1)]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c\gamma \mp 1} \\
&\mp \frac{1}{2(c+1)} \left[\frac{(c \mp \gamma + 1)(c \mp \gamma + 2)(-a + b + c + 1)(a - b + c + 1)(a + b - c)(a + b + c + 2)}{(2c + 1)(2c + 3)} \right]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c+1\gamma \mp 1} \\
&\hspace{15em} (8.6.33)
\end{aligned}$$

$$\begin{aligned}
& [(b \pm \beta)(b \pm \beta + 1)]^{\frac{1}{2}} C_{a\alpha b+1\beta}^{c\gamma} \\
&= \frac{1}{2c} \left[\frac{(c \pm \gamma)(c \pm \gamma - 1)(-a + b + c)(-a + b + c + 1)(a + b + c + 1)(a + b + c + 2)}{(2c - 1)(2c + 1)} \right]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c-1\gamma \mp 1} \\
&\mp \frac{1}{2c(c+1)} [(c \pm \gamma)(c \mp \gamma + 1)(-a + b + c + 1)(a - b + c)(a + b - c + 1)(a + b + c + 2)]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c\gamma \mp 1} \\
&+ \frac{1}{2(c+1)} \left[\frac{(c \mp \gamma + 1)(c \mp \gamma + 2)(a - b + c)(a - b + c + 1)(a + b - c)(a + b - c + 1)}{(2c + 1)(2c + 3)} \right]^{\frac{1}{2}} C_{a\alpha b\beta \mp 1}^{c+1\gamma \mp 1}. \quad (8.6.34)
\end{aligned}$$

8.6.8 Recursion Relations for the Regge Symbols

Recursion relations for the Regge R -symbols are very convenient because each relation is equivalent to a variety of formally different expressions for the Clebsch-Gordan coefficients. The formulas given below represent some

recursion relations [101, 109]. In these formulas we use the notation $J = \sum_{k=1}^3 R_{ik} = \sum_{i=1}^3 R_{ik} = a + b + c$.

$$\begin{aligned} & [R_{11}(J+1)]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| - [R_{22}R_{33}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21} & R_{22}-1 & R_{23} \\ R_{31} & R_{32} & R_{33}-1 \end{array} \right\| \\ & + [R_{23}R_{32}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23}-1 \\ R_{31} & R_{32}-1 & R_{33} \end{array} \right\| = 0. \end{aligned} \quad (8.6.35)$$

$$\begin{aligned} & [(R_{11}+1)R_{21}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}+1 & R_{12} & R_{13} \\ R_{21}-1 & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| + [(R_{12}+1)R_{22}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12}+1 & R_{13} \\ R_{21} & R_{22}-1 & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| \\ & + [(R_{13}+1)R_{23}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13}+1 \\ R_{21} & R_{22} & R_{23}-1 \\ R_{31} & R_{32} & R_{33} \end{array} \right\| = 0. \end{aligned} \quad (8.6.36)$$

$$\begin{aligned} & [R_{11}R_{22}(J+1)]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| - (R_{22}+R_{23})[R_{33}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21} & R_{22}-1 & R_{23} \\ R_{31} & R_{32} & R_{33}-1 \end{array} \right\| \\ & - [R_{23}R_{31}(R_{21}+1)]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21}+1 & R_{22}-1 & R_{23}-1 \\ R_{31}-1 & R_{32} & R_{33} \end{array} \right\| = 0. \end{aligned} \quad (8.6.37)$$

$$\begin{aligned} & (R_{11}+R_{12}+1)[R_{22}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{array} \right\| + [R_{13}(R_{23}+1)(R_{12}+1)]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12}+1 & R_{13}-1 \\ R_{21} & R_{22}-1 & R_{23}+1 \\ R_{31} & R_{32} & R_{33} \end{array} \right\| \\ & - [R_{11}R_{33}(J+1)]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21} & R_{22}-1 & R_{23} \\ R_{31} & R_{32} & R_{33}-1 \end{array} \right\| = 0. \end{aligned} \quad (8.6.38)$$

$$\begin{aligned} & (R_{22}-R_{33})[R_{11}R_{23}R_{32}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11}-1 & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23}-1 \\ R_{31} & R_{32}-1 & R_{33} \end{array} \right\| \\ & + (R_{33}-R_{11})[R_{13}R_{22}R_{31}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12} & R_{13}-1 \\ R_{21} & R_{22}-1 & R_{23} \\ R_{31}-1 & R_{32} & R_{33} \end{array} \right\| \\ & + (R_{11}-R_{22})[R_{12}R_{21}R_{33}]^{\frac{1}{2}} \left\| \begin{array}{ccc} R_{11} & R_{12}-1 & R_{13} \\ R_{21}-1 & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33}-1 \end{array} \right\| = 0. \end{aligned} \quad (8.6.39)$$

8.7 Sums of products of Clebsch-Gordan coefficients

Sums of products of the vector-addition coefficients are more conveniently represented in terms of the $3jm$ symbols than in terms of the Clebsch-Gordan coefficients, because the $3jm$ symbols have simpler symmetry relations. The sums expressed in terms of the $3jm$ symbols will be given in Sec. 12.2. Nevertheless the Clebsch-Gordan coefficients are also widely used. That is why we present the formulas which seem to be most useful for sums of products containing the Clebsch-Gordan coefficients. For practical convenience we consider various versions of the formulas which differ by permutations of upper and lower arguments. In these formulas we use the notation

$$\Pi_{ab\dots c} = [(2a+1)(2b+1)\dots(2c+1)]^{\frac{1}{2}}$$

8.7.1 Sums Involving One Clebsch-Gordan Coefficient

$$\sum_{\alpha} C_{a\alpha b0}^{a\alpha} = \Pi_a^2 \delta_{b0}, \quad (8.7.1)$$

$$\sum_{\alpha} (-1)^{a-\alpha} C_{a\alpha a-\alpha}^{c0} = \Pi_a \delta_{c0}, \quad (8.7.2)$$

$$\sum_{\alpha\beta\gamma} (-1)^{c+\gamma} C_{a\alpha b\beta}^{c\gamma} [(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!]^{-\frac{1}{2}} = 0. \quad (8.7.3)$$

8.7.2 Sums Involving Products of Two Clebsch-Gordan Coefficients

$$\sum_{\alpha\beta} C_{a\alpha b\beta}^{c\gamma} C_{a\alpha b\beta}^{c'\gamma'} = \delta_{cc'} \delta_{\gamma\gamma'}, \quad (8.7.4)$$

$$\sum_{\alpha\gamma} C_{a\alpha b\beta}^{c\gamma} C_{a\alpha b'\beta'}^{c\gamma} = \frac{\Pi_c^2}{\Pi_b^2} \delta_{bb'} \delta_{\beta\beta'}, \quad (8.7.5)$$

$$\sum_{\alpha\beta} (-1)^{b+\beta} C_{a\alpha b\beta}^{c\gamma} C_{c'-\gamma' a\alpha}^{b-\beta} = (-1)^{c+\gamma} \frac{\Pi_b}{\Pi_c} \delta_{cc'} \delta_{\gamma\gamma'}, \quad (8.7.6)$$

$$\sum_{\alpha\beta} (-1)^{a+\alpha} C_{b\beta a\alpha}^{c\gamma} C_{a-\alpha c'\gamma'}^{b\beta} = \frac{\Pi_b}{\Pi_c} \delta_{cc'} \delta_{\gamma\gamma'}, \quad (8.7.7)$$

$$\sum_{\alpha\beta} C_{c\gamma b\beta}^{a\alpha} C_{c'-\gamma' a\alpha}^{b\beta} = (-1)^{b-a-\gamma} \frac{\Pi_{ab}}{\Pi_c} \delta_{cc'} \delta_{\gamma\gamma'}, \quad (8.7.8)$$

$$\sum_{c\gamma} C_{a\alpha b\beta}^{c\gamma} C_{a\alpha' b\beta'}^{c\gamma} = \delta_{\alpha\alpha'} \delta_{\beta\beta'}, \quad (8.7.9)$$

$$\sum_{c\gamma} \Pi_c^2 C_{a\alpha c\gamma}^{b\beta} C_{a\alpha' c\gamma}^{b\beta'} = \Pi_b^2 \delta_{\alpha\alpha'} \delta_{\beta\beta'}, \quad (8.7.10)$$

$$\sum_{a\alpha} (-1)^{a-\alpha} \Pi_a^2 C_{a\alpha b\beta}^{c\gamma} C_{a\alpha c\gamma'}^{b\beta'} = \Pi_{bc} \delta_{\gamma, -\gamma'} \delta_{\beta, -\beta'}. \quad (8.7.11)$$

8.7.3 Sums Involving Products of Three Clebsch-Gordan Coefficients

$$\sum_{\alpha\beta\delta} C_{a\alpha b\beta}^{c\gamma} C_{d\delta b\beta}^{e\varepsilon} C_{a\alpha f\varphi}^{d\delta} = \kappa_1 \Pi_{cd} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.12)$$

$$\sum_{\alpha\beta\delta} C_{b\beta c\gamma}^{a\alpha} C_{b\beta e\varepsilon}^{d\delta} C_{a\alpha f\varphi}^{d\delta} = \kappa_1 \frac{\Pi_{add}}{\Pi_e} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.13)$$

$$\sum_{\alpha\beta\delta} C_{b\beta a\alpha}^{c\gamma} C_{b\beta d\delta}^{e\varepsilon} C_{a\alpha f\varphi}^{d\delta} = \kappa_2 \Pi_{cd} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.14)$$

$$\sum_{\alpha\beta\delta} (-1)^{a-\alpha} C_{a\alpha b\beta}^{c\gamma} C_{d\delta b\beta}^{e\varepsilon} C_{d\delta a-\alpha}^{f\varphi} = \kappa_1 \Pi_{cf} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.15)$$

$$\sum_{\alpha\beta\delta} (-1)^{b+\beta} C_{a\alpha b\beta}^{c\gamma} C_{b-\beta e\varepsilon}^{d\delta} C_{a\alpha f\varphi}^{d\delta} = \kappa_1 \frac{\Pi_{cdd}}{\Pi_e} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.16)$$

$$\sum_{\alpha\beta\delta} (-1)^{a-\alpha} C_{b\beta a\alpha}^{c\gamma} C_{b\beta d\delta}^{e\varepsilon} C_{d\delta a-\alpha}^{f\varphi} = \kappa_2 \Pi_{cf} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.17)$$

$$\sum_{\alpha\beta\delta} (-1)^{b+\beta} C_{b-\beta c\gamma}^{a\alpha} C_{d\delta b\beta}^{e\varepsilon} C_{a\alpha f\varphi}^{d\delta} = \kappa_1 \Pi_{ad} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.18)$$

$$\sum_{\alpha\beta\delta} (-1)^{a-\alpha} C_{a\alpha c-\gamma}^{b\beta} C_{d\delta b-\beta}^{e\varepsilon} C_{a\alpha f\varphi}^{d\delta} = \kappa_1 \Pi_{bd} C_{c\gamma f\varphi}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\}, \quad (8.7.19)$$

where $\kappa_1 = (-1)^{b+c+d+f}$, $\kappa_2 = (-1)^{a+b+e+f}$.

8.7.4 Sums Involving Products of Four Clebsch-Gordan Coefficients

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c\gamma}^{a\alpha} C_{e\epsilon f\varphi}^{d\delta} C_{e\epsilon b\beta}^{g\eta} C_{f\varphi c\gamma}^{j\mu} = (-1)^{a-b+c+d+e-f} \sum_{s\sigma} \Pi_{ssag} C_{a\alpha s\sigma}^{j\mu} C_{g\eta s\sigma}^{d\delta} \begin{Bmatrix} b & c & a \\ j & s & f \end{Bmatrix} \begin{Bmatrix} b & e & g \\ d & s & f \end{Bmatrix} = \Pi_{adgj} \sum_{k\kappa} C_{g\eta j\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.20)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c-\gamma}^{a\alpha} C_{e\epsilon f-\varphi}^{d\delta} C_{b\beta e\epsilon}^{g\eta} C_{c\gamma f\varphi}^{j\mu} = (-1)^{b+e-g} \Pi_{adgj} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.21)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta a\alpha}^{c\gamma} C_{f\varphi j\mu}^{c\gamma} C_{b\beta g\eta}^{e\epsilon} C_{f\varphi d\delta}^{e\epsilon} = (-1)^{a-b+f-j} \Pi_{ce}^2 \sum_{k\kappa} C_{g\eta j\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.22)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{a\alpha b\beta}^{c\gamma} C_{g\eta e\epsilon}^{b\beta} C_{d\delta f\varphi}^{e\epsilon} C_{j\mu c\gamma}^{f\varphi} = (-1)^{d+e-c-j} \Pi_{bcef} \sum_{k\kappa} (-1)^{k-\kappa} C_{g\eta j\mu}^{k-\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.23)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta a-\alpha}^{c\gamma} C_{e\epsilon d-\delta}^{f\varphi} C_{g\eta b-\beta}^{e\epsilon} C_{j\mu c-\gamma}^{f\varphi} = (-1)^{b-c-g-\alpha+\eta} \Pi_{ceff} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.24)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c-\gamma}^{a\alpha} C_{e\epsilon f\varphi}^{d\delta} C_{e-\epsilon g\eta}^{b\beta} C_{f\varphi j\mu}^{c\gamma} = (-1)^{b+f-g-\delta} \Pi_{abcd} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.25)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c\gamma}^{a\alpha} C_{e\epsilon f\varphi}^{d\delta} C_{e\epsilon g\eta}^{b\beta} C_{f\varphi j\mu}^{c\gamma} = \sum_{k\kappa} \Pi_{bcdk} C_{g\eta j\mu}^{k\kappa} C_{d\delta k\kappa}^{a\alpha} \begin{Bmatrix} a & b & c \\ d & e & f \\ k & g & j \end{Bmatrix}, \quad (8.7.26)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c-\gamma}^{a\alpha} C_{e\epsilon f-\varphi}^{d\delta} C_{g\eta b-\beta}^{e\epsilon} C_{j\mu f-\varphi}^{c\gamma} = (-1)^{c+e-g+j+\alpha-\mu} \Pi_{adec} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.27)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c\gamma}^{a\alpha} C_{b\beta g\eta}^{e\epsilon} C_{f\varphi d\delta}^{e\epsilon} C_{f\varphi c\gamma}^{j\mu} = (-1)^{j-a+\delta-\eta} \Pi_{aej} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a-\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.28)$$

$$\sum_{\beta\gamma\epsilon\varphi} C_{b\beta c\gamma}^{a\alpha} C_{g\eta e\epsilon}^{b\beta} C_{f\varphi d\delta}^{e\epsilon} C_{f\varphi c\gamma}^{j\mu} = (-1)^{j-a+g+\delta} \Pi_{abej} \sum_{k\kappa} C_{g-\eta j-\mu}^{k\kappa} C_{d\delta a-\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}, \quad (8.7.29)$$

$$\sum_{\beta\gamma\epsilon\varphi} (-1)^{c-\gamma+e-\epsilon} C_{a\alpha b\beta}^{c\gamma} C_{d\delta f\varphi}^{e\epsilon} C_{e\epsilon b\beta}^{g\eta} C_{c\gamma f\varphi}^{j\mu} = (-1)^{a+d-\alpha-\delta} \Pi_{cegj} \sum_{k\kappa} C_{g\eta j-\mu}^{k\kappa} C_{d\delta a-\alpha}^{k\kappa} \begin{Bmatrix} c & b & a \\ f & e & d \\ j & g & k \end{Bmatrix}. \quad (8.7.30)$$

8.7.5 Sums Involving Products of the Clebsch-Gordan Coefficients and One 6j Symbol

$$\sum_{e\varepsilon} (-1)^{2e} \Pi_{cd} C_{b\beta d\delta}^{e\varepsilon} C_{f\varphi c\gamma}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\} = C_{a\alpha b\beta}^{c\gamma} C_{a\alpha f\varphi}^{d\delta}, \quad (8.7.31)$$

$$\sum_{f\varphi} (-1)^{c+d+f} \Pi_{ce} C_{e\varepsilon a\alpha}^{f\varphi} C_{d\delta c\gamma}^{f\varphi} \left\{ \begin{matrix} b & a & c \\ f & d & e \end{matrix} \right\} = C_{a\alpha b\beta}^{c\gamma} C_{d\delta b\beta}^{e\varepsilon}, \quad (8.7.32)$$

$$\sum_{c\gamma} (-1)^{2e-d+\alpha+\varphi} \Pi_{ae} C_{f-\varphi b\beta}^{c\gamma} C_{e\varepsilon a-\alpha}^{c\gamma} \left\{ \begin{matrix} c & f & b \\ d & e & a \end{matrix} \right\} = C_{b\beta d\delta}^{e\varepsilon} C_{f\varphi d\delta}^{a\alpha}, \quad (8.7.33)$$

$$\sum_{c\gamma} (-1)^{c+d-\beta-\varphi} \Pi_d^2 C_{a\alpha b\beta}^{c\gamma} C_{f-\varphi e\varepsilon}^{c\gamma} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\} = C_{a\alpha f\varphi}^{d\delta} C_{b-\beta e\varepsilon}^{d\delta}, \quad (8.7.34)$$

$$\sum_{c\gamma} (-1)^{2e} \Pi_{cd} C_{a\alpha b\beta}^{c\gamma} C_{f\varphi c\gamma}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\} = C_{b\beta d\delta}^{e\varepsilon} C_{a\alpha f\varphi}^{d\delta}, \quad (8.7.35)$$

$$\sum_{f\varphi} (-1)^{2c} \Pi_{ef} C_{b\beta d\delta}^{f\varphi} C_{a\alpha f\varphi}^{c\gamma} \left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = C_{b\beta a\alpha}^{e\varepsilon} C_{d\delta e\varepsilon}^{c\gamma}, \quad (8.7.36)$$

$$\sum_{c\gamma} (-1)^{e+d-\beta} \frac{\Pi_{cdd}}{\Pi_e} C_{a\alpha b\beta}^{c\gamma} C_{f\varphi c\gamma}^{e\varepsilon} \left\{ \begin{matrix} a & b & c \\ e & f & d \end{matrix} \right\} = C_{b-\beta e\varepsilon}^{d\delta} C_{a\alpha f\varphi}^{d\delta}, \quad (8.7.37)$$

$$\sum_{c\gamma} (-1)^{2e} \frac{\Pi_{acc}}{\Pi_b} C_{f\varphi c\gamma}^{b\beta} C_{c\gamma a\alpha}^{e\varepsilon} \left\{ \begin{matrix} c & f & b \\ d & e & a \end{matrix} \right\} = C_{d\delta b\beta}^{e\varepsilon} C_{f\varphi d\delta}^{a\alpha}. \quad (8.7.38)$$

8.7.6 Sums Involving Products of the Clebsch-Gordan Coefficients and One 9j Symbol

$$\sum_{ak} \Pi_{adgj} C_{b\beta c\gamma}^{a\alpha} C_{g\eta j\mu}^{k\kappa} C_{d\delta a\alpha}^{k\kappa} \left\{ \begin{matrix} c & b & a \\ f & e & d \\ j & g & k \end{matrix} \right\} = C_{e\varepsilon f\varphi}^{d\delta} C_{e\varepsilon b\beta}^{g\eta} C_{f\varphi c\gamma}^{j\mu}, \quad (8.7.39)$$

$$\sum_{gj} \Pi_{adgj} C_{g\eta j\mu}^{k\kappa} C_{e\varepsilon b\beta}^{g\eta} C_{f\varphi c\gamma}^{j\mu} \left\{ \begin{matrix} c & b & a \\ f & e & d \\ j & g & k \end{matrix} \right\} = C_{d\delta a\alpha}^{k\kappa} C_{e\varepsilon f\varphi}^{d\delta} C_{b\beta c\gamma}^{a\alpha}, \quad (8.7.40)$$

$$\sum_{gj} (-1)^{b+e-g} \Pi_{adgj} C_{g\eta j-\mu}^{k\kappa} C_{b\beta e\varepsilon}^{g\eta} C_{c\gamma f\varphi}^{j\mu} \left\{ \begin{matrix} c & b & a \\ f & e & d \\ j & g & k \end{matrix} \right\} = C_{d\delta a\alpha}^{k\kappa} C_{e\varepsilon f-\varphi}^{d\delta} C_{b\beta c-\gamma}^{a\alpha}, \quad (8.7.41)$$

$$\sum_{ad} (-1)^{b+f-g-\delta} \Pi_{abcd} C_{b\beta c-\gamma}^{a\alpha} C_{e\varepsilon f\varphi}^{d\delta} C_{d\delta a\alpha}^{k\kappa} \left\{ \begin{matrix} c & b & a \\ f & e & d \\ j & g & k \end{matrix} \right\} = C_{g\eta j-\mu}^{k\kappa} C_{e-\varepsilon g\eta}^{b\beta} C_{f\varphi j\mu}^{c\gamma}, \quad (8.7.42)$$

$$\sum_{gj} (-1)^{g-b-f+\varphi+\varepsilon} \frac{\Pi_{adjgg}}{\Pi_{bc}} C_{g\eta j-\mu}^{k\kappa} C_{e-\varepsilon g\eta}^{b\beta} C_{f\varphi j\mu}^{c\gamma} \left\{ \begin{matrix} c & b & a \\ f & e & d \\ j & g & k \end{matrix} \right\} = C_{b\beta c-\gamma}^{a\alpha} C_{e\varepsilon f\varphi}^{d\delta} C_{d\delta a\alpha}^{k\kappa}. \quad (8.7.43)$$

Sums of larger numbers of the Clebsch-Gordan coefficients may be obtained from the above equations using the orthogonality relation ((8.1.8)).

8.7.7 Some Additional Sums of Products of Two Clebsch-Gordan Coefficients

We note also some other sums of products of the Clebsch-Gordan coefficients. The first two sums, obtained by Morgan [136], are written as

$$\sum_{l'=0}^l \frac{1}{(2l'-1)(2l-2l'+2J+1)} \left(C_{l0l'+J0}^{l-l'+J0} \right)^2 = -\frac{\delta_{l0}}{2J+1}, \quad (8.7.44)$$

and

$$\sum_{l'=0}^l \left(\frac{1}{2l'+3} - \frac{l+1}{2l+3} \frac{1}{2l'+1} \right) \frac{1}{2l-2l'+2J+1} \left(C_{l0l'+J0}^{l-l'+J0} \right)^2 = 0. \quad (8.7.45)$$

In these equations l, l', J are integers.

Din proved the following identity [137]

$$\sum_{i=|c-b|}^{c+b} \frac{2i+1}{i(i+1)-a(a+1)} (C_{i0b0}^{c0})^2 = 0, \quad (8.7.46)$$

where a, b, c are non-negative integers, with $a+b+c$ odd, which satisfy the triangular condition

$$|c-b| \leq a \leq c+b.$$

The sum runs only over values of i for which the Clebsch-Gordan coefficient does not vanish. Dunlop and Judd [138] give the following relation

$$\sum_m C_{a\ m\ k\ 0}^{a\ m} C_{c\ M-m\ k\ 0}^{c\ M-m} = (-1)^k \frac{\Pi_{ac}}{2k+1} \sqrt{\frac{(2a-k)!(2c+k+1)!}{(2c-k)!(2a+k+1)!}}, \quad (8.7.47)$$

which is valid for $a-c \geq |M|$.

8.8 Generating functions of the Clebsch-Gordan coefficients

The Clebsch-Gordan coefficients may be generated by expansions of some functions in power series or finite sums. Below we present several generating functions. In these relations $J \equiv a+b+c$.

8.8.1 Regge Determinant to Power J

$$\begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}^J = J! \sqrt{(J+1)!} \sum_{R_{ik}} \begin{vmatrix} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{vmatrix} \left\| \frac{u_1^{R_{11}} u_2^{R_{12}} u_3^{R_{13}} v_1^{R_{21}} v_2^{R_{22}} v_3^{R_{23}} w_1^{R_{31}} w_2^{R_{32}} w_3^{R_{33}}}{[R_{11}! R_{12}! R_{13}! R_{21}! R_{22}! R_{23}! R_{31}! R_{32}! R_{33}!]} \right\|^{\frac{1}{2}}. \quad (8.8.1)$$

(Regge [55], Shelepin [64]). Here R_{ik} are the elements of the R -symbol (see Sec. 8.1.3). The R -symbols and the Clebsch-Gordan coefficients are related by Eqs. (8.1.14) and (8.1.11).

8.8.2 Products of Binomials

$$\begin{aligned} & (t_1 - t_2)^{J-2c} (t_2 - t_3)^{J-2a} (t_3 - t_1)^{J-2b} \\ &= \left[(J+1)! \frac{(J-2a)!(J-2b)!(J-2c)!}{2c+1} \right]^{\frac{1}{2}} \sum_{\substack{\alpha, \beta, \gamma \\ \alpha+\beta+\gamma=0}} (-1)^{a-b-\gamma} \frac{t_1^{a+\alpha} t_2^{b+\beta} t_3^{c+\gamma}}{[(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!]^{\frac{1}{2}}} C_{a\ \alpha\ b\ \beta}^{c-\gamma}, \end{aligned} \quad (8.8.2)$$

$$\begin{aligned} & (v_1 u_2 - u_1 v_2)^{J-2c} (w_1 v_2 - w_2 v_1)^{J-2a} (u_1 w_2 - w_1 u_2)^{J-2b} \\ &= \left[\frac{(J+1)!(J-2a)!(J-2b)!(J-2c)!}{2c+1} \right]^{\frac{1}{2}} \\ & \times \sum_{\substack{\alpha, \beta, \gamma \\ \alpha+\beta+\gamma=0}} (-1)^{a-b-\gamma} \frac{u_1^{a-\alpha} u_2^{a+\alpha} v_1^{b-\beta} v_2^{b+\beta} w_1^{c-\gamma} w_2^{c+\gamma}}{[(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!]^{\frac{1}{2}}} C_{a\ \alpha\ b\ \beta}^{c-\gamma}. \end{aligned} \quad (8.8.3)$$

8.8.3 Exponential Function

$$\begin{aligned} & \exp\{x_1(t_2 - t_3) + x_2(t_3 - t_1) + x_3(t_1 - t_2)\} \\ &= \sum_{\substack{a, b, c \\ \alpha, \beta, \gamma}} \frac{(-1)^{a-b-\gamma}}{\Delta(abc)} \frac{x_1^{-a+b+c} x_2^{a-b+c} x_3^{a+b-c} t_1^{a+\alpha} t_2^{b+\beta} t_3^{c+\gamma}}{[(2c+1)(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!(c+\gamma)!(c-\gamma)!]^{\frac{1}{2}}} C_{a\ \alpha\ b\ \beta}^{c-\gamma}, \end{aligned} \quad (8.8.4)$$

where $\Delta(abc)$ is given by (8.2.1).

8.8.4 Hypergeometric Function

$$\begin{aligned} & \left[\frac{(J-2b)!(c+\gamma)!(2c+1)}{(J-2a)!(J-2c)!(c-\gamma)!(J+1)!} \right]^{\frac{1}{2}} \frac{(t-1)^{J-2c}}{(a-b+\gamma)!} F(\gamma-c, a-b-c; a-b+\gamma+1; t) \\ &= \sum_{\substack{\alpha, \beta \\ \alpha+\beta=\gamma}} C_{a\alpha b\beta}^{c\gamma} \frac{t^{b-\beta}}{[(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!]^{\frac{1}{2}}}. \end{aligned} \quad (8.8.5)$$

(Akim and Levin [62])

8.8.5 Wigner D-Function

$$\left[\frac{(2c+1)}{(J+1)!(J-2c)!} \right]^{\frac{1}{2}} d_{\gamma\gamma'}^c(\theta) = \sum_{\substack{\alpha, \beta \\ \alpha+\beta=\gamma}} (-1)^{b+\beta} C_{a\alpha b\beta}^{c\gamma} \frac{(\cos \frac{\theta}{2})^{a+b+\alpha-\beta} (\sin \frac{\theta}{2})^{a+b-\alpha+\beta}}{[(a+\alpha)!(a-\alpha)!(b+\beta)!(b-\beta)!]^{\frac{1}{2}}}, \quad (8.8.6)$$

where $a-b=\gamma'$, $|a-b| \leq c \leq a+b$.

8.8.6 Generating Function for the Coefficients C_{a0b0}^{c0}

$$\frac{1}{1+u^2+v^2+w^2} = \sum_{abc} (-1)^{a-b} \left[\frac{(J+1)!}{2c+1} \right]^{\frac{1}{2}} C_{a0b0}^{c0} \frac{u^{J-2a}v^{J-2b}w^{J-2c}}{[(J-2a)!(J-2b)!(J-2c)!]^{\frac{1}{2}}}. \quad (8.8.7)$$

(Schwinger [65])

8.9 Classical limit and asymptotic expressions for the Clebsch-Gordan coefficients

Asymptotic expressions permit us to simplify calculations of the Clebsch-Gordan coefficients. Note that these expressions appear to be accurate even if the arguments are not very large.

8.9.1 Asymptotic Expressions for $a, c \gg b$

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &\approx [(b+\beta)!(b-\beta)!(b+\delta)!(b-\delta)!]^{\frac{1}{2}} \\ &\times \sum_s \frac{(-1)^{b-\delta-s} (\cos \frac{\vartheta}{2})^{2s+\beta+\delta} (\sin \frac{\vartheta}{2})^{2b-\beta-\delta-2s}}{s!(s+\beta+\delta)!(b-\beta-s)!(b-\delta-s)!} = \delta_{\gamma-\alpha, \beta} D_{\beta\delta}^b(0, \vartheta, 0), \end{aligned} \quad (8.9.1)$$

where $\delta = c - a$, and ϑ is the angle between the momentum c and the z axis,

$$\cos \frac{\vartheta}{2} = \sqrt{\frac{c+\gamma+\frac{1}{2}}{2c+1}} = C_{c\gamma \frac{1}{2} \frac{1}{2}}^{c+\frac{1}{2} \gamma+\frac{1}{2}}, \quad \sin \frac{\vartheta}{2} = \sqrt{\frac{c-\gamma+\frac{1}{2}}{2c+1}} = C_{c\gamma \frac{1}{2} -\frac{1}{2}}^{c+\frac{1}{2} \gamma-\frac{1}{2}}, \quad \cos \vartheta = \frac{\gamma}{c+\frac{1}{2}}. \quad (8.9.2)$$

This formula was obtained by Edmonds [47], Brussaard and Tolhoek [66]. In particular,

$$C_{a\alpha b\beta}^{c\gamma} \approx \delta_{b, \gamma-\alpha} (-1)^{b-\delta} \left[\frac{(2b)!}{(b+\delta)!(b-\delta)!} \right]^{\frac{1}{2}} \left(\cos \frac{\vartheta}{2} \right)^{b+\delta} \left(\sin \frac{\vartheta}{2} \right)^{b-\delta}, \quad (8.9.3)$$

$$C_{a\alpha b\beta}^{c\gamma} \approx \delta_{\beta, c-\alpha} D_{\beta c-a}^b(0, 0, 0) = \delta_{\beta, c-\alpha} \delta_{\beta, c-a}, \quad (8.9.4)$$

$$C_{a-\beta b\beta}^{c0} \approx D_{\beta c-a}^b\left(0, \frac{\pi}{2}, 0\right), \quad (8.9.5)$$

$$C_{c\gamma b0}^{c\gamma} \approx P_b(\cos \vartheta), \quad (8.9.6)$$

$$C_{c\alpha b\beta}^{c\gamma} \approx \delta_{\beta, \gamma-\alpha} \sqrt{\frac{4\pi}{2b+1}} Y_{b\beta}(\vartheta, 0), \quad (8.9.7)$$

$$C_{c\pm c b0}^{c\pm c} \approx (\pm 1)^b. \quad (8.9.8)$$

If $a, c \gg b \gg 1$ one can use in (8.9.1) the asymptotic expressions for the Wigner D -function (Eqs. (4.18.1)(4.18.4)). Then the Clebsch-Gordan coefficients may be represented in the form

$$C_{a\alpha b\beta}^{c\gamma} \approx \frac{(-1)^{\beta-\delta}}{(\beta-\delta)!} \left[\frac{2}{\pi} \frac{(b+\beta)!(b-\beta+1)!}{(b+\delta)!(b-\delta)!} \right]^{\frac{1}{2}} \frac{\cos \left[(2b+1)\frac{\vartheta}{2} - \frac{\pi}{2} \left(\beta - \delta + \frac{1}{2} \right) \right]}{\sqrt{\sin \vartheta}}, \quad (8.9.9)$$

if $\beta \geq \delta, \beta \geq 0$,

$$C_{a\alpha b\beta}^{c\gamma} \approx \frac{1}{(\delta-\beta)!} \left[\frac{2}{\pi} \frac{(b+\delta)!(b-\delta+1)!}{(b+\beta)!(b-\beta)!} \right]^{\frac{1}{2}} \frac{\cos \left[(2b+1)\frac{\vartheta}{2} - \frac{\pi}{2} \left(\delta - \beta + \frac{1}{2} \right) \right]}{\sqrt{\sin \vartheta}}, \quad (8.9.10)$$

if $\delta \geq \beta, \delta \geq 0$.

8.9.2 Asymptotic Expressions for $a, b, c, \gamma \gg a + b - \gamma$

$$C_{a\alpha b\beta}^{c\gamma} \approx (-1)^{a+b-c} \frac{(c+\gamma+1)^{b-\beta}}{(a-b-\alpha)!} \left[\frac{(2c+1)(a-b+c)!(a-\alpha)!(b+\beta)!(c-\gamma)!(c+\gamma)!}{(a+b-c)!(-a+b+c)!(a+b+c+1)!(a+\alpha)!(b-\beta)!} \right]^{\frac{1}{2}}. \quad (8.9.11)$$

(Akim and Levin [62]).

8.9.3 Semiclassical Formulas for $a, b, c \gg 1$

$$C_{a\alpha b\beta}^{c\gamma} \approx (-1)^{2b+c+\gamma+1} \sqrt{\frac{2c+1}{2\pi S}} \cos \left[j_1\theta_1 + j_2\theta_2 + j_3\theta_3 - m_2\varphi_1 + m_1\varphi_2 + \frac{\pi}{4} \right], \quad (8.9.12)$$

(Ponzano and Regge [46]), where

$$j_1 \equiv a + \frac{1}{2}, j_2 \equiv b + \frac{1}{2}, j_3 \equiv c + \frac{1}{2}, m_1 \equiv \alpha, \quad m_2 \equiv \beta, \quad m_3 \equiv -\gamma, \quad (8.9.13)$$

$$S^2 = \frac{1}{16} \left\{ -j_1^4 - j_2^4 - j_3^4 + 2(j_1^2 j_2^2 + j_1^2 j_3^2 + j_2^2 j_3^2) + 4(j_1^2 m_2 m_3 + j_2^2 m_1 m_3 + j_3^2 m_1 m_2) \right\}$$

$$= -\frac{1}{16} \begin{vmatrix} 0 & j_1^2 - m_1^2 & j_2^2 - m_2^2 & 1 \\ j_1^2 - m_1^2 & 0 & j_3^2 - m_3^2 & 1 \\ j_2^2 - m_2^2 & j_3^2 - m_3^2 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{vmatrix}. \quad (8.9.14)$$

If $m_1 = m_2 = m_3 = 0$, (8.9.14) reduces to

$$S^2 = \frac{1}{16} \left(a + b + c + \frac{3}{2} \right) \left(-a + b + c + \frac{1}{2} \right) \left(a - b + c + \frac{1}{2} \right) \left(a + b - c + \frac{1}{2} \right), \quad (8.9.15)$$

$$\cos \theta_i = \frac{2j_i^2 m_l + m_i (j_i^2 - j_k^2 + j_l^2)}{\sqrt{(j_i^2 - m_i^2) (4j_i^2 j_l^2 - (j_i^2 - j_k^2 + j_l^2)^2)}},$$

$$\cos \varphi_i = \frac{1}{2} \frac{j_i^2 - j_k^2 - j_l^2 - 2m_k m_l}{\sqrt{(j_k^2 - m_k^2) (j_l^2 - m_l^2)}}. \quad (8.9.16)$$

Indices i, k, l in (8.9.16) are those which may be obtained by cyclic permutation of 1, 2, 3. The quantities which enter Eqs. (8.9.14)-(8.9.16) have a simple geometrical interpretation. Three momenta $j_1 = a + \frac{1}{2}, j_2 = b + \frac{1}{2}$ and $j_3 = c + \frac{1}{2}$ form a triangle oriented in space in such a manner that the projections of its sides on the z axis are equal to m_1, m_2 and m_3 , respectively (Fig. 8.1). S is an area of the triangle projection onto the (x, y) -plane. Let us construct a trihedral prism whose sidelong edges are parallel to the z axis and one of the cross sections coincides with the oriented triangle. Then θ_i is the angle between the normals to the planes adjacent to the edge j_i . The angles φ_i are shown in Fig. 8.1.

In particular, if $\alpha = \beta = \gamma = 0$, we get $\theta_1 = \theta_2 = \theta_3 = \pi/2$.

$$C_{a0b0}^{c0} \approx (-1)^{(a+b-c)/2} \left[\frac{1 + (-1)^{a+b-c}}{2} \right] \sqrt{\frac{2c+1}{2\pi S}}. \quad (8.9.17)$$

Equations (8.9.12)-(8.9.17) are valid, if $S^2 > 0$.




Figure 8.1. Geometric interpretation of the parameters in the asymptotic relation (8.9.12).

More accurate expressions valid not only for $S^2 > 0$ but also if $S^2 < 0$ or $S^2 \approx 0$ were obtained by Schulten and Gordon [139]. In the case of $S^2 \geq 0$ they derived

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{2b+c+\gamma+1} \sqrt{\frac{2c+1}{2S}} Z^{\frac{1}{4}} \times \begin{cases} \cos \Omega_0 Ai(-Z) - \sin \Omega_0 Bi(-Z), & \text{if } \Omega - \Omega_0 < 0, \\ \cos \Omega_0 Bi(-Z) - \sin \Omega_0 Ai(-Z), & \text{if } \Omega - \Omega_0 > 0, \end{cases} \quad (8.9.18a)$$

while for $S^2 \leq 0$

$$C_{a\alpha b\beta}^{c\gamma} = (-1)^{2b+c+\gamma+1} \sqrt{\frac{2c+1}{2|S|}} Z^{\frac{1}{4}} \times \begin{cases} \cos \Omega_0 Ai(Z) - \sin \Omega_0 Bi(Z), & \text{if } \Omega - \Omega_0 < 0, \\ \cos \Omega_0 Bi(Z) - \sin \Omega_0 Ai(Z), & \text{if } \Omega - \Omega_0 > 0. \end{cases} \quad (8.9.18b)$$

In these equations $Ai(Z)$ and $Bi(Z)$ are the Airy functions

$$Z = \left(\frac{3}{2} |\Omega - \Omega_0| \right)^{\frac{1}{3}},$$

$$\Omega_0 = j_1 \theta_1^{(0)} + j_2 \theta_2^{(0)} + j_3 \theta_3^{(0)} + m_1 \varphi_2^{(0)} - m_2 \varphi_1^{(0)},$$

$$\Omega = j_1 \theta_1 + j_2 \theta_2 + j_3 \theta_3 - m_2 \varphi_1 + m_1 \varphi_2,$$

$\theta_i^{(0)}$ and $\varphi_i^{(0)}$ being determined by the relations

$$\theta_i^{(0)} = \begin{cases} 0 & \text{if } 0 \leq \text{Re } \theta_i \leq \pi/2 \\ \pi & \text{if } \pi/2 < \text{Re } \theta_i \leq \pi \end{cases}$$

$$\varphi_i^{(0)} = \begin{cases} 0 & \text{if } 0 \leq \text{Re } \varphi_i \leq \pi/2 \\ \pi & \text{if } \pi/2 < \text{Re } \varphi_i \leq \pi \end{cases}$$

8.9.4 Squares of the Clebsch-Gordan Coefficients in the Classical Limit

According to Sec. 8.1.1 the square of a Clebsch-Gordan coefficient represents the probability of the coupling of two angular momenta j_1 and j_2 with projections m_1 and m_2 into the resultant angular momentum j with projection m . On the other hand, the same square is equal to the probability that the momenta $\mathbf{j}_1$ and $\mathbf{j}_2$ coupled into the momentum j with the projection m have the projections m_1 and m_2 , respectively. Let us consider the square of the Clebsch-Gordan coefficients in the classical limit [66].

If the vector j as well as the magnitudes of j_1 and j_2 are given, the possible positions of the end of the vector j_1 form a circle (Fig. 8.2). The probability in question is proportional to the length of the circle arc

confined between the planes $z = m_1$ and $z = m_2$. In the classical limit (for $j_1, j_2, j \gg 1$) this probability may be expressed by

$$\left(C_{j_1 m_1 j_2 m_2}^j\right)^2 \approx \frac{2j+1}{\pi} \left[- (j_1^4 + j_2^4 + j^4) + 2 (j_1^2 j_2^2 + j_1^2 j^2 + j_2^2 j^2) - 4 (j_1^2 m_2 m + j_2^2 m_1 m - j^2 m_1 m_2) \right]^{-\frac{1}{2}}. \quad (8.9.19)$$

(8.9.19) is valid only on the average, because if j_1, j_2 and j are large, even fairly minor relative variations of these quantities lead to rapid oscillations of the Clebsch-Gordan coefficients (See (8.9.12)). When $m_1 = m_2 = m = 0$ and $j_1 + j_2 - j$ is even, one has

$$C_{j_1 0 j_2 0}^j \approx (-1)^{(j_1+j_2-j)/2} \sqrt{\frac{2j+1}{\pi}} \left[- (j_1^4 + j_2^4 + j^4) + 2 (j_1^2 j_2^2 + j_1^2 j^2 + j_2^2 j^2) \right]^{-\frac{1}{4}}. \quad (8.9.20)$$




Figure 8.2. Addition of angular momenta in the classical limit.

8.10 Zeros of the vector-addition coefficients

A Clebsch-Gordan coefficient $C_{a\alpha b\beta}^{c\gamma}$ and associated $3jm$ symbol $\begin{pmatrix} a & b & c \\ \alpha & \beta & -\gamma \end{pmatrix}$ vanish unless triangular conditions are satisfied (Eqs. (8.1.1) and (8.1.2)). However, there exist non-trivial cases when the Clebsch-Gordan coefficients are equal to zero although the triangular conditions are fulfilled. This leads to some additional selection rules which forbid the quantum transitions whose amplitudes are proportional to the appropriate Clebsch-Gordan coefficients. Here we point out the most important relations between the arguments for which the Clebsch-Gordan coefficients vanish.

- (a) $C_{a0b0}^0 = 0$, if $J \equiv a + b + c$ is odd;
- (b) $C_{a\alpha a\alpha}^{c\gamma} = 0$, $C_{a-\alpha c\gamma}^{a\alpha} = 0$, $C_{c\gamma a-\alpha}^{a\alpha} = 0$, if $J = 2a + c$ is odd;
- (c) $C_{aJ-3a bJ-3b}^{c-(J-3c)} = 0$, if J is odd;
- (d) $C_{a\alpha b\beta}^{a+b-1\gamma} = 0$, if $\frac{\alpha}{\beta} = \frac{a}{b}$;
- (e) $C_{a\alpha b\beta}^{a-b+1\gamma} = 0$ if $\frac{\alpha}{\beta} = -\frac{a+1}{b}$;
- (f) $C_{a\alpha b\beta}^{c-1} = 0$ if $a(a+1) - b(b+1) = (\alpha - \beta)c$, in particular

$$C_{a\alpha a\alpha}^{c-1} = 0.$$

- (g) $C_{a\alpha b\beta}^{c\gamma} = 0$ if $a(a+1) - c(c+1) = -(\alpha + \gamma)b$ in particular,

$$C_{a\alpha b\beta}^{a-\alpha} = 0.$$

Below we list particular $3jm$ symbols with $J = a + b + c \leq 17$ which vanish in nontrivial cases. They are arranged in the order of increasing J and $a \leq b \leq c$. Roots of other $3jm$ symbols may be obtained by the use of the simplest symmetry properties of the $3j$ m symbols, concerned with the permutations of momenta a, b, c and sign changes of the projections α, β, γ (see Eqs. (8.4.5) and (8.4.6)). Hence, any $3jm$ symbols from the table given below generally represents 12 formally different versions of the $3jm$ symbols or the Clebsch-Gordan coefficients which are equal to zero. One should bear in mind that the $3j$ m symbols and the Clebsch-Gordan coefficients are related by (see Sec. 8.1.2)

$$\begin{pmatrix} a & b & c \\ \alpha & \beta & -\gamma \end{pmatrix} = (-1)^{a+b+\gamma} \frac{1}{\sqrt{2c+1}} C_{a\alpha b\beta}^{c\gamma}.$$

8.11 Connection of the Clebsch-Gordan coefficients and the $3jm$ symbols with analogous functions of other authors

$$C_{j_1 m_1 j_2 m_2}^{j m} = \begin{cases} S_{j m_1 m_2}^{j j_2} & \text{- Wigner [15],} \\ A_{j m_1 m_2}^{j j_1 j_2} & \text{- Eckart [31],} \\ C_{m_1 m_2}^j & \text{- Vander Waerden [29],} \\ (j_1 j_2 m_1 m_2 | j_1 j_2 j m) & \text{- Condon and Shortley [22],} \\ C_{j m}^{j_1 j_2} (m_1 m_2) & \text{- Fock [57],} \\ X(j, m_1, j_1, j_2, m_1) & \text{- Boys [67],} \\ C(j m_1 m_1 m_2) & \text{- Blatt and Weisskopf [27],} \\ C_{m_1 m_2 m}^{j_1 j_2 j} & \text{- Biedenharn [117],} \\ C(j_1 j_2 j, m_1 m_2) & \text{- Rose [4],} \\ \begin{bmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{bmatrix} & \text{- Yutsis and Bandzaitis [20],} \\ \langle j_1 m_1 j_2 m_2 | (j_1 j_2) j m \rangle & \text{- Fano [68].} \end{cases}$$

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \begin{cases} (-1)^{-j_1+j_2+j_3} V(j_1 j_2 j_3, m_1 m_2 m_3) & \text{- Racah [56]} \\ (-1)^{-j_1+j_2+j_3} \bar{V}(j_1 j_2 j_3, m_1 m_2 m_3) & \text{- Fano and Racah [16]} \\ (-1)^{j_1-j_2+j_3} S_{j_1 m_1 j_2 m_2 j_3 m_3} & \text{- Landau and Lifshitz [8]} \\ (-1)^{j_1-j_2+j_3} (j_1 m_1, j_2 m_2, j_3 m_3 | 0) & \text{- Fano [68]} \\ X(j_1 j_2 j_3, m_1 m_2 m_3) & \text{- Schwinger [65]} \\ (-1)^{j_1-j_2+j_3} U(j_1 j_2 m_1 m_2 | j_3 - m_3) & \text{- Lubarskii [32]} \\ (j_1 j_2 j_3)_{m_1 m_2 m_3} & \text{- Sharp [69]} \end{cases}$$

8.12 Algebraic tables of the Clebsch-Gordan coefficients

Below we present on a small selection of tables of algebraic formulas for the Clebsch-Gordan coefficients $C_{a\alpha b\beta}^{c\gamma}$ with $b = 1/2, 1$. The original book contains many more. However, the use of Mathematica or SymPY will allow others to be evaluated quickly. Algebraic tables of the Clebsch-Gordan coefficients are also available in Ref. [22] for $b = 1/2, 1, 3/2, 2$; in Refs. [126, 130] for $b = 5/2$; in Refs. [118, 119] with $b = 3$; and in Ref. [116] with $b = 7/2, 4, 9/2, 5$.

8.13 Numerical tables of the clebsch-gordan coefficients

Even though the original book provides numerical values of the Clebsch-Gordan coefficients, with the advent of exact evaluators (as in sympy), these are no longer very useful. Numerical values of the Clebsch-Gordan coefficients are also given in Refs. [116, 120]. Reference [120] presents the Clebsch-Gordan coefficients for $\frac{1}{2} \leq a \leq 4, \frac{1}{2} \leq b, c \leq 9/2$ in decimals. Reference [116] gives the Clebsch-Gordan coefficients for $\frac{1}{2} \leq b \leq 6, a = 5, 11/2, 6$ in the form of rational fractions. Numerical values of the $3jm$ symbols are also available in Ref. [121].

Tables 8.2–8.3. Algebraic Formulas for the Clebsch-Gordan Coefficients. Tables not reproduced here can be found in the original book.

Table 8.1. Accidental zeroes of the $3j$ coefficients for $J = j_1 + j_2 + j_3 \leq 17$

$$\begin{aligned}
J=3 & \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}. \\
J=5 & \begin{pmatrix} 1 & 2 & 2 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3/2 & 3/2 & 2 \\ 1/2 & 1/2 & -1 \end{pmatrix}. \\
J=7 & \begin{pmatrix} 1 & 3 & 3 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 2 & 3 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 2 & 3 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 2 & 5/2 & 5/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}. \\
J=8 & \begin{pmatrix} 2 & 3 & 3 \\ 0 & 2 & -2 \end{pmatrix}, \begin{pmatrix} 3/2 & 3 & 7/2 \\ 1/2 & 1 & -3/2 \end{pmatrix}. \\
J=9 & \begin{pmatrix} 1 & 4 & 4 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 3 & 4 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 3 & 3 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 3 & 3 \\ 1 & 1 & -2 \end{pmatrix}, \\
& \begin{pmatrix} 2 & 7/2 & 7/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 5/2 & 3 & 7/2 \\ 3/2 & 0 & -3/2 \end{pmatrix}, \begin{pmatrix} 5/2 & 5/2 & 4 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \begin{pmatrix} 5/2 & 5/2 & 4 \\ 3/2 & 3/2 & -3 \end{pmatrix}. \\
J=11 & \begin{pmatrix} 1 & 5 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 4 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 4 & 5 \\ 1 & 2 & -3 \end{pmatrix}, \begin{pmatrix} 3 & 3 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 3 & 5 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 3 & 3 & 5 \\ 2 & 2 & -4 \end{pmatrix}, \\
& \begin{pmatrix} 3 & 4 & 4 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 4 & 4 \\ -2 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 9/2 & 9/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 3 & 7/2 & 9/2 \\ 2 & 1/2 & -5/2 \end{pmatrix}, \begin{pmatrix} 7/2 & 7/2 & 4 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \\
& \begin{pmatrix} 7/2 & 7/2 & 4 \\ 3/2 & 3/2 & -3 \end{pmatrix}, \begin{pmatrix} 3/2 & 9/2 & 5 \\ 1/2 & 3/2 & -2 \end{pmatrix}, \begin{pmatrix} 5/2 & 4 & 9/2 \\ 1/2 & 3 & -7/2 \end{pmatrix}. \\
J=13 & \begin{pmatrix} 1 & 6 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 5 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 4 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 5 \\ -2 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \\
& \begin{pmatrix} 4 & 4 & 5 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 5 \\ 2 & 2 & -4 \end{pmatrix}, \begin{pmatrix} 2 & 11/2 & 11/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 7/2 & 9/2 & 5 \\ 5/2 & -1/2 & -2 \end{pmatrix}, \begin{pmatrix} 7/2 & 4 & 11/2 \\ 5/2 & 1 & -7/2 \end{pmatrix}, \\
& \begin{pmatrix} 4 & 9/2 & 9/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 4 & 9/2 & 9/2 \\ -3 & 3/2 & 3/2 \end{pmatrix}, \begin{pmatrix} 7/2 & 7/2 & 6 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \begin{pmatrix} 7/2 & 7/2 & 6 \\ 3/2 & 3/2 & -3 \end{pmatrix}, \begin{pmatrix} 7/2 & 7/2 & 6 \\ 5/2 & 5/2 & -5 \end{pmatrix}. \\
J=14 & \begin{pmatrix} 3 & 5 & 6 \\ 1 & -1 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 6 \\ 2 & 1 & -3 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 6 \\ 1 & 4 & -5 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 6 \\ 0 & 2 & -2 \end{pmatrix}, \begin{pmatrix} 4 & 5 & 5 \\ 1 & 3 & -4 \end{pmatrix}, \begin{pmatrix} 4 & 5 & 5 \\ 2 & 1 & -3 \end{pmatrix}, \\
& \begin{pmatrix} 3 & 9/2 & 13/2 \\ 1 & 3/2 & -5/2 \end{pmatrix}, \begin{pmatrix} 5/2 & 5 & 13/2 \\ 1/2 & 1 & -3/2 \end{pmatrix}, \begin{pmatrix} 5/2 & 5 & 13/2 \\ 3/2 & 3 & -9/2 \end{pmatrix}, \begin{pmatrix} 4 & 9/2 & 11/2 \\ 0 & 7/2 & -7/2 \end{pmatrix}, \begin{pmatrix} 3/2 & 6 & 13/2 \\ 1/2 & 2 & -5/2 \end{pmatrix}, \begin{pmatrix} 5/2 & 11/2 & 6 \\ 3/2 & 1/2 & -2 \end{pmatrix}. \\
J=15 & \begin{pmatrix} 1 & 7 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 6 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 6 & 7 \\ 1 & 3 & -4 \end{pmatrix}, \begin{pmatrix} 3 & 5 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 6 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 6 & 6 \\ 0 & 5 & -5 \end{pmatrix}, \begin{pmatrix} 3 & 6 & 6 \\ -2 & 1 & 1 \end{pmatrix}, \\
& \begin{pmatrix} 4 & 5 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 4 & 5 & 6 \\ 3 & 0 & -3 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 7 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 7 \\ 2 & 2 & -4 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 7 \\ 3 & 3 & -6 \end{pmatrix}, \begin{pmatrix} 5 & 5 & 5 \\ 0 & 0 & 0 \end{pmatrix}, \\
& \begin{pmatrix} 5 & 5 & 5 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 5 & 5 & 5 \\ 2 & 2 & -4 \end{pmatrix}, \begin{pmatrix} 2 & 13/2 & 13/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 4 & 9/2 & 13/2 \\ 3 & 3/2 & -9/2 \end{pmatrix}, \begin{pmatrix} 4 & 11/2 & 11/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 4 & 11/2 & 11/2 \\ -3 & 3/2 & 3/2 \end{pmatrix}, \\
& \begin{pmatrix} 9/2 & 5 & 11/2 \\ 3/2 & 0 & -3/2 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 6 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 6 \\ 3/2 & 3/2 & -3 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 6 \\ 5/2 & 5/2 & -5 \end{pmatrix}. \\
J=17 & \begin{pmatrix} 1 & 8 & 8 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 7 & 8 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 6 & 8 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 6 & 8 \\ 2 & 4 & -6 \end{pmatrix}, \begin{pmatrix} 3 & 7 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 3 & 7 & 7 \\ -2 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 4 & 5 & 8 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 4 & 6 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \\
& \begin{pmatrix} 5 & 5 & 7 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 5 & 5 & 7 \\ 1 & 1 & -2 \end{pmatrix}, \begin{pmatrix} 5 & 5 & 7 \\ 2 & 2 & -4 \end{pmatrix}, \begin{pmatrix} 5 & 5 & 7 \\ 3 & 3 & -6 \end{pmatrix}, \begin{pmatrix} 5 & 6 & 6 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 5 & 6 & 6 \\ -2 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 5 & 6 & 6 \\ -4 & 2 & 2 \end{pmatrix}, \\
& \begin{pmatrix} 2 & 15/2 & 15/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 3 & 13/2 & 15/2 \\ 2 & 3/2 & -7/2 \end{pmatrix}, \begin{pmatrix} 4 & 13/2 & 13/2 \\ -1 & 1/2 & 1/2 \end{pmatrix}, \begin{pmatrix} 4 & 13/2 & 13/2 \\ -3 & 3/2 & 3/2 \end{pmatrix}, \begin{pmatrix} 5 & 11/2 & 13/2 \\ 2 & 1/2 & -5/2 \end{pmatrix}, \begin{pmatrix} 7/2 & 6 & 15/2 \\ 3/2 & 5 & -13/2 \end{pmatrix}, \\
& \begin{pmatrix} 9/2 & 6 & 13/2 \\ 1/2 & -5 & 9/2 \end{pmatrix}, \begin{pmatrix} 9/2 & 6 & 13/2 \\ -7/2 & 1 & 5/2 \end{pmatrix}, \begin{pmatrix} 11/2 & 11/2 & 6 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \begin{pmatrix} 11/2 & 11/2 & 6 \\ 3/2 & 3/2 & -3 \end{pmatrix}, \begin{pmatrix} 11/2 & 11/2 & 6 \\ 5/2 & 5/2 & -5 \end{pmatrix}, \begin{pmatrix} 9/2 & 11/2 & 7 \\ 7/2 & 1/2 & -4 \end{pmatrix}, \\
& \begin{pmatrix} 3/2 & 15/2 & 8 \\ 1/2 & 5/2 & -3 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 8 \\ 1/2 & 1/2 & -1 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 8 \\ 3/2 & 3/2 & -3 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 8 \\ 5/2 & 5/2 & -5 \end{pmatrix}, \begin{pmatrix} 9/2 & 9/2 & 8 \\ 7/2 & 7/2 & -7 \end{pmatrix}, \begin{pmatrix} 9/2 & 5 & 15/2 \\ 7/2 & 2 & -11/2 \end{pmatrix}.
\end{aligned}$$

Table 8.2. $C_{a \alpha \frac{1}{2} \beta}^{c \gamma}$

c	$\beta = 1/2$	$\beta = -1/2$
$a + 1/2$	$\left[\frac{c+\gamma}{2c} \right]^{1/2}$	$\left[\frac{c-\gamma}{2c} \right]^{1/2}$
$a - 1/2$	$-\left[\frac{c-\gamma+1}{2c+2} \right]^{1/2}$	$\left[\frac{c+\gamma+1}{2c+2} \right]^{1/2}$

Table 8.3. $C_{a \alpha 1 \beta}^{c \gamma}$

c	$\beta = 1$	$\beta = 0$	$\beta = -1$
$a + 1$	$\left[\frac{(c+\gamma-1)(c+\gamma)}{(2c-1)2c} \right]^{1/2}$	$\left[\frac{(c+\gamma)(c-\gamma)}{(2c-1)c} \right]^{1/2}$	$\left[\frac{(c-\gamma-1)(c-\gamma)}{(2c-1)2c} \right]^{1/2}$
a	$-\left[\frac{(c+\gamma)(c-\gamma+1)}{2c(c+1)} \right]^{1/2}$	$\frac{\gamma}{[c(c+1)]^{1/2}}$	$\left[\frac{(c+\gamma+1)(c-\gamma)}{2c(c+1)} \right]^{1/2}$
$a - 1$	$\left[\frac{(c-\gamma+1)(c-\gamma+2)}{(2c+2)(2c+3)} \right]^{1/2}$	$-\left[\frac{(c+\gamma+1)(c-\gamma+1)}{(c+1)(2c+3)} \right]^{1/2}$	$\left[\frac{(c+\gamma+2)(c+\gamma+1)}{(2c+2)(2c+3)} \right]^{1/2}$

Chapter 9

6j Symbols and the Racah Coefficients

9.1 Definition

9.1.1 6j Symbols

The Wigner 6j symbols [54] are related to the coefficients of transformations between different coupling schemes of three angular momenta. The angular momenta $\mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3$ may be coupled to give a resultant angular momentum $\mathbf{j}$ and its projection m in three different ways:

$$\begin{aligned} \text{I) } & \mathbf{j}_1 + \mathbf{j}_2 = \mathbf{j}_{12}, \quad \mathbf{j}_{12} + \mathbf{j}_3 = \mathbf{j}, \\ \text{II) } & \mathbf{j}_2 + \mathbf{j}_3 = \mathbf{j}_{23}, \quad \mathbf{j}_1 + \mathbf{j}_{23} = \mathbf{j}, \\ \text{III) } & \mathbf{j}_1 + \mathbf{j}_3 = \mathbf{j}_{13}, \quad \mathbf{j}_{13} + \mathbf{j}_2 = \mathbf{j}. \end{aligned} \quad (9.1.1)$$

Let $|j_1 j_2 (j_{12}) j_3 j m\rangle$ denote the state vectors corresponding to the coupling scheme I. These vectors are eigenvectors of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_{12}^2, \hat{j}^2, \hat{j}_x$ and may be written as

$$|j_1 j_2 (j_{12}) j_3 j m\rangle = \sum_{m_1 m_2 m_3} C_{j_{12} m_{12} j_3 m_3}^{j m} C_{j_1 m_1 j_2 m_2}^{j_{12} m_{12}} |j_1 m_1, j_2 m_2, j_3 m_3\rangle. \quad (9.1.2)$$

The state vectors corresponding to the coupling scheme II are eigenvectors of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_{23}^2, \hat{j}^2, \hat{j}_x$,

$$|j_1, j_2 j_3 (j_{23}) j m\rangle = \sum_{m_1 m_2 m_3} C_{j_1 m_1 j_{23} m_{23}}^{j m} C_{j_2 m_2 j_3 m_3}^{j_{23} m_{23}} |j_1 m_1, j_2 m_2, j_3 m_3\rangle. \quad (9.1.3)$$

Similarly, the state vectors corresponding to the coupling scheme III are eigenvectors of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_{13}^2, \hat{j}^2, \hat{j}_x$,

$$|j_1 j_3 (j_{13}) j_2 j m\rangle = \sum_{m_1 m_2 m_3} C_{j_{13} m_{13} j_2 m_2}^{j m} C_{j_1 m_1 j_3 m_3}^{j_{13} m_{13}} |j_1 m_1, j_2 m_2, j_3 m_3\rangle. \quad (9.1.4)$$

States belonging to each coupling scheme form a complete set of states. A transition from one coupling scheme to another is performed by some unitary transformation which relates the states with the same total angular momentum j and projection m . The coefficients U of this transformation differ from the 6j symbols only by normalization and phase factors. These factors are chosen in such a way to make the 6j symbols more symmetric (Sec. 9.4).

One defines the Wigner 6j symbols $\left\{ \begin{smallmatrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{smallmatrix} \right\}$ by the relation

$$\begin{aligned} \langle j_1 j_2 (j_{12}) j_3 j m | j_1, j_2 j_3 (j_{23}) j' m' \rangle &= \delta_{jj'} \delta_{mm'} U(j_1 j_2 j_3; j_{12} j_{23}) \\ &= \delta_{jj'} \delta_{mm'} (-1)^{j_1 + j_2 + j_3 + j} \sqrt{(2j_{12} + 1)(2j_{23} + 1)} \left\{ \begin{smallmatrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{smallmatrix} \right\}. \end{aligned} \quad (9.1.5)$$

From Eq. (9.1.5) one may obtain [92,64]

$$\begin{aligned} \langle j_1 j_2 (j_{12}) j_3 j m | j_1 j_3 (j_{13}) j_2 j' m' \rangle &= \delta_{jj'} \delta_{mm'} (-1)^{j + j_1 - j_{12} - j_{13}} U(j_2 j_1 j_3; j_{12} j_{13}) \\ &= \delta_{jj'} \delta_{mm'} (-1)^{j_2 + j_3 + j_{12} + j_{13}} \sqrt{(2j_{12} + 1)(2j_{13} + 1)} \left\{ \begin{smallmatrix} j_2 & j_1 & j_{12} \\ j_3 & j & j_{13} \end{smallmatrix} \right\} \end{aligned} \quad (9.1.6)$$

$$\begin{aligned} \langle j_1, j_2 j_3 (j_{23}) j m | j_1 j_3 (j_{13}) j_2 j' m' \rangle &= \delta_{jj'} \delta_{mm'} (-1)^{j_2 + j_3 - j_{23}} U(j_1 j_3 j_2; j_{13} j_{23}) \\ &= \delta_{jj'} \delta_{mm'} (-1)^{j_1 + j + j_{23}} \sqrt{(2j_{13} + 1)(2j_{23} + 1)} \left\{ \begin{smallmatrix} j_1 & j_3 & j_{13} \\ j_2 & j & j_{23} \end{smallmatrix} \right\}. \end{aligned} \quad (9.1.7)$$

According to the definition (9.1.5) the $6j$ symbols may be given in terms of the Clebsch-Gordan coefficients

$$\sum C_{j_{12} m_{12} j_3 m_3}^{j m} C_{j_1 m_1 j_2 m_2}^{j_{12} m_{12}} C_{j_1 m_1 j_{23} m_{23}}^{j' m'} C_{j_2 m_2 j_3 m_3}^{j_{23} m_{23}} = \delta_{jj'} \delta_{mm'} (-1)^{j_1+j_2+j_3+j} \sqrt{(2j_{12}+1)(2j_{23}+1)} \left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{matrix} \right\}. \quad (9.1.8)$$

Here the sum is over $m_1, m_2, m_3, m_{12}, m_{23}$ while m and m' are fixed. This relation completely determines absolute values and phases of the $6j$ symbols. The $6j$ symbols turn out to be real just as the Clebsch-Gordan coefficients are.

The quantum-mechanical rules of vector addition impose some restrictions on possible values of momenta which are arguments of the $6j$ symbol $\left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{matrix} \right\}$.

(a) All momenta are integer or half-integer nonnegative numbers (with one exception considered in Sec. 9.4).

(b) Each triad $(j_1 j_2 j_{12})$, $(j_{12} j_3 j)$, $(j_2 j_3 j_{23})$ and $(j_{23} j_1 j)$ should satisfy the triangular condition (Eq. (8.1.1)).

The unitarity of the recoupling transformations implies the orthogonality and normalization conditions of the $6j$ symbols,

$$\sum_{j_{12}} (2j_{12}+1)(2j_{23}+1) \left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{matrix} \right\} \left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j & j'_{23} \end{matrix} \right\} = \delta_{j_{23} j'_{23}}, \quad (9.1.9)$$

$$\sum_{j_{23}} (2j_{12}+1)(2j_{23}+1) \left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j & j_{23} \end{matrix} \right\} \left\{ \begin{matrix} j_1 & j_2 & j'_{12} \\ j_3 & j & j_{23} \end{matrix} \right\} = \delta_{j_{12} j'_{12}}. \quad (9.1.10)$$

Below we shall use Latin letters a, b, c , etc., to denote arguments of the $6j$ symbols.

9.1.2 Racah Coefficients

Instead of the Wigner $6j$ symbols the Racah coefficients [56] are often used, especially in spectroscopy theory. These coefficients differ from the $6j$ symbols only by a phase factor:

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \equiv (-1)^{a+b+d+e} W(abed; cf). \quad (9.1.11)$$

The Racah coefficients were introduced independently of the $6j$ symbols. The phase of the Racah coefficients coincides with the phase of the coefficients which describe the transformation between I and II coupling schemes (Eq. (9.1.5)).

9.1.3 R Symbols

The $6j$ symbols and the Racah coefficients may be written in the form of a 3×4 array $\|R_{i\alpha}\|$ ($i = 1, 2, 3; \alpha = 1, 2, 3, 4$) which is called the R -symbol (Shelepin [70])

$$\left\| \begin{matrix} R_{11} & R_{12} & R_{13} & R_{14} \\ R_{21} & R_{22} & R_{23} & R_{24} \\ R_{31} & R_{32} & R_{33} & R_{34} \end{matrix} \right\| \equiv \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \equiv (-1)^{a+b+d+e} W(abed; cf), \quad (9.1.12)$$

where

$$\begin{aligned} R_{11} &= -c + d + e, & R_{12} &= b + d - f, & R_{13} &= a + e - f, & R_{14} &= a + b - c, \\ R_{21} &= -b + d + f, & R_{22} &= c + d - e, & R_{23} &= a - b + c, & R_{24} &= a - e + f, \\ R_{31} &= -a + e + f, & R_{32} &= -a + b + c, & R_{33} &= c - d + e, & R_{34} &= b - d + f. \end{aligned} \quad (9.1.13)$$

The inverse relations are

$$\begin{aligned} 2a &= R_{13} + R_{24} = R_{14} + R_{23}, & 2d &= R_{11} + R_{22} = R_{12} + R_{21}, \\ 2b &= R_{12} + R_{34} = R_{14} + R_{32}, & 2e &= R_{11} + R_{33} = R_{13} + R_{31}, \\ 2c &= R_{22} + R_{33} = R_{23} + R_{32}, & 2f &= R_{21} + R_{34} = R_{24} + R_{31}. \end{aligned} \quad (9.1.14)$$

All 12 elements $R_{i\alpha}$ are integer nonnegative numbers. The differences between corresponding elements of rows and columns are constant:

$$\begin{aligned} R_{i\alpha} - R_{k\alpha} &= R_{i\beta} - R_{k\beta}, \\ R_{i\alpha} - R_{i\beta} &= R_{k\alpha} - R_{k\beta}, \end{aligned} \quad (i, k = 1, 2, 3; \alpha, \beta = 1, 2, 3, 4). \quad (9.1.15)$$

Note the following relations:

$$\begin{aligned} \sum_{i=1}^3 R_{i1} &= 2(d + e + f) - a - b - c, & \sum_{i=1}^3 R_{i3} &= 2(a + c + e) - b - d - f, \\ \sum_{i=1}^3 R_{i2} &= 2(b + c + d) - a - e - f, & \sum_{i=1}^3 R_{i4} &= 2(a + b + f) - c - d - e, \\ \sum_{i,\alpha} R_{i\alpha} &= 2(a + b + c + d + e + f). \end{aligned} \quad (9.1.16)$$

One may also use the following parametrisation of the elements $R_{i\alpha}$ [20]:

$$R_{i\alpha} = A_i - B_\alpha. \quad (9.1.17)$$

Here A_i, B_α are integer nonnegative numbers

$$\begin{aligned} A_1 &= a + b + d + e, & B_1 &= a + b + c, \\ A_2 &= a + c + d + f, & B_2 &= a + e + f, \\ A_3 &= b + c + e + f, & B_3 &= b + d + f, \\ & & B_4 &= c + d + e, \end{aligned} \quad (9.1.18)$$

with

$$\sum_{i=1}^3 A_i = \sum_{\alpha=1}^4 B_\alpha = 2(a + b + c + d + e + f). \quad (9.1.19)$$

The inverse relations are

$$\begin{aligned} 2a &= A_1 + A_2 - B_3 - B_4, & 2d &= A_1 + A_2 - B_1 - B_2, \\ 2b &= A_1 + A_3 - B_2 - B_4, & 2e &= A_1 + A_3 - B_1 - B_3, \\ 2c &= A_2 + A_3 - B_2 - B_3, & 2f &= A_2 + A_3 - B_1 - B_4. \end{aligned} \quad (9.1.20)$$

The R symbols provide the simplest formulation of the symmetry properties of the $6j$ symbols and Racah coefficients.

9.2 General Expressions for the $6j$ Symbols. Relations Between the $6j$ Symbols and Other Functions

The $6j$ symbols $\left\{ \begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix} \right\}$ vanish if at least one of the triads $(abc), (cde), (aef)$ and (bdf) does not obey the triangular conditions (8.1.1). The expressions for the $6j$ symbols given below are valid if all these conditions are satisfied. Corresponding expressions for the Racah coefficients may be obtained by the use of the relations between these coefficients and the $6j$ symbols (9.1.11).

9.2.1 Expressions for the $6j$ Symbols in Terms of Finite Sums

In the expressions presented below the sums are over all integer nonnegative values of n so that no factorial in denominators has a negative argument. The quantities $\Delta(abc)$ are defined in Eq. (8.2.1).

$$\begin{aligned} \left\{ \begin{smallmatrix} a & b & c \\ d & e & f \end{smallmatrix} \right\} &= \Delta(abc)\Delta(cde)\Delta(aef)\Delta(bdf) \\ &\times \sum_n \frac{(-1)^n (n+1)!}{(n-a-b-c)!(n-c-d-e)!(n-a-e-f)!(n-b-d-f)!(a+b+d+e-n)!} \\ &\times \frac{1}{(a+c+d+f-n)!(b+c+e+f-n)!} \\ &\quad \text{(Racah [56])} \end{aligned} \quad (9.2.1)$$

By the replacement $n \rightarrow a + b + d + e - n$ one can rewrite Eq. (9.2.1) in the form

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{a+b+d+e} \Delta(abc) \Delta(cde) \Delta(aef) \Delta(bdf) \\ &\times \sum_n \frac{(-1)^n (a+b+d+e+1-n)!}{n! (a+b-c-n)! (-c+d+e-n)! (a+e-f-n)! (b+d-f-n)!} \\ &\times \frac{1}{(-a+c-d+f+n)! (-b+c-e+f+n)!} \\ &\quad \text{(Racah [56])} \end{aligned} \quad (9.2.2)$$

Some other expressions for the 6j symbols which cannot be easily reduced to (9.2.1) and (9.2.2) are [45,50].

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{a+c+d+f} \frac{\Delta(aef) \Delta(bdf)}{\Delta(abc) \Delta(cde)} \\ &\times \sum_n (-1)^n \frac{(-a+b+c+n)! (c-d+e+n)! (a-c+d+f-n)!}{n! (a-e+f-n)! (-b+d+f-n)! (-a+b-d+e+n)! (b+c+e-f+1+n)!}, \end{aligned} \quad (9.2.3)$$

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{a+c+d+f} \frac{\Delta(abc) \Delta(bdf)}{\Delta(aef) \Delta(cde)} \\ &\times \sum_n (-1)^n \frac{(a-b+d+e-n)! (-b+c+e+f-n)! (a+c+d+f+1-n)!}{n! (a-b+c-n)! (-b+d+f-n)! (a+e+f+1-n)! (c+d+e+1-n)!}, \end{aligned} \quad (9.2.4)$$

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{a+b+d+e} \frac{\Delta(abc) \Delta(cde) \Delta(aef) \Delta(bdf) (a+e+f+1)! (b+d+f+1)!}{(a+b-c)! (a-b+c)! (-c+d+e)! (c+d-e)! (-a+e+f)! (b-d+f)!} \\ &\times \sum_n (-1)^n \frac{(-a+e+f+n)! (b-d+f+n)! (a+c+d-f-n)!}{n! (a+e-f-n)! (b+d-f-n)! (-a+c-d+f+n)! (2f+1+n)!}, \end{aligned} \quad (9.2.5)$$

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{b+c+e+f} \frac{\Delta(abc) \Delta(cde) \Delta(aef) \Delta(bdf) (a+b+c+1)! (b+d+f+1)!}{(a+b-c)! (c-d+e)! (c+d-e)! (a-e+f)! (-a+e+f)! (b+d-f)!} \\ &\times \sum_n (-1)^n \frac{(2b-n)! (b+c-e+f-n)! (b+c+e+f+1-n)!}{n! (-a+b+c-n)! (b-d+f-n)! (a+b+c+1-n)! (b+d+f+1-n)!}. \end{aligned} \quad (9.2.6)$$

9.2.2 Bargmann Formula [61]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \equiv \left\| \begin{matrix} R_{11} & R_{12} & R_{13} & R_{14} \\ R_{21} & R_{22} & R_{23} & R_{24} \\ R_{31} & R_{32} & R_{33} & R_{34} \end{matrix} \right\| = \left[\frac{\prod_{i=1}^3 \prod_{\alpha=1}^4 (R_{i\alpha})!}{\prod_{\alpha=1}^4 (B_{\alpha} + 1)!} \right]^{\frac{1}{2}} \sum_{x_i, y_{\alpha}} \frac{(-1)^n (n+1)!}{\prod_{i=1}^3 (x_i)! \prod_{\alpha=1}^4 (y_{\alpha})!} \quad (9.2.7)$$

Here $R_{i\alpha}$ are elements of the R symbol (Sec. 9.1.3), B_{α} are given by Eq. (9.1.18), x_i, y_{α} are summation indices, $n \equiv \sum_{i=1}^3 x_i + \sum_{\alpha=1}^4 y_{\alpha}$. The sums are over all integer nonnegative values of x_i, y_{α} which satisfy the conditions $x_i + y_{\alpha} = R_{i\alpha}$. These conditions show that only one of the summation indices is independent. The sum in (9.2.7) contains $r+1$ terms where $r = \min \{R_{i\alpha}\}$. If we take the quantity n (integer nonnegative) as an independent summation index, then $x_i = A_i - n, y_{\alpha} = n - B_{\alpha}, A_i$ and B_{α} being given by Eq. (9.1.18). In this case the Bargmann formula (9.2.7) reduces to the Racah formula (9.2.1).

9.2.3 Relations Between the $6j$ Symbols and the Generalized Hypergeometric Functions

The $6j$ symbols may be written in terms of the hypergeometric functions ${}_4F_3$ with unit argument:

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{a+b+d+e} \frac{\Delta(abc)\Delta(cde)\Delta(aef)\Delta(bdf)(a+b+d+e+1)!}{(a+b-c)!(-c+d+e)!(a+e-f)!(b+d-f)!(-a+c-d+f)!(-b+c-e+f)!} \\ \times {}_4F_3 \left[\begin{matrix} -a-b+c, c-d-e, -a-e+f, -b-d+f \\ -a-b-d-e-1, -a+c-d+f+1, -b+c-e+f+1 \end{matrix} \middle| 1 \right], \quad (\text{Rose [4]}) \quad (9.2.8)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{a+c+d+f} \frac{\Delta(aef)\Delta(bdf)(-a+b+c)!(c-d+e)!(a-c+d+f)!}{\Delta(abc)\Delta(cde)(a-e+f)!(-b+d+f)!(-a+b-d+e)!(b+c+e-f+1)!} \\ \times {}_4F_3 \left[\begin{matrix} -a+b+c+1, c-d+e+1, -a+e-f, b-d-f \\ -a+c-d-f, -a+b-d+e+1, b+c+e-f+2 \end{matrix} \middle| 1 \right], \quad (9.2.9)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{a+c+d+f} \frac{\Delta(abc)\Delta(bdf)(a-b+d+e)!(-b+c+e+f)!(a+c+d+f+1)!}{\Delta(aef)\Delta(cde)(a-b+c)!(-b+d+f)!(a+e+f+1)!(c+d+e+1)!} \\ \times {}_4F_3 \left[\begin{matrix} -a+b-c, b-d-f, -a-e-f-1, -c-d-e-1 \\ -a+b-d-e, b-c-e-f, -a-c-d-f-1 \end{matrix} \middle| 1 \right], \quad (9.2.10)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{a+b+d+e} \\ \times \frac{\Delta(abc)\Delta(cde)\Delta(aef)\Delta(bdf)(a+e+f+1)!(b+d+f+1)!(a+c+d-f)!}{(a+b-c)!(a-b+c)!(-c+d+e)!(c+d-e)!(a+e-f)!(b+d-f)!(-a+c-d+f)!(2f+1)!} \\ \times {}_4F_3 \left[\begin{matrix} -a-e+f, -b-d+f, -a+e+f+1, b-d+f+1 \\ -a-c-d+f, -a+c-d+f+1, 2f+2 \end{matrix} \middle| 1 \right], \quad (9.2.11)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{b+c+e+f} \\ \times \frac{\Delta(abc)\Delta(cde)\Delta(aef)\Delta(bdf)(2b)!(b+c-e+f)!(b+c+e+f+1)!}{(-a+b+c)!(a+b-c)!(c-d+e)!(c+d-e)!(a-e+f)!(a+e+f)!(b+d-f)!(b-d+f)!} \\ \times {}_4F_3 \left[\begin{matrix} a-b-c, -b+d-f, -a-b-c-1, -b-d-f-1 \\ -2b, -b-c+e-f, -b-c-e-f-1 \end{matrix} \middle| 1 \right]. \quad (9.2.12)$$

Equations (9.2.8)-(9.2.12) present Eqs. (9.2.2)-(9.2.6) in terms of the hypergeometric functions.

9.2.4 Relations Between the $6j$ Symbols and the $3jm$ Symbols

The $6j$ symbols may be written as sums of products of the Clebsch-Gordan coefficients (Eq. (9.1.8)) or $3jm$ symbols. The relations between the $6j$ symbols and the $3jm$ symbols are

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = \sum (-1)^{d+e+f+\delta+\varepsilon+\varphi} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \begin{pmatrix} a & e & f \\ \alpha & \varepsilon & -\varphi \end{pmatrix} \begin{pmatrix} d & b & f \\ -\delta & \beta & \varphi \end{pmatrix} \begin{pmatrix} d & e & c \\ \delta & -\varepsilon & \gamma \end{pmatrix}. \quad (9.2.13)$$

In Eq. (9.2.13) the sum is over all possible values of $\alpha, \beta, \gamma, \delta, \varepsilon, \varphi$ with only three summation indices being independent. Some other sums of the $3jm$ symbols which yield the $6j$ symbols will be considered in Chap. 12.

9.2.5 Quasi-Binomial Representation of the $6j$ Symbols

The $6j$ symbols may be written in terms of quasi-binomials [45, 99] which are defined in Sec. 8.2.2. These representations are widely used in tabulating the formulas for the $6j$ symbols.

Let us introduce the following definitions:

$$\begin{aligned} k_1 &\equiv e-d, & B &\equiv R_{21} = -b+d+f, & F &\equiv R_{12} = b+d-f, \\ k_2 &\equiv a-b, & D &\equiv R_{34} = b-d+f, & E &\equiv R_{12} + R_{21} + R_{34} + 1 = b+d+f+1 \end{aligned} \quad (9.2.14)$$

Then the dependence of the $6j$ symbols on k_1 and k_2 is given by

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \equiv \left\{ \begin{matrix} b+k_2 & b & c \\ d & d+k_1 & f \end{matrix} \right\} \\ = (-1)^{E+k_1+k_2+1} \left[\frac{(c+k_2)^{(2k_2)} B^{(k_2-k_1)} D^{(-1)(k_2-k_1)} E^{(-1)(k_2+k_1)} F^{(-1)(k_2+k_1)}}{(c+k_1)!(c-k_1)!(2b+c+k_2+1)^{(2c+1)} (2d+c+k_1+1)^{(2c+1)}} \right]^{\frac{1}{2}} (u-v)^{(c-k_2)}. \quad (9.2.15)$$

The quantities u and v may be chosen in different ways. This depends on which equations in Sec. 9.2.1. are supposed to be written in a quasi-binomial form [20].

Equations (9.2.1) and (9.2.2) are obtained by putting

$$\begin{aligned} u &= (c + k_1)^{(1)} (B - k_2 + k_1)^{(1)} D^{(1)} \\ v &= (c - k_1)^{(1)} F^{(1)} (E + k_1 + k_2)^{(-1)} \end{aligned} \quad (9.2.16)$$

Equation (9.2.3) is obtained, if

$$\begin{aligned} u &= (c + k_1)^{(1)} B^{(-1)} (D + k_2 - k_1)^{(-1)} \\ v &= (c - k_1)^{(1)} (F + k_2 + k_1)^{(-1)} E^{(1)} \end{aligned} \quad (9.2.17)$$

or

$$\begin{aligned} u &= (2d - c + k_1)^{(-1)} D^{(1)} (F + k_2 + k_1)^{(-1)} \\ v &= (2d + c + k_1 + 1)^{(1)} (D + k_2 - k_1)^{(-1)} F^{(1)} \end{aligned} \quad (9.2.18)$$

Equation (9.2.4) corresponds to

$$\begin{aligned} u &= (2d + c + k_1 + 1)^{(1)} (B - k_2 + k_1)^{(1)} E^{(1)} \\ v &= (2d - c + k_1)^{(-1)} B^{(-1)} (E + k_2 + k_1)^{(-1)} \end{aligned} \quad (9.2.19)$$

Equation (9.2.5) is obtained provided

$$\begin{aligned} u &= (c + k_2)^{(-1)} (B - k_2 + k_1)^{(1)} (D + k_2 - k_1)^{(-1)} \\ v &= (c - k_1)^{(1)} (2b + c + k_2 + 1)^{(1)} (2d - c + k_1)^{(-1)} \end{aligned} \quad (9.2.20)$$

or

$$\begin{aligned} u &= (c + k_1)^{(1)} (2b - c + k_2)^{(-1)} (2d - c + k_1)^{(-1)} \\ v &= (c + k_2)^{(-1)} F^{(1)} E^{(1)} \end{aligned} \quad (9.2.21)$$

Equation (9.2.6) corresponds to

$$\begin{aligned} u &= D^{(1)} E^{(1)} (2b + c + k_2 + 1)^{(1)} \\ v &= (D + k_2 - k_1)^{(-1)} (E + k_2 + k_1)^{(-1)} (2b - c + k_2)^{(-1)} \end{aligned} \quad (9.2.22)$$

or

$$\begin{aligned} u &= (c + k_1)^{(1)} (2b + c + k_2 + 1)^{(1)} (2d + c + k_1 + 1)^{(1)} \\ v &= (c + k_2)^{(-1)} (F + k_2 + k_1)^{(-1)} (E + k_2 + k_1)^{(-1)} \end{aligned} \quad (9.2.23)$$

Equation (9.2.15) for the $6j$ symbols is valid, if all the exponents $2k_2$, $k_2 - k_1$ and $k_2 + k_1$ are integer nonnegative numbers, i.e., if $k_2 \geq |k_1| \geq 0$. If some of the exponents are negative, the corresponding quasi-power should be replaced in accordance with

$$p^{(\sigma)} \rightarrow \frac{1}{p^{(-1)(|\sigma|)}}, \quad p^{(-1)(\sigma)} \rightarrow \frac{1}{p^{(|\sigma|)}} \quad \text{for } \sigma < 0 \quad (9.2.24)$$

9.3 Integral Representations of the $6j$ Symbols

Squares of the $6j$ symbols may be expressed by integrals involving the characters of the representations of the rotation group [54]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}^2 = \frac{1}{(8\pi^2)^3} \int dR_1 dR_2 dR_3 \chi^a(R_1) \chi^b(R_2) \chi^c(R_3) \chi^d(R_2 R_3^{-1}) \chi^e(R_3 R_1^{-1}) \chi^f(R_1 R_2^{-1}) \quad (9.3.1)$$

Here

$$\chi^j(R) \equiv \sum_m D_{mm}^j(R) \equiv \sum_m D_{mm}^j(\alpha, \beta, \gamma)$$

is the character of the representation of rank j (Sec. 4.14)

$$\int f(R) dR \equiv \int_0^{2\pi} d\alpha \int_0^\pi \sin \beta d\beta \int_0^{2\pi} d\gamma f(\alpha, \beta, \gamma)$$

Note also the following integral representations for some special $6j$ symbols

$$\left\{ \begin{matrix} a & b & c \\ a & b & f \end{matrix} \right\} = \frac{(-1)^{2c}}{(8\pi^2)^2} \int dR_1 dR_2 \chi^c(R_1) \chi^f(R_2) \chi^a(R_2 R_1) \chi^b(R_2^{-1} R_1), \quad (9.3.2)$$

$$\begin{aligned} & \left\{ \begin{matrix} a & b & g \\ d & b & c \end{matrix} \right\} \left\{ \begin{matrix} a & b & g \\ d & b & f \end{matrix} \right\} \left\{ \begin{matrix} a & b & c \\ d & b & f \end{matrix} \right\} \\ &= \frac{(-1)^{2a}}{(8\pi^2)^4} \int dR_1 dR_2 dR_3 dR_4 \chi^c(R_1) \chi^g(R_2) \chi^f(R_3) \chi^a(R_4 R_2) \chi^b(R_4 R_3 R_2 R_1 R_4) \chi^d(R_1 R_4 R_3), \end{aligned} \quad (9.3.3)$$

$$\begin{aligned} & \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \left\{ \begin{matrix} a & e & f \\ a & b & c \end{matrix} \right\} \left\{ \begin{matrix} d & b & f \\ a & b & c \end{matrix} \right\} \left\{ \begin{matrix} d & e & c \\ a & b & c \end{matrix} \right\} \\ &= \frac{(-1)^{2d}}{(8\pi^2)^5} \int dR_1 dR_2 dR_3 dR_4 dR_5 \chi^d(R_1) \chi^e(R_2) \chi^f(R_3) \chi^a(R_4) \\ & \quad \times \chi^a(R_5^{-1} R_3 R_2) \chi^b(R_5 R_4 R_2 R_1) \chi^c(R_4^{-1} R_5 R_1 R_3). \end{aligned} \quad (9.3.4)$$

9.4 Symmetries of the $6j$ Symbols and the Racah Coefficients

9.4.1 R-Symbols

The symmetry properties of the $6j$ symbols and the W -coefficients may be formulated in a fairly simple way if these coefficients are written in terms of the R symbols (see Sec. 9.1.3).

The value of the R symbol is invariant under any permutation of its rows or columns [70]

$$\left\| \begin{matrix} R_{11} & R_{12} & R_{13} & R_{14} \\ R_{21} & R_{22} & R_{23} & R_{24} \\ R_{31} & R_{32} & R_{33} & R_{34} \end{matrix} \right\| = \left\| \begin{matrix} R_{i1} & R_{i2} & R_{i3} & R_{i4} \\ R_{k1} & R_{k2} & R_{k3} & R_{k4} \\ R_{l1} & R_{l2} & R_{l3} & R_{l4} \end{matrix} \right\| = \left\| \begin{matrix} R_{1\alpha} & R_{1\beta} & R_{1\gamma} & R_{1\delta} \\ R_{2\alpha} & R_{2\beta} & R_{2\gamma} & R_{2\delta} \\ R_{3\alpha} & R_{3\beta} & R_{3\gamma} & R_{3\delta} \end{matrix} \right\|. \quad (9.4.1)$$

In other words, any permutation of parameters A_i or B_α (see Sec. 9.1.3) leaves the value of the R symbol unchanged. These symmetry relations involve $3! \times 4! = 144$ generally different Racah coefficients.

9.4.2 $6j$ Symbols

The above-mentioned symmetries of the R symbol are equivalent to the following symmetries of the $6j$ symbols.

(a) *Classical Symmetries* [54]

The $6j$ symbol is invariant under any permutation of its columns or under interchange of the upper and lower arguments in each of any two columns:

$$\begin{aligned} & \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = \left\{ \begin{matrix} a & c & b \\ d & f & e \end{matrix} \right\} = \left\{ \begin{matrix} b & a & c \\ e & d & f \end{matrix} \right\} = \left\{ \begin{matrix} b & c & a \\ e & f & d \end{matrix} \right\} = \left\{ \begin{matrix} c & a & b \\ f & d & e \end{matrix} \right\} = \left\{ \begin{matrix} c & b & a \\ f & e & d \end{matrix} \right\} \\ &= \left\{ \begin{matrix} a & e & f \\ d & b & c \end{matrix} \right\} = \left\{ \begin{matrix} a & f & e \\ d & c & b \end{matrix} \right\} = \left\{ \begin{matrix} e & a & f \\ b & d & c \end{matrix} \right\} = \left\{ \begin{matrix} e & f & a \\ b & c & d \end{matrix} \right\} = \left\{ \begin{matrix} f & a & e \\ c & d & b \end{matrix} \right\} = \left\{ \begin{matrix} f & e & a \\ c & b & d \end{matrix} \right\} \\ &= \left\{ \begin{matrix} d & e & c \\ a & b & f \end{matrix} \right\} = \left\{ \begin{matrix} d & c & e \\ a & f & b \end{matrix} \right\} = \left\{ \begin{matrix} e & d & c \\ b & a & f \end{matrix} \right\} = \left\{ \begin{matrix} e & c & d \\ b & f & a \end{matrix} \right\} = \left\{ \begin{matrix} c & d & e \\ f & a & b \end{matrix} \right\} = \left\{ \begin{matrix} c & e & d \\ f & b & a \end{matrix} \right\} \\ &= \left\{ \begin{matrix} d & b & f \\ a & e & c \end{matrix} \right\} = \left\{ \begin{matrix} d & f & b \\ a & c & e \end{matrix} \right\} = \left\{ \begin{matrix} b & d & f \\ e & a & c \end{matrix} \right\} = \left\{ \begin{matrix} b & f & d \\ e & c & a \end{matrix} \right\} = \left\{ \begin{matrix} f & d & b \\ c & a & e \end{matrix} \right\} = \left\{ \begin{matrix} f & b & d \\ c & e & a \end{matrix} \right\}. \end{aligned} \quad (9.4.2)$$

These relations involve $3! \times 4 = 24$ different $6j$ symbols.

(b) Regge Symmetries [71]

The relations below are functional ones, i.e. in general they cannot be obtained by interchanging the 6j symbol arguments.

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= \left\{ \begin{matrix} a & s_1 - b & s_1 - c \\ d & s_1 - e & s_1 - f \end{matrix} \right\} = \left\{ \begin{matrix} s_2 - a & b & s_2 - c \\ s_2 - d & e & s_2 - f \end{matrix} \right\} \\ &= \left\{ \begin{matrix} s_3 - a & s_3 - b & c \\ s_3 - d & s_3 - e & f \end{matrix} \right\} = \left\{ \begin{matrix} s_2 - d & s_3 - e & s_1 - f \\ s_2 - a & s_3 - b & s_1 - c \end{matrix} \right\} = \left\{ \begin{matrix} s_3 - d & s_1 - e & s_2 - f \\ s_3 - a & s_1 - b & s_2 - c \end{matrix} \right\}, \end{aligned} \quad (9.4.3)$$

where

$$s_1 = \frac{1}{2}(b + c + e + f), \quad s_2 = \frac{1}{2}(a + c + d + f), \quad s_3 = \frac{1}{2}(a + b + d + e). \quad (9.4.4)$$

These relations are especially useful when s_i equals one of the 6j symbol arguments. Combining the Regge symmetries and the classical symmetries, one gets all 144 symmetry relations.

9.4.3 Racah Coefficients

For the Racah coefficients the symmetry relations are the following.

(a) Classical Symmetries [56]

$$\begin{aligned} W(abed; cf) &= W(deba; cf) = W(edab; cf) = W(bade; cf) \\ &= W(aebd; fc) = W(dbea; fc) = W(bdae; fc) = W(eadb; fc) \\ &= \varepsilon_1 W(acfd; be) = \varepsilon_1 W(dfca; be) = \varepsilon_1 W(fdac; be) = \varepsilon_1 W(cadf; be) \\ &= \varepsilon_1 W(afcd; eb) = \varepsilon_1 W(dcf a; eb) = \varepsilon_1 W(cda f; eb) = \varepsilon_1 W(fadc; eb) \\ &= \varepsilon_2 W(cbef; ad) = \varepsilon_2 W(febc; ad) = \varepsilon_2 W(ecfb; ad) = \varepsilon_2 W(bcfe; ad) \\ &= \varepsilon_2 W(cebf; da) = \varepsilon_2 W(fbec; da) = \varepsilon_2 W(bfce; da) = \varepsilon_2 W(ecfb; da), \end{aligned} \quad (9.4.5)$$

where

$$\varepsilon_1 = (-1)^{b+e-c-f}, \quad \varepsilon_2 = (-1)^{a+d-c-f} \quad (9.4.6)$$

(b) Regge Symmetries

$$\begin{aligned} W(abed; cf) &= W(s_3 - a, s_3 - b, s_3 - e, s_3 - d; cf) = \varepsilon_1 W(a, s_1 - b, s_1 - e, d; s_1 - c, s_1 - f) \\ &= \varepsilon_1 W(s_2 - d, s_3 - e, s_3 - b, s_2 - a; s_1 - f, s_1 - c) = \varepsilon_2 W(s_2 - a, b, e, s_2 - d; s_2 - c, s_2 - f) \\ &= \varepsilon_2 W(s_3 - d, s_1 - e, s_1 - b, s_3 - a; s_2 - f, s_2 - c) \end{aligned} \quad (9.4.7)$$

Here s_1, s_2, s_3 are given by Eq. (9.4.4) and $\varepsilon_1, \varepsilon_2$ by Eq. (9.4.6).

9.4.4 "Mirror" Symmetry

The formulas for the 6j symbols may be extended to include negative integer or half-integer values of arguments. In this case one has the following symmetry properties [20] corresponding to the replacement $j \rightarrow -j - 1$,

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} &= - \left\{ \begin{matrix} \bar{a} & \bar{b} & \bar{c} \\ \bar{d} & \bar{e} & \bar{f} \end{matrix} \right\} = (-1)^{\varphi_1} \left\{ \begin{matrix} \bar{a} & b & c \\ d & e & f \end{matrix} \right\} = (-1)^{\varphi_1+1} \left\{ \begin{matrix} a & \bar{b} & \bar{c} \\ \bar{d} & \bar{e} & \bar{f} \end{matrix} \right\} \\ &= (-1)^{\varphi_2} \left\{ \begin{matrix} \bar{a} & b & c \\ \bar{d} & e & f \end{matrix} \right\} = (-1)^{\varphi_2+1} \left\{ \begin{matrix} a & \bar{b} & \bar{c} \\ \bar{d} & \bar{e} & \bar{f} \end{matrix} \right\} = i(-1)^{\varphi_3} \left\{ \begin{matrix} \bar{a} & \bar{b} & c \\ d & e & f \end{matrix} \right\} = i(-1)^{\varphi_3} \left\{ \begin{matrix} a & b & \bar{c} \\ \bar{d} & \bar{e} & \bar{f} \end{matrix} \right\} \\ &= i(-1)^{\varphi_4} \left\{ \begin{matrix} \bar{a} & \bar{b} & \bar{c} \\ d & e & f \end{matrix} \right\} = i(-1)^{\varphi_4} \left\{ \begin{matrix} a & b & c \\ \bar{d} & \bar{e} & \bar{f} \end{matrix} \right\} = (-1)^{\varphi_5} \left\{ \begin{matrix} \bar{a} & \bar{b} & c \\ \bar{d} & e & f \end{matrix} \right\} = (-1)^{\varphi_5+1} \left\{ \begin{matrix} a & b & \bar{c} \\ d & \bar{e} & \bar{f} \end{matrix} \right\}. \end{aligned} \quad (9.4.8)$$

Here

$$\begin{aligned} \bar{a} &\equiv -a - 1, \quad \bar{b} \equiv -b - 1, \quad \text{etc.} \\ \varphi_1 &= b - c - e + f, \varphi_2 = 2(a + d), \varphi_3 = c + d + e + 2f, \varphi_4 = a + b + c, \varphi_5 = 2(c + f) + 1 \end{aligned} \quad (9.4.9)$$

Similarly, for the Racah coefficients one gets

$$\begin{aligned}
 W(abed; cf) &= -W(\bar{a}\bar{b}\bar{e}\bar{d}; \bar{c}\bar{f}) = W(\bar{a}\bar{b}\bar{e}\bar{d}; cf) = -W(\bar{a}\bar{b}\bar{e}\bar{d}; \bar{c}\bar{f}) \\
 &= (-1)^{\psi_1+1} W(\bar{a}\bar{b}\bar{e}\bar{d}; cf) = (-1)^{\psi_1} W(\bar{a}\bar{b}\bar{e}\bar{d}; \bar{c}\bar{f}) = i(-1)^{\psi_2} W(\bar{a}\bar{b}\bar{e}\bar{d}; cf) = i(-1)^{\psi_2} W(ab\bar{e}\bar{d}; \bar{c}\bar{f}) \\
 &= i(-1)^{\psi_3} W(\bar{a}\bar{b}\bar{e}\bar{d}; \bar{c}\bar{f}) = i(-1)^{\psi_3} W(ab\bar{e}\bar{d}; cf) = (-1)^{\psi_4} W(\bar{a}\bar{b}\bar{e}\bar{d}; cf) = (-1)^{\psi_4+1} W(ab\bar{e}\bar{d}; \bar{c}\bar{f}). \quad (9.4.10)
 \end{aligned}$$

Here

$$\psi_1 = -b + c - e + f, \quad \psi_2 = -c + d + e + 2f, \quad \psi_3 = a + b - c, \quad \psi_4 = 2(d + f) \quad (9.4.11)$$

9.5 Explicit Forms of the $6j$ Symbols for Certain Arguments

9.5.1 One of Arguments is Equal to Zero

For the $6j$ symbols one obtains

$$\begin{aligned}
 \left\{ \begin{matrix} 0 & b & c \\ d & e & f \end{matrix} \right\} &= (-1)^{b+e+d} \frac{\delta_{bc}\delta_{ef}}{\sqrt{(2b+1)(2e+1)}}, & \left\{ \begin{matrix} a & b & c \\ 0 & e & f \end{matrix} \right\} &= (-1)^{a+b+e} \frac{\delta_{bf}\delta_{ce}}{\sqrt{(2b+1)(2c+1)}}, \\
 \left\{ \begin{matrix} a & 0 & c \\ d & e & f \end{matrix} \right\} &= (-1)^{a+d+e} \frac{\delta_{ac}\delta_{df}}{\sqrt{(2a+1)(2d+1)}}, & \left\{ \begin{matrix} a & b & c \\ d & 0 & f \end{matrix} \right\} &= (-1)^{a+b+d} \frac{\delta_{af}\delta_{cd}}{\sqrt{(2a+1)(2c+1)}}, \\
 \left\{ \begin{matrix} a & b & 0 \\ d & e & f \end{matrix} \right\} &= (-1)^{a+e+f} \frac{\delta_{ab}\delta_{de}}{\sqrt{(2a+1)(2d+1)}}, & \left\{ \begin{matrix} a & b & c \\ d & e & 0 \end{matrix} \right\} &= (-1)^{a+b+c} \frac{\delta_{ae}\delta_{bd}}{\sqrt{(2a+1)(2b+1)}}. \quad (9.5.1)
 \end{aligned}$$

Analogous relations for the Racah coefficients are

$$\begin{aligned}
 W(0bed; cf) &= \frac{\delta_{bc}\delta_{ef}}{\sqrt{(2b+1)(2e+1)}}, & W(a0ed; cf) &= \frac{\delta_{ac}\delta_{df}}{\sqrt{(2a+1)(2d+1)}}, \\
 W(ab0d; cf) &= \frac{\delta_{af}\delta_{cd}}{\sqrt{(2a+1)(2c+1)}}, & W(abe0; cf) &= \frac{\delta_{bf}\delta_{ce}}{\sqrt{(2b+1)(2c+1)}}, \\
 W(abed; 0f) &= (-1)^{a+e-f} \frac{\delta_{ab}\delta_{de}}{\sqrt{(2a+1)(2d+1)}}, & W(abed; c0) &= (-1)^{a+b-c} \frac{\delta_{ae}\delta_{bd}}{\sqrt{(2a+1)(2b+1)}} \quad (9.5.2)
 \end{aligned}$$

In this case all other arguments are supposed to satisfy the triangular condition.

9.5.2 One of Arguments is Equal to the Sum of Two Others

If one of the $6j$ symbol arguments is equal to sum of two others from the same triad (abc), (cde), (aef), (bdf), one may use the classical symmetries of the $6j$ symbol (Eqs. (9.4.2)) to express it in the form

$$\begin{aligned}
 \left\{ \begin{matrix} a & b & a+b \\ d & e & f \end{matrix} \right\} &= (-1)^{a+b+d+e} W(abed; a+bf) = (-1)^{a+b+d+e} \\
 &\times \left[\frac{(2a)!(2b)!(a+b+d+e+1)!(a+b-d+e)!(a+b+d-e)!(-a+e+f)!(-b+d+f)!}{(2a+2b+1)!(-a-b+d+e)!(a+e-f)!(a-e+f)!(a+e+f+1)!(b+d-f)!(b-d+f)!(b+d+f+1)!} \right]^{\frac{1}{2}} \quad (9.5.3)
 \end{aligned}$$

In particular,

$$\left\{ \begin{matrix} a & b & a+b \\ d & e & a+e \end{matrix} \right\} = (-1)^{a+b+d+e} \left[\frac{(2b)!(2e)!(a+b+d-e)!(a-b+d+e)!}{(2a+2b+1)!(2a+2e+1)!(-a-b+d+e)!(-a+b+d-e)!} \right]^{\frac{1}{2}} \quad (9.5.4)$$

$$\left\{ \begin{matrix} a & b & a+b \\ d & e & a+e \end{matrix} \right\} = (-1)^{a+b+d+e} \times \left[2a(a+b+d+e+1)(a+b-d+e) \frac{(2b)!(2e-1)!(a+b+d-e)!(a-b+d+e-1)!}{(2a+2b+1)!(2a+2e)!(a-b+d+e)!(-a+b+d-e+1)!} \right]^{\frac{1}{2}}, \quad (9.5.5)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & e & a+b \end{matrix} \right\} = (-1)^{2a+b+e} \frac{(2a)!(b+e)!}{(2a+2b+1)!(-b+e)!}, \quad (9.5.6)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & e & a+b-1 \end{matrix} \right\} = (-1)^{2a+b+e} \frac{(2a-1)!(b+e-1)!}{(2a+2b)!(-b+e)!} \left[\frac{2a \cdot 2b(b+e)(2a+b-e)(2a+b+e+1)}{(2a+2b+1)(-b+e+1)} \right]^{\frac{1}{2}} \quad (9.5.7)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & b & f \end{matrix} \right\} = (-1)^{2a+2b} \frac{(2a)!(2b)!}{(a+b-f)!(a+b+f+1)!}, \quad (9.5.8)$$

$$\left\{ \begin{matrix} a & b & a+b \\ b & a & f \end{matrix} \right\} = (-1)^{2a+2b} \frac{(2a)!(2b)!}{[(2a-f)!(2a+f+1)!(2b-f)!(2b+f+1)!]^{\frac{1}{2}}}, \quad (9.5.9)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & b & a+b \end{matrix} \right\} = (-1)^{2a+2b} \frac{(2a)!(2b)!}{(2a+2b+1)!}, \quad (9.5.10)$$

$$\left\{ \begin{matrix} a & b & a+b \\ b & b & a+b-1 \end{matrix} \right\} = (-1)^{2a+2b} \frac{(2a)!(2b)!}{(2a+2b)!}, \quad (9.5.11)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & b & a-b \end{matrix} \right\} = (-1)^{2a+2b} \frac{1}{2a+1}, \quad (a \geq b), \quad (9.5.12)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a & b & a-b+1 \end{matrix} \right\} = (-1)^{2a+2b} \frac{2b}{(2a+1)(2a+2)}, \quad (a \geq b-1), \quad (9.5.13)$$

$$\left\{ \begin{matrix} a & b & a+b \\ a+b+e & e & a+e \end{matrix} \right\} = (-1)^{2(a+b+e)} \frac{1}{[(2a+2b+1)(2a+2e+1)]^{\frac{1}{2}}}. \quad (9.5.14)$$

9.5.3 One of Arguments is Smaller by Unity than the Sum of Two Others

If the 6j symbol has one argument which is one less than the sum of two others from the same triad (abc) , (cde) , (aef) , (bdf) , one may use the classical symmetries of the 6j symbol (Eqs. (9.4.2)) to bring it into the form

$$\begin{aligned} & \left\{ \begin{matrix} a & b & a+b-1 \\ d & e & f \end{matrix} \right\} = (-1)^{a+b+d+e} W(abed; a+b-1f) \\ & = (-1)^{a+b+d+e} 2\{ab(a+b) + (a+b)f(f+1) - ad(d+1) - be(e+1)\} \\ & \times \left[\frac{(2a-1)!(2b-1)!(a+b+d+e)!(a+b-d+e-1)!(a+b+d-e-1)!(-a+e+f)!(-b+d+f)!}{(2a+2b)!(-a-b+d+e+1)!(a+e-f)!(a-e+f)!(a+e+f+1)!(b+d-f)!(b-d+f)!(b+d+f+1)!} \right]^{\frac{1}{2}} \end{aligned} \quad (9.5.15)$$

In particular

$$\left\{ \begin{matrix} a & b & a+b-1 \\ d & e & a+e-1 \end{matrix} \right\} = (-1)^{a+b+d+e} 2 \{ a(a+b+e-1)(a+b+e) - ad(d+1) - 2be \} \\ \times \left[\frac{(2b-1)!(2e-1)!(a+b+d-e-1)!(a-b+d+e-1)!}{(2a+2b)!(2a+2e)!(-a-b+d+e+1)!(-a+b+d-e+1)!} \right]^{\frac{1}{2}}, \quad (9.5.16)$$

$$\left\{ \begin{matrix} a & b & a+b-1 \\ a & e & a+b-1 \end{matrix} \right\} = (-1)^{2a+b+e} 2 \{ b(2a+b-1)(2a+b) - be(e+1) - 2a^2 \} \frac{(2a-1)!(b+e-1)!}{(2a+2b)!(-b+e+1)!}, \quad (9.5.17)$$

$$\left\{ \begin{matrix} a & b & a+b-1 \\ a & b & f \end{matrix} \right\} = (-1)^{2a+2b} 2 \{ ab(a+b) + (a+b)f(f+1) - a^2(a+1) \\ - b^2(b+1) \} \frac{(2a-1)!(2b-1)!}{(a+b-f)!(a+b+f+1)!}, \quad (9.5.18)$$

$$\left\{ \begin{matrix} a & b & a+b-1 \\ b & a & f \end{matrix} \right\} = (-1)^{2a+2b} 2 \{ (a+b)f(f+1) - 2ab \} \frac{(2a-1)!(2b-1)!}{[(2a-f)!(2a+f+1)!(2b-f)!(2b+f+1)!]^{\frac{1}{2}}}; \quad (9.5.19)$$

$$\left\{ \begin{matrix} a & b & a+b-1 \\ a+b+e-1 & e & a+e-1 \end{matrix} \right\} = (-1)^{2(a+b+e)} \left[\frac{2b \cdot 2e}{(2a+2b)(2a+2b-1)(2a+2e)(2a+2e-1)} \right]^{\frac{1}{2}}. \quad (9.5.20)$$

Some other cases are given by Eqs. (9.5.7), (9.5.9), (9.5.11).

9.5.4 Arguments a, b, d, e are Equal in Pairs

If $a = b$ and $d = e$ or $a = e$ and $b = d$, the Wigner $6j$ symbol may be rewritten as [72]

$$\left\{ \begin{matrix} a & a & c \\ b & b & f \end{matrix} \right\} = \left\{ \begin{matrix} a & b & f \\ b & a & c \end{matrix} \right\} = (-1)^{2a+2b} W(aabb; cf) = (-1)^{2a+2b} W(abab; fc) \\ = (-1)^{a+b+c+f} \left[\frac{(2a-c)!(2b-c)!}{(2a+c+1)!(2b+c+1)!} \right]^{\frac{1}{2}} V_c(a, f, b) \quad (9.5.21)$$

where c is integer, and $V_c(a, f, b) = V_c(b, f, a)$. According to Eq. (9.6.6), the quantities V_c satisfy the recursion relation

$$V_{c+1} = \frac{2c+1}{c+1} V_1 V_c - c(2c+1) V_c - \frac{c}{c+1} [4a(a+1) + 1 - c^2] [4b(b+1) + 1 - c^2] V_{c-1} \quad (9.5.22)$$

Let us denote

$$\tilde{a} \equiv a(a+1), \quad \tilde{b} \equiv b(b+1), \quad x \equiv f(f+1) - a(a+1) - b(b+1) = \tilde{f} - \tilde{a} - \tilde{b} \quad (9.5.23)$$

Then for some special values of c the functions V_c are given by

$$V_0(a, f, b) = 1, \quad (9.5.24)$$

$$V_1(a, f, b) = -2x, \quad (9.5.25)$$

$$V_2(a, f, b) = 6x^2 + 6x - 8\tilde{a}\tilde{b}, \quad (9.5.26)$$

$$V_3(a, f, b) = -20x^3 - 80x^2 - 16x[3 + \tilde{a} + \tilde{b} - 3\tilde{a}\tilde{b}] + 80\tilde{a}\tilde{b}, \quad (9.5.27)$$

$$V_4(a, f, b) = 70x^4 + 700x^3 + 40x^2[39 + 5\tilde{a} + 5\tilde{b} - 6\tilde{a}\tilde{b}] \\ + 80x[9 + 6\tilde{a} + 6\tilde{b} - 17\tilde{a}\tilde{b}] - 48\tilde{a}\tilde{b}[27 + 4\tilde{a} + 4\tilde{b} - 2\tilde{a}\tilde{b}], \quad (9.5.28)$$

$$V_{2a-1}(a, f, b) = (-1)^{1+a-b+f} 2 \left\{ a^2 + \tilde{f} - \tilde{b} \right\} \frac{(2a)!(2a-1)!(2a+2b)!(-a+b+f)!}{(2b-2a+1)!(a+b-f)!(a-b+f)!(a+b+f+1)!} \quad (9.5.29)$$

$$\left(a \leq b + \frac{1}{2} \right),$$

$$V_{2b-1}(a, f, b) = (-1)^{1-a+b+f} 2 \left\{ b^2 + \tilde{f} - \tilde{a} \right\} \frac{(2b)!(2b-1)!(2a+2b)!(a-b+f)!}{(2a-2b+1)!(a+b-f)!(a-b+f)!(a+b+f+1)!} \quad (9.5.30)$$

$$\left(b \leq a + \frac{1}{2} \right),$$

$$V_{2a}(a, f, b) = (-1)^{a-b+f} \frac{(2a)!(2a)!(2a+2b+1)!(-a+b+f)!}{(2b-2a)!(a+b-f)!(a-b+f)!(a+b+f+1)!} \quad (9.5.31)$$

$$(a \leq b),$$

$$V_{2b}(a, f, b) = (-1)^{-a+b+f} \frac{(2b)!(2b)!(2a+2b+1)!(a-b+f)!}{(2a-2b)!(a+b-f)!(a-b+f)!(a+b+f+1)!} \quad (9.5.32)$$

$$(b \leq a).$$

For special values of f one has

$$V_c(a, a-b, b) = \frac{(2b)!(2a+c+1)!}{(2a+1)!(2b-c)!}, \quad (a \geq b) \quad (9.5.33)$$

$$V_c(a, b-a, b) = \frac{(2a)!(2b+c+1)!}{(2b+1)!(2a-c)!}, \quad (a \leq b) \quad (9.5.34)$$

$$V_c(a, a-b+1, b) = 2\{2b(a+1) - (a-b+1)c(c+1)\} \frac{(2b-1)!(2a+c+1)!}{(2a+2)!(2b-c)!}, \quad (9.5.35)$$

$$(a \geq b-1)$$

$$V_c(a, b-a+1, b) = 2\{2a(b+1) + (a-b-1)c(c+1)\} \frac{(2a-1)!(2b+c+1)!}{(2b+2)!(2a-c)!}, \quad (9.5.36)$$

$$(b \geq a-1),$$

$$V_c(a, a+b-1, b) = (-1)^{c+1} 2\{(a+b)c(c+1) - 2ab\} \frac{(2a-1)!(2b-1)!}{(2a-c)!(2b-c)!}, \quad (9.5.37)$$

$$V_c(a, a+b, b) = (-1)^c \frac{(2a)!(2b)!}{(2a-c)!(2b-c)!} \quad (9.5.38)$$

See also Eqs. (9.5.9) and (9.5.19).

9.6 Recursion Relations

9.6.1 Relations in Which Arguments are Changed by $1/2$

$$\begin{aligned}
& [(a+b+c+1)(-a+b+c)(c+d+e+1)(c+d-e)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \\
& = -2c[(b+d+f+1)(b+d-f)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b-\frac{1}{2} & c-\frac{1}{2} \\ d-\frac{1}{2} & e & f \end{matrix} \right\} \\
& + [(a+b-c+1)(a-b+c)(-c+d+e+1)(c-d+e)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c-1 \\ d & e & f \end{matrix} \right\}, \tag{9.6.1} \\
& (a-b-d+e)[(a+b+c+1)(c+d+e+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}
\end{aligned}$$

$$\begin{aligned}
& = -[(a-b+c)(c-d+e)(a+e-f)(a+e+f+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a-\frac{1}{2} & b & c-\frac{1}{2} \\ d & e-\frac{1}{2} & f \end{matrix} \right\} \\
& + [(-a+b+c)(c+d-e)(b+d-f)(b+d+f+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b-\frac{1}{2} & c-\frac{1}{2} \\ d-\frac{1}{2} & e & f \end{matrix} \right\}, \tag{9.6.2}
\end{aligned}$$

$$\begin{aligned}
& [(-a+b+c)(a-b+c+1)(a+e-f+1)(b+d+f+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}, \\
& = [(c+d-e)(c-d+e+1)(a+e+f+2)(b+d-f)]^{\frac{1}{2}} \left\{ \begin{matrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d-\frac{1}{2} & e+\frac{1}{2} & f \end{matrix} \right\} \\
& + (a-b-d+e+1)[(-a+b+c)(b-d+f)]^{\frac{1}{2}} \left\{ \begin{matrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f-\frac{1}{2} \end{matrix} \right\}, \tag{9.6.3}
\end{aligned}$$

Note. Relation (9.6.3), as given in the original source, does not hold identically: a direct evaluation of both sides shows that they differ for generic physical values of the arguments. For example, at $a = b = \frac{1}{2}$, $c = 1$, $d = \frac{1}{2}$, $e = \frac{3}{2}$, $f = 1$ the left-hand side equals $-\frac{2\sqrt{3}}{3}$ whereas the right-hand side equals $-\frac{\sqrt{6}}{3}$. The equation therefore appears to contain a misprint whose correct form we have been unable to reconstruct; the neighbouring recursions (9.6.1), (9.6.2), (9.6.4) and (9.6.5) have all been verified numerically.

$$\begin{aligned}
& (2d+1)(2f+1)[(a+b+c+1)(a-b+c)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \\
& = -[(a+e+f+1)(a-e+f)(b+d+f+1)(-b+d+f)(c+d+e+1)(c+d-e)]^{\frac{1}{2}} \left\{ \begin{matrix} a-\frac{1}{2} & b & c-\frac{1}{2} \\ d-\frac{1}{2} & e & f-\frac{1}{2} \end{matrix} \right\} \\
& - [(-a+e+f+1)(a+e-f)(b-d+f+1)(b+d-f)(c+d+e+1)(c+d-e)]^{\frac{1}{2}} \left\{ \begin{matrix} a-\frac{1}{2} & b & c-\frac{1}{2} \\ d-\frac{1}{2} & e & f+\frac{1}{2} \end{matrix} \right\} \\
& - [(a+e+f+1)(a-e+f)(b+d-f+1)(b-d+f)(-c+d+e+1)(c-d+e)]^{\frac{1}{2}} \left\{ \begin{matrix} a-\frac{1}{2} & b & c-\frac{1}{2} \\ d+\frac{1}{2} & e & f-\frac{1}{2} \end{matrix} \right\} \\
& + [(-a+e+f+1)(a+e-f)(b+d+f+2)(-b+d+f+1)(-c+d+e+1)(c-d+e)]^{\frac{1}{2}} \\
& \times \left\{ \begin{matrix} a-\frac{1}{2} & b & c-\frac{1}{2} \\ d+\frac{1}{2} & e & f+\frac{1}{2} \end{matrix} \right\}. \tag{9.6.4}
\end{aligned}$$

9.6.2 Relations in Which Arguments are Changed by 1

$$\begin{aligned}
& (2c+1)\{2[a(a+1)d(d+1)+b(b+1)e(e+1)-c(c+1)f(f+1)] \\
& -[a(a+1)+b(b+1)-c(c+1)][d(d+1)+e(e+1)-c(c+1)]\} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \\
& = -c[(a+b+c+2)(-a+b+c+1)(a-b+c+1)(a+b-c) \\
& \times (d+e+c+2)(-d+e+c+1)(d-e+c+1)(d+e-c)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c+1 \\ d & e & f \end{matrix} \right\} \\
& - (c+1)[(a+b+c+1)(-a+b+c)(a-b+c)(a+b-c+1) \\
& \times (d+e+c+1)(-d+e+c)(d-e+c)(d+e-c+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c-1 \\ d & e & f \end{matrix} \right\} \tag{9.6.5}
\end{aligned}$$

In particular

$$\begin{aligned}
& (2c+1)\{-2a(a+1)-2b(b+1)+2f(f+1)+c(c+1)\}\left\{\begin{matrix} a & a & c \\ b & b & f \end{matrix}\right\} \\
& = (c+1)[(2a+c+2)(2a-c)(2b+c+2)(2b-c)]^{\frac{1}{2}}\left\{\begin{matrix} a & a & c+1 \\ b & b & f \end{matrix}\right\} \\
& + c[(2a+c+1)(2a-c+1)(2b+c+1)(2b-c+1)]^{\frac{1}{2}}\left\{\begin{matrix} a & a & c-1 \\ b & b & f \end{matrix}\right\}
\end{aligned} \tag{9.6.6}$$

$$\begin{aligned}
& (2c+1)\{[a(a+1)+b(b+1)-c(c+1)]^2-2[a^2(a+1)^2+b^2(b+1)^2-c(c+1)f(f+1)]\}\left\{\begin{matrix} a & b & c \\ a & b & f \end{matrix}\right\} \\
& = c(a+b+c+2)(-a+b+c+1)(a-b+c+1)(a+b-c)\left\{\begin{matrix} a & b & c+1 \\ a & b & f \end{matrix}\right\} \\
& + (c+1)(a+b+c+1)(-a+b+c)(a-b+c)(a+b-c+1)\left\{\begin{matrix} a & b & c-1 \\ a & b & f \end{matrix}\right\}
\end{aligned} \tag{9.6.7}$$

9.7 Generating Function

The $6j$ symbols turn out to be the coefficients of a power series expansion of the generating function $f(\tau_{i\alpha})$ [73], which depends on 12 variables $\tau_{i\alpha}$ ($i = 1, 2, 3; \alpha = 1, 2, 3, 4$),

$$f(r_{i\alpha}) \equiv \left[1 + \sum_{i=1}^3 \prod_{\alpha=1}^4 r_{i\alpha} + \sum_{\alpha=1}^4 \prod_{i=1}^3 r_{i\alpha}\right]^{-2} = \sum_{R_{i\alpha}} N(R_{i\alpha}) \left\{\begin{matrix} a & b & c \\ d & e & f \end{matrix}\right\} \prod_{i=1}^3 \prod_{\alpha=1}^4 (r_{i\alpha})^{R_{i\alpha}} \tag{9.7.1}$$

The exponents $R_{i\alpha}$ are elements of the R symbol. The relation between $R_{i\alpha}$ and arguments a, b, c , etc. is given by Eqs. (9.1.13)-(9.1.14). The normalization factors N are

$$N(R_{i\alpha}) = \left[\frac{\prod_{\alpha=1}^4 (B_{\alpha} + 1)!}{\prod_{i=1}^3 \prod_{\alpha=1}^4 (R_{i\alpha})!} \right]^{\frac{1}{2}}, \tag{9.7.2}$$

where B_{α} are given by Eqs. (9.1.18).

9.8 Sums Involving the $6j$ Symbols

In this section only the most important sums involving products of the $6j$ symbols are presented. We shall use the notation $\{abc\}$ for the symbol which is equal to 1 if a, b, c satisfy the triangular conditions (8.1.1) and is zero otherwise. In the equations below the sum is over all possible values (integer or half-integer) of X which

obey all triangular conditions.

$$\sum_X (2X+1) \left\{ \begin{matrix} a & b & X \\ a & b & c \end{matrix} \right\} = (-1)^{2c} \{abc\}, \quad (9.8.1)$$

$$\sum_X (-1)^{a+b+X} (2X+1) \left\{ \begin{matrix} a & b & X \\ b & a & c \end{matrix} \right\} = \delta_{c0} \sqrt{(2a+1)(2b+1)}, \quad (9.8.2)$$

$$\sum_X (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ c & d & q \end{matrix} \right\} = \delta_{pq} \frac{\{adp\}\{bcp\}}{2p+1}, \quad (9.8.3)$$

$$\sum_X (-1)^{p+q+X} (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ d & c & q \end{matrix} \right\} = \left\{ \begin{matrix} a & c & q \\ b & d & p \end{matrix} \right\}, \quad (9.8.4)$$

$$\sum_X (-1)^{2X} (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ a & b & r \end{matrix} \right\} = \left\{ \begin{matrix} a & f & r \\ d & q & e \\ p & c & b \end{matrix} \right\}, \quad (9.8.5)$$

$$\sum_X (-1)^{R+X} (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ b & a & r \end{matrix} \right\} = \left\{ \begin{matrix} p & q & r \\ e & a & d \end{matrix} \right\} \left\{ \begin{matrix} p & q & r \\ f & b & c \end{matrix} \right\}, \quad (9.8.6)$$

$$(R = a + b + c + d + e + f + p + q + r)$$

$$\sum_X (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ g & h & r \end{matrix} \right\} \left\{ \begin{matrix} g & h & X \\ a & b & s \end{matrix} \right\} = (-1)^{p-s-q+r} \left\{ \begin{matrix} - & h & a & s \\ c & - & p & b \\ f & r & - & g \\ q & e & d & - \end{matrix} \right\}, \quad (9.8.7)$$

$$\sum_X (-1)^{T-X} (2X+1) \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ g & h & r \end{matrix} \right\} \left\{ \begin{matrix} g & h & X \\ b & a & s \end{matrix} \right\} = \left\{ \begin{matrix} a & d & e & h \\ p & q & r & s \\ b & c & f & g \end{matrix} \right\}, \quad (9.8.8)$$

$$(T = a + b + c + d + e + f + g + h + p + q + r + s).$$

Many other sums, involving the $6j$ symbols as well as the $3jm$ symbols and the $9j$ symbols, will be given in Chap. 12.

9.9 Asymptotics of the $6j$ Symbols for Large Angular Momenta

9.9.1 Asymptotic Relations Between the $6j$ Symbols and the Clebsch-Gordan Coefficients

If $R \gg 1$ and a, b, c etc. are arbitrary, one gets the following asymptotic relation

$$\left\{ \begin{matrix} a & b & c \\ d+R & e+R & f+R \end{matrix} \right\} \approx \frac{(-1)^{a+b+d+e}}{\sqrt{2R(2c+1)}} C_{a\alpha b\beta}^{c\gamma} \quad (9.9.1)$$

where $\alpha = f - e, \beta = d - f, \gamma = d - e$.

For the $6j$ symbols and the $3jm$ symbols this relation assumes the form [74]

$$\left\{ \begin{matrix} a & b & c \\ d+R & e+R & f+R \end{matrix} \right\} \approx \frac{(-1)^{a+b+c+2(d+e+f)}}{\sqrt{2R}} \left(\begin{matrix} a & b & c \\ e-f & f-d & d-e \end{matrix} \right). \quad (9.9.2)$$

The asymptotic relation between the R symbols which correspond to the $6j$ symbols and the Clebsch-Gordan coefficients is written as

$$\begin{aligned} & \left\| \begin{matrix} -c+d+e+2R & b+d-f & a+e-f & a+b-c \\ -b+d+f+2R & c+d-e & a-b+c & a-e+f \\ -a+e+f+2R & -a+b+c & c-d+e & b-d+f \end{matrix} \right\| \\ & \approx \frac{(-1)^{a+b+c+2(d+e+f)}}{\sqrt{2R}} \left\| \begin{matrix} -a+b+c & a-b+c & a+b-c \\ a+e-f & b-d+f & c+d-e \\ a-e+f & b+d-f & c-d+e \end{matrix} \right\|. \end{aligned} \quad (9.9.3)$$

In particular, when $d = e = f = 0$ one obtains [66]

$$\left\{ \begin{matrix} a & b & c \\ R & R & R \end{matrix} \right\} \approx \frac{(-1)^c}{\sqrt{2R(2c+1)}} C_{a0b0}^{c0} \quad (9.9.4)$$

9.9.2 Asymptotic Expressions for the $6j$ Symbols

The asymptotic behaviour of the $6j$ symbols $\begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix}$ for large angular momenta is closely associated with geometric properties of the tetrahedron whose edges are $a + \frac{1}{2}, b + \frac{1}{2}, c + \frac{1}{2}, d + \frac{1}{2}, e + \frac{1}{2}, f + \frac{1}{2}$, etc. (Fig. 9.1).




Figure 9.1. The tetrahedron associated with asymptotic behaviour of the $6j$ symbols.

(a) *The Ponzano-Regge Formula [46] (semiclassical approximation to the $6j$ symbols)*

If $a, b, c, d, e, f \gg 1$, then

$$\begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix} \approx \frac{1}{\sqrt{12\pi V}} \cos \left(\sum_{i,k=1}^4 j_{ik} \Theta_{ik} + \frac{\pi}{4} \right), \quad (V^2 > 0) \quad (9.9.5)$$

Here

$$\begin{aligned} j_{12} &= a + \frac{1}{2}, & j_{13} &= b + \frac{1}{2}, & j_{14} &= c + \frac{1}{2}, \\ j_{23} &= f + \frac{1}{2}, & j_{24} &= e + \frac{1}{2}, & j_{34} &= d + \frac{1}{2}, \\ j_{ik} &= j_{ki}, & j_{ii} &= 0. \end{aligned} \quad (9.9.6)$$

V is the volume of the tetrahedron, Θ_{ik} is the angle between two external normals to the planes adjacent to the edge j_{ik} .

The tetrahedron volume is equal to

$$V^2 = \frac{1}{2^3(3!)^2} \begin{vmatrix} 0 & j_{34}^2 & j_{24}^2 & j_{23}^2 & 1 \\ j_{34}^2 & 0 & j_{14}^2 & j_{13}^2 & 1 \\ j_{24}^2 & j_{14}^2 & 0 & j_{12}^2 & 1 \\ j_{23}^2 & j_{13}^2 & j_{12}^2 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{vmatrix} \quad (9.9.7)$$

The angles Θ_{ik} are given by

$$S_i S_k \sin \Theta_{ik} = \frac{3}{2} V j_{ik}, \quad (i \neq k) \quad (9.9.8)$$

Here S_i is the area of the triangle opposite to the vertex p_i (Fig. 9.1). One can evaluate S_i , using the standard formulas. For example,

$$S_1^2 = \frac{1}{16} (j_{12} + j_{13} + j_{14}) (j_{12} + j_{13} - j_{14}) (j_{12} - j_{13} + j_{14}) (-j_{12} + j_{13} + j_{14}) = -\frac{1}{16} \begin{vmatrix} 0 & j_{12}^2 & j_{13}^2 & 1 \\ j_{12}^2 & 0 & j_{14}^2 & 1 \\ j_{13}^2 & j_{14}^2 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{vmatrix}. \quad (9.9.9)$$

The asymptotic expressions (9.9.5) are valid only if $V^2 > 0$ (classically allowed domain). If $V^2 < 0$ (classically forbidden domain, when an associated tetrahedron does not exist), the asymptotic expression becomes

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \approx \frac{1}{2\sqrt{12\pi|V|}} \cos \Phi \exp \left\{ - \left| \sum_{i,k=1}^4 j_{ik} \operatorname{Im} \Theta_{ik} \right| \right\}, \quad (V^2 < 0) \quad (9.9.10)$$

where

$$\Phi = \sum_{i,k=1}^4 \left(j_{ik} - \frac{1}{2} \right) \operatorname{Re} \Theta_{ik} \quad (9.9.11)$$

In this case the $6j$ symbols are exponentially small even if the triangular condition is satisfied.

Near the classical domain boundary, where $V^2 \approx 0$, Eqs. (9.9.5), (9.9.10) are of little use, although one may use the improved expressions [46]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \approx 2^{-\frac{4}{3}} \left(\prod_{i=1}^4 S_i \right)^{-\frac{1}{6}} \{ \cos \Phi \operatorname{Ai}(z) + \sin \Phi \operatorname{Bi}(z) \} \quad (9.9.12)$$

$\operatorname{Ai}(z)$ and $\operatorname{Bi}(z)$ being the Airy functions [25],

$$z = \begin{cases} -(3V)^2 \left(4 \prod_{i=1}^4 S_i \right)^{-\frac{2}{3}} & \text{if } V^2 > 0, \\ (3|V|)^2 \left(4 \prod_{i=1}^4 S_i \right)^{-\frac{2}{3}} & \text{if } V^2 < 0. \end{cases} \quad (9.9.13)$$

Note that the Ponzano-Regge approximation is sufficiently accurate even at comparatively small angular momenta a, b, c etc. Asymptotic expressions similar to (9.9.12) but extended over the entire domain of angular momenta are obtained in Ref. [139].

(b) *The Edmonds' Formula* [33]

If f, m, n are arbitrary integers or half-integers and $a, b, c \gg f, m, n$, then

$$\left\{ \begin{matrix} a & b & c \\ b+m & a+n & f \end{matrix} \right\} \approx \frac{(-1)^{a+b+c+f+m}}{\sqrt{(2a+1)(2b+1)}} d_{mn}^f(\theta), \quad (9.9.14)$$

where $d_{mn}^f(\Theta)$ is the rotation matrix (Chap. 4), Θ is an angle between the tetrahedron edges $a+n+\frac{1}{2}$ and $b+m+\frac{1}{2}$ (Fig. 9.2)

$$\cos \theta = \frac{a(a+1) + b(b+1) - c(c+1)}{2\sqrt{a(a+1)b(b+1)}}. \quad (9.9.15)$$




Figure 9.2. The angle Θ which enters Eq. (9.9.14).

In particular, if $m = n = 0, a, b, c \gg f$ and f is an arbitrary integer, then (9.9.14) turns into the Racah formula [75],

$$\left\{ \begin{matrix} a & b & c \\ b & a & f \end{matrix} \right\} \approx \frac{(-1)^{a+b+c+f}}{\sqrt{(2a+1)(2b+1)}} P_f(\cos \theta), \quad (9.9.16)$$

where P_f is the Legendre polynomial. If, in addition, f is large ($a, b, c \gg f \gg 1$), one can substitute into (9.9.16) asymptotic expressions for the Legendre polynomials to obtain

$$\left\{ \begin{matrix} a & b & c \\ b & a & f \end{matrix} \right\} \approx (-1)^{a+b+c+f} \left[\frac{4}{\pi(2a+1)(2b+1)(2f+1)\sin\theta} \right]^{\frac{1}{3}} \cos \left[\left(f + \frac{1}{2} \right) \theta - \frac{\pi}{4} \right]. \quad (9.9.17)$$

This expression may also be written in the form (cf. Eq. (9.9.5))

$$\left\{ \begin{matrix} a & b & c \\ b & a & f \end{matrix} \right\} \approx \frac{(-1)^{a+b+c+f}}{\sqrt{12\pi V}} \cos \left[\left(f + \frac{1}{2} \right) \Theta - \frac{\pi}{4} \right], \quad (9.9.18)$$

where the tetrahedron volume is given by

$$V = \frac{1}{6} \left(a + \frac{1}{2} \right) \left(b + \frac{1}{2} \right) \left(f + \frac{1}{2} \right) \sin \Theta. \quad (9.9.19)$$

(c) $a, b, c, d, e, f \gg 1, m, n, p$

In this case, one has

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} \left\{ \begin{matrix} a & b & c \\ d+m & e+n & f+p \end{matrix} \right\} \approx \frac{\Theta(V^2)}{24\pi V} \cos(m\Theta_1 + n\Theta_2 + p\Theta_3). \quad (9.9.20)$$

where V is the tetrahedron volume (9.9.7)

$$\Theta(V^2) = \begin{cases} 1, & \text{if } V^2 > 0, \\ 0, & \text{if } V^2 < 0, \end{cases} \quad (9.9.21)$$

Θ_i is the angle between two external normals to the planes adjacent to the edge $l_i + \frac{1}{2}$ (Fig. 9.3), with $l_1 \equiv d, l_2 \equiv e, l_3 \equiv f$. The angles Θ_i can be evaluated from

$$\cos \Theta_i = \frac{\cos \varphi_{ik} \cos \varphi_{il} - \cos \varphi_{kl}}{\sin \varphi_{ik} \sin \varphi_{il}} \quad (9.9.22)$$

where combinations i, k, l are obtained by cyclic permutations of 1, 2, 3.




Figure 9.3. Geometrical interpretation of the angles in Eqs. (9.9.20)-(9.9.23).

In Eq. (9.9.22) $\varphi_{ik} \equiv \varphi_{ki}$ is the angle between the edges $l_i + \frac{1}{2}$ and $l_k + \frac{1}{2}$ of the tetrahedron (Fig. 9.3)

$$\cos \varphi_{ik} = \frac{l_i(l_i + 1) + l_k(l_k + 1) - j_l(j_l + 1)}{2\sqrt{l_i(l_i + 1)l_k(l_k + 1)}} \quad (9.9.23)$$

with $i \neq k \neq l$ and $j_1 \equiv a, j_2 \equiv b, j_3 \equiv c$.

(d) $m = n = p = 0$

In particular, for $m = n = p = 0$ Eq. (9.9.20) yields the Wigner formula [15]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}^2 \approx \frac{\Theta(V^2)}{24\pi V}, \quad (a, b, c, d, e, f \gg 1) \quad (9.9.24)$$

This formula is valid only on the average because the $6j$ symbols oscillate rapidly with momentum variations in the region of large angular momenta.

(e) $R \rightarrow \infty$

If a, b, c , etc. are fixed and $R \rightarrow \infty$, one has [46]

$$\begin{aligned} \left\{ \begin{matrix} a & b+R & c+R \\ d & e+R & f+R \end{matrix} \right\} &\approx (-1)^\varphi \left[\frac{(a-b+c)!(a-e+f)!(c+d-e)!(-b+d+f)!}{(a+b-c)!(a+e-f)!(-c+d+e)!(b+d-f)!} \right]^{\frac{1}{2} \text{sign}(c+f-b-e)} \\ &\times \frac{(2R)^{-1-|b+e-c-f|}}{|b+e-c-f|!} \left[1 + O\left(\frac{1}{R^2}\right) \right]. \end{aligned} \quad (9.9.25)$$

In this case

$$\begin{aligned} \varphi &= a + d + \min\{b + e, c + f\}, \\ \text{sign } x &= \begin{cases} 1, & \text{if } x \geq 0 \\ -1, & \text{if } x < 0 \end{cases} \end{aligned} \quad (9.9.26)$$

9.10 Relations Between the Wigner $6j$ Symbols and Analogous Functions of Other Authors

Racah [56]:

$$W(abed; cf) = (-1)^{a+b+d+e} \left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}$$

Jahn [76]:

$$U(abed; cf) = [(2c+1)(2f+1)]^{\frac{1}{2}} W(abed; cf)$$

Biedenharn, Blatt and Rose [72]:

$$Z(abed; cf) = i^{-a+e+f} [(2a+1)(2b+1)(2d+1)(2e+1)]^{\frac{1}{2}} C_{a0e0}^{f0} W(abed; cf)$$

9.11 Tables of Algebraic Expressions for the $6j$ Symbols

Algebraic expressions for the $6j$ symbols $\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\}$ with $d = \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}, 3, \frac{7}{2}, 4$ are presented in Tables 9.1-9.8.

We use the following notations

$$\begin{aligned} s &= a + b + c \\ X &= -a(a+1) + b(b+1) + c(c+1) \end{aligned}$$

The tables of algebraic formulas for the $6j$ symbols and Racah coefficients are also given in Refs. [10, 20, 72].

Chapter 10

9j and 12j Symbols

10.1 Definition of the 9j Symbols

10.1.1 9j Symbols as Recoupling Coefficients

The Wigner 9j symbols [54] (or Fano coefficients [68]) are associated with the coefficients of unitary transformations which connect state vectors corresponding to different coupling schemes of four angular momenta.

Let us consider an addition of four angular momenta $\mathbf{j}_1, \mathbf{j}_2, \mathbf{j}_3$ and $\mathbf{j}_4$ to form a resultant angular momentum j with projection m . There exist different coupling schemes of these angular momenta. We pay attention to the following schemes¹

$$\begin{aligned} \text{I) } & \mathbf{j}_1 + \mathbf{j}_2 = \mathbf{j}_{12}, \quad \mathbf{j}_3 + \mathbf{j}_4 = \mathbf{j}_{34}, \quad \mathbf{j}_{12} + \mathbf{j}_{34} = \mathbf{j}; \\ \text{II) } & \mathbf{j}_1 + \mathbf{j}_3 = \mathbf{j}_{13}, \quad \mathbf{j}_2 + \mathbf{j}_4 = \mathbf{j}_{24}, \quad \mathbf{j}_{13} + \mathbf{j}_{24} = \mathbf{j}; \\ \text{III) } & \mathbf{j}_1 + \mathbf{j}_4 = \mathbf{j}_{14}, \quad \mathbf{j}_2 + \mathbf{j}_3 = \mathbf{j}_{23}, \quad \mathbf{j}_{14} + \mathbf{j}_{23} = \mathbf{j}; \end{aligned} \quad (10.1.1)$$

Let $|j_1 j_2 (j_{12}) j_3 j_4 (j_{34}) j m\rangle$ be a state vector which corresponds to scheme I. This vector is an eigenvector of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_4^2, \hat{j}_{12}^2, \hat{j}_{34}^2, \hat{j}^2$ and $\hat{j}_x$ and may be written as

$$|j_1 j_2 (j_{12}) j_3 j_4 (j_{34}) j m\rangle = \sum_{\substack{m_1 m_2 m_3 m_4 \\ m_{12} m_{34}}} C_{j_{12} m_{12} j_{34} m_{34}}^{j m} C_{j_1 m_1 j_2 m_2}^{j_{12} m_{12}} C_{j_3 m_3 j_4 m_4}^{j_{34} m_{34}} |j_1 m_1, j_2 m_2, j_3 m_3, j_4 m_4\rangle. \quad (10.1.2)$$

Here $|j_1 m_1, j_2 m_2, j_3 m_3, j_4 m_4\rangle$ is eigenvector of the operators $\hat{j}_i^2$ and $\hat{j}_{ix} (i = 1, 2, 3, 4)$. A state vector which corresponds to scheme II is the eigenvector of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_4^2, \hat{j}_{13}^2, \hat{j}_{24}^2, \hat{j}^2$ and $\hat{j}_z$. It can be represented in the form

$$|j_1 j_3 (j_{13}) j_2 j_4 (j_{24}) j m\rangle = \sum_{\substack{m_1 m_2 m_3 m_4 \\ m_{13} m_{24}}} C_{j_{13} m_{13} j_{24} m_{24}}^{j m} C_{j_1 m_1 j_3 m_3}^{j_{13} m_{13}} C_{j_2 m_2 j_4 m_4}^{j_{24} m_{24}} |j_1 m_1, j_2 m_2, j_3 m_3, j_4 m_4\rangle \quad (10.1.3)$$

Similarly, a state vector, which corresponds to scheme III, is the eigenvector of the operators $\hat{j}_1^2, \hat{j}_2^2, \hat{j}_3^2, \hat{j}_4^2, \hat{j}_{14}^2, \hat{j}_{23}^2, \hat{j}^2$ and $\hat{j}_x$. This state vector is given by

$$|j_1 j_4 (j_{14}) j_2 j_3 (j_{23}) j m\rangle = \sum_{\substack{m_1 m_2 m_3 m_4 \\ m_{14} m_{23}}} C_{j_{14} m_{14} j_{23} m_{23}}^{j m} C_{j_1 m_1 j_4 m_4}^{j_{14} m_{14}} C_{j_2 m_2 j_3 m_3}^{j_{23} m_{23}} |j_1 m_1, j_2 m_2, j_3 m_3, j_4 m_4\rangle \quad (10.1.4)$$

The state vectors associated with each of these coupling schemes form a complete set of functions. Any transformation which connects the state vectors in different coupling schemes, is determined by a unitary matrix. Such a matrix relates the state vectors with the same resultant angular momentum j and its projection m . It may be shown that the transformation matrix is independent of m . The Wigner 9j symbols are proportional to the matrix elements (transformation coefficient). Thus, the Wigner 9j symbols (or, equivalently, the Fano coefficients)

$$\left\{ \begin{matrix} j_1 & j_2 & j_{12} \\ j_3 & j_4 & j_{34} \\ j_{13} & j_{24} & j \end{matrix} \right\}$$

¹Along with the coupling schemes (10.1.1) there exist some other ones, e.g., $\mathbf{j}_1 + \mathbf{j}_2 = \mathbf{j}_{12}, \mathbf{j}_{12} + \mathbf{j}_3 = \mathbf{j}_{123}, \mathbf{j}_{123} + \mathbf{j}_4 = \mathbf{j}$. However, the coefficients of transformation which connects a state vector in this coupling scheme with vectors in coupling schemes, I, II or III are related to the products of two 6j symbols rather than to the 9j symbols.

are related to the coefficients of the transformation between state vectors in schemes I and II by

$$\begin{aligned} & \langle j_1 j_2 (j_{12}) j_3 j_4 (j_{34}) j m | j_1 j_3 (j_{13}) j_2 j_4 (j_{24}) j' m' \rangle \\ &= \delta_{jj'} \delta_{mm'} [(2j_{12} + 1) (2j_{13} + 1) (2j_{24} + 1) (2j_{34} + 1)]^{\frac{1}{2}} \begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j_4 & j_{34} \\ j_{13} & j_{24} & j \end{Bmatrix} \end{aligned} \quad (10.1.5)$$

Using this definition, we may also obtain

$$\begin{aligned} & \langle j_1 j_2 (j_{12}) j_3 j_4 (j_{34}) j m | j_1 j_4 (j_{14}) j_2 j_3 (j_{23}) j' m' \rangle \\ &= \delta_{jj'} \delta_{mm'} (-1)^{j_3 + j_4 - j_{34}} [(2j_{12} + 1) (2j_{14} + 1) (2j_{23} + 1) (2j_{34} + 1)]^{\frac{1}{2}} \begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_4 & j_3 & j_{34} \\ j_{14} & j_{23} & j \end{Bmatrix}, \end{aligned} \quad (10.1.6)$$

$$\begin{aligned} & \langle j_1 j_3 (j_{13}) j_2 j_4 (j_{24}) j m | j_1 j_4 (j_{14}) j_2 j_3 (j_{23}) j' m' \rangle \\ &= \delta_{jj'} \delta_{mm'} (-1)^{j_3 - j_4 - j_{23} + j_{24}} [(2j_{13} + 1) (2j_{14} + 1) (2j_{24} + 1) (2j_{23} + 1)]^{\frac{1}{2}} \begin{Bmatrix} j_1 & j_3 & j_{13} \\ j_4 & j_2 & j_{24} \\ j_{14} & j_{23} & j \end{Bmatrix}. \end{aligned} \quad (10.1.7)$$

As follows from definition (10.1.5) a 9j symbol can be represented as a sum of products of the Clebsch-Gordan coefficients:

$$\begin{aligned} & \sum_{\substack{m_1 m_2 m_3 m_4 \\ m_{12} m_{34} m_{13} m_{24}}} C_{j_1 m_1 j_2 m_2}^{j_{12} m_{12}} C_{j_3 m_3 j_4 m_4}^{j_{34} m_{34}} C_{j_{12} m_{12} j_{34} m_{34}}^{j m} C_{j_1 m_1 j_3 m_3}^{j_{13} m_{13}} C_{j_2 m_2 j_4 m_4}^{j_{24} m_{24}} C_{j_{13} m_{13} j_{24} m_{24}}^{j' m'} \\ &= \delta_{jj'} \delta_{mm'} [(2j_{12} + 1) (2j_{13} + 1) (2j_{24} + 1) (2j_{34} + 1)]^{\frac{1}{2}} \begin{Bmatrix} j_1 & j_2 & j_{12} \\ j_3 & j_4 & j_{34} \\ j_{13} & j_{24} & j \end{Bmatrix} \end{aligned} \quad (10.1.8)$$

This equation unambiguously fixes the absolute value and phase of the 9j symbol. The 9j symbols are real.

Below we shall also use Latin letters (a, b, c , etc.) to denote arguments of the 9j symbols.

In accordance with the quantum mechanical rules of angular momentum addition, arguments of the 9j symbol

$$\begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}$$

satisfy the following conditions.

- (a) Arguments $a, b, c, \dots, j$ are integer or half-integer non-negative numbers (generalization of the 9j symbols to the case of negative arguments will be considered in Sec. 10.4.3).
- (b) A 9j symbol vanishes unless the triangular conditions (see Sec. 8.1.1) are fulfilled for triads (abc) , (def) , (ghj) , (adg) , (beh) and (cfj) .

The 9j symbols satisfy the orthogonality and normalization relations

$$\sum_{gh} (2g + 1)(2h + 1) \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \begin{Bmatrix} a & b & c' \\ d & e & f' \\ g & h & j \end{Bmatrix} = \delta_{cc'} \delta_{ff'} \{abc\} \{def\} \{cfj\} \frac{1}{(2c + 1)(2f + 1)}, \quad (10.1.9)$$

$$\sum_{cf} (2c + 1)(2f + 1) \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \begin{Bmatrix} a & b & c \\ d & e & f \\ g' & h' & j \end{Bmatrix} = \delta_{gg'} \delta_{hh'} \{adg\} \{beh\} \{ghj\} \frac{1}{(2g + 1)(2h + 1)}. \quad (10.1.10)$$

These relations follow from the unitarity of transformations which relate state vectors in different coupling schemes.

10.1.2 9j Symbol and r Symbol

A 9j symbol may be represented as a 3×6 array $\|r_{ik}, r'_{ik}\|$ (Shelepin [70], Wu [77]) which is called the r -symbol:

$$\begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \equiv \left\| \begin{array}{cccccc} r_{11} & r_{12} & r_{13} & r'_{11} & r'_{12} & r'_{13} \\ r_{21} & r_{22} & r_{23} & r'_{21} & r'_{22} & r'_{23} \\ r_{31} & r_{32} & r_{33} & r'_{31} & r'_{32} & r'_{33} \end{array} \right\|, \quad (10.1.11)$$

²The array $\|r_{ik}, r'_{ik}\|$ under discussion (Wu [77]) differs from the r -symbol introduced by Shelepin [70].

where

$$\begin{aligned}
 r_{11} &= -a + b + c, & r_{12} &= a - b + c, & r_{13} &= a + b - c, \\
 r_{21} &= -d + e + f, & r_{22} &= d - e + f, & r_{23} &= d + e - f, \\
 r_{31} &= -g + h + j, & r_{32} &= g - h + j, & r_{33} &= g + h - j, \\
 r'_{11} &= -a + d + g, & r'_{12} &= -b + e + h, & r'_{13} &= -c + f + j, \\
 r'_{21} &= a - d + g, & r'_{22} &= b - e + h, & r'_{23} &= c - f + j, \\
 r'_{31} &= a + d - g, & r'_{32} &= b + e - h, & r'_{33} &= c + f - j.
 \end{aligned} \tag{10.1.12}$$

The inverse relations are as follows

$$\begin{aligned}
 2a &= r_{12} + r_{13} = r'_{21} + r'_{31}, & 2d &= r_{22} + r_{23} = r'_{11} + r'_{31}, & 2g &= r_{32} + r_{33} = r'_{11} + r'_{21}, \\
 2b &= r_{11} + r_{13} = r'_{22} + r'_{32}, & 2e &= r_{21} + r_{23} = r'_{12} + r'_{32}, & 2h &= r_{31} + r_{33} = r'_{12} + r'_{22}, \\
 2c &= r_{11} + r_{12} = r'_{23} + r'_{33}, & 2f &= r_{21} + r_{22} = r'_{13} + r'_{33}, & 2j &= r_{31} + r_{32} = r'_{13} + r'_{23}.
 \end{aligned} \tag{10.1.13}$$

All eighteen elements r_{ik}, r'_{ik} are integer and non-negative. Only nine of them are linearly independent. The elements r_{ik} and r'_{ik} satisfy the following relations

$$\begin{aligned}
 \sum_{i=1}^3 r_{1i} &= a + b + c, & \sum_{i=1}^3 r_{2i} &= d + e + f, & \sum_{i=1}^3 r_{3i} &= g + h + j, \\
 \sum_{i=1}^3 r_{i1} &= R - 2(a + d + g), & \sum_{i=1}^3 r_{i2} &= R - 2(b + e + h), & \sum_{i=1}^3 r_{i3} &= R - 2(c + f + j), \\
 \sum_{i=1}^3 r'_{1i} &= R - 2(a + b + c), & \sum_{i=1}^3 r'_{2i} &= R - 2(d + e + f), & \sum_{i=1}^3 r'_{3i} &= R - 2(g + h + j), \\
 \sum_{i=1}^3 r'_{i1} &= a + d + g, & \sum_{i=1}^3 r'_{i2} &= b + e + h, & \sum_{i=1}^3 r'_{i3} &= c + f + j, \\
 \sum_{i,k=1}^3 r_{ik} &= R, & \sum_{i,k=1}^3 r'_{ik} &= R,
 \end{aligned} \tag{10.1.14}$$

where

$$R = a + b + c + d + e + f + g + h + j. \tag{10.1.15}$$

10.2 Explicit Forms of the $9j$ Symbols and Their Relations to Other Functions

The expressions given below represent the $9j$ symbols in the forms of algebraic sums and sums involving products of the Clebsch-Gordan coefficients, $3jm$ and $6j$ symbols. Unlike the $3jm$ and $6j$ symbols the expressions for the $9j$ symbols in terms of the generalized hypergeometric functions are still unknown. Convenient expressions for the $9j$ symbols in terms of quasi-binomials can be obtained only in some special cases.

10.2.1 Expressions for the $9j$ Symbols in the Forms of Algebraic Sums

$$\begin{aligned}
 & \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = \frac{\Delta(abc)\Delta(def)\Delta(beh)\Delta(ghj)}{\Delta(adg)\Delta(cfj)} \\
 & \times \frac{(a + d - g)!(c + f - j)!(g + h + j + 1)!}{(a + d + g + 1)!(a - b + c)!(-a + b + c)!(d - e + f)!(-d + e + f)!(b - e + h)!(-b + e + h)!} \\
 & \times \sum_{xyzt} (-1)^{a-c+e-g+j+x+y+z+t} \frac{(2a-x)!(2b-y)!(2d-z)!(2e-t)!}{x!y!z!t!} \\
 & \times \frac{(-a + b + c + x)!(-b + e + h + y)!(-d + e + f + z)!(b - e + g - j + t)!(-a - e + f + g + x + t)!}{(a + b - c - x)!(b + e - h - y)!(d + e - f - z)!(b + e - g + j - t)!(a + d - g - x - z)!(e - b + h + y - t)!} \\
 & \times \frac{(c - d + e + j + z - t)!}{(-d + e + f + z - t)!(b - e + g - j - y + t)!(-a + c - e + g - j + x + t)!(-a + c - d + f + g + j + 1 + x + z)!}.
 \end{aligned} \tag{10.2.1}$$

In Eq. (10.2.1) the sums are over all integer values of x, y, z, t for which the factorial arguments are greater than or equal to zero. An expression for the $9j$ symbols in the form of triple sum was proposed by Alisauskas and Yutsis [140].

10.2.2 Wu Formulas [77]

$$\left\{ \begin{matrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{matrix} \right\} = \Delta(j_{11}j_{12}j_{13}) \Delta(j_{21}j_{22}j_{23}) \Delta(j_{31}j_{32}j_{33}) \Delta(j_{11}j_{21}j_{31}) \\ \times \Delta(j_{12}j_{22}j_{32}) \Delta(j_{13}j_{23}j_{33}) \sum_{\nu, \omega} (-1)^{\omega_4 + \omega_8 + \omega_6} \frac{(n+1)!}{\prod_{p,q=1}^3 \nu_{pq}! \prod_{\alpha=1}^6 \omega_{\alpha}!} \quad (10.2.2)$$

where

$$n = \sum_{p,q=1}^3 \nu_{pq} + \sum_{\alpha=1}^6 \omega_{\alpha} \quad (10.2.3)$$

In Eq. (10.2.3) the summation is over 15 integer and non-negative variables $\nu_{pq}(p, q = 1, 2, 3), \omega_{\alpha}(\alpha = 1, 2, \dots, 6)$. These variables satisfy the following conditions owing to which only six variables are independent:

$$\begin{aligned} \omega_1 + \omega_4 + \nu_{21} + \nu_{31} &= r_{11}, & \omega_1 + \omega_4 + \nu_{12} + \nu_{13} &= r'_{11} \\ \omega_3 + \omega_6 + \nu_{22} + \nu_{32} &= r_{12}, & \omega_3 + \omega_6 + \nu_{11} + \nu_{13} &= r'_{12} \\ \omega_2 + \omega_5 + \nu_{23} + \nu_{33} &= r_{13}, & \omega_2 + \omega_5 + \nu_{11} + \nu_{12} &= r'_{13} \\ \omega_3 + \omega_5 + \nu_{11} + \nu_{31} &= r_{21}, & \omega_3 + \omega_5 + \nu_{22} + \nu_{23} &= r'_{21} \\ \omega_2 + \omega_4 + \nu_{12} + \nu_{32} &= r_{22}, & \omega_2 + \omega_4 + \nu_{21} + \nu_{23} &= r'_{22} \\ \omega_1 + \omega_6 + \nu_{13} + \nu_{33} &= r_{23}, & \omega_1 + \omega_6 + \nu_{21} + \nu_{22} &= r'_{23} \\ \omega_2 + \omega_6 + \nu_{11} + \nu_{21} &= r_{31}, & \omega_2 + \omega_6 + \nu_{32} + \nu_{33} &= r'_{31} \\ \omega_1 + \omega_5 + \nu_{12} + \nu_{22} &= r_{32}, & \omega_1 + \omega_5 + \nu_{31} + \nu_{33} &= r'_{32} \\ \omega_3 + \omega_4 + \nu_{13} + \nu_{23} &= r_{33}, & \omega_3 + \omega_4 + \nu_{31} + \nu_{32} &= r'_{33} \end{aligned} \quad (10.2.4)$$

ν_{pq} and ω_{α} are related to arguments of the $9j$ symbols by

$$\sum_{q=1}^3 \nu_{pq} = n - \sum_{q=1}^3 r_{pq}, \quad \sum_{q=1}^3 \nu_{qp} = n - \sum_{q=1}^3 r'_{qp}. \quad (10.2.5)$$

$$\begin{aligned} n + \omega_1 + \omega_4 - \nu_{11} &= j_{12} + j_{13} + j_{21} + j_{31}, & n + \omega_3 + \omega_5 - \nu_{21} &= j_{22} + j_{23} + j_{11} + j_{31}, \\ n + \omega_3 + \omega_6 - \nu_{12} &= j_{11} + j_{13} + j_{22} + j_{32}, & n + \omega_2 + \omega_4 - \nu_{22} &= j_{21} + j_{23} + j_{12} + j_{32}, \\ n + \omega_2 + \omega_5 - \nu_{13} &= j_{11} + j_{12} + j_{23} + j_{33}, & n + \omega_1 + \omega_6 - \nu_{23} &= j_{21} + j_{22} + j_{13} + j_{33}, \\ n + \omega_2 + \omega_6 - \nu_{31} &= j_{32} + j_{33} + j_{11} + j_{21}, \\ n + \omega_1 + \omega_5 - \nu_{32} &= j_{31} + j_{33} + j_{12} + j_{22}, \\ n + \omega_3 + \omega_4 - \nu_{33} &= j_{31} + j_{32} + j_{13} + j_{23}. \end{aligned} \quad (10.2.6)$$

A choice of independent summation variables can be made in many ways, leading to formally different representations of the $9j$ symbols. Some of them are given below.

(a)

$$\left\{ \begin{matrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{matrix} \right\} = \Delta(j_{11}j_{12}j_{13}) \Delta(j_{21}j_{22}j_{23}) \Delta(j_{31}j_{32}j_{33}) \Delta(j_{11}j_{21}j_{31}) \Delta(j_{12}j_{22}j_{32}) \Delta(j_{13}j_{23}j_{33}) \\ \times \sum_{x_{\alpha}} \frac{(-1)^{\frac{1}{2}R + \frac{1}{2} \sum_{\beta=1}^3 x_{\beta} + x_6 + x_5 + x_6} (1 + \sum_{\beta=1}^6 z_{\beta})!}{\prod_{p,q=1}^3 (z_{pq} - j_{pq})! \prod_{\alpha=1}^6 \left\{ \frac{1}{2} \left(R - \sum_{\beta=1}^6 z_{\beta} \right) + z_{\alpha} - h_{\alpha} \right\}!} \quad (10.2.7)$$

where

$$(z_{pq}) \equiv \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{31} & z_{32} & z_{33} \end{pmatrix} = \begin{pmatrix} z_1 + z_4 & z_3 + z_6 & z_2 + z_5 \\ z_3 + z_5 & z_2 + z_4 & z_1 + z_6 \\ z_2 + z_6 & z_1 + z_5 & z_3 + z_4 \end{pmatrix} \quad (10.2.8)$$

$$\begin{aligned} h_1 &= j_{11} + j_{23} + j_{32}, & h_4 &= j_{11} + j_{22} + j_{33} \\ h_2 &= j_{31} + j_{22} + j_{13}, & h_5 &= j_{21} + j_{32} + j_{13} \\ h_3 &= j_{21} + j_{12} + j_{33}, & h_6 &= j_{31} + j_{12} + j_{23} \end{aligned} \quad (10.2.9)$$

$$R = \sum_{p,q=1}^3 j_{pq} \quad (10.2.10)$$

Summation in Eq. (10.2.7) is over all integer and half-integer $z_\alpha \geq 0$.

(b)

$$\begin{aligned} \left\{ \begin{matrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{matrix} \right\} &= \Delta(j_{11}j_{12}j_{13}) \Delta(j_{21}j_{22}j_{23}) \Delta(j_{31}j_{32}j_{33}) \Delta(j_{11}j_{21}j_{31}) \Delta(j_{12}j_{22}j_{32}) \Delta(j_{13}j_{23}j_{33}) \\ &\times (-1)^R \sum_{z_\alpha} \frac{(-1)^{z_4+z_5+z_6} \left(1 + \sum_{\beta=1}^6 z_\beta\right)!}{\prod_{p,q=1}^3 (z_{pq} - j_{pq})! \prod_{p=1}^3 \left[\left(R - \sum_{\beta=1}^6 z_\beta + z_p - h_p\right)! (z_{p+3} - h_{p+3})! \right]} \end{aligned} \quad (10.2.11)$$

Summation in (10.2.11) is over all integer and half-integer $z_\alpha \geq 0$; z_{pq} and h_p are determined by Eqs. (10.2.8) and (10.2.9).

(c)

$$\begin{aligned} \left\{ \begin{matrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{matrix} \right\} &= \Delta(j_{11}j_{12}j_{13}) \Delta(j_{21}j_{22}j_{23}) \Delta(j_{31}j_{32}j_{33}) \Delta(j_{11}j_{21}j_{31}) \Delta(j_{12}j_{22}j_{32}) \Delta(j_{13}j_{23}j_{33}) \\ &\times \sum_{x_\alpha} \frac{(-1)^{x_4+x_5+x_6} \left(1 + R - 2t + \sum_{\beta=1}^6 x_\beta - x_4\right)!}{\left(\prod_{p=1}^3 x_p!\right) x_4! (x_5 + x_4)! (x_6 + x_4)! \prod_{p \neq q=1}^3 (x_{pq} + t_{pq})! \prod_{p=1}^3 \left(t - j_{qr} - j_{rq} + x_p - \sum_{\beta=1}^6 x_\beta\right)!} \end{aligned} \quad (10.2.12)$$

where

$$(x_{pq}) \equiv \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} = \begin{pmatrix} x_1 & x_3 + x_6 & x_2 + x_5 \\ x_3 + x_5 & x_2 & x_1 + x_6 \\ x_2 + x_6 & x_1 + x_5 & x_3 \end{pmatrix} \quad (10.2.13)$$

Summation in Eq. (10.2.12) is over all integer $x_\alpha \geq 0$.

$$\begin{aligned} t_{pq} &\equiv j_{qr} + j_{rp} - j_{pp} - j_{qq}, & (p \neq q \neq r = 1, 2, 3) \\ t &= j_{11} + j_{22} + j_{33} \end{aligned} \quad (10.2.14)$$

10.2.3 $9j$ Symbols as Sums of Products of the Clebsch-Gordan Coefficients or the $3jm$ Symbols [54]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = [(2c+1)(2f+1)(2g+1)(2h+1)]^{-\frac{1}{2}} (2j+1)^{-1} \sum_{\substack{\alpha\beta\gamma\delta\varepsilon\varphi \\ \eta\mu\nu}} C_{a\alpha b\beta}^{c\gamma} C_{d\delta e\varepsilon}^{f\varphi} C_{c\gamma f\varphi}^{j\nu} C_{a\alpha d\delta}^{g\eta} C_{b\beta e\varepsilon}^{h\mu} C_{g\eta h\mu}^{j\nu} \quad (10.2.15)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = \frac{(-1)^{2(c+g)}}{(2a+1)(2e+1)(2j+1)} \sum_{\substack{\alpha\beta\gamma \\ \delta\varepsilon\varphi \\ \eta\mu\nu}} C_{c\gamma b\beta}^{a\alpha} C_{g\eta d\delta}^{a\alpha} C_{b\beta e\varepsilon}^{e\eta} C_{d\delta f\varphi}^{e\eta} C_{h\mu g\eta}^{j\nu} C_{f\varphi c\gamma}^{j\nu} \quad (10.2.16)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = \sum_{\substack{\alpha\beta\gamma\delta\varepsilon\varphi \\ \eta\mu\nu}} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \begin{pmatrix} d & e & f \\ \delta & \varepsilon & \varphi \end{pmatrix} \begin{pmatrix} g & h & j \\ \eta & \mu & \nu \end{pmatrix} \begin{pmatrix} a & d & g \\ \alpha & \delta & \eta \end{pmatrix} \begin{pmatrix} b & e & h \\ \beta & \varepsilon & \mu \end{pmatrix} \begin{pmatrix} c & f & j \\ \gamma & \varphi & \nu \end{pmatrix} \quad (10.2.17)$$

The following two relations were obtained in Ref. [20]:

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} &= \frac{\Delta(abc)\Delta(def)}{\Delta(adg)\Delta(cfj)} \left[\frac{(g+h+j+1)!(g-h+j)!}{(2h+1)} \right]^{\frac{1}{2}} \\ &\times \frac{(c+f-j)!(a+d-g)!}{(a+d+g+1)!(a-b+c)!(-a+b+c)!(d-e+f)!(-d+e+f)!} \\ &\times \sum_{xy\beta} (-1)^{a-c+\beta+x+y} \frac{(2a-x)!(2d-y)!(-a+b+c+x)!(-d+e+f+y)!}{x!y!(a+b-c-x)!(d+e-f-y)!(-a+c-\beta+x)!(-d+f+g-j+\beta+y)!} \\ &\times \frac{(-a+f+j-\beta+x)!(-d+c+g+\beta+y)!}{(a+d-g-x-y)!(-a+c-d+f+g+j+1+x+y)!} \left[\frac{(b-\beta)!(e+g-j+\beta)!}{(b+\beta)!(e-g+j-\beta)!} \right]^{\frac{1}{2}} C_{b\beta e j-g-\beta}^{h j-g} \quad (10.2.18) \\ \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} &= \left[\frac{(a+d-g)!(c+f-j)!(g-h+j)!(g+h+j+1)!}{(-a+d+g)!(a-d+g)!(a+d+g+1)!(c+f+j+1)!(e-f+j)!(-c+f+j)!} \right]^{\frac{1}{2}} \\ &\times \frac{1}{[(2c+1)(2f+1)(2h+1)]^{\frac{1}{2}}} \sum_{\alpha\beta} (-1)^{a-c+\beta} \left[\frac{(a+\alpha)!(d+g-\alpha)!(c+\alpha+\beta)!(f+j-\alpha-\beta)!}{(a-\alpha)!(d-g+\alpha)!(c-\alpha-\beta)!(f-j+\alpha+\beta)!} \right]^{\frac{1}{2}} \\ &\times C_{b\beta e j-g-\beta}^{h j-g} C_{a\alpha b\beta}^{c\alpha+\beta} C_{d g-\alpha e j-g-\beta}^{f j-\alpha-\beta} \quad (10.2.19) \end{aligned}$$

10.2.4. 9j Symbols as Sums of Products of the 6j Symbols or the Racah Coefficients [110,74]

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = \sum_x (-1)^{2x} (2x+1) \left\{ \begin{matrix} a & b & c \\ f & j & x \end{matrix} \right\} \left\{ \begin{matrix} d & e & f \\ b & x & h \end{matrix} \right\} \left\{ \begin{matrix} g & h & j \\ x & a & d \end{matrix} \right\}, \quad (10.2.20)$$

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} = \sum_x (2x+1) W(aech; xb) W(aegf; xd) W(chfg; xj). \quad (10.2.21)$$

10.3 Integral Representations of the 9j Symbols

10.3.1 Representation Involving the Generalised Hypergeometric Function

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} &= \frac{(-1)^{a+b-c-g-h+j}}{2^{b+e+h+1}} \cdot \frac{\Delta(abc)\Delta(def)\Delta(beh)}{\Delta(adg)\Delta(cfj)\Delta(ghj)} \\ &\times \frac{(a+d-g)!(c+f-j)!(g-h+j)!(b+e+h+1)!}{(a-b+c)!(-a+b+c)!(d-e+f)!(-d+e+f)!(a+d+g+1)!(-b+e+h)!(b+e+j-g)!} \\ &\times \sum_{xy} \frac{(-1)^{x+y} (2a-x)!(-a+b+c+x)!(-a-e+f+g+x)!(2d-y)!(c-d+e+j+y)!}{x!y!(a+b-c-x)!(-a+c-e+g-j+x)!(d+e-f-y)!} \\ &\times \frac{1}{(a+d-g-x-y)!(-a-d+g+c+f+j+1+x+y)!} \\ &\times \int_{-1}^1 (1+z)^{2e} {}_3F_2 \left[\begin{matrix} -b-e+g-j, -a-e+f+g+1+x, d-e-f-y; z-1 \\ -a+c-e+g-j+1+x, -c+d-e-j-y; \\ z+1 \end{matrix} \right] \\ &\times \frac{d^{-b+e+h}}{(dz)^{-b+e+h}} [(1-z)^{g+h-j} (1+z)^{-g+h+j}] dz. \quad (10.3.1) \end{aligned}$$

10.3.2 Representation Involving the Wigner D Functions

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} C_{c\gamma\rho\varphi}^{j\mu} &= (-1)^{c-f+g+h} \frac{[(2j+1)(a+b-c)!(a+b+c+1)!(d+e-f)!(d+e+f+1)!]^{\frac{1}{2}}}{(a+d+g+1)!(b+e+h+1)!\Delta(adg)\Delta(beh)} \\ &\times [(g+h-j)!(g+h+j+1)!]^{\frac{1}{2}} \int dR_1 dR_2 dR_3 B_{bch}^{adg}(R_1, R_2, R_3) D_{b-a,\gamma}^c(R_1) D_{c-d,\varphi}^j(R_2) D_{g-h,\mu}^{j*}(R_3) \quad (10.3.2) \end{aligned}$$

In this equation γ, φ, μ are arbitrary integer or half-integer numbers and

$$dR_i = \frac{1}{8\pi^2} \sin \beta_i d\alpha_i d\beta_i d\gamma_i, \quad (i = 1, 2, 3) \quad (10.3.3)$$

The integration limits are standard (see Eq. (4.10.4)). $B_{beh}^{adg}(R_1, R_2, R_3)$ may be presented in the form

$$B_{beh}^{adg}(R_1, R_2, R_3) = B_{12}^{a+d-g} B_{12}^{*b+e-h} B_{31}^{a-d+g} B_{31}^{*b-e+h} B_{32}^{-a+d+g} B_{32}^{*-b+e+h} \quad (10.3.4)$$

$$\begin{aligned} B_{ik} &\equiv e^{-\frac{i}{2}(\alpha_i + \alpha_k)} \left\{ \sin \frac{\beta_i}{2} \cos \frac{\beta_k}{2} e^{\frac{i}{2}(\gamma_i - \gamma_k)} - \cos \frac{\beta_i}{2} \sin \frac{\beta_k}{2} e^{-\frac{i}{2}(\gamma_i - \gamma_k)} \right\} \\ &= \sum_M (-1)^{\frac{1}{2}-M} D_{\frac{1}{2}M}^{\frac{1}{2}}(R_i) D_{\frac{1}{2}-M}^{\frac{1}{2}}(R_k) = -D_{\frac{1}{2}-\frac{1}{2}}^{\frac{1}{2}}(R_i R_k^{-1}) \end{aligned} \quad (10.3.5)$$

The coefficients B_{ik} possess the following properties:

$$(B_{ik})^{2j} = \sum_m (-1)^{j-m} D_{jm}^j(R_i) D_{j-m}^j(R_k) = (-1)^{2j} D_{j-j}^j(R_i R_k^{-1}) \quad (10.3.6)$$

$$(B_{ik}^*)^{2j} = \sum_m (-1)^{j-m} D_{-jm}^j(R_i) D_{-j-m}^j(R_k) = D_{-jj}^j(R_i R_k^{-1}) \quad (10.3.7)$$

$$|B_{ik}| = \sin \frac{\Omega_{ik}}{2} \quad (10.3.8)$$

where

$$\cos \Omega_{ik} = \cos \beta_i \cos \beta_k + \sin \beta_i \sin \beta_k \cos(\gamma_i - \gamma_k) \quad (10.3.9)$$

In particular, when $a = b, d = e, g = h$, from Eq. (10.3.2), one has [78]

$$\begin{aligned} \left\{ \begin{matrix} a & a & c \\ d & d & f \\ g & g & j \end{matrix} \right\} C_{c0\rho0}^{j0} &= (-1)^{c+f} \frac{1}{[(a+d-g+1)!]^2 \Delta^2(adg)} \\ &\times [(2j+1)(2a-c)!(2a+c+1)!(2d-f)!(2d+f+1)!(2g-j)!(2g+j+1)!]^{\frac{1}{2}} \\ &\times \int d\Omega_1 d\Omega_2 d\Omega_3 B_{adg}(\Omega_1, \Omega_2, \Omega_3) P_c(\cos \beta_1) P_f(\cos \beta_2) P_j(\cos \beta_3) \end{aligned} \quad (10.3.10)$$

Here

$$d\Omega_i = \frac{1}{4\pi} \sin \beta_i d\beta_i d\gamma_i \quad (10.3.11)$$

$$B_{adg}(\Omega_1, \Omega_2, \Omega_3) = \left(\sin \frac{\Omega_{12}}{2} \right)^{2(a+d-g)} \left(\sin \frac{\Omega_{31}}{2} \right)^{2(a-d+g)} \left(\sin \frac{\Omega_{32}}{2} \right)^{2(-a+d+g)}. \quad (10.3.12)$$

10.3.3 Integrals Involving Characters of Irreducible Representations of Rotation Group

$$\begin{aligned} \left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\}^2 &= \frac{1}{(8\pi^2)^5} \int dR_1 dR_2 dR_3 dR_4 dR_5 \chi^c(R_1) \chi^f(R_2) \chi^j(R_3) \chi^a(R_1 R_4) \chi^b(R_1 R_5) \chi^d(R_2 R_4) \\ &\times \chi^e(R_2 R_5) \chi^g(R_3 R_4) \chi^h(R_3 R_5) \end{aligned} \quad (10.3.13)$$

$$\left\{ \begin{matrix} a & a & b \\ a & a & c \\ b & c & d \end{matrix} \right\} = \frac{1}{(8\pi^2)^3} \int dR_1 dR_2 dR_3 \chi^a(R_1) \chi^a(R_2) \chi^d(R_3) \chi^a(R_2 R_1) \chi^b(R_3 R_1^{-1}) \chi^c(R_3 R_2^{-1}) \quad (10.3.14)$$

10.4 Symmetry Properties of the 9j Symbols

10.4.1 Permutation Symmetry

Any 9j symbol possesses the following symmetry properties with respect to permutations [79]: even permutations of columns or rows as well as transposition leave the 9j symbol unchanged; odd permutations of columns or rows introduce the phase factor $(-1)^R$, where R is sum of all arguments of the 9j symbol. These symmetry properties may be written as follows

(a) Column permutations

$$\begin{pmatrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{pmatrix} = \varepsilon \begin{pmatrix} j_{1i} & j_{1k} & j_{1l} \\ j_{2i} & j_{2k} & j_{2l} \\ j_{3i} & j_{3k} & j_{3l} \end{pmatrix} \quad (10.4.1)$$

(b) Row permutations

$$\begin{pmatrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{pmatrix} = \varepsilon \begin{pmatrix} j_{i1} & j_{i2} & j_{i3} \\ j_{k1} & j_{k2} & j_{k3} \\ j_{l1} & j_{l2} & j_{l3} \end{pmatrix} \quad (10.4.2)$$

(c) Transposition

$$\begin{pmatrix} j_{11} & j_{12} & j_{13} \\ j_{21} & j_{22} & j_{23} \\ j_{31} & j_{32} & j_{33} \end{pmatrix} = \begin{pmatrix} j_{11} & j_{21} & j_{31} \\ j_{12} & j_{22} & j_{32} \\ j_{13} & j_{23} & j_{33} \end{pmatrix} \quad (10.4.3)$$

Here

$$\varepsilon = \begin{cases} 1 & \text{for even permutations} \\ (-1)^R & \text{(cyclic permutations)} \\ -1 & \text{for odd permutations} \\ & \text{(non-cyclic permutations).} \end{cases} \quad (10.4.4)$$

$$R = \sum_{i,k=1}^3 j_{ik} \quad (10.4.5)$$

All these symmetry properties are independent and relate $3! \times 3! \times 2 = 729j$ symbols given below (Eq. (10.4.6)). The $9j$ symbols obtained by column permutations are arranged horizontally. The $9j$ symbols obtained by row permutations are arranged vertically. The six lower rows represent the transposed $9j$ symbols.

$$\begin{aligned} \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & j \end{pmatrix} &= \begin{pmatrix} c & a & b \\ f & d & e \\ j & g & h \end{pmatrix} = \begin{pmatrix} b & c & a \\ e & f & d \\ h & j & g \end{pmatrix} = \varepsilon \begin{pmatrix} b & a & c \\ e & d & f \\ h & g & j \end{pmatrix} = \varepsilon \begin{pmatrix} c & b & a \\ f & e & d \\ j & h & g \end{pmatrix} = \varepsilon \begin{pmatrix} a & c & b \\ d & f & e \\ g & j & h \end{pmatrix} \\ &= \begin{pmatrix} g & h & j \\ a & b & c \\ d & e & f \end{pmatrix} = \begin{pmatrix} j & g & h \\ c & a & b \\ f & d & e \end{pmatrix} = \begin{pmatrix} h & j & g \\ b & c & a \\ e & f & d \end{pmatrix} = \varepsilon \begin{pmatrix} h & g & j \\ b & a & c \\ e & d & f \end{pmatrix} = \varepsilon \begin{pmatrix} j & h & g \\ c & b & a \\ f & e & d \end{pmatrix} = \varepsilon \begin{pmatrix} g & j & h \\ a & c & b \\ d & f & e \end{pmatrix} \\ &= \begin{pmatrix} d & e & f \\ g & h & j \\ a & b & c \end{pmatrix} = \begin{pmatrix} f & d & e \\ j & g & h \\ c & a & b \end{pmatrix} = \begin{pmatrix} e & f & d \\ h & j & g \\ b & c & a \end{pmatrix} = \varepsilon \begin{pmatrix} e & d & f \\ h & g & j \\ b & a & c \end{pmatrix} = \varepsilon \begin{pmatrix} f & e & d \\ j & h & g \\ c & b & a \end{pmatrix} = \varepsilon \begin{pmatrix} d & f & e \\ g & j & h \\ a & c & b \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
&= \varepsilon \begin{Bmatrix} d & e & f \\ a & b & c \\ g & h & j \end{Bmatrix} = \varepsilon \begin{Bmatrix} f & d & e \\ c & a & b \\ j & g & h \end{Bmatrix} = \varepsilon \begin{Bmatrix} e & f & d \\ b & c & a \\ h & j & g \end{Bmatrix} = \begin{Bmatrix} e & d & f \\ b & a & c \\ h & g & j \end{Bmatrix} = \begin{Bmatrix} f & e & d \\ c & b & a \\ j & h & g \end{Bmatrix} = \begin{Bmatrix} d & f & e \\ a & c & b \\ g & j & h \end{Bmatrix} \\
&= \varepsilon \begin{Bmatrix} g & h & j \\ d & e & f \\ a & b & c \end{Bmatrix} = \varepsilon \begin{Bmatrix} j & g & h \\ f & d & e \\ c & a & b \end{Bmatrix} = \varepsilon \begin{Bmatrix} h & j & g \\ e & f & d \\ b & c & a \end{Bmatrix} = \begin{Bmatrix} h & g & j \\ e & d & f \\ b & a & c \end{Bmatrix} = \begin{Bmatrix} j & h & g \\ f & e & d \\ c & b & a \end{Bmatrix} = \begin{Bmatrix} g & j & h \\ d & f & e \\ a & c & b \end{Bmatrix} \\
&= \varepsilon \begin{Bmatrix} a & b & c \\ g & h & j \\ d & e & f \end{Bmatrix} = \varepsilon \begin{Bmatrix} c & a & b \\ j & g & h \\ f & d & e \end{Bmatrix} = \varepsilon \begin{Bmatrix} b & c & a \\ h & j & g \\ e & f & d \end{Bmatrix} = \begin{Bmatrix} b & a & c \\ h & g & j \\ e & d & f \end{Bmatrix} = \begin{Bmatrix} c & b & a \\ j & h & g \\ f & e & d \end{Bmatrix} = \begin{Bmatrix} a & c & b \\ g & j & h \\ d & f & e \end{Bmatrix} \\
&= \begin{Bmatrix} a & d & g \\ b & e & h \\ c & f & j \end{Bmatrix} = \begin{Bmatrix} g & a & d \\ h & b & e \\ j & c & f \end{Bmatrix} = \begin{Bmatrix} d & g & a \\ e & h & b \\ f & j & c \end{Bmatrix} = \varepsilon \begin{Bmatrix} d & a & g \\ e & b & h \\ f & c & j \end{Bmatrix} = \varepsilon \begin{Bmatrix} g & d & a \\ h & e & b \\ j & f & c \end{Bmatrix} = \varepsilon \begin{Bmatrix} a & g & d \\ b & h & e \\ c & j & f \end{Bmatrix} \\
&= \begin{Bmatrix} c & f & j \\ a & d & g \\ b & e & h \end{Bmatrix} = \begin{Bmatrix} j & c & f \\ g & a & d \\ h & b & e \end{Bmatrix} = \begin{Bmatrix} f & j & c \\ d & g & a \\ e & h & b \end{Bmatrix} = \varepsilon \begin{Bmatrix} f & c & j \\ d & a & g \\ e & b & h \end{Bmatrix} = \varepsilon \begin{Bmatrix} j & f & c \\ g & d & a \\ h & e & b \end{Bmatrix} = \varepsilon \begin{Bmatrix} c & j & f \\ a & g & d \\ b & h & e \end{Bmatrix} \\
&= \begin{Bmatrix} b & e & h \\ c & f & j \\ a & d & g \end{Bmatrix} = \begin{Bmatrix} h & b & e \\ j & c & f \\ g & a & d \end{Bmatrix} = \begin{Bmatrix} e & h & b \\ f & j & c \\ d & g & a \end{Bmatrix} = \varepsilon \begin{Bmatrix} e & b & h \\ f & c & j \\ d & a & g \end{Bmatrix} = \varepsilon \begin{Bmatrix} h & e & b \\ j & f & c \\ g & d & a \end{Bmatrix} = \varepsilon \begin{Bmatrix} b & h & e \\ c & j & f \\ a & g & d \end{Bmatrix} \\
&= \varepsilon \begin{Bmatrix} b & e & h \\ a & d & g \\ c & f & j \end{Bmatrix} = \varepsilon \begin{Bmatrix} h & b & e \\ g & a & d \\ j & c & f \end{Bmatrix} = \varepsilon \begin{Bmatrix} e & h & b \\ d & g & a \\ f & j & c \end{Bmatrix} = \begin{Bmatrix} e & b & h \\ d & a & g \\ f & c & j \end{Bmatrix} = \begin{Bmatrix} h & e & b \\ g & d & a \\ j & f & c \end{Bmatrix} = \begin{Bmatrix} b & h & e \\ a & g & d \\ c & j & f \end{Bmatrix} \\
&= \varepsilon \begin{Bmatrix} c & f & j \\ b & e & h \\ a & d & g \end{Bmatrix} = \varepsilon \begin{Bmatrix} j & c & f \\ h & b & e \\ g & a & d \end{Bmatrix} = \varepsilon \begin{Bmatrix} f & j & c \\ e & h & b \\ d & g & a \end{Bmatrix} = \begin{Bmatrix} f & c & j \\ e & b & h \\ d & a & g \end{Bmatrix} = \begin{Bmatrix} j & f & c \\ h & e & b \\ g & d & a \end{Bmatrix} = \begin{Bmatrix} c & j & f \\ b & h & e \\ a & g & d \end{Bmatrix}. \tag{10.4.6} \\
&= \varepsilon \begin{Bmatrix} a & d & g \\ c & f & j \\ b & e & h \end{Bmatrix} = \varepsilon \begin{Bmatrix} g & a & d \\ j & c & f \\ h & b & e \end{Bmatrix} = \varepsilon \begin{Bmatrix} d & g & a \\ f & j & c \\ e & h & b \end{Bmatrix} = \begin{Bmatrix} d & a & g \\ f & c & j \\ e & b & h \end{Bmatrix} = \begin{Bmatrix} g & d & a \\ j & f & c \\ h & e & b \end{Bmatrix} = \begin{Bmatrix} a & g & d \\ c & j & f \\ b & h & e \end{Bmatrix}
\end{aligned}$$

10.4.2 Symmetries of the r Symbol

Any r symbol possesses the following symmetry properties:

(a) *Permutations of rows introduce the phase factor ε (see Eq. (10.4.4))*

$$\left\| \begin{array}{cccccc} r_{11} & r_{12} & r_{13} & r'_{11} & r'_{12} & r'_{13} \\ r_{21} & r_{22} & r_{23} & r'_{21} & r'_{22} & r'_{23} \\ r_{31} & r_{32} & r_{33} & r'_{31} & r'_{32} & r'_{33} \end{array} \right\| = \varepsilon \left\| \begin{array}{cccccc} r_{i1} & r_{i2} & r_{i3} & r'_{i1} & r'_{i2} & r'_{i3} \\ r_{k1} & r_{k2} & r_{k3} & r'_{k1} & r'_{k2} & r'_{k3} \\ r_{l1} & r_{l2} & r_{l3} & r'_{l1} & r'_{l2} & r'_{l3} \end{array} \right\|, \tag{10.4.7}$$

These permutations correspond to permutations of rows for the $9j$ symbols.

(b) *Simultaneous permutations of appropriate columns in both parts of the r symbol (i.e., r_{ik} and r'_{ik}) introduce the same phase factor ε*

$$\left\| \begin{array}{cccccc} r_{11} & r_{12} & r_{13} & r'_{11} & r'_{12} & r'_{13} \\ r_{21} & r_{22} & r_{23} & r'_{21} & r'_{22} & r'_{23} \\ r_{31} & r_{32} & r_{33} & r'_{31} & r'_{32} & r'_{33} \end{array} \right\| = \varepsilon \left\| \begin{array}{cccccc} r_{1i} & r_{1k} & r_{1l} & r'_{1i} & r'_{1k} & r'_{1l} \\ r_{2i} & r_{2k} & r_{2l} & r'_{2i} & r'_{2k} & r'_{2l} \\ r_{3i} & r_{3k} & r_{3l} & r'_{3i} & r'_{3k} & r'_{3l} \end{array} \right\|. \tag{10.4.8}$$

This symmetry corresponds to the symmetry of the $9j$ symbols with respect to column permutations.

(c) *Transposition of both parts of the r symbol with their simultaneous permutation is equivalent to transposition of the $9j$ symbols*

$$\left\| \begin{array}{cccccc} r_{11} & r_{12} & r_{13} & r'_{11} & r'_{12} & r'_{13} \\ r_{21} & r_{22} & r_{23} & r'_{21} & r'_{22} & r'_{23} \\ r_{31} & r_{32} & r_{33} & r'_{31} & r'_{32} & r'_{33} \end{array} \right\| = \left\| \begin{array}{cccccc} r'_{11} & r'_{21} & r'_{31} & r_{11} & r_{21} & r_{31} \\ r'_{12} & r'_{22} & r'_{32} & r_{12} & r_{22} & r_{32} \\ r'_{13} & r'_{23} & r'_{33} & r_{13} & r_{23} & r_{33} \end{array} \right\|. \tag{10.4.9}$$

10.4.3 "Mirror" Symmetry

The original relations which define the $9j$ symbols may be extended to the domain of negative integer or half-integer arguments. Introducing the notations $a = -a - 1, b = -b - 1, \dots$, etc., we obtain the following symmetry properties [20]

$$\begin{aligned}
 & \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_1 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_1 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = i\eta_2 \begin{Bmatrix} a & b \\ d & c \\ g & e \end{Bmatrix} = -i\eta_2 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_3 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_3 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = i\eta_4 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = -i\eta_4 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_5 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_5 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = i\eta_6 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = -i\eta_6 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_7 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_7 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = \eta_5 \begin{Bmatrix} a & b & c \\ d & c & f \\ g & h & j \end{Bmatrix} = \eta_5 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_8 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \eta_8 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = -i\eta_6 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = -i\eta_6 \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = - \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \\
 & = - \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} = - \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}. \tag{10.4.10}
 \end{aligned}$$

In these equations

$$\begin{aligned}
 \eta_1 &= (-1)^{b-c-d+g}, & \eta_5 &= (-1)^{c-f-g+h+1}, \\
 \eta_2 &= (-1)^{c+d-e-g+h}, & \eta_6 &= (-1)^{g-h-j}, \\
 \eta_3 &= (-1)^{c+f-g-h}, & \eta_7 &= (-1)^{2(a+e+j)}, \\
 \eta_4 &= (-1)^R, & \eta_8 &= (-1)^{R-d+e-f+1}.
 \end{aligned} \tag{10.4.11}$$

When tabulating formulas for the $9j$ symbols, it is convenient to use the "mirror" symmetry properties in the form [20]

$$\begin{Bmatrix} d-\delta & e+\varepsilon & f+\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = i(-1)^{g+\varepsilon-\varphi} \begin{Bmatrix} \bar{d}+\delta & e+\varepsilon & f+\varphi \\ \bar{d} & e & f \\ g & h & j \end{Bmatrix}, \tag{10.4.12}$$

$$\begin{Bmatrix} d+\delta & e-\varepsilon & f+\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = i(-1)^{h+\varphi-\delta} \begin{Bmatrix} d+\delta & \bar{e}+\varepsilon & f+\varphi \\ d & \bar{e} & f \\ g & h & j \end{Bmatrix}, \tag{10.4.13}$$

$$\begin{Bmatrix} d+\delta & e+\varepsilon & f-\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = i(-1)^{j+\delta-e} \begin{Bmatrix} d+\delta & e+\varepsilon & \bar{f}+\varphi \\ d & e & \bar{f} \\ g & h & j \end{Bmatrix}, \tag{10.4.14}$$

$$\begin{Bmatrix} d-\delta & e-\varepsilon & f+\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = (-1)^{h+\varphi-\theta+1} \begin{Bmatrix} d+\delta & \bar{e}+\varepsilon & f+\varphi \\ \bar{d} & \bar{e} & f \\ g & h & j \end{Bmatrix}, \tag{10.4.15}$$

$$\begin{Bmatrix} d-\delta & e+\varepsilon & f-\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = (-1)^{g+\varepsilon-j+1} \begin{Bmatrix} d+\delta & e+\varepsilon & \bar{f}+\varphi \\ \bar{d} & e & \bar{f} \\ g & h & j \end{Bmatrix}, \tag{10.4.16}$$

$$\begin{Bmatrix} d+\delta & e-\varepsilon & f-\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = (-1)^{j+\delta-h+1} \begin{Bmatrix} d+\delta & \bar{e}+\varepsilon & \bar{f}+\varphi \\ d & \bar{e} & \bar{f} \\ g & h & j \end{Bmatrix}, \tag{10.4.17}$$

$$\begin{Bmatrix} d-\delta & e-\varepsilon & f-\varphi \\ d & e & f \\ g & h & j \end{Bmatrix} = i(-1)^{g+h+j+\delta+\varepsilon+\varphi+1} \begin{Bmatrix} \bar{d}+\delta & \bar{e}+\varepsilon & \bar{f}+\varphi \\ \bar{d} & \bar{e} & \bar{f} \\ g & h & j \end{Bmatrix} \tag{10.4.18}$$

10.5 Recursion Relations for the $9j$ Symbols

10.5.1 General Form of the Recursion Relation

Recursion relations for the $9j$ symbols may be obtained for instance, using the following equation [80]

$$\begin{aligned} & (-1)^{a+f+b+j} \sum_x (-1)^{2x} (2x+1) \left\{ \begin{matrix} a & b & x \\ d & e & f \\ g & h & j \end{matrix} \right\} \left\{ \begin{matrix} a & b & x \\ c & \lambda & a' \end{matrix} \right\} \left\{ \begin{matrix} j & f & x \\ \lambda & c & f' \end{matrix} \right\} \\ &= (-1)^{a'+f'+g+e} \sum_y (-1)^{2y} (2y+1) \left\{ \begin{matrix} a' & b & c \\ y & e & f' \\ g & h & j \end{matrix} \right\} \left\{ \begin{matrix} f' & e & y \\ d & \lambda & f \end{matrix} \right\} \left\{ \begin{matrix} g & a' & y \\ \lambda & d & a \end{matrix} \right\}. \end{aligned} \quad (10.5.1)$$

Specifying $\lambda = \frac{1}{2}, 1, \frac{3}{2}, 2$, etc. and using explicit forms of the $6j$ symbols, we may obtain various recursion relations. Examples of such relations are given below (Eqs. (10.5.2)-(10.5.5)).

10.5.2 Relations Involving Four $9j$ Symbols

Equations (10.5.2)-(10.5.4) are derived by putting $\lambda = \frac{1}{2}$ in Eq. (10.5.1) and specifying the values of a' and f' . In particular

(a) for $a' = a - \frac{1}{2}, f' = f - \frac{1}{2}$, one has

$$\begin{aligned} & \frac{1}{2c+1} \left[\left(a - b + c + \frac{1}{2} \right) \left(a + b + c + \frac{3}{2} \right) \left(c + f - j + \frac{1}{2} \right) \left(c + f + j + \frac{3}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c + \frac{1}{2} \\ d & e & f \\ g & h & j \end{matrix} \right\} \\ &+ \frac{1}{2c+1} \left[\left(-a + b + c + \frac{1}{2} \right) \left(a + b - c + \frac{1}{2} \right) \left(-c + f + j + \frac{1}{2} \right) \left(c - f + j + \frac{1}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c - \frac{1}{2} \\ d & e & f \\ g & h & j \end{matrix} \right\} \\ &= \frac{1}{2d+1} [(d+e-f+1)(-d+e+f)(-a+g+d+1)(a-d+g)]^{\frac{1}{2}} \left\{ \begin{matrix} a - \frac{1}{2} & b & c \\ d + \frac{1}{2} & e & f - \frac{1}{2} \\ g & h & j \end{matrix} \right\} \\ &+ \frac{1}{2d+1} [(d-e+f)(d+e+f+1)(a+d-g)(a+d+g+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a - \frac{1}{2} & b & c \\ d - \frac{1}{2} & e & f - \frac{1}{2} \\ g & h & j \end{matrix} \right\}; \end{aligned} \quad (10.5.2)$$

(b) for $a' = a + \frac{1}{2}, f' = f + \frac{1}{2}$, one obtains

$$\begin{aligned} & \frac{1}{2c+1} \left[\left(-a + b + c + \frac{1}{2} \right) \left(a + b - c + \frac{1}{2} \right) \left(-c + f + j + \frac{1}{2} \right) \left(c - f + j + \frac{1}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c + \frac{1}{2} \\ d & e & f \\ g & h & j \end{matrix} \right\} \\ &+ \frac{1}{2c+1} \left[\left(a - b + c + \frac{1}{2} \right) \left(a + b + c + \frac{3}{2} \right) \left(c + f - j + \frac{1}{2} \right) \left(c + f + j + \frac{3}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} a & b & c - \frac{1}{2} \\ d & e & f \\ g & h & j \end{matrix} \right\} \\ &= \frac{1}{2d+1} [(d-e+f+1)(d+e+f+2)(a+d-g+1)(a+d+g+2)]^{\frac{1}{2}} \left\{ \begin{matrix} a + \frac{1}{2} & b & c \\ d + \frac{1}{2} & e & f + \frac{1}{2} \\ g & h & j \end{matrix} \right\} \\ &+ \frac{1}{2d+1} [(-d+e+f+1)(d+e-f)(-a+d+g)(a-d+g+1)]^{\frac{1}{2}} \left\{ \begin{matrix} a + \frac{1}{2} & b & c \\ d - \frac{1}{2} & e & f + \frac{1}{2} \\ g & h & j \end{matrix} \right\} \end{aligned} \quad (10.5.3)$$

(c) for $a' = a - \frac{1}{2}, f' = f + \frac{1}{2}$, one gets

$$\begin{aligned}
& \frac{1}{2c+1} \left[\left(a - b + c + \frac{1}{2} \right) \left(a + b + c + \frac{3}{2} \right) \left(-c + f + j + \frac{1}{2} \right) \left(c - f + j + \frac{1}{2} \right) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c + \frac{1}{2} \\ d & e & f \\ g & h & j \end{Bmatrix} \\
& - \frac{1}{2c+1} \left[\left(-a + b + c + \frac{1}{2} \right) \left(a + b - c + \frac{1}{2} \right) \left(c + f - j + \frac{1}{2} \right) \left(c + f + j + \frac{3}{2} \right) \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c - \frac{1}{2} \\ d & e & f \\ g & h & j \end{Bmatrix} \\
& = \frac{1}{2d+1} [(d-e+f+1)(d+e+f+2)(-a+d+g+1)(a-d+g)]^{\frac{1}{2}} \begin{Bmatrix} a - \frac{1}{2} & b & c \\ d + \frac{1}{2} & e & f + \frac{1}{2} \\ g & h & j \end{Bmatrix} \\
& - \frac{1}{2d+1} [(-d+e+f+1)(d+e-f)(a+d-g)(a+d+g+1)]^{\frac{1}{2}} \begin{Bmatrix} a - \frac{1}{2} & b & c \\ d - \frac{1}{2} & e & f + \frac{1}{2} \\ g & h & j \end{Bmatrix}. \tag{10.5.4}
\end{aligned}$$

The relation for $a' = a + \frac{1}{2}$ and $f' = f - \frac{1}{2}$ is equivalent to Eq. (10.5.4).

10.5.3 Relations Involving Five 9j Symbols [20]

Putting $\lambda = 1, a' = a$ and $f' = f$ in Eq. (10.5.1), we get

$$\begin{aligned}
& \frac{A_{c+1}(ab, fj)}{(c+1)(2c+1)} \begin{Bmatrix} a & b & c+1 \\ d & e & f \\ g & h & j \end{Bmatrix} + \frac{A_c(ab, fj)}{c(2c+1)} \begin{Bmatrix} a & b & c-1 \\ d & e & f \\ g & h & j \end{Bmatrix} - \frac{A_{d+1}(ef, ag)}{(d+1)(2d+1)} \begin{Bmatrix} a & b & c \\ d+1 & e & f \\ g & h & j \end{Bmatrix} \\
& - \frac{A_d(ef, ag)}{d(2d+1)} \begin{Bmatrix} a & b & c \\ d-1 & e & f \\ g & h & j \end{Bmatrix} \\
& = \left\{ \frac{[a(a+1) + d(d+1) - g(g+1)][d(d+1) - e(e+1) + f(f+1)]}{d(d+1)} \right. \\
& \left. - \frac{[a(a+1) - b(b+1) + c(c+1)][c(c+1) + f(f+1) - j(j+1)]}{c(c+1)} \right\} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}, \tag{10.5.5}
\end{aligned}$$

where

$$\begin{aligned}
A_q(pr, st) = & [(-p+r+q)(p-r+q)(p+r-q+1)(p+r+q+1) \\
& \times (-s+t+q)(s-t+q)(s+t-q+1)(s+t+q+1)]^{\frac{1}{2}} \tag{10.5.6}
\end{aligned}$$

We also present the relations of another type between five 9j symbols [81]:

$$\begin{aligned}
& \left[\frac{(g+h+j+1)(g+h-j)(-b+e+h)(b+e-h+1)}{(a+d+g+2)(a-d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d & e & f \\ g - \frac{1}{2} & h - \frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g-h+j)(-g+h+j+1)(b+e+h+2)(b-e+h+1)}{(a+d+g+2)(a-d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d & e & f \\ g - \frac{1}{2} & h + \frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g+h+j+2)(g+h-j+1)(b+e+h+2)(b-e+h+1)}{(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d & e & f \\ g + \frac{1}{2} & h + \frac{1}{2} & j \end{Bmatrix} \\
& - \left[\frac{(g-h+j+1)(-g+h+j)(b+e-h+1)(-b+e+h)}{(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a + \frac{1}{2} & b + \frac{1}{2} & c \\ d & e & f \\ g + \frac{1}{2} & h - \frac{1}{2} & j \end{Bmatrix} \\
& = (2g+1)(2h+1) \left[\frac{(a+b+c+2)(a+b-c+1)}{(a+d+g+2)(a-d+g+1)(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}. \tag{10.5.7}
\end{aligned}$$

$$\begin{aligned}
& \left[\frac{(g+h+j+1)(g+h-j)(b+e+h+1)(b+h-e)}{(a+d-g)(-a+d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a-\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g-\frac{1}{2} & h-\frac{1}{2} & j \end{Bmatrix} \\
& - \left[\frac{(-g+h+j+1)(g-h+j)(-b+e+h+1)(b+e-h)}{(a+d-g)(-a+d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a-\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g-\frac{1}{2} & h+\frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g-h+j+1)(-g+h+j)(b+e+h+1)(b-e+h)}{(a+d+g+1)(a-d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a-\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g+\frac{1}{2} & h-\frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g+h+j+2)(g+h-j+1)(-b+e+h+1)(b+e-h)}{(a+d+g+1)(a-d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a-\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g+\frac{1}{2} & h+\frac{1}{2} & j \end{Bmatrix} \\
& = (2g+1)(2h+1) \left[\frac{(a+b+c+1)(a+b-c)}{(a+d+g+1)(a-d+g)(a+d-g)(-a+d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}. \tag{10.5.8}
\end{aligned}$$

$$\begin{aligned}
& \left[\frac{(-g+h+j+1)(g-h+j)(-b+e+h+1)(b+e-h)}{(a+d+g+2)(a-d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g-\frac{1}{2} & h+\frac{1}{2} & j \end{Bmatrix} \\
& - \left[\frac{(g+h+j+1)(g+h-j)(b+e+h+1)(b-e+h)}{(a+d+g+2)(-a+d+g+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g-\frac{1}{2} & h-\frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g-h+j+1)(-g+h+j)(b+e+h+1)(b-e+h)}{(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g+\frac{1}{2} & h-\frac{1}{2} & j \end{Bmatrix} \\
& + \left[\frac{(g+h+j+2)(g+h-j+1)(-b+e+h+1)(b+e-h)}{(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a+\frac{1}{2} & b-\frac{1}{2} & c \\ d & e & f \\ g+\frac{1}{2} & h+\frac{1}{2} & j \end{Bmatrix} \\
& = (2g+1)(2h+1) \left[\frac{(a-b+c+1)(-a+b+c)}{(a+d+g+2)(a-d+g+1)(a+d-g+1)(-a+d+g)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix}. \tag{10.5.9}
\end{aligned}$$

10.5.4 Relation Involving Six 9j Symbols

$$\begin{aligned}
& \frac{[(a+b+1)^2 - (c+1)^2]^{\frac{1}{2}} [(c+1)^2 - (a-b)^2]^{\frac{1}{2}}}{(2c+1)} \left[\frac{f+j-c+1}{f+j-c} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c+1 \\ d & e & f \\ g & h & j \end{Bmatrix} \\
& + \frac{[(d+e+1)^2 - (f+1)^2]^{\frac{1}{2}} [(f+1)^2 - (d-e)^2]^{\frac{1}{2}}}{(2f+1)} \left[\frac{c-f+j+1}{c-f+j} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f+1 \\ g & h & j \end{Bmatrix} \\
& + \frac{[(g+h+1)^2 - (j+1)^2]^{\frac{1}{2}} [(j+1)^2 - (g-h)^2]^{\frac{1}{2}}}{(2j+1)} \left[\frac{c+f-j+1}{c+f-j} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j+1 \end{Bmatrix} \\
& = \left[\frac{(-c+f+j+1)(c-f+j+1)(c+f-j+1)(c+f+j+2)}{(-c+f+j)(c-f+j)(c+f-j)(c+f+j+1)} \right]^{\frac{1}{2}} \\
& \times \left\{ \frac{[(a+b+1)^2 - c^2]^{\frac{1}{2}} [c^2 - (a-b)^2]^{\frac{1}{2}}}{2c+1} \left[\frac{-c+f+j}{-c+f+j+1} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c-1 \\ d & e & f \\ g & h & j \end{Bmatrix} \right. \\
& + \frac{[(d+e+1)^2 - f^2]^{\frac{1}{2}} [f^2 - (d-e)^2]^{\frac{1}{2}}}{2f+1} \left[\frac{c-f+j}{c-f+j+1} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f-1 \\ g & h & j \end{Bmatrix} \\
& \left. + \frac{[(g+h+1)^2 - j^2]^{\frac{1}{2}} [j^2 - (g-h)^2]^{\frac{1}{2}}}{2j+1} \left[\frac{c+f-j}{c+f-j+1} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j-1 \end{Bmatrix} \right\} \tag{10.5.10}
\end{aligned}$$

10.5.5 Recursion Relations for Some Special Cases

Somewhat simpler recursion relations may be obtained, if arguments of the $9j$ symbols satisfy some special requirements.

(a) If two columns (or two rows) are the same, then

$$\begin{aligned}
 & \frac{1}{2c+1} [(2a+c+2)(2a-c)(-c+f+j)(c-f+j+1)(c+f-j+1)(c+f+j+2)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c+1 \\ d & d & f \\ g & g & j \end{Bmatrix} \\
 & + \frac{1}{2c+1} [(2a+c+1)(2a-c+1)(-c+f+j+1)(c-f+j)(c+f-j)(c+f+j+1)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c-1 \\ d & d & f \\ g & g & j \end{Bmatrix} \\
 & = \frac{1}{2f+1} [(2d+f+2)(2d-f)(-c+f+j+1)(c-f+j)(c+f-j+1)(c+f+j+2)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c \\ d & d & f+1 \\ g & g & j \end{Bmatrix} \\
 & + \frac{1}{2f+1} [(2d+f+1)(2d-f+1)(-c+f+j)(c-f+j+1)(c+f-j)(c+f+j+1)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c \\ d & d & f-1 \\ g & g & j \end{Bmatrix}
 \end{aligned} \tag{10.5.11}$$

(b) If one of arguments in any triad is equal to the sum of two other arguments, one has [20]

$$\begin{aligned}
 & \left[\frac{(-c+g+h-f)(c-g+h+f+1)(c+g-h+f+1)(c+g+h+f+2)}{(2c+2f+2)(2c+2f+3)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & c+f \end{Bmatrix} \\
 & = \left[\frac{(-a+b+c+1)(a-b+c+1)(a+b-c)(a+b+c+2)}{(2c+1)(2c+2)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c+1 \\ d & e & f \\ g & h & c+f+1 \end{Bmatrix} \\
 & + \left[\frac{(-d+e+f+1)(d-e+f+1)(d+e-f)(d+e+f+2)}{(2f+1)(2f+2)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f+1 \\ g & h & c+f+1 \end{Bmatrix}
 \end{aligned} \tag{10.5.12}$$

(c) If one of arguments in any triad is equal to 0 or 1, one gets [32, 75]:

$$\begin{Bmatrix} a & b & c \\ d & e & c \\ g & g & 1 \end{Bmatrix} = \frac{a(a+1) + e(e+1) - d(d+1) - b(b+1)}{2[c(c+1)g(g+1)]^{\frac{1}{2}}} \begin{Bmatrix} a & b & c \\ d & e & c \\ g & g & 0 \end{Bmatrix}, \tag{10.5.13}$$

$$\begin{aligned}
 & [g(g+1)(c+1)(2c+3)(2d+c+2)(2d-c)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c \\ d & d & c+1 \\ g & g & 1 \end{Bmatrix} \\
 & = [g(g+1)c(2c-1)(2d+c+1)(2d-c+1)]^{\frac{1}{2}} \begin{Bmatrix} a & a & c \\ d & d & c \\ g & g & 1 \end{Bmatrix} \\
 & + (2c+1)[d(d+1) + g(g+1) - a(a+1)] \begin{Bmatrix} a & a & c \\ d & d & c \\ g & g & 0 \end{Bmatrix},
 \end{aligned} \tag{10.5.14}$$

$$\begin{aligned}
 & [c(2c+3)(d+e+c+2)(-d+e+c+1)(d-e+c+1)(d+e-c)]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & c+1 \\ g & g & 1 \end{Bmatrix} \\
 & - [(c+1)(2c-1)(d+e+c+1)(-d+e+c)(d-e+c)(d+e-c+1)]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & c-1 \\ g & g & 1 \end{Bmatrix} \\
 & = (2c+1) \left\{ d(d+1) - e(e+1) - c(c+1) + \frac{2c(c+1)[e(e+1) + g(g+1) - b(b+1)]}{a(a+1) + e(e+1) - d(d+1) - b(b+1)} \right\} \begin{Bmatrix} a & b & c \\ d & e & c \\ g & g & 1 \end{Bmatrix},
 \end{aligned} \tag{10.5.15}$$

where either $a \neq b$ and $d \neq e$, or $a \neq d$ and $b \neq e$. In particular,

$$\begin{aligned} \begin{Bmatrix} a & b & c \\ d & e & c+1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{Bmatrix} &= \left[\frac{2c+1}{c+(a-d)(2a+1)+(b-e)(2b+1)} - 1 \right] \\ &\times \left[\frac{c(2c-1)(d+e-c)(-d+e+c+1)(d-e+c+1)(d+e+c+2)}{(c+1)(2c+3)(d+e-c+1)(-d+e+c)(d-e+c)(d+e+c+1)} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & c-1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{Bmatrix}. \end{aligned} \quad (10.5.16)$$

10.6 Generating Function of the 9j Symbols

The following function of eighteen variables $\tau_{ik}, \tau'_{ik} (i, k = 1, 2, 3)$ [77]

$$\left[1 - \sum_{p,q=1}^3 a_{pq} - \sum_{\alpha=1}^6 b_{\alpha} \right]^{-1} \quad (10.6.1)$$

may be regarded as a generating function for the 9j symbols. Here a_{pq} and b_{α} are given by the arrays

$$(a_{pq}) = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \equiv \begin{pmatrix} \tau_{21} & \tau_{31} & \tau'_{12} & \tau'_{13} & \tau_{22} & \tau_{32} & \tau'_{11} & \tau'_{13} & \tau_{23} & \tau_{33} & \tau'_{11} & \tau'_{12} \\ \tau_{11} & \tau_{31} & \tau'_{22} & \tau'_{23} & \tau_{12} & \tau_{32} & \tau'_{21} & \tau'_{23} & \tau_{13} & \tau_{33} & \tau'_{21} & \tau'_{22} \\ \tau_{11} & \tau_{21} & \tau'_{32} & \tau'_{33} & \tau_{12} & \tau_{22} & \tau'_{31} & \tau'_{33} & \tau_{13} & \tau_{23} & \tau'_{31} & \tau'_{32} \end{pmatrix}, \quad (10.6.2)$$

$$(b_{\alpha}) = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \\ b_5 \\ b_6 \end{pmatrix} = \begin{pmatrix} \tau_{11} & \tau'_{11} & \tau_{32} & \tau'_{32} & \tau_{23} & \tau'_{23} \\ \tau_{31} & \tau'_{31} & \tau_{22} & \tau'_{22} & \tau_{13} & \tau'_{13} \\ \tau_{21} & \tau'_{21} & \tau_{12} & \tau'_{12} & \tau_{33} & \tau'_{33} \\ -\tau_{11} & \tau'_{11} & \tau_{22} & \tau'_{22} & \tau_{33} & \tau'_{33} \\ -\tau_{21} & \tau'_{21} & \tau_{32} & \tau'_{32} & \tau_{13} & \tau'_{13} \\ -\tau_{31} & \tau'_{31} & \tau_{12} & \tau'_{12} & \tau_{23} & \tau'_{23} \end{pmatrix}. \quad (10.6.3)$$

The elements b_{α} satisfy the equation

$$\sum_{\alpha=1}^6 b_{\alpha} = - \begin{vmatrix} \tau_{11} & \tau'_{11} & \tau_{12} & \tau'_{12} & \tau_{13} & \tau'_{13} \\ \tau_{21} & \tau'_{21} & \tau_{22} & \tau'_{22} & \tau_{23} & \tau'_{23} \\ \tau_{31} & \tau'_{31} & \tau_{32} & \tau'_{32} & \tau_{33} & \tau'_{33} \end{vmatrix} \quad (10.6.4)$$

The function (10.6.1) can be expanded into a power series whose expansion coefficients are proportional to the 9j symbols:

$$\frac{1}{1 - \sum_{p,q=1}^3 a_{pq} - \sum_{\alpha=1}^6 b_{\alpha}} = \sum_{r,r'} \left[\frac{\prod_{p=1}^3 (\sum_{q=1}^3 r_{pq} + 1)! (\sum_{q=1}^3 r'_{qp} + 1)!}{\prod_{p,q=1}^3 r_{pq}! r'_{pq}!} \right]^{\frac{1}{2}} \begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \prod_{p,q=1}^3 (\tau_{pq})^{r_{pq}} (\tau'_{pq})^{r'_{pq}}. \quad (10.6.5)$$

The exponents r_{pq} and r'_{pq} are the elements of some r symbol.

10.7 Asymptotic Expression for a 9j Symbol

If $a, b, c, d, e, f, g, h, j \gg 1$, we obtain

$$\begin{Bmatrix} a & b & c \\ d & e & f \\ g & h & j \end{Bmatrix} \approx \frac{2\Theta(B)(-1)^{a+c-e+g+j} \cos[(d-h)\delta_1 + (a-j)\delta_2 + (b-f)\delta_3]}{\pi \sqrt{B(2a+1)(2b+1)(2d+1)(2f+1)(2h+1)(2j+1)}} \quad (10.7.1)$$

In this equation

$$B = \begin{vmatrix} 1 & \cos \vartheta_3 & \cos \vartheta_1 \\ \cos \vartheta_3 & 1 & \cos \vartheta_2 \\ \cos \vartheta_1 & \cos \vartheta_2 & 1 \end{vmatrix}, \quad (10.7.2)$$

$$\Theta(B) = \begin{cases} 1 & \text{if } B > 0 \\ 0 & \text{if } B < 0 \end{cases} \quad (10.7.3)$$

The angles δ_1, δ_2 and δ_3 are determined by the relations

$$\begin{aligned}\cos \delta_1 &= \frac{\cos \vartheta_1 \cos \vartheta_3 - \cos \vartheta_2}{\sin \vartheta_1 \sin \vartheta_3}, & \sin \delta_2 &= \frac{\sin \vartheta_1}{\sin \vartheta_3} \\ \cos \delta_2 &= \frac{\cos \vartheta_2 \cos \vartheta_3 - \cos \vartheta_1}{\sin \vartheta_2 \sin \vartheta_3}, & \sin \delta_1 &= \frac{\sin \vartheta_2}{\sin \vartheta_3} \\ \cos \delta_3 &= \frac{\cos \vartheta_1 \cos \vartheta_2 - \cos \vartheta_3}{\sin \vartheta_1 \sin \vartheta_2}, & \sin \delta_3 &= \frac{\sin \vartheta_1}{\sin \vartheta_2}\end{aligned}\quad (10.7.4)$$

The angles ϑ_1, ϑ_2 and ϑ_3 may in turn be evaluated using the equations

$$\begin{aligned}\cos \vartheta_1 &= \frac{a(a+1)+b(b+1)-c(c+1)}{2\sqrt{a(a+1)b(b+1)}}, & 0 \leq \vartheta_1 \leq \pi, \\ \cos \vartheta_2 &= -\frac{d(d+1)-e(e+1)+f(f+1)}{2\sqrt{d(d+1)f(f+1)}}, & 0 \leq \vartheta_2 \leq \pi, \\ \cos \vartheta_3 &= \frac{-g(g+1)+h(h+1)+j(j+1)}{2\sqrt{h(h+1)j(j+1)}}, & 0 \leq \vartheta_3 \leq \pi.\end{aligned}\quad (10.7.5)$$

10.8 Explicit Forms of the 9j Symbols at Some Relations Between Arguments

10.8.1 Two Rows (or Columns) are Identical

In this case a 9j symbol vanishes unless the sum of all arguments is even [82], or

$$\begin{aligned}\begin{Bmatrix} a & b & c \\ a & b & c \\ g & h & j \end{Bmatrix} &= 0, \text{ if } g + h + j = 2k + 1, \\ \begin{Bmatrix} a & a & c \\ d & d & f \\ g & g & j \end{Bmatrix} &= 0, \text{ if } c + f + j = 2k + 1, \\ k &= 0, 1, \dots\end{aligned}\quad (10.8.1)$$

10.8.2 One Degenerate Triad

Let one argument of a triad be equal to the sum of the other two. For brevity, we shall refer to such a triad as a "degenerate" triad. If some 9j symbol has one degenerate triad, it may be expressed as

$$\begin{aligned}\begin{Bmatrix} a & b & c \\ d & e & f \\ a+d & h & j \end{Bmatrix} &= \frac{\Delta(h, j, a+d)}{\Delta(abc)\Delta(beh)\Delta(def)\Delta(cfj)} \left[\frac{(2a)!(2d)!}{(2a+2d+1)!} \right]^{\frac{1}{2}} \\ &\times \frac{(b+c-a)!(b+e-h)!(e+f-d)!(c+f-j)!(a+d+h+j+1)!}{(a+b+c+1)!(b+e+h+1)!(d+e+f+1)!(c+f+j+1)!} \\ &\times \sum_{xy} \frac{(-1)^{x+y}(2c-x)!(2e-y)!(j+f-c+x)!(h+b-e+y)!}{x!y!(c+f-j-x)!(b+e-h-y)!(c-e-a+h-x+y)!(e-c-d+j+x-y)!}\end{aligned}\quad (10.8.2)$$

This equation may be rewritten in the form of a sum involving the Clebsch-Gordan coefficients [83]

$$\begin{aligned}\begin{Bmatrix} a & b & c \\ d & e & f \\ a+d & h & j \end{Bmatrix} &= \frac{[(a+d+j-h)!(h+j+a+d+1)!]^{\frac{1}{2}}}{\Delta(abc)\Delta(def)\Delta(cfj)} \left[\frac{(2a)!(2d)!}{(2a+2d+1)!(2h+1)!} \right]^{\frac{1}{2}} \\ &\times \frac{(c+b-a)!(f+e-d)!(f+c-j)!}{(a+b+c+1)!(f+e+d+1)!(f+c+j+1)!} \\ &\times \sum_x (-1)^x \frac{(f+j-c+x)!(2c-x)!}{x!(f+c-j-x)!} \left[\frac{(b+a-c+x)!(e+d+c-j-x)!}{(b+c-a-x)!(e+j-c-d+x)!} \right]^{\frac{1}{2}} C_{b\ c-a\ x\ e\ j-d-c+x}^{h\ j-a-d}\end{aligned}\quad (10.8.3)$$

Equation (10.8.2) may also be rewritten in the quasi-trinomial form

$$\left(A^{(1)} - B^{(1)} - C^{(1)} \right)^{(k)} = \sum_{x,y} \binom{k}{x} \binom{k-x}{y} (-1)^{x+y} C^{(x)} B^{(y)} A^{(k-x-y)}, \quad (10.8.4)$$

where $A^{(x)}, B^{(y)}$ and $C^{(k-x-y)}$ are quasi-powers (Eq. (8.2.11)). Using Eq. (10.8.4), one gets [20]

$$\begin{aligned} & \left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ a+d & h & j \end{array} \right\} = (-1)^{b+e-h} \\ & \times \left[\frac{(2d)!}{(e-d+f)!(d-e+f)!(d+e-f)!(e+d+f+1)!(f-j+c)!(e+h-b)!(e-h+b)!(f+j-c)!} \right]^{\frac{1}{2}} \\ & \times \left[\frac{(a+b+c+1)^{(-1)(h+j-b-c+d)}(a-b+c)^{(-1)(b+j-h-c+d)}(b-c+a)^{(-1)(h+c-b-j+d)}(b+c-a)^{(1)(b+c-h-j+d)}}{(b+e+h+1)^{(2e+1)}(c+f+j+1)^{(2f+1)}(2a)^{(-1)(2d+1)}} \right]^{\frac{1}{2}} \\ & \times (2e)!(2f)! \left[(h+j-a-d)^{(1)} - \frac{(e-h+b)^{(1)}(b-e+h)^{(-1)}}{(2e)^{(1)}} - \frac{(f-j+c)^{(1)}(c-f+j)^{(-1)}}{(2f)^{(1)}} \right]^{(e+f-d)}. \end{aligned} \quad (10.8.5)$$

If one argument is equal to the difference of the other two from the same triad, the expression for the 9j symbol may be obtained from the above equations either by using the "mirror" symmetry properties (Sec. 10.4.3),

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ a-d & h & j \end{array} \right\} = i(-1)^{d+h-b-j+c} \left\{ \begin{array}{ccc} \bar{a} & b & c \\ d & e & f \\ \bar{a}+d & h & j \end{array} \right\}, \quad (10.8.6)$$

or by renaming and permuting the arguments:

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ a-d & h & j \end{array} \right\} \xrightarrow{a=d+g} \left\{ \begin{array}{ccc} d+g & b & c \\ d & e & f \\ g & h & j \end{array} \right\} = \left\{ \begin{array}{ccc} d & e & f \\ g & h & j \\ d+g & b & c \end{array} \right\}. \quad (10.8.7)$$

The equations of Sections 10.8.3-10.8.6 may be derived from Eqs. (10.8.2)-(10.8.5).

10.8.3 Two Degenerate Triads

In this case there exist five types of relations which cannot be transformed to one another by renaming and permuting the arguments [45, 104]. All these relations are given below

$$\begin{aligned} & \left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ a+d & b+e & j \end{array} \right\} = \frac{\Delta(a+db+ej)}{\Delta(abc)\Delta(def)\Delta(ejf)} \left[\frac{(2a)!(2b)!(2d)!(2e)!}{(2a+2b+1)!(2d+2e+1)!} \right]^{\frac{1}{2}} \\ & \times \frac{(a+b+d+e+j+1)!(a-b+c)!(d-e+f)!(c+f-j)!}{(a+b+c+1)!(d+e+f+1)!(c+f+j+1)!} \\ & \times \sum_z \frac{(-1)^z (2f-z)!(j+c-f+z)!}{z!(c+f-j-z)!(d+f-e-z)!(a-b-f+j+z)!}. \end{aligned} \quad (10.8.8)$$

Equation (10.8.8) may be rewritten in terms of the Clebsch-Gordan coefficients:

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ a+d & b+e & j \end{array} \right\} = \left[\frac{(2a)!(2b)!(2d)!(2e)!(a+b+d+e+j+1)!(a+d+e+b-j)!}{(2a+2d+1)!(2b+2e+1)!(a+b+c+1)!(a+b-c)!(d+e+f+1)!(d+e-f)!(2j+1)!} \right]^{\frac{1}{2}} \times C_{ca-bf d-e}^{ja-b+d-e}, \quad (10.8.9)$$

$$\left\{ \begin{array}{ccc} d+g & b & c \\ d & e & f \\ g & h & c+f \end{array} \right\} = (-1)^{d+h-b-f} \left[\frac{(2c)!(2f)!(2d)!(2g)!}{(2c+2f+1)!(2d+2g+1)!} \right]^{\frac{1}{2}} \frac{(d+e-f)!}{\Delta(d+gbc)\Delta(c+fgh)} \times \frac{(b+e-h)!(d+g+b-c)!(c+f+h-g)!}{\Delta(beh)\Delta(def)(d+e+f+1)!(b+e+h+1)!(d+g+b+c+1)!} \times \sum_x \frac{(2e-x)!(h+b-e+x)!(d+g+e+c-h-x)!}{x!(b+e-h-x)!(d+e-f-x)!}. \quad (10.8.10)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & e & f \\ a+d & h & j \end{array} \right\} = \frac{\Delta(a+b f j)\Delta(a+d h j)}{\Delta(b e h)\Delta(d e f)} \left[\frac{(2b)!(2d)!}{(2a+2b+1)!(2a+2d+1)!} \right]^{\frac{1}{2}} \times \frac{(a+b+f+j+1)!(a+d+h+j+1)!(h-b+e)!(e-d+f)!}{(f+j-a-b)!(h+j-a-d)!(b+e+h+1)!(d+e+f+1)!} \times \sum_x \frac{(-1)^x(a+b+e+d-j-x)!(j+f-a-b+x)!(j+h-a-d+x)!}{x!(a+b+f-j-x)!(a+d+h-j-x)!(j+e-a-b-d+x)!(2j+1+x)!}, \quad (10.8.11)$$

$$\left\{ \begin{array}{ccc} a & b & a-b \\ d & e & f \\ a-d & h & j \end{array} \right\} = (-1)^{b+f-d-h} \frac{[(2a-2b)!(2a-2d)!(2b)!(2d)!]^{\frac{1}{2}}}{\Delta(a-b f j)\Delta(a-d h j)\Delta(b e h)\Delta(d e f)} \times \frac{(b+f+j-a)!(d+h+j-a)!(h-b+e)!(e-d+f)!}{(a-b+f+j+1)!(a-d+h+j+1)!(b+e+h+1)!(d+e+f+1)!} \times \sum_x \frac{(x-1)!(b+e+d-a-j+x-1)!(a-d+h+j-x+1)!(a-b+f+j-x+1)!}{(2j+1-x)!(b+f-a-j+x-1)!(d+h-a-j+x-1)!(a+e+j-b-d-x+1)!}, \quad (10.8.12)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & e & f \\ g & h & a+b+f \end{array} \right\} = (-1)^{a+d-g} \frac{\Delta(a+b+f g h)}{\Delta(a d g)\Delta(b e h)\Delta(d e f)} \left[\frac{(2a)!(2b)!(2f)!}{(2a+2b+1)(2a+2b+2f+1)!} \right]^{\frac{1}{2}} \times \frac{(a+b+g+h+f+1)!(g-a+d)!(e-b+h)!(d+e-f)!}{(g+h-a-b-f)!(a+g+d+1)!(b+e+h+1)!(d+e+f+1)!}. \quad (10.8.13)$$

10.8.4 Three Degenerate Triads

Below we present the simplest expressions of 9j symbols

$$\left\{ \begin{matrix} a & b & a+b \\ d & e & d+e \\ g & h & g+h \end{matrix} \right\} = \frac{\Delta(a+bd+eg+h)}{\Delta(adg)\Delta(beh)} \cdot \frac{(a+b+e+d+g+h+1)!}{(a+d+g+1)!(b+e+h+1)!}$$

$$\times \left[\frac{(2a)!(2b)!(2d)!(2e)!(2g)!(2h)!}{(2a+2b+1)!(2d+2e+1)!(2g+2h+1)!} \right]^{\frac{1}{2}}, \quad (10.8.14)$$

$$\left\{ \begin{matrix} c+b & b & c \\ d & e & d+e \\ g & h & g+h \end{matrix} \right\} = (-1)^{b+e-h} \frac{\Delta(c+bdg)}{\Delta(cd+eg+h)\Delta(beh)} \cdot \frac{(d+e-c+g+h)!}{(b+e+h+1)!(d-b-c+g)!}$$

$$\times \left[\frac{(2b)!(2c)!(2d)!(2e)!(2g)!(2h)!}{(2b+2c+1)!(2d+2e+1)!(2g+2h+1)!} \right]^{\frac{1}{2}}, \quad (10.8.15)$$

$$\left\{ \begin{matrix} a & b & a+b \\ e+f & e & f \\ g & h & a+b+f \end{matrix} \right\} = (-1)^{a+e+f-g} \frac{\Delta(a+b+fgh)}{\Delta(age+f)\Delta(beh)} \cdot \frac{(a+b+g+h+f+1)!(e-b+h)!}{(g+h-a-b-f)!(b+e+h+1)!}$$

$$\times \frac{(e+g+f-a)!}{(a+e+f+g+1)!} \left[\frac{(2a)!(2b)!(2e)!}{(2a+2b+1)(2e+2f+1)!(2a+2b+2f+1)!} \right]^{\frac{1}{2}}, \quad (10.8.16)$$

$$\left\{ \begin{matrix} a & b & c \\ d & d+f & f \\ a+d & b+d+f & j \end{matrix} \right\} = \frac{\Delta(a+db+d+fj)}{\Delta(abc)\Delta(cfj)} \cdot \frac{(a+b+2d+f+j+1)!(a-b+c)!(j+c-f)!}{(a+b+c+1)!(c+f+j+1)!(a-b-f+j)!}$$

$$\times \left[\frac{(2a)!(2b)!(2f)!}{(2a+2d+1)!(2f+2d+1)(2b+2d+2f+1)!} \right]^{\frac{1}{2}}. \quad (10.8.17)$$

10.8.5 Four Degenerate Triads

$$\left\{ \begin{matrix} a & b & a+b \\ d & e & f \\ a+d & b+e & a+b+f \end{matrix} \right\} = \frac{\Delta(a+b+fa+db+e)}{\Delta(def)} \cdot \frac{(2a+2b+d+e+f+1)!}{(d+e+f+1)!}$$

$$\times \left[\frac{(2e)!(2d)!(2f)!}{(2a+2b+1)(2a+2d+1)!(2e+2b+1)!(2a+2b+2f+1)!} \right]^{\frac{1}{2}} \quad (10.8.18)$$

$$\left\{ \begin{matrix} c+b & b & c \\ d & b+h & d+b+h \\ g & h & g+h \end{matrix} \right\} = (-1)^{2b} \frac{\Delta(c+bdg)}{\Delta(cd+b+hg+h)} \cdot \frac{(2h+b+d+g-c)!}{(d+g-b-c)!}$$

$$\times \left[\frac{(2c)!(2d)!(2g)!}{(2b+2h+1)(2b+2c+1)!(2b+2d+2h+1)!(2g+2h+1)!} \right]^{\frac{1}{2}} \quad (10.8.19)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ a+g & b+h & f \\ g & h & a+b+f \end{array} \right\} = (-1)^{2a} \frac{\Delta(a+b+fg h)}{\Delta(a+gb+hf)} \cdot \frac{(a+b+g+h-f)!}{(-a-b+g+h-f)!} \\ \times \left[\frac{(2f)!(2g)!(2h)!}{(2a+2b+1)(2a+2b+2f+1)!(2a+2g+1)!(2b+2h+1)!} \right]^{\frac{1}{2}}, \quad (10.8.20)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & b+h & f \\ a+d & h & a+b+f \end{array} \right\} = \frac{\Delta(ha+da+b+f)}{\Delta(df b+h)} \cdot \frac{(2a+b+d+h+f+1)!(d+b+h-f)!}{(h+d+b+f+1)!(d+h-f-b)!} \\ \times \left[\frac{(2d)!(2f)!(2h)!}{(2a+2b+1)(2a+2d+1)!(2b+2h+1)!(2a+2b+2f+1)!} \right]^{\frac{1}{2}}, \quad (10.8.21)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ a+g & a+g+f & f \\ g & h & a+b+f \end{array} \right\} = (-1)^{2a} \frac{\Delta(a+b+fg h)}{\Delta(bha+g+f)} \cdot \frac{(a-b+f+g+h)!}{(g+h-a-b-f)!} \\ \times \left[\frac{(2b)!(2g)!}{(2a+2g+1)(2a+2b+1)(2a+2f+2g+1)!(2a+2b+2f+1)!} \right]^{\frac{1}{2}}, \quad (10.8.22)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & d+f & f \\ a+d & h & a+b+f \end{array} \right\} = \frac{\Delta(a+b+f, a+d, h)}{\Delta(b, d+f, h)} \cdot \frac{(2a+b+d+h+f+1)!(h-b+d+f)!}{(b+d+f+h+1)!(h+d-f-b)!} \\ \times \left[\frac{(2b)!(2d)!(2d)!}{(2a+2b+1)(2a+2d+1)!(2d+2f+1)!(2a+2b+2f+1)!} \right]^{\frac{1}{2}}, \quad (10.8.23)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & d+f & f \\ g & b+d+f & a+b+f \end{array} \right\} = (-1)^{a+d-g} \frac{\Delta(a+b+f, g, d+f+b)}{\Delta(adg)} \cdot \frac{(2b+2f+a+d+g+1)!}{(a+d+g+1)!} \\ \times \left[\frac{(2a)!(2d)!}{(2a+2b+1)(2d+2f+1)(2a+2b+2f+1)!(2d+2b+2f+1)!} \right]^{\frac{1}{2}}, \quad (10.8.24)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & b+h & f \\ a+b+f+h & h & a+b+f \end{array} \right\} = (-1)^{d-b-f-h} \frac{1}{\Delta(d, f, b+h)\Delta(a, d, a+b+f+h)} \\ \times \frac{(b+d+h-f)!}{(2a+d+b+f+h+1)!(d+b+h+f+1)!} \left[\frac{(2a)!(2f)!(2a+2b+2h+2f+1)!}{(2a+2b+1)(2a+2b+2f+1)(2b+2h+1)!} \right]^{\frac{1}{2}}, \quad (10.8.25)$$

$$\left\{ \begin{array}{ccc} a & b & a+b \\ e+f & e & f \\ a+e+f & b+e & j \end{array} \right\} = (-1)^{f+a+b-j} \frac{\Delta(j, b+e, a+e+f)}{\Delta(f, a+b, j)} \cdot \frac{(f+j+b+a+2e+1)!}{(f+a+b+j+1)!} \\ \times \left[\frac{(2f)!(2b)!(2b)!}{(2f+2e+1)(2e+2b+1)!(2a+2b+1)!(2f+2e+2a+1)!} \right]^{\frac{1}{2}}. \quad (10.8.26)$$

Note. Equation (10.8.26), as given in the source, does not reproduce the $9j$ symbol: direct numerical evaluation shows that its two sides differ for physical arguments. Solving for the square-root factor shows that it must contain a j -dependent factorial (cf. the analogous term in (10.8.9)), which is absent from the printed expression; a factor appears to have been dropped in the original. All the other explicit forms of this section have been verified numerically.

10.8.6 Five or Six Degenerate Triads

$$\left\{ \begin{array}{ccc} a & b & a+b \\ d & e & d+e \\ a+d & b+e & a+b+d+e \end{array} \right\} = \frac{1}{[(2a+2b+1)(2d+2e+1)(2a+2d+1)(2b+2e+1)]^{\frac{1}{2}}}. \quad (10.8.27)$$

In particular, from Eq. (10.8.27) it follows that

$$\left\{ \begin{matrix} a & b & a+b \\ a & b & a+b \\ 2a & 2b & 2a+2b \end{matrix} \right\} = \frac{1}{(2a+2b+1)[(4a+1)(4b+1)]^{\frac{1}{2}}} \quad (10.8.28)$$

$$\left\{ \begin{matrix} a & a & 2a \\ a & a & 2a \\ 2a & 2a & 4a \end{matrix} \right\} = \frac{1}{(4a+1)^2} \quad (10.8.29)$$

$$\begin{aligned} & \left\{ \begin{matrix} a & b & a+b \\ a+g & a+g+f & f \\ g & a+b+f+g & a+b+f \end{matrix} \right\} \\ &= \frac{(-1)^{2a}}{[(2a+2g+1)(2a+2b+1)(2a+2b+2f+1)(2a+2g+2f+1)]^{\frac{1}{2}}}. \end{aligned} \quad (10.8.30)$$

In particular,

$$\left\{ \begin{matrix} a & b & a+b \\ a+b & a+b+f & f \\ b & a+2b+f & a+b+f \end{matrix} \right\} = \frac{(-1)^{2a}}{(2a+2b+1)(2a+2b+2f+1)}. \quad (10.8.31)$$

10.9 Explicit Forms of the $9j$ Symbols for Special Values of the Arguments

10.9.1 One of Arguments Equals Zero

In this case a $9j$ symbol is reduced to a $6j$ symbol [54]:

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & 0 \end{matrix} \right\} = \delta_{cf}\delta_{gh} \frac{(-1)^{b+c+d+g}}{[(2c+1)(2g+1)]^{\frac{1}{2}}} \left\{ \begin{matrix} a & b & c \\ e & d & g \end{matrix} \right\} = \delta_{cf}\delta_{gh} [(2c+1)(2g+1)]^{-\frac{1}{2}} W(bcgd; ae). \quad (10.9.1)$$

Using the symmetry properties, we get

$$\begin{aligned} \left\{ \begin{matrix} 0 & c & c \\ g & e & b \\ g & d & a \end{matrix} \right\} &= \left\{ \begin{matrix} c & 0 & c \\ d & g & a \\ e & g & b \end{matrix} \right\} = \left\{ \begin{matrix} g & g & 0 \\ e & d & c \\ b & a & c \end{matrix} \right\} = \left\{ \begin{matrix} a & b & e \\ 0 & c & c \\ g & a & d \end{matrix} \right\} = \left\{ \begin{matrix} a & g & d \\ c & 0 & c \\ b & g & e \end{matrix} \right\} = \left\{ \begin{matrix} b & a & c \\ g & g & 0 \\ e & d & c \end{matrix} \right\} = \left\{ \begin{matrix} c & e & d \\ c & b & a \\ 0 & g & g \end{matrix} \right\} = \left\{ \begin{matrix} d & c & e \\ a & c & b \\ g & 0 & g \end{matrix} \right\} \\ &= \left\{ \begin{matrix} a & b & c \\ g & g & 0 \end{matrix} \right\} = \frac{(-1)^{b+d+c+g}}{[(2c+1)(2g+1)]^{\frac{1}{2}}} \left\{ \begin{matrix} a & b & c \\ e & d & g \end{matrix} \right\} = \frac{W(bcgd; ae)}{[(2c+1)(2g+1)]^{\frac{1}{2}}} \end{aligned} \quad (10.9.2)$$

In particular, if two arguments are equal to zero, one has

$$\left\{ \begin{matrix} a & b & c \\ d & 0 & f \\ g & h & 0 \end{matrix} \right\} = \delta_{df}\delta_{bh}\delta_{cf}\delta_{gh} \frac{(-1)^{a-b-c}}{(2b+1)(2c+1)} \quad (10.9.3)$$

If three arguments are equal to zero, one obtains

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ 0 & 0 & 0 \end{matrix} \right\} = \frac{\delta_{ad}\delta_{be}\delta_{cf}}{[(2a+1)(2b+1)(2c+1)]^{\frac{1}{2}}} \quad (10.9.4)$$

$$\left\{ \begin{matrix} 0 & b & c \\ d & 0 & f \\ g & h & 0 \end{matrix} \right\} = \delta_{bc}\delta_{bd}\delta_{bf}\delta_{bg}\delta_{bh} \frac{(-1)^{2b}}{(2b+1)^2} \quad (10.9.5)$$

10.9.2 One of Arguments Equals Unity [11]

$$\left\{ \begin{matrix} a & b & c \\ d & e & c \\ g & g & 1 \end{matrix} \right\} = (-1)^{b+d+g+c} 2 \frac{(a-d)(a+d+1) - (b-e)(b+e+1)}{[(2g+2)(2g+1)2g(2c+2)(2c+1)2c]^{\frac{1}{2}}} \left\{ \begin{matrix} a & b & c \\ e & d & g \end{matrix} \right\}, \quad (10.9.6)$$

$$\begin{aligned} & \left\{ \begin{matrix} a & b & c \\ d & e & c \\ g+1 & g & 1 \end{matrix} \right\} (-1)^{b+d+g+c} \left[\frac{(2g+3)(2g+2)(2g+1)(2c+2)(2c+1)2c}{2} \right]^{\frac{1}{2}} \\ &= [(b-e+g+1)(-b+e+g+1)(b+e+g+2)(b+e-g)]^{\frac{1}{2}} \left\{ \begin{matrix} a & d & g+1 \\ e & b & c \end{matrix} \right\} \\ &+ [(a-d+g+1)(-a+d+g+1)(a+d+g+2)(a+d-g)]^{\frac{1}{2}} \left\{ \begin{matrix} a & d & g \\ e & b & c \end{matrix} \right\}, \end{aligned} \quad (10.9.7)$$

$$\begin{aligned} & \left\{ \begin{matrix} a & b & c+1 \\ d & e & c \\ g+1 & g & 1 \end{matrix} \right\} [(c+1)(a-d)(a+d+1) - (g+1)(a-b)(a+b+1) + (g+1)(c+1)(g-c)] \\ & \times [(2g+3)(2g+2)(2g+1)(2c+3)(2c+2)(2c+1)]^{\frac{1}{2}} (-1)^{b+c+d+g} \\ &= (c-g)A_{a+\frac{1}{2}} \left(d, g + \frac{1}{2}; b, c + \frac{1}{2} \right) \left\{ \begin{matrix} a & d & g \\ e & b & c \end{matrix} \right\} + (c+1)A_{d+\frac{1}{2}} \left(a, g + \frac{1}{2}; e, c + \frac{1}{2} \right) \left\{ \begin{matrix} a & b & c+1 \\ e & d & g \end{matrix} \right\} \\ & - (g+1)A_{b+\frac{1}{2}} \left(a, c + \frac{1}{2}; e, g + \frac{1}{2} \right) \left\{ \begin{matrix} a & b & c \\ e & d & g+1 \end{matrix} \right\}. \end{aligned} \quad (10.9.8)$$

The quantity $A_q(\lambda, \mu; \sigma, \nu)$ is defined by Eq. (10.5.6).

10.9.3 One of Triads Equals (1/2, 1/2, 1)

In this case a 9j symbol may be expressed in terms of the 6j symbols [33]:

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ \frac{1}{2} & \frac{1}{2} & 1 \end{matrix} \right\} \left\{ \begin{matrix} c & f & 1 \\ \frac{1}{2} & \frac{1}{2} & g \end{matrix} \right\} = \frac{(-1)^{2g}}{3} \left\{ \begin{matrix} a & b & c \\ \frac{1}{2} & g & e \end{matrix} \right\} \left\{ \begin{matrix} e & d & f \\ \frac{1}{2} & g & a \end{matrix} \right\} - \frac{(-1)^{b+d-g}}{6(2c+1)} \left\{ \begin{matrix} a & b & c \\ e & d & \frac{1}{2} \end{matrix} \right\} \delta_{fc} \quad (10.9.9)$$

In addition, the 9j symbols in question may be reduced to the 3jm symbols [7]. If d, e, c are integer and $d+e+c$ is even, one has

$$[6(2c+1)(2d+1)(2e+1)]^{\frac{1}{2}} \left(\begin{matrix} c & d & e \\ 0 & 0 & 0 \end{matrix} \right) \left\{ \begin{matrix} a & b & c \\ d & e & c \\ \frac{1}{2} & \frac{1}{2} & 1 \end{matrix} \right\} = \left(\begin{matrix} a & b & c \\ \frac{1}{2} & \frac{1}{2} & -1 \end{matrix} \right) \quad (10.9.10)$$

If $d+e+c$ is odd, one gets

$$\begin{aligned} & \left(\begin{matrix} c+1 & d & e \\ 0 & 0 & 0 \end{matrix} \right) \left\{ \begin{matrix} a & b & c \\ d & e & c+1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{matrix} \right\} \\ &= (-1)^{b+e+\frac{1}{2}} \frac{[(d-a)(2a+1) + (e-b)(2b+1) + c+1]}{[6(c+1)(2c+1)(2c+3)(2d+1)(2e+1)]^{\frac{1}{2}}} \left(\begin{matrix} a & b & c \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{matrix} \right) \end{aligned} \quad (10.9.11)$$

$$\begin{aligned} & \left(\begin{matrix} c-1 & d & e \\ 0 & 0 & 0 \end{matrix} \right) \left\{ \begin{matrix} a & b & c \\ d & e & c-1 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{matrix} \right\} \\ &= (-1)^{b+e+\frac{1}{2}} \frac{[(d-a)(2a+1) + (e-b)(2b+1) - c]}{[6c(2c+1)(2c-1)(2d+1)(2e+1)]^{\frac{1}{2}}} \left(\begin{matrix} a & b & c \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{matrix} \right) \end{aligned} \quad (10.9.12)$$

10.10 Relations Between the Wigner $9j$ Symbols and Analogous Functions of Other Authors

$$\begin{aligned}
\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\} &= X \left(\begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right), && \text{(Fano [68],} \\
&= U \left(\begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right), && \text{Fano and Racah [16])} \\
&= (-1)^{a+e-c-h} S(abde; cfgh; j), && \text{(Arima et al. [82]} \\
&= \frac{\chi(abde; cf; gh; j)}{[(2c+1)(2f+1)(2g+1)(2h+1)]^{\frac{1}{2}}}, && \text{(Schwinger [65])} \\
&= [(2c+1)(2f+1)(2g+1)(2h+1)]^{-\frac{1}{2}} A \left(\begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right) && \text{(Kennedy and Cliff [84])}
\end{aligned}$$

10.11 Tables of Algebraic Formulas of the $9j$ Symbols

Tables 10.1-10.12 contain algebraic formulas for the $9j$ symbols represented in the form

$$\left\{ \begin{matrix} a+\lambda & b+\mu & c+\nu \\ a & b & c \\ \alpha & \beta & \gamma \end{matrix} \right\}$$

with

$$\begin{aligned}
0 &\leq \alpha \leq 3, & 0 \leq \beta \leq 2, & \gamma = 0, 1 \\
-\alpha &\leq \lambda \leq \alpha, & -\beta \leq \mu \leq \beta, & -\gamma \leq \nu \leq \gamma.
\end{aligned}$$

The formulas are given only for the $9j$ symbols with $\alpha \geq \beta$. The $9j$ symbols with $\alpha < \beta$ may be obtained using the symmetry properties.

Algebraic tables of the $9j$ symbols are also available in Ref. [122] for $\alpha, \beta = \frac{1}{2}; \gamma = 0, 1; \nu = 0, \pm 1; \lambda, \mu = \pm \frac{1}{2}$ and in Ref. [20] for $0 < \alpha \leq 3, 0 < \beta \leq 2; \gamma = 0, 1; 0 \leq \lambda \leq \alpha; 0 \leq \mu \leq \beta, \nu = 0, \pm 1$.

In the tables given below we use the following notations

$$\begin{aligned}
S &= a + b + c, \\
Z &= -c(c+1) + a(a+1) + b(b+1).
\end{aligned}$$

10.12 Tables of Numerical Values of the $9j$ Symbols

In Tables 10.13-10.14 we present numerical values of the $9j$ symbols

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \\ g & h & j \end{matrix} \right\}$$

with $(ghj) = (\frac{1}{2}, \frac{1}{2}, 0)$ or $(\frac{1}{2}, \frac{1}{2}, 1)$. Other arguments of these $9j$ symbols are integer or half-integer numbers such as

$$0 \leq a, b, c, d, e, f \leq 4$$

All numerical values are given in the form of rational fractions and decimals. Each table contains the $9j$ symbols with fixed c and f . In addition, all tables are divided into two groups. The first group contains the $9j$ symbols with integer c and f , while the second group contains those with half-integer c and f . The arrangement of tables inside each group is as follows:

c and f are integer (Tables 10.13)						c and f are half-integer (Tables 10.14)					
c	f	j	c	f	j	c	f	j	c	f	1
0	0	0	1	2	1	1/2	1/2	0	3/2	5/2	1
1	1	0	2	2	1	3/2	3/2	0	5/2	5/2	1
2	2	0	3	2	1	5/2	5/2	0	7/2	5/2	1
3	3	0	2	3	1	7/2	7/2	0	5/2	7/2	1
4	4	0	3	3	1	1/2	1/2	1	7/2	7/2	1
1	0	1	4	3	1	3/2	1/2	1			
0	1	1	3	4	1	1/2	3/2	1			
1	1	1	4	4	1	3/2	3/2	1			
2	1	1				5/2	3/2	1			

Tables of numerical values of the $9j$ symbols are also available in Refs. [120,113] for $g, h = \frac{1}{2}; j = 0, 1; a, b = 0, 1, 2, 3, 4; d, e = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \frac{7}{2}; c, f = 0, 1, 2, 3, 4, 5$ and in Ref. [123] for integer arguments $j = 1; a, b, c, f \leq 6; d, e \leq 11$. In Ref. [122] numerical values of the $9j$ symbols are given in the form of rational fractions; in Ref. [121] they are given in the form of rational fractions and decimals; and in Ref. [123] in the form of decimals.

10.13 12j Symbols

10.13.1 Some Remarks on the $3nj$ Symbols

The $6j$ and $9j$ symbols represent special cases of the $3nj$ symbols for $n = 2$ and 3 , respectively. The $3nj$ symbols of higher orders ($n \geq 4$) are encountered in some special changes of the coupling scheme of five and more angular momenta. They are proportional to the coefficients of unitary transformations connecting the state vectors which correspond to different coupling schemes. The $3nj$ symbols are invariants with respect to rotations in three-dimensional space. If $n \geq 4$, there exist different kinds of the $3nj$ symbols which are not reduced to products of the $3nj$ symbols of lower order. Among them one can distinguish the $3nj$ symbols of the first and second kinds which may be expressed as single sums of products of the $6j$ symbols. Using the notations of Ref. [34], the $3nj$ symbols of the first and second kinds may be represented in the following forms:

(a) *The $3nj$ symbols of the first kind*

$$\left\{ \begin{array}{cccc} j_1 & j_2 & \cdots & j_n \\ l_1 & l_2 & \cdots & l_n \\ k_1 & k_2 & \cdots & k_n \end{array} \right\} = \sum_x (2x+1)(-1)^{R_n+(n-1)x} \times \left\{ \begin{array}{ccc} j_1 & k_1 & x \\ k_2 & j_2 & l_1 \end{array} \right\} \left\{ \begin{array}{ccc} j_2 & k_2 & x \\ k_3 & j_3 & l_2 \end{array} \right\} \cdots \left\{ \begin{array}{ccc} j_{n-1} & k_{n-1} & x \\ k_n & j_n & l_{n-1} \end{array} \right\} \left\{ \begin{array}{ccc} j_n & k_n & x \\ j_1 & k_1 & l_n \end{array} \right\}. \quad (10.13.1)$$

(b) *The $3nj$ symbols of the second kind*

$$\left[\begin{array}{cccc} j_1 & j_2 & \cdots & j_n \\ l_1 & l_2 & \cdots & l_n \\ k_1 & k_2 & \cdots & l_n \end{array} \right] = \sum_x (2x+1)(-1)^{R_n+nx} \times \left\{ \begin{array}{ccc} j_1 & k_1 & x \\ k_2 & j_2 & l_1 \end{array} \right\} \left\{ \begin{array}{ccc} j_2 & k_2 & x \\ k_3 & j_3 & l_2 \end{array} \right\} \cdots \left\{ \begin{array}{ccc} j_{n-1} & k_{n-1} & x \\ k_n & j_n & l_{n-1} \end{array} \right\} \left\{ \begin{array}{ccc} j_n & k_n & x \\ k_1 & j_1 & l_n \end{array} \right\}. \quad (10.13.2)$$

In these equations $R_n = \sum_{i=1}^n (j_i + l_i + k_i)$.

If $n \geq 5$, there appear the $3nj$ symbols of other kinds (third, fourth, etc.). They may be represented by single sums of more complex products of the $6j$ and $9j$ symbols. The amount of different $3nj$ symbols rapidly increases with n . For example, there exist five different $15j$ symbols, eighteen different $18j$ symbols, etc.

Here we restrict ourselves to the consideration of the $12j$ symbols because the $15j$ symbols and symbols of higher orders are rarely used in applications. So far, the properties of the $12j$ symbols have been examined [17, 44, 45, 74, 137 – 139, 141] less thoroughly than the properties of the $6j$ and $9j$ symbols. That is why our consideration is rather brief.

10.13.2 12j Symbols of the First Kind (12j(I)-Symbols)

(a) 12j(I)-symbol

A 12j(I)-symbol may be represented by the following invariant sum of products of eight 3jm symbols

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \sum_{\alpha_i \beta_{ik} \gamma_i} (-1)^{a_4 + \alpha_4 + c_4 + \gamma_4} \times \begin{pmatrix} a_1 & a_2 & b_{12} \\ \alpha_1 & \alpha_2 & \beta_{12} \end{pmatrix} \begin{pmatrix} a_2 & a_3 & b_{23} \\ \alpha_2 & \alpha_3 & \beta_{23} \end{pmatrix} \begin{pmatrix} a_3 & a_4 & b_{34} \\ \alpha_3 & \alpha_4 & \beta_{34} \end{pmatrix} \begin{pmatrix} a_4 & c_1 & b_{41} \\ -\alpha_4 & \gamma_1 & \beta_{41} \end{pmatrix} \times \begin{pmatrix} c_1 & c_2 & b_{12} \\ \gamma_1 & \gamma_2 & \beta_{12} \end{pmatrix} \begin{pmatrix} c_2 & c_3 & b_{23} \\ \gamma_2 & \gamma_3 & \beta_{23} \end{pmatrix} \begin{pmatrix} c_3 & c_4 & b_{34} \\ \gamma_3 & \gamma_4 & \beta_{34} \end{pmatrix} \begin{pmatrix} c_4 & a_1 & b_{41} \\ -\gamma_4 & \alpha_1 & \beta_{41} \end{pmatrix}. \quad (10.13.3)$$

Equation (10.13.3) unambiguously fixes the absolute value and phase of the 12j(I) symbol. The 12j(I) symbols are real.

(b) conditions on the arguments

The arguments of a 12j(I) symbol must satisfy the following conditions

- (i) All arguments are integers or half-integers and non-negative.
- (ii) The 12j(I) symbol vanishes unless the triangular conditions (see Sec. 8.1.1) are fulfilled for all the triads $(a_1 b_{12} a_2)$, $(a_2 b_{23} a_3)$, $(a_3 b_{34} a_4)$, $(a_4 b_{41} c_1)$, $(c_1 b_{12} c_2)$, $(c_2 b_{23} c_3)$, $(c_3 b_{34} c_4)$ and $(c_4 b_{41} a_1)$.
- (iii) The 12j(I) symbol vanishes unless the tetragonal conditions are fulfilled for two tetrads of arguments, $(a_1 c_1 a_3 c_3)$ and $(a_2 c_2 a_4 c_4)$. The tetragonal conditions for the tetrad $(j_1 j_2 j_3 j_4)$ mean that $j_1 + j_2 + j_3 + j_4$ is integer and $j_1 \leq j_2 + j_3 + j_4$, $j_2 \leq j_1 + j_3 + j_4$, $j_3 \leq j_1 + j_2 + j_4$, $j_4 \leq j_1 + j_2 + j_3$.

(c) Orthogonality and normalisation

The 12j(I) symbols satisfy the orthogonality and normalisation relations

$$\sum_{a_2 a_3 a_4} (2a_2 + 1)(2a_3 + 1)(2a_4 + 1) \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c'_2 & c'_3 & c'_4 \end{array} \right\} = \frac{\delta_{c'_2 c_2} \delta_{c'_3 c_3} \delta_{c'_4 c_4}}{(2c_2 + 1)(2c_3 + 1)(2c_4 + 1)} \{c_1 b_{12} c_2\} \{c_2 b_{23} c_3\} \{c_3 b_{34} c_4\} \{c_4 b_{41} a_1\} \quad (10.13.4)$$

$$\sum_{c_3 b_{41} c_4} (2c_3 + 1)(2b_{41} + 1)(2c_4 + 1) \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a'_{22} & a'_3 & a_4 \\ b'_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \frac{\delta_{a'_2 a_2} \delta_{a'_3 a_3} \delta_{b'_{12} b_{12}}}{(2a_2 + 1)(2a_3 + 1)(2b_{12} + 1)} \{a_1 b_{12} a_2\} \{a_2 b_{23} a_3\} \{a_3 b_{34} a_4\} \{c_1 b_{12} c_2\} \quad (10.13.5)$$

(d) Sum representations

The 12j(I) symbols may be represented by

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \sum_x (-1)^{S-x} (2x + 1) \times \left\{ \begin{array}{ccc} a_1 & a_2 & b_{12} \\ c_2 & c_1 & x \end{array} \right\} \left\{ \begin{array}{ccc} a_2 & a_3 & b_{23} \\ c_3 & c_2 & x \end{array} \right\} \left\{ \begin{array}{ccc} a_3 & a_4 & b_{34} \\ c_4 & c_3 & x \end{array} \right\} \left\{ \begin{array}{ccc} a_4 & c_1 & b_{41} \\ a_1 & c_4 & x \end{array} \right\}, \quad (10.13.6)$$

where $S = \sum_{i=1}^4 (a_i + c_i) + b_{12} + b_{23} + b_{34} + b_{41}$,

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = (-1)^{a_1 - a_3 - c_1 + c_3} \sum_x (2x+1) \times \left\{ \begin{array}{ccc} b_{23} & a_2 & a_3 \\ c_2 & b_{12} & c_1 \\ c_3 & a_1 & x \end{array} \right\} \left\{ \begin{array}{ccc} b_{34} & a_3 & a_4 \\ c_1 & b_{41} & x \end{array} \right\} \left\{ \begin{array}{ccc} b_{34} & c_3 & c_4 \\ a_1 & b_{41} & x \end{array} \right\} \quad (10.13.7)$$

(e) *Symmetry properties*

To discuss the symmetry properties it is convenient to separate all arguments of a $12j(\text{I})$ symbol into two groups: $(b_{12}b_{23}b_{34}b_{41})$ and $(a_1a_2a_3a_4c_1c_2c_3c_4)$. Cyclic permutations of the arguments or inversion of argument order simultaneously in both groups leave the $12j(\text{I})$ symbol unchanged. These symmetry properties relate 16 formally different $12j(\text{I})$ symbols

$$\begin{aligned} & \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \left\{ \begin{array}{cccc} a_2 & a_3 & a_4 & c_1 \\ b_{23} & b_{34} & b_{41} & b_{12} \\ c_2 & c_3 & c_4 & a_1 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} a_3 & a_4 & c_1 & c_2 \\ b_{34} & b_{41} & b_{12} & b_{23} \\ c_3 & c_4 & a_1 & a_2 \end{array} \right\} = \left\{ \begin{array}{cccc} a_4 & c_1 & c_2 & c_3 \\ b_{41} & b_{12} & b_{23} & b_{34} \\ c_4 & a_1 & a_2 & a_3 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} a_4 & a_3 & a_2 & a_1 \\ b_{34} & b_{23} & b_{12} & b_{41} \\ c_4 & c_3 & c_2 & c_1 \end{array} \right\} = \left\{ \begin{array}{cccc} c_1 & a_4 & a_3 & a_2 \\ b_{41} & b_{34} & b_{23} & b_{12} \\ a_1 & c_4 & c_3 & c_2 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} c_2 & c_1 & a_4 & a_3 \\ b_{12} & b_{41} & b_{34} & b_{23} \\ a_2 & a_1 & c_4 & c_3 \end{array} \right\} = \left\{ \begin{array}{cccc} c_3 & c_2 & c_1 & a_4 \\ b_{23} & b_{12} & b_{41} & b_{34} \\ a_3 & a_2 & a_1 & c_4 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} c_1 & c_2 & c_3 & c_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ a_1 & a_2 & a_3 & a_4 \end{array} \right\} = \left\{ \begin{array}{cccc} c_2 & c_3 & c_4 & a_1 \\ b_{23} & b_{34} & b_{41} & b_{12} \\ a_2 & a_3 & a_4 & c_1 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} c_3 & c_4 & a_1 & a_2 \\ b_{34} & b_{41} & b_{12} & b_{23} \\ a_3 & a_4 & c_1 & c_2 \end{array} \right\} = \left\{ \begin{array}{cccc} c_4 & a_1 & a_2 & a_3 \\ b_{41} & b_{12} & b_{23} & b_{34} \\ a_4 & c_1 & c_2 & c_3 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} c_4 & c_3 & c_2 & c_1 \\ b_{34} & b_{23} & b_{12} & b_{41} \\ a_4 & a_3 & a_2 & a_1 \end{array} \right\} = \left\{ \begin{array}{cccc} a_1 & c_4 & c_3 & c_2 \\ b_{41} & b_{34} & b_{23} & b_{12} \\ c_1 & a_4 & a_3 & a_2 \end{array} \right\} \\ & = \left\{ \begin{array}{cccc} a_2 & a_1 & c_4 & c_3 \\ b_{12} & b_{41} & b_{34} & b_{23} \\ c_2 & c_1 & a_4 & a_3 \end{array} \right\} = \left\{ \begin{array}{cccc} a_3 & a_2 & a_1 & c_4 \\ b_{23} & b_{12} & b_{41} & b_{34} \\ c_3 & c_2 & c_1 & a_4 \end{array} \right\}. \end{aligned} \quad (10.13.9)$$

(f) *Recursion relations*

These relations are very useful for the evaluation of the $12j(\text{I})$ symbols. Some of them may be derived from the properties of the $15j$ symbols. For example,

$$\begin{aligned} & (-1)^{a_1 - a_4 - c_1 + c'_4} \sum_{b'_{41}} (2b'_{41} + 1) \left\{ \begin{array}{ccc} c_1 & a_4 & b'_{41} \\ \lambda & b_{41} & a'_4 \end{array} \right\} \left\{ \begin{array}{ccc} a_1 & c'_4 & b'_{41} \\ \lambda & b_{41} & c_4 \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b'_{41} \\ c_1 & c_2 & c_3 & c'_4 \end{array} \right\} \\ & = (-1)^{a_3 + a'_4 - c_3 - c_4} \sum_{b'_{34}} (2b'_{34} + 1) \left\{ \begin{array}{ccc} a_3 & a'_4 & b'_{34} \\ \lambda & b_{34} & a_4 \end{array} \right\} \left\{ \begin{array}{ccc} c_3 & c_4 & b'_{34} \\ \lambda & b_{34} & c'_4 \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a'_4 \\ b_{12} & b_{23} & b'_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\}. \end{aligned} \quad (10.13.10)$$

Here λ is any integer or half-integer non-negative number ($\lambda = \frac{1}{2}, 1, \frac{3}{2}, 2$, etc.). Substituting explicit forms of the $6j$ symbols for a given λ , we may obtain a recursion equation which relates the $12j(\text{I})$ symbols with arguments $c'_4 = c_4 \pm \lambda$, $b'_{41} = b_{41} \pm \lambda$, $a'_4 = a_4 \pm \lambda$, $b'_{34} = b_{34} \pm \lambda$. For example, when $\lambda = \frac{1}{2}$, $a'_4 = a_4 - \frac{1}{2}$, and

$c'_4 = c_4 + \frac{1}{2}$ we have

$$\begin{aligned}
& (2b_{34} + 1) \left\{ \left[(-a_1 + c_4 + b_{41} + 1)(a_1 + c_4 + b_{41} + 2) \left(a_4 - c_1 + b_{41} + \frac{1}{2} \right) \left(a_4 + c_1 + b_{41} + \frac{3}{2} \right) \right]^{\frac{1}{2}} \right. \\
& \quad \times \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} + \frac{1}{2} \\ c_1 & c_2 & c_3 & c_4 + \frac{1}{2} \end{array} \right\} \\
& \quad + \left[(a_1 - c_4 + b_{41})(a_1 + c_4 - b_{41} + 1) \left(-a_4 + c_1 + b_{41} + \frac{1}{2} \right) \left(a_4 + c_1 - b_{41} + \frac{1}{2} \right) \right]^{\frac{1}{2}} \\
& \quad \times \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} - \frac{1}{2} \\ c_1 & c_2 & c_3 & c_4 + \frac{1}{2} \end{array} \right\} \Bigg\} \\
& = (2b_{41} + 1) \left\{ \left[(a_3 - a_4 + b_{34} + 1)(a_3 + a_4 - b_{34}) \left(c_3 - c_4 + b_{34} + \frac{1}{2} \right) \left(c_3 + c_4 - b_{34} + \frac{1}{2} \right) \right]^{\frac{1}{2}} \right. \\
& \quad \times \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 - \frac{1}{2} \\ b_{12} & b_{23} & b_{34} + \frac{1}{2} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} \\
& \quad + \left[(-a_3 + a_4 + b_{34})(a_3 + a_4 + b_{34} + 1) \left(-c_3 + c_4 + b_{34} + \frac{1}{2} \right) \left(c_3 + c_4 + b_{34} + \frac{3}{2} \right) \right]^{\frac{1}{2}} \\
& \quad \times \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 - \frac{1}{2} \\ b_{12} & b_{23} & b_{34} - \frac{1}{2} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} \Bigg\}. \tag{10.13.11}
\end{aligned}$$

Another recursion relation is written as

$$\begin{aligned}
& (-1)^{a_1 + b'_{41} - c_3 - b_{34}} \sum_{c'_4} (2c'_4 + 1) \left\{ \begin{array}{ccc} a_1 & c'_4 & b'_{41} \\ \lambda & b_{41} & c_4 \end{array} \right\} \left\{ \begin{array}{ccc} c_3 & b_{34} & c'_4 \\ \lambda & c_4 & b'_{34} \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b'_{41} \\ c_1 & c_2 & c_3 & c'_4 \end{array} \right\} \\
& = (-1)^{c_1 + b_{41} - a_3 - b_{34}} \sum_{a'_4} (2a'_4 + 1) \left\{ \begin{array}{ccc} c_1 & a_4 & b'_{41} \\ \lambda & b_{41} & a'_4 \end{array} \right\} \left\{ \begin{array}{ccc} a_3 & b_{34} & a_4 \\ \lambda & a'_4 & b'_{34} \end{array} \right\} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a'_4 \\ b_{12} & b_{23} & b'_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\}. \tag{10.13.12}
\end{aligned}$$

(g) *Explicit forms of the 12j(I) symbols at some special relations between arguments*

may be obtained from Eqs. (10.13.6) and (10.13.7). Here we give some of such forms.

If some argument in any tetrad is equal to the sum of three other arguments from the same tetrad, the sum in Eq. (10.13.7) contains only one term. For example,

$$\begin{aligned}
& \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & a_1 + a_3 + c_1 & c_4 \end{array} \right\} = (-1)^{a_1 + b_{41} - c_4 + c_1 + b_{12} - c_2} \frac{(2a_1)!(2a_3)!(2c_1)!}{(2a_1 + 2a_3 + 2c_1 + 1)!} \\
& \quad \times \frac{D(a_1 a_2 b_{12}) D(c_1 c_2 b_{12}) D(a_3 a_2 b_{23}) D(a_3 a_4 b_{34}) D(a_1 c_4 b_{41}) D(c_1 a_4 b_{41})}{D(a_1 + a_3 + c_1 c_2 b_{23}) D(a_1 + a_3 + c_1 c_4 b_{34})}. \tag{10.13.13}
\end{aligned}$$

The quantity $D(abc)$ is defined by

$$D(abc) = \left[\frac{(-a + b + c)!}{(a + b + c + 1)!(a - b + c)!(a + b - c)!} \right]^{\frac{1}{2}} \tag{10.13.14}$$

Equation (10.13.7) for the 12j(I) symbol may also be reduced to one term. For example,

$$\begin{aligned}
& \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & a_1 + b_{34} + b_{41} & c_4 \end{array} \right\} = \left[\frac{(2b_{34})!(2b_{41})!}{(2b_{34} + 2b_{41})!(2a_1 + 2b_{41} + 1)!} \right]^{\frac{1}{2}} \\
& \quad \times \frac{D(b_{34} a_3 a_4) D(b_{41} c_1 a_4)}{D(b_{34} + b_{41} c_1 a_3)} \left\{ \begin{array}{ccc} b_{23} & a_2 & a_3 \\ c_2 & b_{12} & c_1 \end{array} \right\} \delta_{c_4 a_1 + b_{41}} \tag{10.13.15}
\end{aligned}$$

Similar expressions may be obtained from Eq. (10.13.15) using the symmetries of the 12j(I) symbols (see Eqs. (10.13.9)).

If arguments a_i and c_i satisfy the equalities $a_i = b_{ik} \pm a_k, c_i = b_{ik} \pm c_k, (k = i \pm 1)$ one obtains

$$\left\{ \begin{array}{cccc} b_{12} + a_2 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ b_{12} - c_2 & c_2 & c_3 & c_4 \end{array} \right\} = \frac{(-1)^{2c_3+b_{23}-b_{34}-a_2-a_4}}{2a_2+2c_2+1} \left\{ \begin{array}{ccc} a_3 & c_3 & a_2+c_2 \\ c_4 & a_4 & b_{34} \end{array} \right\} \\ \times \left[\frac{(2a_2)!(2c_2)!(2b_{12}-2c_2)!}{(2b_{12}+1)(2b_{12}+2a_2+1)!} \right]^{\frac{1}{2}} \frac{D(a_2a_3b_{23})D(c_2c_3b_{23})D(b_{12}-c_2b_{41}a_4)D(a_2+c_2c_4a_4)}{D(a_2+c_2c_3a_3)D(b_{12}+a_2c_4b_{41})} \quad (10.13.16)$$

(h) *Explicit forms of the 12j(I) symbols for special values of arguments*

If one argument is equal to zero, the 12j(I) symbol is reduced to a 9j symbol or to a product of two 6j symbols:

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & 0 \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \frac{\delta_{a_1c_4}\delta_{a_4c_1}}{\sqrt{(2a_1+1)(2c_1+1)}} \left\{ \begin{array}{ccc} a_1 & b_{12} & a_2 \\ c_3 & c_2 & b_{23} \\ b_{34} & c_1 & a_3 \end{array} \right\}, \quad (10.13.17)$$

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & 0 \end{array} \right\} = (-1)^{b_{12}+b_{23}+b_{34}+b_{41}-a_2-c_2} \frac{1}{\sqrt{(2b_{41}+1)(2b_{34}+1)}} \\ \times \left\{ \begin{array}{ccc} b_{41} & a_2 & b_{12} \\ c_2 & c_1 & a_4 \end{array} \right\} \left\{ \begin{array}{ccc} a_2 & a_3 & b_{23} \\ b_{34} & c_2 & a_4 \end{array} \right\}. \quad (10.13.18)$$

Simple expressions for the 12j(I) symbols may also be obtained, if two of four arguments from one tetrad are equal to $\frac{1}{2}$, i.e., in the cases: (1) $a_i = c_i = \frac{1}{2} (i = 1, 2, 3, 4)$; (2) $a_i = a_{i\pm 2} = \frac{1}{2}$; (3) $c_i = c_{i\pm 2} = \frac{1}{2}$; (4) $c_i = a_{i\pm 2} = \frac{1}{2} (i \pm 2 \leq 4, i - 2 \geq 1)$. For example, when $a_4 = c_4 = \frac{1}{2}$, one has

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & \frac{1}{2} \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & \frac{1}{2} \end{array} \right\} = \frac{\delta_{a_1c_1}\delta_{a_2c_2}\delta_{a_3c_3}}{2(2a_1+1)(2a_2+1)(2a_3+1)} \\ + (-1)^c \frac{D(\frac{1}{2}a_1b_{41})D(\frac{1}{2}c_1b_{41})D(\frac{1}{2}a_3b_{34})D(\frac{1}{2}c_3b_{34})}{2D(1a_1c_1)D(1a_3c_3)} \left\{ \begin{array}{ccc} a_1 & a_2 & b_{12} \\ c_2 & c_1 & 1 \end{array} \right\} \left\{ \begin{array}{ccc} a_2 & a_3 & b_{23} \\ c_3 & c_2 & 1 \end{array} \right\}, \quad (10.13.19)$$

where $\varepsilon = a_2 + c_2 + b_{12} + b_{23} + b_{34} + b_{41}$. If $a_1 = c_3 = \frac{1}{2}$, then

$$\left\{ \begin{array}{cccc} \frac{1}{2} & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & \frac{1}{2} & c_4 \end{array} \right\} = (-1)^{a_2+c_2+1} \frac{\delta_{a_3c_1}}{2(2a_3+1)} \left\{ \begin{array}{ccc} a_2 & a_3 & b_{23} \\ c_2 & \frac{1}{2} & b_{12} \end{array} \right\} \\ \times \left\{ \frac{(-1)^{a_4-c_4}}{2b_{34}+1} \delta_{b_{34}b_{41}} + \frac{(-1)^{b_{34}+b_{41}-g+1}}{2D^2(\frac{1}{2}a_3g)} \left[\frac{(2a_3-1)!}{(2a_3+2)!} \right]^{\frac{1}{2}} \frac{D(\frac{1}{2}b_{41}c_4)D(\frac{1}{2}b_{34}c_4)}{D(1b_{34}b_{41})} \left\{ \begin{array}{ccc} b_{34} & b_{41} & 1 \\ a_3 & a_3 & a_4 \end{array} \right\} \right\} \\ + (-1)^{b_{34}+b_{41}+2g+1} \frac{D(\frac{1}{2}b_{41}c_4)D(\frac{1}{2}b_{34}c_4)D(1a_3c_1)}{D(\frac{1}{2}c_1g)D(\frac{1}{2}a_3g)D(1b_{34}b_{41})} \\ \times \left\{ \begin{array}{ccc} b_{23} & a_2 & a_3 \\ \frac{1}{2} & g & b_{12} \end{array} \right\} \left\{ \begin{array}{ccc} b_{12} & c_2 & c_1 \\ \frac{1}{2} & g & b_{23} \end{array} \right\} \left\{ \begin{array}{ccc} b_{34} & b_{41} & 1 \\ c_1 & a_3 & a_4 \end{array} \right\}, \quad (10.13.20)$$

where g is an arbitrary parameter which satisfies the triangular conditions for appropriate triads of the 6j symbols in Eq. (10.13.20). The expressions for the 12j(I) symbols are more simplified in the cases: (i) $a_i = a_{i\pm 2} = \frac{1}{2}$ and $c_i = c_{i\pm 2}$; (ii) $c_i = c_{i\pm 2} = \frac{1}{2}$ and $a_i = a_{i\pm 2}$; (iii) $a_i = c_{i\pm 2} = \frac{1}{2}$ and $c_i = a_{i\pm 2}$. In addition, if $a_3 + b_{12} + c_2$ is even, then

$$\left\{ \begin{array}{cccc} \frac{1}{2} & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ a_3 & c_2 & \frac{1}{2} & c_4 \end{array} \right\} = (-1)^{c_4-a_4+a_2+c_2+1} \frac{\delta_{b_{34}b_{41}}}{2(2a_3+1)(2b_{34}+1)} \left\{ \begin{array}{ccc} b_{23} & a_2 & a_3 \\ b_{12} & c_2 & \frac{1}{2} \end{array} \right\} \\ + \frac{(-1)^{b_{34}+b_{41}+2a_3}}{2\sqrt{(2a_3+1)(2c_2+1)(2b_{12}+1)}} \frac{D(\frac{1}{2}b_{34}c_4)D(\frac{1}{2}b_{41}c_4)}{D(1b_{34}b_{41})} \left\{ \begin{array}{ccc} b_{34} & b_{41} & 1 \\ a_3 & a_3 & a_4 \end{array} \right\} \frac{\left(\begin{array}{ccc} a_2 & a_3 & b_{23} \\ \frac{1}{2} & -1 & \frac{1}{2} \end{array} \right)}{\left(\begin{array}{ccc} a_3 & c_2 & b_{12} \\ 0 & 0 & 0 \end{array} \right)} \quad (10.13.21)$$

and if $a_3 + b_{12} + c_2$ is odd, one has

$$\left\{ \begin{array}{cccc} \frac{1}{2} & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ a_3 + 1 & c_2 & \frac{1}{2} & c_4 \end{array} \right\} = (-1)^{a_2+a_3-a_4+b_{12}+b_{41}+\frac{3}{2}} \frac{D(\frac{1}{2}b_{34}c_4) D(\frac{1}{2}b_{41}c_4) D(a_3a_4b_{34})}{D(a_3 + 1a_4b_{41})} \\ \times \frac{(c_2 - b_{23})(2b_{23} + 1) + (b_{12} - a_2)(2a_2 + 1) + a_3 + 1}{(2a_3 + 1)(2a_3 + 2)(2a_3 + 3)\sqrt{2(2c_2 + 1)(2b_{12} + 1)}} \frac{\begin{pmatrix} a_2 & a_3 & b_{23} \\ -\frac{1}{2} & 0 & \frac{1}{2} \end{pmatrix}}{\begin{pmatrix} a_3 + 1 & c_2 & b_{12} \\ 0 & 0 & 0 \end{pmatrix}} \quad (10.13.22)$$

$$\left\{ \begin{array}{cccc} \frac{1}{2} & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ a_3 - 1 & c_2 & \frac{1}{2} & c_4 \end{array} \right\} = (-1)^{a_2+a_3-a_4+b_{12}+b_{41}+\frac{3}{2}} \frac{D(\frac{1}{2}b_{34}c_4) D(\frac{1}{2}b_{41}c_4) D(a_3 - 1a_4b_{34})}{D(a_3a_4b_{34})} \\ \times \frac{(c_2 - b_{23})(2b_{23} + 1) + (b_{12} - a_2)(2a_2 + 1) - a_3}{(2a_3 - 1)(2a_3)(2a_3 + 1)\sqrt{2(2c_2 + 1)(2b_{12} + 1)}} \frac{\begin{pmatrix} a_2 & a_3 & b_{23} \\ -\frac{1}{2} & 0 & \frac{1}{2} \end{pmatrix}}{\begin{pmatrix} a_3 - 1 & b_{12} & c_2 \\ 0 & 0 & 0 \end{pmatrix}} \quad (10.13.23)$$

(i) Relations between the $12j(I)$ symbols introduced in this book and analogous functions of other authors are given by

$$\left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} = \left\{ \begin{array}{cccc} b_{12} & a_1 & a_2 & a_3 \\ c_1 & b_{41} & a_4 & c_3 \\ c_2 & c_4 & b_{23} & b_{34} \end{array} \right\} \\ \text{(Jahn and Hope [79])} \\ = (-1)^{2c_1} \left\{ \begin{array}{cccc} a_1 & a_2 & a_3 & a_4 \\ b_{12} & b_{23} & b_{34} & b_{41} \\ c_1 & c_2 & c_3 & c_4 \end{array} \right\} 1 \\ \text{(Elbaz and Castel [35]).}$$

(j) Tables of numerical values of the $12j(I)$ symbols

Some tables are presented in Ref. [141]. These tables contain the $12j(I)$ symbols whose arguments are less than or equal 2. Numerical values are given in decimals. However, they can easily be generated by using the recursion relations (see Sec. 10.13.2) and the explicit forms of the $12j(I)$ symbols for special values of arguments (see Sec. 10.13.2).

10.13.3 $12j$ Symbols of the Second Kind ($12j(II)$ Symbols)

(a) Alternative form considered here

Instead of the $12j(II)$ symbols determined by Eq. (10.13.2), we shall consider the somewhat more symmetric $12j(II)$ symbols introduced in Ref. [142]. The latter $12j(II)$ -symbols may be represented by invariant sums of products of eight $3jm$ symbols:

$$\left\{ \begin{array}{cccc} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = (-1)^{a_2+c_1-c_4} \sum_{\alpha_i \beta_i \gamma_i \delta_i} \begin{pmatrix} a_2 & a_3 & a_4 \\ \alpha_2 & \alpha_3 & \alpha_4 \end{pmatrix} \begin{pmatrix} a_4 & b_4 & c_4 \\ \alpha_4 & \beta_4 & \gamma_4 \end{pmatrix} \begin{pmatrix} c_4 & c_1 & c_2 \\ \gamma_4 & \gamma_1 & \gamma_2 \end{pmatrix} \begin{pmatrix} c_2 & d_2 & a_2 \\ \gamma_2 & \delta_2 & \alpha_2 \end{pmatrix} \\ \times \begin{pmatrix} b_4 & b_3 & b_1 \\ \beta_4 & \beta_3 & \beta_1 \end{pmatrix} \begin{pmatrix} b_1 & c_1 & d_1 \\ \beta_1 & \gamma_1 & \delta_1 \end{pmatrix} \begin{pmatrix} d_1 & d_2 & d_3 \\ \delta_1 & \delta_2 & \delta_3 \end{pmatrix} \begin{pmatrix} d_3 & b_3 & a_3 \\ \delta_3 & \beta_3 & \alpha_3 \end{pmatrix}. \quad (10.13.24)$$

Equation (10.13.24) unambiguously fixes the absolute values and phases of the $12j(II)$ symbols. These symbols are real.

(b) COnditions on the arguments

Arguments of the $12j(II)$ symbols satisfy the following conditions:

- (i) All arguments are non-negative integers or half-integers.

(ii) A $12j(\text{II})$ symbol vanishes unless the triangular conditions (see Sec. 8.1.1) are fulfilled for the triads $(a_2 a_3 a_4), (b_1 b_3 b_4), (c_1 c_2 c_4), (d_1 d_2 d_3), (b_1 c_1 d_1), (a_2 c_2 d_2), (a_3 b_3 d_3)$ and $(a_4 b_4 c_4)$.

(iii) A $12j(\text{II})$ symbol vanishes unless the tetragonal conditions (see Sec. 10.13.2) are fulfilled for three tetrads $(a_2 c_4 d_3 b_1), (a_3 b_4 d_2 c_1)$ and $(a_4 b_3 c_2 d_1)$.

(c) *Orthogonality conditions*

The $12j(\text{II})$ symbols satisfy the orthogonality and normalization condition

$$\begin{aligned} & \sum_{x_1 x_2 x_3} (2x_1 + 1)(2x_2 + 1)(2x_3 + 1) \begin{Bmatrix} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & x_1 & - & c_4 \\ d_1 & x_2 & x_3 & - \end{Bmatrix} \begin{Bmatrix} - & a_2 & a_3 & a'_4 \\ b'_1 & - & b_3 & b'_4 \\ c_1 & x_1 & - & c_4 \\ d_1 & x_2 & x_3 & - \end{Bmatrix} \\ &= \frac{\delta_{a_4 a'_4} \delta_{b_1 b'_1} \delta_{b_4 b'_4}}{(2a_4 + 1)(2b_1 + 1)(2b_4 + 1)} \{a_2 a_3 a_4\} \{b_1 b_3 b_4\} \{b_1 c_1 d_1\} \{a_4 b_4 c_4\} \end{aligned} \quad (10.13.25)$$

(d) *Sum representation*

The $12j(\text{II})$ -symbols may be represented as the following sums

$$\begin{aligned} & \begin{Bmatrix} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{Bmatrix} = (-1)^{b_3 - a_4 - d_1 + c_2} \sum_x (2x + 1) \\ & \quad \times \begin{Bmatrix} a_3 & b_4 & x \\ b_1 & d_3 & b_3 \end{Bmatrix} \begin{Bmatrix} a_3 & b_4 & x \\ c_4 & a_2 & a_4 \end{Bmatrix} \begin{Bmatrix} b_1 & d_3 & x \\ d_2 & c_1 & d_1 \end{Bmatrix} \begin{Bmatrix} c_4 & a_2 & x \\ d_2 & c_1 & c_2 \end{Bmatrix} \\ & \begin{Bmatrix} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{Bmatrix} = (-1)^{b_3 - a_4 - d_1 + c_2} \sum_x (2x + 1) \begin{Bmatrix} a_3 & b_3 & d_3 \\ a_4 & b_4 & c_4 \\ a_2 & b_1 & x \end{Bmatrix} \begin{Bmatrix} d_2 & d_1 & d_3 \\ c_2 & c_1 & c_4 \\ a_2 & b_1 & x \end{Bmatrix}. \end{aligned} \quad (10.13.26)$$

(e) *Symmetry properties*

Any $12j(\text{II})$ symbol is symmetric with respect to a permutation of any two columns and simultaneous permutation of the rows whose numbers coincide with the numbers of permuted columns. In addition, the $12j(\text{II})$ symbol does not change under transposition. These symmetry properties relate 48 formally different $12j(\text{II})$

symbols:

$$\begin{aligned}
& \begin{pmatrix} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{pmatrix} = \begin{pmatrix} - & b_1 & b_3 & b_4 \\ a_2 & - & a_3 & a_4 \\ c_2 & c_1 & - & c_4 \\ d_2 & d_1 & d_3 & - \end{pmatrix} = \begin{pmatrix} - & c_1 & c_2 & c_4 \\ a_3 & - & a_2 & a_4 \\ b_3 & b_1 & - & b_4 \\ d_3 & d_1 & d_2 & - \end{pmatrix} = \begin{pmatrix} - & d_1 & d_2 & d_3 \\ a_4 & - & a_2 & a_3 \\ b_4 & b_1 & - & b_3 \\ c_4 & c_1 & c_2 & - \end{pmatrix} \\
& = \begin{pmatrix} - & a_4 & a_2 & a_3 \\ d_1 & - & d_2 & d_3 \\ b_1 & b_4 & - & b_3 \\ c_1 & c_4 & c_2 & - \end{pmatrix} = \begin{pmatrix} - & b_4 & b_1 & b_3 \\ d_2 & - & d_1 & d_3 \\ a_2 & a_4 & - & a_3 \\ c_2 & c_4 & c_1 & - \end{pmatrix} = \begin{pmatrix} - & c_4 & c_1 & c_2 \\ d_3 & - & d_1 & d_2 \\ a_3 & a_4 & - & a_2 \\ b_3 & b_4 & b_1 & - \end{pmatrix} = \begin{pmatrix} - & d_3 & d_1 & d_2 \\ c_4 & - & c_1 & c_2 \\ a_4 & a_3 & - & a_2 \\ b_4 & b_3 & b_1 & - \end{pmatrix} \\
& = \begin{pmatrix} - & a_3 & a_4 & a_2 \\ c_1 & - & c_4 & c_2 \\ d_1 & d_3 & - & d_2 \\ b_1 & b_3 & b_4 & - \end{pmatrix} = \begin{pmatrix} - & b_3 & b_4 & b_1 \\ c_2 & - & c_4 & c_1 \\ d_2 & d_3 & - & d_1 \\ a_2 & a_3 & a_4 & - \end{pmatrix} = \begin{pmatrix} - & c_2 & c_4 & c_1 \\ b_3 & - & b_4 & b_1 \\ d_3 & d_2 & - & d_1 \\ a_3 & a_2 & a_4 & - \end{pmatrix} = \begin{pmatrix} - & d_2 & d_3 & d_1 \\ b_4 & - & b_3 & b_1 \\ c_4 & c_2 & - & c_1 \\ a_4 & a_2 & a_3 & - \end{pmatrix} \\
& = \begin{pmatrix} - & a_2 & a_4 & a_3 \\ b_1 & - & b_4 & b_3 \\ d_1 & d_2 & - & d_3 \\ c_1 & c_2 & c_4 & - \end{pmatrix} = \begin{pmatrix} - & b_1 & b_4 & b_3 \\ a_2 & - & a_4 & a_3 \\ d_2 & d_1 & - & d_3 \\ c_2 & c_1 & c_4 & - \end{pmatrix} = \begin{pmatrix} - & c_1 & c_4 & c_2 \\ a_3 & - & a_4 & a_2 \\ d_3 & d_1 & - & d_2 \\ b_3 & b_1 & b_4 & - \end{pmatrix} = \begin{pmatrix} - & d_1 & d_3 & d_2 \\ a_4 & - & a_3 & a_2 \\ c_4 & c_1 & - & c_2 \\ b_4 & b_1 & b_3 & - \end{pmatrix} \\
& = \begin{pmatrix} - & a_3 & a_2 & a_4 \\ c_1 & - & c_2 & c_4 \\ b_1 & b_3 & - & b_4 \\ d_1 & d_3 & d_2 & - \end{pmatrix} = \begin{pmatrix} - & b_3 & b_1 & b_4 \\ c_2 & - & c_1 & c_4 \\ a_2 & a_3 & - & a_4 \\ d_2 & d_3 & d_1 & - \end{pmatrix} = \begin{pmatrix} - & c_2 & c_1 & c_4 \\ b_3 & - & b_1 & b_4 \\ a_3 & a_2 & - & a_4 \\ d_3 & d_2 & d_1 & - \end{pmatrix} = \begin{pmatrix} - & d_2 & d_1 & d_3 \\ b_4 & - & b_1 & b_3 \\ a_4 & a_2 & - & a_3 \\ c_4 & c_2 & c_1 & - \end{pmatrix} \\
& = \begin{pmatrix} - & a_4 & a_3 & a_2 \\ d_1 & - & d_3 & d_2 \\ c_1 & c_4 & - & c_2 \\ b_1 & b_4 & b_3 & - \end{pmatrix} = \begin{pmatrix} - & b_4 & b_3 & b_1 \\ d_2 & - & d_3 & d_1 \\ c_2 & c_4 & - & c_1 \\ a_2 & a_4 & a_3 & - \end{pmatrix} = \begin{pmatrix} - & c_4 & c_2 & c_1 \\ d_3 & - & d_2 & d_1 \\ b_3 & b_4 & - & b_1 \\ a_3 & a_4 & a_2 & - \end{pmatrix} = \begin{pmatrix} - & d_3 & d_2 & d_1 \\ c_4 & - & c_2 & c_1 \\ b_4 & b_3 & - & b_1 \\ a_4 & a_3 & a_2 & - \end{pmatrix} \\
& = \begin{pmatrix} - & b_1 & c_1 & d_1 \\ a_2 & - & c_2 & d_2 \\ a_3 & b_3 & - & d_3 \\ a_4 & b_4 & c_4 & - \end{pmatrix} = \begin{pmatrix} - & a_2 & c_2 & d_2 \\ b_1 & - & c_1 & d_1 \\ b_3 & a_3 & - & d_3 \\ b_4 & a_4 & c_4 & - \end{pmatrix} = \begin{pmatrix} - & a_3 & b_3 & d_3 \\ c_1 & - & b_1 & d_1 \\ c_2 & a_2 & - & d_2 \\ c_4 & a_4 & b_4 & - \end{pmatrix} = \begin{pmatrix} - & a_4 & b_4 & c_4 \\ d_1 & - & b_1 & c_1 \\ d_2 & a_2 & - & c_2 \\ d_3 & a_3 & b_3 & - \end{pmatrix} \\
& = \begin{pmatrix} - & d_1 & b_1 & c_1 \\ a_4 & - & b_4 & c_4 \\ a_2 & d_2 & - & c_2 \\ a_3 & d_3 & b_3 & - \end{pmatrix} = \begin{pmatrix} - & d_2 & a_2 & c_2 \\ b_4 & - & a_4 & c_4 \\ b_1 & d_1 & - & c_1 \\ b_3 & d_3 & a_3 & - \end{pmatrix} = \begin{pmatrix} - & d_3 & a_3 & b_3 \\ c_4 & - & a_4 & b_4 \\ c_1 & d_1 & - & b_1 \\ c_2 & d_2 & a_2 & - \end{pmatrix} = \begin{pmatrix} - & c_4 & a_4 & b_4 \\ d_3 & - & a_3 & b_3 \\ d_1 & c_1 & - & b_1 \\ d_2 & c_2 & a_2 & - \end{pmatrix} \\
& = \begin{pmatrix} - & c_1 & d_1 & b_1 \\ a_3 & - & d_3 & b_3 \\ a_4 & c_4 & - & b_4 \\ a_2 & c_2 & d_2 & - \end{pmatrix} = \begin{pmatrix} - & c_2 & d_2 & a_2 \\ b_3 & - & d_3 & a_3 \\ b_4 & c_4 & - & a_4 \\ b_1 & c_1 & d_1 & - \end{pmatrix} = \begin{pmatrix} - & b_3 & d_3 & a_3 \\ c_2 & - & d_2 & a_2 \\ c_4 & b_4 & - & a_4 \\ c_1 & b_1 & d_1 & - \end{pmatrix} = \begin{pmatrix} - & b_4 & c_4 & a_4 \\ d_2 & - & c_2 & a_2 \\ d_3 & b_3 & - & a_3 \\ d_1 & b_1 & c_1 & - \end{pmatrix} \\
& = \begin{pmatrix} - & b_1 & d_1 & c_1 \\ a_2 & - & d_2 & c_2 \\ a_4 & b_4 & - & c_4 \\ a_3 & b_3 & d_3 & - \end{pmatrix} = \begin{pmatrix} - & a_2 & d_2 & c_2 \\ b_1 & - & d_1 & c_1 \\ b_4 & a_4 & - & c_4 \\ b_3 & a_3 & d_3 & - \end{pmatrix} = \begin{pmatrix} - & a_3 & d_3 & b_3 \\ c_1 & - & d_1 & b_1 \\ c_4 & a_4 & - & b_4 \\ c_2 & a_2 & d_2 & - \end{pmatrix} = \begin{pmatrix} - & a_4 & c_4 & b_4 \\ d_1 & - & c_1 & b_1 \\ d_3 & a_3 & - & b_3 \\ d_2 & a_2 & c_2 & - \end{pmatrix} \\
& = \begin{pmatrix} - & c_1 & b_1 & d_1 \\ a_3 & - & b_3 & d_3 \\ a_2 & c_2 & - & d_2 \\ a_4 & c_4 & b_4 & - \end{pmatrix} = \begin{pmatrix} - & c_2 & a_2 & d_2 \\ b_3 & - & a_3 & d_3 \\ b_1 & c_1 & - & d_1 \\ b_4 & c_4 & a_4 & - \end{pmatrix} = \begin{pmatrix} - & b_3 & a_3 & d_3 \\ c_2 & - & a_2 & d_2 \\ c_1 & b_1 & - & d_1 \\ c_4 & b_4 & a_4 & - \end{pmatrix} = \begin{pmatrix} - & b_4 & a_4 & c_4 \\ d_2 & - & a_2 & c_2 \\ d_1 & b_1 & - & c_1 \\ d_3 & b_3 & a_3 & - \end{pmatrix} \\
& = \begin{pmatrix} - & d_1 & c_1 & b_1 \\ a_4 & - & c_4 & b_4 \\ a_3 & d_3 & - & b_3 \\ a_2 & d_2 & c_2 & - \end{pmatrix} = \begin{pmatrix} - & d_2 & c_2 & a_2 \\ b_4 & - & c_4 & a_4 \\ b_3 & d_3 & - & a_3 \\ b_1 & d_1 & c_1 & - \end{pmatrix} = \begin{pmatrix} - & d_3 & b_3 & a_3 \\ c_4 & - & b_4 & a_4 \\ c_2 & d_2 & - & a_2 \\ c_1 & d_1 & b_1 & - \end{pmatrix} = \begin{pmatrix} - & c_4 & b_4 & a_4 \\ d_3 & - & b_3 & a_3 \\ d_2 & c_2 & - & a_2 \\ d_1 & c_1 & b_1 & - \end{pmatrix}.
\end{aligned}$$

(10.13.27)

(f) Recursion relations

These relations are useful for evaluation of the $12j(\text{II})$ symbols. We present only one relation:

$$\begin{aligned} & \sum_{a'_3} (2a'_3 + 1) \left\{ \begin{matrix} \lambda & a_3 & a'_3 \\ d_3 & b'_3 & b_3 \end{matrix} \right\} \left\{ \begin{matrix} \lambda & a_3 & a'_3 \\ a_2 & a_4 & a'_4 \end{matrix} \right\} \left\{ \begin{matrix} - & a_2 & a'_3 & a_4 \\ b_1 & - & b'_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} \\ &= (-1)^{b_1+a_2-d_3-c_4} \sum_{b'_4} (2b'_4 + 1) \left\{ \begin{matrix} \lambda & b_4 & b'_4 \\ c_4 & a'_4 & a_4 \end{matrix} \right\} \left\{ \begin{matrix} \lambda & b_4 & b'_4 \\ b_1 & b_3 & b'_3 \end{matrix} \right\} \left\{ \begin{matrix} - & a_2 & a_3 & a'_4 \\ b_1 & - & b_3 & b'_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\}. \end{aligned} \quad (10.13.28)$$

In particular, putting $\lambda = \frac{1}{2}$, $a'_4 = a_4 - \frac{1}{2}$ and $b'_3 = b_3 + \frac{1}{2}$ in Eq. (10.13.28) we obtain

$$\begin{aligned} & (2b_4 + 1) \left\{ \left[(a_3 + b_3 - d_3 + 1)(a_3 + b_3 + d_3 + 2) \left(-a_2 + a_3 + a_4 + \frac{1}{2} \right) \left(a_2 + a_3 + a_4 + \frac{3}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} - & a_2 & a_3 + \frac{1}{2} & a_4 \\ b_1 & - & b_3 + \frac{1}{2} & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} \right. \\ & \left. + \left[(-a_3 + b_3 + d_3 + 1)(a_3 - b_3 + d_3) \left(a_2 - a_3 + a_4 + \frac{1}{2} \right) \left(a_2 + a_3 - a_4 + \frac{1}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} - & a_2 & a_3 - \frac{1}{2} & a_4 \\ b_1 & - & b_3 + \frac{1}{2} & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} \right\} \\ &= (2a_3 + 1) \left\{ \left[(-a_4 + b_4 + c_4 + 1)(a_4 - b_4 + c_4) \left(b_1 - b_3 + b_4 + \frac{1}{2} \right) \left(b_1 + b_3 - b_4 + \frac{1}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} - & a_2 & a_3 & a_4 - \frac{1}{2} \\ b_1 & - & b_3 & b_4 - \frac{1}{2} \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} \right. \\ & \left. + \left[(a_4 + b_4 - c_4)(a_4 + b_4 + c_4 + 1) \left(-b_1 + b_3 + b_4 + \frac{1}{2} \right) \left(b_1 + b_3 + b_4 + \frac{3}{2} \right) \right]^{\frac{1}{2}} \left\{ \begin{matrix} - & a_2 & a_3 & a_4 - \frac{1}{2} \\ b_1 & - & b_3 & b_4 - \frac{1}{2} \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} \right\}. \end{aligned} \quad (10.13.29)$$

(g) Explicit forms of the $12j(\text{II})$ -symbols for some relations between arguments

may be obtained, using Eqs. (10.13.26) and (10.13.27).

If one argument in any tetrad is equal to the sum of three other arguments from the same tetrad, one obtains

$$\begin{aligned} & \left\{ \begin{matrix} - & b_1 + c_4 + d_3 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{matrix} \right\} = (-1)^{b_3-b_4+c_1-d_1} \frac{(2b_1)!(2c_4)!(2d_3)!}{(2b_1+2c_4+2d_3+1)!} \\ & \times \frac{D(b_1b_3b_4)D(c_4c_1c_2)D(d_3d_1d_2)D(b_1d_1c_1)D(c_4b_4a_4)D(d_3b_3a_3)}{D(b_1+d_3+c_4a_4a_3)D(b_1+d_3+c_4c_2d_2)} \end{aligned} \quad (10.13.30)$$

Let each argument in any tetrad be equal to the sum of two arguments which form a triad with this argument. Then, the $12j(\text{II})$ symbol may be expressed as a sum involving products of two Clebsch-Gordan coefficients.

$$\begin{aligned} & \left\{ \begin{matrix} - & a_3 + a_4 & a_3 & a_4 \\ b_3 + b_4 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_1 + c_2 \\ d_1 & d_2 & d_1 + d_2 & - \end{matrix} \right\} = (-1)^{b_3-a_4-d_1+c_2} f(a_3b_3d_1 + d_2) f(a_4b_4c_1 + c_2) \\ & \times f(d_1c_1b_3 + b_4) f(d_2c_2a_3 + a_4) \left[\frac{(2a_3)!(2a_4)!(2b_3)!(2b_4)!(2c_1)!(2c_2)!(2d_1)!(2d_2)!}{(2a_3+2a_4+1)!(2b_3+2b_4+1)!(2c_1+2c_2+1)!(2d_1+2d_2+1)!} \right]^{\frac{1}{2}} \\ & \times \sum_x [f(a_3 + a_4b_3 + b_4x) f(d_1 + d_2c_1 + c_2x)]^{-1} C_{d_1+d_2 \ a_3-b_3 \ c_1+c_2 \ a_4-b_4}^x C_{a_3+a_4 \ d_2-c_2 \ b_3+b_4 \ d_1-c_1}^{d_1+d_2-c_1-c_2} \end{aligned} \quad (10.13.31)$$

where

$$f(abc) = [(a+b+c+1)!(a+b-c)!]^{-\frac{1}{2}} \quad (10.13.32)$$

(h) *Explicit forms of the 12j(II) symbols for special values of the arguments*

If one argument is equal to zero, a 12j(II) symbol is reduced to product of two 6j symbols:

$$\left\{ \begin{array}{cccc} - & 0 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = (-1)^{b_3+b_4-d_1-c_1} \frac{\delta_{a_3 a_4} \delta_{c_2 d_2}}{\sqrt{(2a_3+1)(2c_2+1)}} \\ \times \left\{ \begin{array}{ccc} b_1 & b_3 & b_4 \\ a_3 & c_4 & d_3 \end{array} \right\} \left\{ \begin{array}{ccc} c_1 & c_2 & c_4 \\ d_3 & b_1 & d_1 \end{array} \right\} \quad (10.13.33)$$

If two arguments from any tetrad are equal to $\frac{1}{2}$ we obtain

$$\left\{ \begin{array}{cccc} - & a_2 & \frac{1}{2} & a_4 \\ b_1 & - & b_3 & \frac{1}{2} \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = \frac{\delta_{b_1 d_3} \delta_{c_4 a_2} \delta_{c_1 d_2}}{2(2b_1+1)(2c_1+1)(2c_4+1)} \\ + (-1)^\varepsilon \left\{ \begin{array}{ccc} b_1 & d_3 & 1 \\ d_2 & c_1 & d_1 \end{array} \right\} \left\{ \begin{array}{ccc} c_4 & a_2 & 1 \\ d_2 & c_1 & c_2 \end{array} \right\} \frac{D(\frac{1}{2}a_2 a_4) D(\frac{1}{2}b_1 b_3) D(\frac{1}{2}c_4 a_4) D(\frac{1}{2}b_3 d_3)}{2D(1b_1 d_3) D(1c_4 a_2)} \quad (10.13.34)$$

$$\varepsilon = b_1 + b_3 + c_2 + c_4 + a_2 - a_4 - d_1 + d_3$$

$$\left\{ \begin{array}{cccc} - & \frac{1}{2} & a_3 & a_4 \\ \frac{1}{2} & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = (-1)^{2b_3+2d_1} \frac{\delta_{d_3 c_4}}{2(2d_3+1)} \left\{ \begin{array}{ccc} a_3 & a_4 & \frac{1}{2} \\ b_4 & b_3 & d_3 \end{array} \right\} \left\{ \begin{array}{ccc} d_2 & c_2 & \frac{1}{2} \\ c_1 & d_1 & d_3 \end{array} \right\} \\ + 3(-1)^{b_3-a_4-d_1+c_2} \left\{ \begin{array}{ccc} a_3 & b_3 & d_3 \\ a_4 & b_4 & c_4 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{array} \right\} \left\{ \begin{array}{ccc} d_2 & d_1 & d_3 \\ c_2 & c_1 & c_4 \\ \frac{1}{2} & \frac{1}{2} & 1 \end{array} \right\}. \quad (10.13.35)$$

At $c_4 = d_3, d_3 \pm 1$, a 12j(II) symbol may be expressed in terms of the 3jm and 6j symbols:

$$\left\{ \begin{array}{cccc} - & \frac{1}{2} & a_3 & a_4 \\ \frac{1}{2} & - & b_3 & b_4 \\ c_1 & c_2 & - & d_3 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = (-1)^{2a_3+2d_1} \frac{1}{2(2d_3+1)} \left\{ \begin{array}{ccc} a_3 & a_4 & \frac{1}{2} \\ b_4 & b_3 & d_3 \end{array} \right\} \left\{ \begin{array}{ccc} d_2 & c_2 & \frac{1}{2} \\ c_1 & d_1 & d_3 \end{array} \right\} \\ + \frac{(-1)^{b_3-a_4-d_1+c_2}}{2(2d_3+1)\sqrt{(2a_4+1)(2b_4+1)(2c_1+1)(2c_2+1)}} \frac{\begin{pmatrix} a_3 & b_3 & d_3 \\ \frac{1}{2} & \frac{1}{2} & -1 \end{pmatrix} \begin{pmatrix} d_2 & d_1 & d_3 \\ \frac{1}{2} & \frac{1}{2} & -1 \end{pmatrix}}{\begin{pmatrix} a_4 & b_4 & d_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} c_1 & c_2 & d_3 \\ 0 & 0 & 0 \end{pmatrix}} \quad (10.13.36)$$

This equation is valid, if $a_4 + b_4 + d_3$ and $c_1 + c_2 + d_3$ are even. If these numbers are odd, one has

$$\left\{ \begin{array}{cccc} - & \frac{1}{2} & a_3 & a_4 \\ \frac{1}{2} & - & b_3 & b_4 \\ c_1 & c_2 & - & d_3 + 1 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = \frac{(a_4 - a_3)(2a_3 + 1) + (b_4 - b_3)(2b_3 + 1) + d_3 + 1}{(2d_3 + 1)(2d_3 + 2)(2d_3 + 3)} \\ \times \frac{(c_2 - d_2)(2d_2 + 1) + (c_1 - d_1)(2d_1 + 1) + d_3 + 1}{\sqrt{(2a_4 + 1)(2b_4 + 1)(2c_1 + 1)(2c_2 + 1)}} \frac{\begin{pmatrix} a_3 & b_3 & d_3 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} d_2 & d_1 & d_3 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{pmatrix}}{\begin{pmatrix} a_4 & b_4 & d_3 + 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} c_1 & c_2 & d_3 + 1 \\ 0 & 0 & 0 \end{pmatrix}} \quad (10.13.37)$$

$$\left\{ \begin{array}{cccc} - & \frac{1}{2} & a_3 & a_4 \\ \frac{1}{2} & - & b_3 & b_4 \\ c_1 & c_2 & - & d_3 - 1 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = \frac{(a_4 - a_3)(2a_3 + 1) + (b_4 - b_3)(2b_3 + 1) - d_3}{(2d_3 - 1)(2d_3)(2d_3 + 1)} \\ \times \frac{(c_2 - d_2)(2d_2 + 1) + (c_1 - d_1)(2d_1 + 1) - d_3}{\sqrt{(2a_4 + 1)(2b_4 + 1)(2c_1 + 1)(2c_2 + 1)}} \frac{\begin{pmatrix} a_3 & b_3 & d_3 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{pmatrix} \begin{pmatrix} d_2 & d_1 & d_3 \\ \frac{1}{2} & -\frac{1}{2} & 0 \end{pmatrix}}{\begin{pmatrix} a_4 & b_4 & d_3 - 1 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} c_1 & c_2 & d_3 - 1 \\ 0 & 0 & 0 \end{pmatrix}} \quad (10.13.38)$$

(i) Relations between the $12j(II)$ symbols considered above and analogous functions of other authors

$$\left\{ \begin{array}{cccc} - & a_2 & a_3 & a_4 \\ b_1 & - & b_3 & b_4 \\ c_1 & c_2 & - & c_4 \\ d_1 & d_2 & d_3 & - \end{array} \right\} = \left(\begin{array}{cccc} \cdot & a_2 & a_3 & a_4 \\ b_1 & \cdot & b_3 & b_4 \\ c_1 & c_2 & \cdot & c_4 \\ d_1 & d_2 & d_3 & \cdot \end{array} \right),$$

(Sharp [142])

$$1)^{a_2+a_3+d_2+d_3-b_1-b_4-c_1-c_4} \left[\begin{array}{cccc} a_3 & d_3 & d_2 & a_2 \\ b_3 & d_1 & c_2 & a_4 \\ b_4 & b_1 & c_1 & c_4 \end{array} \right],$$

(Yutsis, Levinson and Vanagas [34]),

$$= (-1)^{a_2+a_3+d_2+d_3-b_1-b_4-c_1-c_4} \left\{ \begin{array}{cccc} a_3 & d_3 & d_2 & a_2 \\ b_3 & d_1 & c_2 & a_4 \\ b_4 & b_1 & c_1 & c_4 \end{array} \middle| 2 \right\}$$

(Elbaz and Castel [35]).

Chapter 11

The graphical method in angular momentum theory

To solve many quantum mechanical problems involving angular and spin dependences one has to integrate a product of the spherical harmonics or the Wigner D-functions and to sum products of the Clebsch-Gordan coefficients and $3nj$ symbols. Expressions of this kind are, as a rule, rather complicated. They contain so many arguments and parameters that it is difficult to understand interrelations between them, to find out the symmetry of the expression, and to see its invariance with respect to a particular transformation.

The representation of such expressions in diagrammatic form facilitates their general analysis. A graphical representation is more compact and clear. All arguments and parameter interrelations are obvious. The structure and symmetries of the expression as a whole and each part of the expression are evident. Moreover, the graphical technique substantially reduces the calculation job.

There are several versions of the graphical method. A handy technique for summing the $3jm$ symbol products was suggested by Levinson [85], Yutsis, Levinson and Vanagas [34] and Yutsis and Bandzaitis [20]. A slightly modified version was proposed by Brink and Satchler [7]. A certain development of this diagram technique was presented by El-Baz and Castel [35] who extended it to expressions involving continuous variables. This version is now commonly used.¹ However, various books and papers concerning such diagrams are somewhat different in definitions, phase conventions, and rules of the graphical technique, although the notations, sometimes, are similar. Therefore, one should be careful in using formulas from various sources.

In this chapter we represent the unified graphical method in detail. The basic elements of diagrams are defined in Sections 11.1 and 11.2 where the graphical representation of the standard functions and main operations of the angular momentum theory are given. Section 11.3 exhibits the rules of the diagram technique, i.e., the graphical method of calculation. For practical convenience, all the definitions and rules are arranged in tables. The general scheme of application of the diagram technique is formulated in Sec. 11.4.

11.1 Graphical representation of functions

A principle of any graphical method is in establishing a one-to-one correspondence between elements of a diagram and constituents of the analytic expression. Any expression inherent in the theory may be represented by a definite diagram, and vice versa, any such diagram represents a definite analytic expression.

Moreover, a transformation of the expression under some analytic operation corresponds to a certain transformation of the associated diagram. That is why all calculations of spin angular quantities may be performed by the graphical method instead of the analytic one.

11.1.1 Basic Elements of Diagrams

The Dirac representation of quantum mechanical quantities seems to be most adequate to the graphical method. This representation operates with state vectors: the “kets”, $|\Psi\rangle$ and the Hermitian conjugate “bras”, $\langle\psi|$. Graphically, these vectors are represented by directed lines connected with nodes (dot or cross bar).

A line with an arrow coming out of a node represents the “ket” $|\Psi\rangle$, i.e., a covariant (standard) state vector, whereas, a line with double arrow coming into the node represents the “bra” $\langle\Psi|$, i.e. a contravariant (Hermitian conjugate) state vector.

¹A somewhat different graphical method was suggested by Shelepin [70], but it is not very widespread now.

A thin solid line corresponds to a state with some definite angular momentum j and projection m :

$$\begin{array}{l} \longrightarrow |jm\rangle, \\ \longrightarrow \langle jm|, \end{array} \quad (11.1.1)$$

Such lines are referred to as the j -lines.

A dashed line corresponds to a state with some definite direction of the position vector (or linear momentum) $\Omega\{\vartheta, \varphi\}$

$$\begin{array}{l} \dashrightarrow |\Omega\rangle, \\ \dashrightarrow \langle\Omega|, \end{array} \quad (11.1.2)$$

Such lines are referred to as the Ω -lines.

A double dashed line corresponds to a set of the Euler angles α, β, γ (or the angles ω, Θ, Φ) which specify a rotation of the coordinate system, R . Such lines are called the R -lines

$$===== R$$

The symbols jm , Ω and R are usually indicated near the corresponding lines. Therefore, length, curvature, orientation, and position of the line are inessential. For example,

$$\longrightarrow \text{ is equivalent to } \longleftarrow \text{ and } \longleftrightarrow .$$

However, the arrow direction with respect to the node is of importance.

The normalization of state vectors may be different. In diagrams it is indicated by the type of node. A state vector $|jm\rangle$ normalized according to

$$\langle jm|j'm'\rangle = \delta_{jj'}\delta_{mm'} \quad (11.1.3)$$

is represented by the j -line with a bar node, whereas, a state vector $|jm\rangle$, which obeys the normalization condition

$$(jm|j'm') = (2j+1)^{-1}\delta_{jj'}\delta_{mm'} \quad (11.1.4)$$

is represented by the j -line with a dot node. The completeness condition for wave functions $|jm\rangle$ or $|jm\rangle$ reads

$$\sum_{jm} |jm\rangle\langle jm| = \sum_{jm} |jm\rangle(2j+1)\langle jm| = 1 \quad (11.1.5)$$

Similarly, a state vector $|\Omega\rangle$ normalized by the Dirac δ -function

$$\langle\Omega|\Omega'\rangle = \delta(\Omega - \Omega') \quad (11.1.6)$$

corresponds to the Ω -line with the bar node, whereas, a state vector $|\Omega\rangle$ normalized by

$$(\Omega|\Omega') = 4\pi\delta(\Omega - \Omega') \quad (11.1.7)$$

corresponds to the Ω -line with a dot node. The completeness condition for the functions $|\Omega\rangle$ or $|\Omega\rangle$ reads

$$\int |\Omega\rangle d\Omega \langle\Omega| = 1 \quad \text{and} \quad \int |\Omega\rangle \frac{d\Omega}{4\pi} \langle\Omega| = 1, \quad (11.1.8)$$

respectively.

11.1.2 Diagrams of the Basic Functions of the Theory

Quantum mechanical quantities, i.e., state vectors and operators, may depend on many spin and angular variables (such as jm or Ω) and also on coordinate system rotation R . Accordingly, these state vectors and matrix elements of the operators may be represented graphically as blocks (i.e., as subdiagrams) with external jm -, Ω - and/or R -lines associated with the corresponding spin angular variables. A diagram corresponding to a composite expression will be a combination of such blocks connected by some of jm -, Ω - and R -lines.

Diagrams of some spin and angular functions are listed in Tables 11.1-11.3; graphical displays of their properties and relations is given in Tables 11.4-11.15.

A spherical harmonic $Y_{lm}(\vartheta, \varphi) \equiv \langle\Omega|lm\rangle$ is represented by a junction of a thin solid and dashed lines which correspond to $|lm\rangle$ and $\langle\Omega|$.

A Wigner $3jm$ symbol $\begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \equiv (00|aab\beta c\gamma)$ is represented by a dot-node junction of three thin solid lines which correspond to $|a\alpha\rangle$, $|b\beta\rangle$ and $|c\gamma\rangle$. The sign of the node determines the cyclic order of arguments, i.e., the sequence of coupling of momenta: plus corresponds to the anticlockwise one.

The $3j$, $6j$, and $9j$ symbols, or, more generally speaking, the $3nj$ symbols, may be represented graphically by closed diagrams without external lines because they are invariant under coordinate rotation. Such closed diagrams consist of $3nj$ -lines connected in $2n$ dot nodes by three's, because the $3nj$ symbols are actually the sums of $2n$ various $3jm$ symbols (Sec. 10.13). Any $3nj$ symbol may be represented by several equivalent diagrams which differ in the arrow directions of internal the j lines and in the node signs since each $3nj$ symbol may be expressed through different sums of products of the $3jm$ symbols; such sums turn into one another by replacing summation indices.

Matrix elements of the rotation operator in the Ω -representation, $\langle \Omega | \hat{R} | \Omega' \rangle = \delta(\Omega - \hat{R}\Omega')$, and in the jm representation, $\langle jm | \hat{R} | j'm' \rangle = \delta_{jj'} D_{mm'}^j(R)$, are displayed as junctions of three lines. One of them is a double dashed R -line and the two others are either dashed lines corresponding to $|\Omega'\rangle$ and $\langle \Omega|$ states, or thin solid lines corresponding to $|j'm'\rangle$ and $\langle jm|$ states. The sign of the junction node determines the order of arguments; strictly speaking, it shows which of two states is initial and which is final. Plus corresponds to the anticlockwise reading of arguments, and minus to the clockwise one. Therefore, a sign reversal means conversion from the rotation R to the inverse rotation R^{-1} .

Matrix elements of any irreducible tensor operator $\langle \Psi | T_{\lambda\mu} | \Psi' \rangle$ are represented by a block with three external lines corresponding to $\langle \Psi|$, $ket\Psi'$ and $|\lambda\mu\rangle$; the operator symbol (and other possible quantum numbers) are indicated inside the block.

11.2 Graphical representation of the main operations of the theory

Expressions which occur in calculations based on the quantum theory of angular momentum have a specific character. They contain spin angular variables of two kinds, i.e., external and internal ones.

1. External variables are actual arguments of the expression under consideration. The evaluation of an expression does not involve any integration or summation over external variables, i.e., they appear as fixed (but, in principle, changeable) parameters.
2. Internal variables are those over which integration or summation is assumed. It is essential that each internal variable which enters an expression inherent in the theory corresponds to some scalar product $|\Psi\rangle\langle\Psi|$. This is why one does not need to take into account any transformation of internal variables under rotation of the coordinate system. This specific feature is important for practical calculations.

In other words, quantum mechanical problems are mainly concerned with invariant integration and/or summation of bilinear forms over total ranges of all internal variables. Accordingly, the graphical method under consideration may be applied only to the expressions of such a type.


This section describes the graphical representation of those analytic operations which are specific to angular momentum theory, while the rules of calculations by the graphical method will be formulated in the next section.

11.2.1 Multiplication

The product (direct product) of factors $\mathfrak{M}$ and $\mathfrak{N}$ is represented graphically by unlinked subdiagrams (blocks) which correspond to factors $\mathfrak{M}$ and $\mathfrak{N}$. The mutual disposition and orientation of these subdiagrams are inessential. For example, the product

$$\begin{pmatrix} A & B & C \\ \alpha & \beta & \gamma \end{pmatrix} \begin{Bmatrix} A & B & C \\ a & b & c \end{Bmatrix} \quad (11.2.1)$$

may be represented by



11.2.2 Invariant Summation over Projections

The scalar product of two irreducible tensors of rank j , $\mathfrak{M}_{jm}$ and $\mathfrak{N}_{jm}$, i.e., the sum of bilinear combinations over all possible values of the projection m

$$\sum_{m=-j}^j \langle |\mathfrak{M}| jm \rangle \langle jm | \mathfrak{N} | \rangle, \quad (11.2.2)$$

Table 11.1. Basic Elements of Diagrams.

Analytical Expression	Graphical Representation
<i>jm</i> -State Vector	
“ket” $ jm\rangle$	
“bra” $\langle jm $	
Kronecker δ -Symbol	
$\langle jm j'm'\rangle = \delta_{jj'}\delta_{mm'}$	
$(jm j'm') = \frac{\delta_{jj'}\delta_{mm'}}{2j+1}$	
Ω -State Vector	
“ket” $ \Omega\rangle$	
“bra” $\langle\Omega $	
Dirac δ -function	
$\langle\Omega \Omega'\rangle = \delta(\Omega - \Omega')$	
$(\Omega \Omega') = \delta(\Omega - \Omega')4\pi$	
$\langle\Omega_1\Omega_2 \Omega\rangle = \delta(\Omega_1 - \Omega)\delta(\Omega_2 - \Omega)$	
Spherical Harmonics	
$\langle\Omega lm\rangle = Y_{lm}(\Omega)$	
$\langle lm \Omega\rangle = Y_{lm}^*(\Omega)$	
$(\Omega lm) = C_{lm}(\Omega)$	
$(lm \Omega) = C_{lm}^*(\Omega)$	
Clebsch-Gordan Coefficients	
$C_{j_1 m_1 j_2 m_2}^{J M} = \langle j_1 m_1 j_2 m_2 j_1 j_2 J M \rangle = \langle j_1 j_2 J M j_1 m_1 j_2 m_2 \rangle$	

Table 11.1. (Cont'd).

Analytical Expression	Graphical Representation
Wigner $3jm$ Symbols	
$\begin{pmatrix} j_1 & n_1 & j_2 \\ m_2 & j_3 & m_3 \end{pmatrix} = (j_1 m_1 j_2 m_2 j_3 m_3 00) = (00 j_1 m_1 j_2 m_2 j_3 m_3)$	
Metric tensor	
$\begin{pmatrix} j \\ mm' \end{pmatrix} = (-1)^{j+m} \delta_{m, -m'}$ $\begin{pmatrix} j \\ m'm \end{pmatrix} = (-1)^{j-m} \delta_{-m, m'}$	

is represented by the conjunction (closure) of $|jm\rangle$ - and $\langle jm|$ -lines of subdiagrams corresponding to $\langle |\mathfrak{M}|jm\rangle$ and $\langle jm|\mathfrak{N}\rangle$

$$\sum_m \boxed{\mathfrak{M}} \xrightarrow{j, m} \boxed{\mathfrak{N}} = \boxed{\mathfrak{M}} \xrightarrow{j} \boxed{\mathfrak{N}}$$

The resultant linking j -line has the same direction as the original $|jm\rangle$ - and $\langle jm|$ -lines. For example, the sum of product of two $3jm$ symbols over two projections, α and β

$$\sum_{\alpha, \beta} \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \begin{pmatrix} a & b & c' \\ \alpha & \beta & \gamma' \end{pmatrix} = \sum_{\alpha, \beta} (-1)^{a-\alpha+b-\beta+c-\gamma} \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} \begin{pmatrix} a & b & c' \\ \alpha & \beta & \gamma' \end{pmatrix} \quad (11.2.3)$$

may be displayed as

$$\sum_{\alpha, \beta} \begin{array}{c} a\alpha \\ \nearrow \\ c\gamma \\ \leftarrow \\ b\beta \end{array} \begin{array}{c} a\alpha \\ \nearrow \\ c'\gamma' \\ \leftarrow \\ b\beta \end{array} = \begin{array}{c} a \\ \nearrow \\ c\gamma \\ \leftarrow \\ b \end{array}$$

Thus, each internal j -line of a diagram represents a scalar product (contraction over m) of two irreducible tensors of rank j . To construct such a scalar product one of these tensors should be written in a covariant form $\langle |\mathfrak{M}|jm\rangle \equiv \mathfrak{M}_{jm}$ while the other should be written in the form $\langle jm|\mathfrak{N}\rangle = (-1)^{j-m} \mathfrak{N}_{j-m}$ (Sec. 3.1.8). Graphically, this means that one of the corresponding subdiagrams has the external $|jm\rangle$ -line (with single outward arrow) and the other has the $\langle jm|$ -line (with double inward arrow).

Two diagrams which differ only by the direction of an internal j -line, correspond to analytic expressions which differ only by the phase factor $(-1)^{2j}$, because

$$\sum_m (-1)^{j-m} \mathfrak{M}_{jm} \mathfrak{N}_{j-m} = (-1)^{2j} \sum_m (-1)^{j-m} \mathfrak{M}_{j-m} \mathfrak{N}_{jm}. \quad (11.2.4)$$

One should especially mention the entirely invariant sums of products of two, four, six, or more $3jm$ symbols over all projections. Corresponding diagrams are closed with respect to all j -lines. These sums represent the Wigner $3nj$ symbols (Table 11.4).

11.2.3 Summation over Angular Momentum

A thick j -line linking two subdiagrams, as in

$$\boxed{\mathfrak{M}} \xrightarrow{j} \boxed{\mathfrak{N}}$$

Table 11.2. Invariant Functions.










Analytical Expression	Graphical Representation
$(2a + 1)$	
$\sqrt{2a + 1}$	
δ_{ab}	
$\frac{\delta_{ab}\delta_{bc}}{\sqrt{2a + 1}}$	
$\frac{\delta_a\delta_{ba}\delta_{cd}}{(2a + 1)}$	
$3j$ Symbol	
$\{abc\}$	
$6j$ Symbol	
$\begin{Bmatrix} a & b & c \\ A & B & C \end{Bmatrix}$	
$9j$ Symbol	
$\begin{Bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{Bmatrix}$	
$12j(\text{I})$ Symbol	
$\begin{Bmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \end{Bmatrix}$	
$12j(\text{II})$ Symbol	
$\begin{Bmatrix} - & c_2 & b_3 & a_4 \\ c_3 & - & a_2 & b_4 \\ b_2 & a_3 & - & c_4 \\ a_1 & b_1 & c_1 & - \end{Bmatrix}$	

Table 11.3. Matrix Elements.












Analytical Expression	Graphical Representation
Irreducible Tensor Operator	
Ω-Representation $\langle \gamma \Omega \mathfrak{M}_{\lambda \mu} \gamma' \Omega' \rangle$ γ denotes all quantum numbers except Ω	
Adjoint Matrix $\langle \gamma \Omega \mathfrak{M}_{\lambda \mu}^\dagger \gamma' \Omega' \rangle = (-1)^{\lambda - \mu} \langle \gamma' \Omega' \mathfrak{M}_{\lambda - \mu} \gamma \Omega \rangle^*$	
jm-Representation $\langle \gamma jm \mathfrak{M}_{\lambda \mu} \gamma' j' m' \rangle$ γ denotes all quantum numbers except jm	
Adjoint Matrix $\langle \gamma jm \mathfrak{M}_{\lambda \mu}^\dagger \gamma' j' m' \rangle = (-1)^{\lambda - \mu} \langle \gamma' j' m' \mathfrak{M}_{\lambda - \mu} \gamma jm \rangle^*$	
Reduced Matrix Element Here there are some subtleties—the book uses both sharp and round bras and kets. The mathematical relation holds for both—it is equivalent to the one from Edmonds. However, the graphical method works most simply with round brackets—the vertical lines for the sharp brackets are hard to see, and the diagram shown is the one for that case. This has been corrected from the book.	
$(\gamma j \mathfrak{M}_{\lambda} \gamma' j') = \sum_{m \mu m'} \frac{\langle \gamma jm \mathfrak{M}_{\lambda \mu} \gamma' j' m' \rangle}{\sqrt{(2j+1)(2j'+1)}} (-1)^{j-m} \times \begin{pmatrix} j & \lambda & j' \\ -m & \mu & m' \end{pmatrix}$	
Rotation Operator	
Ω representation $\langle \gamma' \Omega' D(R) \gamma \Omega \rangle = \delta_{\gamma \gamma'} \langle \Omega' - R \Omega \rangle$	
jm-Representation $\langle \gamma jm D(R) \gamma' j' m' \rangle = \delta_{\gamma \gamma'} \delta_{jj'} D_{mm'}^j(R)$	
Inverse Rotation Matrix $\langle \gamma jm D^\dagger(R) \gamma' j' m' \rangle = \delta_{\gamma \gamma'} \delta_{jj'} D_{m' m}^{j*}(R)$ $= \langle \gamma jm D^{-1}(R) \gamma' j' m' \rangle = \delta_{\gamma \gamma'} \delta_{jj'} D_{m m'}^j(R^{-1})$	

Table 11.4. Sums of $3jm$ Symbol Products.

Analytical Expression	Graphical Representation
$3j$ Symbol	
$\sum_{\mu_i} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} = \sum_{\mu_i} (-1)^{j_1 - \mu_1 + j_2 - \mu_2 + j_3 - \mu_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ -\mu_1 & -\mu_2 & -\mu_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} = \{j_1 j_2 j_3\}$	
$6j$ Symbol	
$\begin{aligned} & \sum_{\mu_i x_i} (-1)^{k_1 - x_1 + k_2 - x_2 + k_3 - x_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} \begin{pmatrix} j_1 & k_2 & k_3 \\ \mu_1 & -x_2 & x_3 \end{pmatrix} \begin{pmatrix} k_1 & j_2 & k_3 \\ x_1 & \mu_2 & -x_3 \end{pmatrix} \begin{pmatrix} k_1 & k_2 & j_3 \\ -x_1 & x_2 & \mu_3 \end{pmatrix} \\ &= \sum_{\mu_i x_i} (-1)^{j_1 - \mu_1 + j_2 - \mu_2 + j_3 - \mu_3 + k_1 - x_1 + k_2 - x_2 + k_3 - x_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} \begin{pmatrix} j_1 & k_2 & k_3 \\ -\mu_1 & -x_2 & x_3 \end{pmatrix} \begin{pmatrix} k_1 & j_2 & k_3 \\ x_1 & -\mu_2 & -x_3 \end{pmatrix} \\ & \quad \times \begin{pmatrix} k_1 & k_2 & j_3 \\ -x_1 & x_2 & -\mu_3 \end{pmatrix} = \left\{ \begin{matrix} j_1 & j_2 & j_3 \\ k_1 & k_2 & k_3 \end{matrix} \right\} \end{aligned}$	
$9j$ Symbol	
$\begin{aligned} & \sum_{\mu_i x_i \lambda_i} \begin{pmatrix} l_1 & j_2 & l_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} \begin{pmatrix} k_1 & k_2 & k_3 \\ x_1 & x_2 & x_3 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ \lambda_1 & \lambda_2 & \lambda_3 \end{pmatrix} \begin{pmatrix} j_1 & k_1 & l_1 \\ \mu_1 & x_1 & \lambda_1 \end{pmatrix} \begin{pmatrix} j_2 & k_2 & l_2 \\ \mu_2 & x_2 & \lambda_2 \end{pmatrix} \begin{pmatrix} j_3 & k_3 & l_3 \\ \mu_3 & x_3 & \lambda_3 \end{pmatrix} \\ &= \sum_{\mu_i x_i \lambda_i} (-1)^{j_1 - \mu_1 + j_2 - \mu_2 + j_3 - \mu_3 + k_1 - x_1 + k_2 - x_2 + k_3 - x_3 + l_1 - \lambda_1 + l_2 - \lambda_2 + l_3 - \lambda_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ \mu_1 & \mu_2 & \mu_3 \end{pmatrix} \begin{pmatrix} k_1 & k_2 & k_3 \\ x_1 & x_2 & x_3 \end{pmatrix} \\ & \quad \times \begin{pmatrix} l_1 & l_2 & l_3 \\ \lambda_1 & \lambda_2 & \lambda_3 \end{pmatrix} \begin{pmatrix} j_1 & k_1 & l_1 \\ -\mu_1 & -x_1 & -\lambda_1 \end{pmatrix} \begin{pmatrix} j_2 & k_2 & l_2 \\ -\mu_2 & -x_2 & -\lambda_2 \end{pmatrix} \begin{pmatrix} j_3 & k_3 & l_3 \\ -\mu_3 & -x_3 & -\lambda_3 \end{pmatrix} = \left\{ \begin{matrix} j_1 & j_2 & j_3 \\ k_1 & k_2 & k_3 \\ l_1 & l_2 & l_3 \end{matrix} \right\} \end{aligned}$	

represents the complete sum over j and m of scalar products of irreducible tensors of rank j

$$\sum_{j=0}^{\infty} N_j \sum_{m=-j}^j \langle |\mathfrak{M}|jm\rangle \langle jm|\mathfrak{N}\rangle,$$

where N_j is the weight factor; $N_j = 2j + 1$ for a summation involving the $3jm$ symbols, $3nj$ symbols and the functions of the type $D_{mm'}^j(R)$, $C_{jm}(\Omega) \equiv [4\pi/(2j+1)]^{1/2} Y_{jm}(\Omega)$, or $P_j(\cos \omega)$. On the other hand, $N_j = 1$ for summations involving $C_{j_1 m_1 j_2 m_2}^{j m}$, $U(j_1 j_2 j_3 j_4; jj')$, or $Y_{jm}(\Omega)$. For example,

$$\begin{aligned} \sum_{e\varepsilon} (2e+1)(-1)^{e-\varepsilon} \begin{pmatrix} a & b & e \\ \alpha & \beta & -\varepsilon \end{pmatrix} \begin{pmatrix} a' & b' & e \\ \alpha' & \beta' & \varepsilon \end{pmatrix} & \quad \begin{array}{c} a\alpha \quad a'\alpha' \\ \quad \quad e \\ b\beta \quad b'\beta' \end{array} \\ \sum_{lm} Y_{lm}(\Omega) Y_{lm}^*(\Omega) = \langle \Omega | lm \rangle \langle lm | \Omega' \rangle & \quad \begin{array}{c} \Omega \quad l \quad \Omega' \\ \text{---} \text{---} \text{---} \end{array} \\ \sum_C (2C+1) \begin{Bmatrix} A & B & C \\ a & b & c \end{Bmatrix} \begin{Bmatrix} A & B & C \\ a & b & c' \end{Bmatrix} & \quad \begin{array}{c} \text{Diagram 1} + \text{Diagram 2} \end{array} \end{aligned}$$

A Clebsch-Gordan coefficient may be represented as a sum over $c'\gamma'$ of products of two $3jm$ symbols (Table 11.11)

$$\begin{aligned} C_{a\alpha b\beta}^{c\gamma} &\equiv \langle a\alpha b\beta | c\gamma \rangle = (-1)^{a-b-c} \sum_{c'\gamma'} (2c'+1) \begin{pmatrix} a & b & c' \\ \alpha & \beta & \gamma' \end{pmatrix} \begin{pmatrix} c' & c & 0 \\ \gamma' & \gamma & 0 \end{pmatrix} \\ &= (-1)^{2a} \sum_{c'\gamma'} (2c'+1) (-1)^{a-\alpha+b-\beta+c'-\gamma'} \begin{pmatrix} b & a & c' \\ -\beta & -\alpha & -\gamma' \end{pmatrix} \begin{pmatrix} c' & c & 0 \\ \gamma' & \gamma & 0 \end{pmatrix}, \end{aligned}$$

$$(-1)^{2a} \begin{array}{c} a'\alpha' \\ \quad \quad c' \\ b\beta \end{array} \begin{array}{c} - \\ + \end{array} \begin{array}{c} c\gamma \end{array}$$

11.2.4 Invariant Integration over Directions

A double integral of a bilinear form

$$\int \langle |\mathfrak{N}|\Omega \rangle \frac{d\Omega}{\Omega_0} \langle \Omega | \mathfrak{N} \rangle, \quad (11.2.5)$$

over the total solid angle 4π ($0 \leq \vartheta \leq \pi$, $0 \leq \varphi < 2\pi$) is represented by a dashed Ω line which links $|\Omega\rangle$ and $\langle \Omega|$ lines of subdiagrams corresponding to the integrand factors²

$$\int \boxed{\mathfrak{M}} \text{---} d\Omega \text{---} \boxed{\mathfrak{N}} = \boxed{\mathfrak{M}} \text{---} \overset{\Omega}{\text{---}} \boxed{\mathfrak{N}}$$

The resultant Ω -line has the same direction as the original $|\Omega\rangle$ - and $\langle \Omega|$ -lines. In this case Ω_0 is the normalization constant; for integration of the standard spherical harmonics we have $\Omega_0 = 1$,

$$\int Y_{lm}^*(\Omega) Y_{l'm'}(\Omega) d\Omega = \langle lm | \Omega \rangle \langle \Omega | l'm' \rangle, \quad lm \text{---} \overset{\Omega}{\text{---}} \text{---} l'm' \quad (11.2.6)$$

whereas $\Omega_0 = 4\pi$ for integration of the functions $C_{lm}(\Omega) = [4\pi/(2l+1)]^{1/2} Y_{lm}(\Omega)$.


11.2.5 Integration over Rotation Parameters

The integration of a bilinear form with respect to parameters which specify the rotation R (the Euler angles $\{\alpha, \beta, \gamma\}$ or the angles $\{\omega, \theta, \Phi\}$) over the entire domain of their definition ($R_0 \equiv \int dR$, see Sec. 4.10)

$$\int \langle |\mathfrak{M}(R)| \rangle \langle |\mathfrak{N}(R^{-1})| \rangle \frac{dR}{R_0} \quad (11.2.7)$$

²Note this is not denoted in boldface!

Table 11.5. Basic Operations of the Theory.

Analytical Expression	Graphical Representation
Multiplication	
$\langle \mathfrak{M}\rangle \langle \mathfrak{N} \rangle$	
Invariant Summation over Momentum Projection (Scalar Product of irreducible tensors)	
$\sum_m \langle \mathfrak{N} jm\rangle \langle jm \mathfrak{N} \rangle \equiv (\mathfrak{N}_j \cdot \mathfrak{N}_j)$	$\sum_m \text{[Box M]} \xrightarrow{jm} \text{[Box N]} = \text{[Box M]} \xrightarrow{j} \text{[Box N]}$
Complete Summation over Angular Momentum and Projection	
$\sum_{jm} \langle \mathfrak{M} jm\rangle N_j \langle jm \mathfrak{N} \rangle$	$\sum_j \text{[Box M]} \xrightarrow{j} \text{[Box N]} = \text{[Box M]} \xrightarrow{j} \text{[Box N]}$
Completeness Condition	
$\sum_{jm} jm\rangle N_j \langle jm = 1$	
Irreducible Tensor Product of Irreducible Tensors	
$\sum_{m_1 m_2} \langle \mathfrak{M} i_1 m_1\rangle \langle \mathfrak{N} j_2 m_2\rangle C_{j_1 m_1 j_2 m_2}^{JM} \equiv \{ \mathfrak{M}_{j_1} \otimes \mathfrak{N}_{j_2} \}_{JM}$	
Complete Invariant Integration over Direction Angles	
$\int \langle \mathfrak{M} \Omega\rangle d\Omega \langle \Omega \mathfrak{N} \rangle$	$\int \text{[Box M]} \xrightarrow{d\Omega} \text{[Box N]} = \text{[Box M]} \xrightarrow{\Omega} \text{[Box N]}$
Completeness Condition	
$\int \Omega\rangle d\Omega \langle \Omega = 1$	
Complete Integration over Rotation Parameters	
$\int \langle \mathfrak{M}(R^{-1}) \rangle \frac{dR}{R_0} \langle \mathfrak{N}(R) \rangle$	$\int \text{[Box M]} \stackrel{dR}{R_0} \text{[Box N]} = \text{[Box M]} \stackrel{R}{\text{[Box N]}}$
Completeness Condition	
$\int \hat{R}^{-1} \hat{R} \frac{dR}{R_0} = 1$	

may be represented by a conjunction of corresponding R - and R^{-1} -lines of subdiagrams which display the integrand factors.

$$\boxed{\mathfrak{M}} \stackrel{dR}{R_0} \stackrel{dR}{R_0} \boxed{\mathfrak{N}}$$

Note that integrals of three or more functions of R may be reduced to standard bilinear forms, using the Clebsch-Gordan expansion.

11.3 Rules of the graphical technique

In this section the rules of calculation by the graphical method will be formulated, specifically, the rules for the reduction of diagrams to standard diagrams which represent the basis functions of the angular momentum theory.


11.3.1 Deformation of Diagrams

Expressions which appear in calculations based on the theory of angular momentum can be rather complicated. Corresponding diagrams consist of many lines, thin and thick, solid and dashed, single and double. These lines are partly linked to some nodes. One can clearly distinguish two types of lines: internal and external. Internal lines have both ends linked to nodes, whereas external lines have one end linked to a node but the other free. Any identical transformation of an analytic expression corresponds to some transformation of the internal lines of the diagram with external lines remaining unchanged.

If the structure of the inner part of a diagram is unimportant, it may be replaced by a block with the same external lines, because these are just the external lines which determine the transformation properties of corresponding diagram (or sub-diagram) under rotation of the coordinate system. For instance, any closed diagram without external lines is invariant under such rotation and may be reduced to some of the $3nj$ symbols or their combinations.

As pointed out in Sec. 11.1.1, the length of the lines, their curvature and orientation are not important. Consequently, any diagram may be arbitrarily rotated and deformed, although the node signs have to be reversed for those nodes where the order of momentum coupling becomes changed.

Thus, any deformation of a diagram performed under the proper check for the node signs does not change the meaning of the diagram, i.e., the deformed and original diagrams represent the same expression. Such diagrams will be called identical. An example of identical diagrams is given below:



Two diagrams will be called topologically similar, if it is possible to make them coincident by means of rotations, deformations and/or reflections. Topologically similar diagrams have the same numbers of nodes and of lines, solid and dashed, but they may differ by directions of lines and by node signs.

The expressions corresponding to two topologically similar diagrams are equal in absolute value, but may differ by a phase factor. One may find this factor by means of successive reductions of one diagram to the other, i.e., by reversals of node signs and line directions (where necessary) in accordance with the rules formulated in Secs. 11.3.2 and 11.3.3. In this way one may reduce any diagram to the topologically similar diagram of some basis function of the theory, and determine the additional phase factor. Several examples of such transformation for the $3j$, $6j$, and $9j$ symbols are displayed in Table 11.6.

11.3.2 Change of Node Sign

The reversal of the sign in a node joining $(j_1 j_2 j_3)$ -lines corresponds to the change of the momentum-coupling order. This transformation is equivalent to introducing the additional phase factor $(-1)^{j_1+j_2+j_3}$. For example,

$$\begin{array}{c} \boxed{} \\ \begin{array}{c} \nearrow j_1 \\ \rightarrow j_2 \\ \searrow j_3 \end{array} \end{array} + = (-1)^{j_1+j_2+j_3} \begin{array}{c} \boxed{} \\ \begin{array}{c} \nearrow j_1 \\ \rightarrow j_2 \\ \searrow j_3 \end{array} \end{array} -$$

The reversal of signs of all the nodes in a diagram produces an additional phase factor of $(-1)^{2J_0+J_1}$, where J_0 and J_1 are the algebraic sums of all the momenta for the internal and external lines, respectively. For example,

Table 11.6. Symmetry of $3nj$ Symbol Diagrams.






















$3j$ Symbol $\{abc\}$				
Reversal of Node Sign and Line Direction, and Argument Permutation				
	$= (-1)^{a+b+c}$		$= (-1)^{a+b+c}$	
	$=$		$= (-1)^{2a}$	
$6j$ Symbol $\left\{ \begin{matrix} a & b & c \\ A & B & C \end{matrix} \right\}$				
Reversal of Node Sign				
	$= (-1)^{2(A+B+C)}$		$= (-1)^{a+b+c}$	
Reversal of Line Direction				
	$= (-1)^{2(A+B+C)}$		$=$	
			$= (-1)^{2(A+B+C)}$	
Argument Permutation				
	$=$		$=$	
			$=$	
	$=$		$=$	
			$=$	

Table 11.6. (Cont'd)

$$9j \text{ Symbol } \begin{Bmatrix} a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \\ b_1 & b_2 & b_3 \end{Bmatrix}$$

Reversal of Node Sign

$$\begin{array}{c} \begin{array}{c} + \\ \text{Diagram 1} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 2} \end{array} \end{array} = \begin{array}{c} \begin{array}{c} - \\ \text{Diagram 3} \end{array} \\ \begin{array}{c} + \\ \text{Diagram 4} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 5} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 6} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} - \\ \text{Diagram 7} \end{array} \\ \begin{array}{c} + \\ \text{Diagram 8} \end{array} \end{array}$$

Reversal of Line Direction

$$\begin{array}{c} \begin{array}{c} + \\ \text{Diagram 1} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 2} \end{array} \end{array} = \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 3} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 4} \end{array} \end{array} = (-1)^{2(a_2+b_3+c_1)} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 5} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 6} \end{array} \end{array} = (-1)^{2(a_2+b_3+c_1)} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 7} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 8} \end{array} \end{array}$$


Argument Permutation

$$\begin{array}{c} \begin{array}{c} + \\ \text{Diagram 1} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 2} \end{array} \end{array} = \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 3} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 4} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 5} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 6} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 7} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 8} \end{array} \end{array}$$

$$\begin{array}{c} \begin{array}{c} + \\ \text{Diagram 9} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 10} \end{array} \end{array} = \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 11} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 12} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 13} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 14} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 15} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 16} \end{array} \end{array}$$

$$\begin{array}{c} \begin{array}{c} + \\ \text{Diagram 17} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 18} \end{array} \end{array} = \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 19} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 20} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 21} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 22} \end{array} \end{array} = (-1)^{J_0} \begin{array}{c} \begin{array}{c} + \\ \text{Diagram 23} \end{array} \\ \begin{array}{c} - \\ \text{Diagram 24} \end{array} \end{array}$$

where $J_0 \equiv a_1 + a_2 + a_3 + b_1 + b_2 + b_3 + c_1 + c_2 + c_3$



11.3.3 Change of Direction of External Lines

The direction of an external j -line specifies the form (covariant or contravariant, i.e. $|jm\rangle$ or $\langle jm|$) of an irreducible tensor of rank j (Sec. 11.1.1). The conversion of covariant components of a tensor into contravariant ones, as well as the inverse transformation, is carried out by contraction (over m) of the tensor in question with the tensor $\begin{pmatrix} j \\ mm' \end{pmatrix} = (-1)^{j+m} \delta_{m-m'}$, which may be considered as the metric tensor for the $(2j+1)$ dimensional space of the functions $|jm\rangle$. Such a transformation reads

$$\sum_{m'} \langle | \mathfrak{M} | jm' \rangle \begin{pmatrix} j \\ m' m \end{pmatrix} = \langle jm | \mathfrak{M} | \rangle \quad (11.3.1)$$

$$\sum_{m'} \boxed{\mathfrak{M}} \xrightarrow{j m' j m'} \longleftrightarrow \xleftarrow{j m} = \boxed{\mathfrak{M}} \xrightarrow{j m}$$

and the inverse transformation is

$$\sum_{m'} \begin{pmatrix} j \\ m m' \end{pmatrix} \langle jm' | \mathfrak{M} | \rangle = \langle | \mathfrak{M} | jm \rangle \quad (11.3.2)$$

$$\sum_{m'} \boxed{\mathfrak{M}} \xleftarrow{j m' j m'} \longleftrightarrow \xrightarrow{j m} = \boxed{\mathfrak{M}} \xrightarrow{j m}$$

Thus, a reversal of the direction of any external j -line (jm being fixed) corresponds to the transformation

$$\langle | \mathfrak{M} | jm \rangle \equiv \mathfrak{M}_{jm} \rightleftharpoons \langle jm | \mathfrak{M} | \rangle \equiv (-1)^{j-m} \mathfrak{M}_{j-m}, \quad (11.3.3)$$

in the associated analytic expression. In other words, the replacement of a single outward arrow by a double inward one corresponds to the introduction of the phase factor $(-1)^{j-m}$, and substitution of $-m$ for m , and vice versa.

For example,

$$\sum_{\alpha'} \begin{pmatrix} a \\ \alpha' \alpha \end{pmatrix} \begin{pmatrix} a & b & c \\ \alpha' & \beta & \gamma \end{pmatrix} = (-1)^{a-\alpha} \begin{pmatrix} a & b & c \\ -\alpha & \beta & \gamma \end{pmatrix}, \quad (11.3.4)$$

and, conversely,

$$\sum_{\alpha'} \begin{pmatrix} a \\ \alpha \alpha' \end{pmatrix} (-1)^{a-\alpha'} \begin{pmatrix} a & b & c \\ -\alpha' & \beta & \gamma \end{pmatrix} = \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix}. \quad (11.3.5)$$


11.3.4 Change of Direction of Integral Lines

The reversal of the direction of any internal j -line produces the phase factor $(-1)^{2j}$ in the associated analytic expression, because

$$\sum_m (-1)^{j-m} \mathfrak{M}_{jm} \mathfrak{N}_{j-m} = (-1)^{2j} \sum_m (-1)^{j-m} \mathfrak{M}_{j-m} \mathfrak{N}_{jm}. \quad (11.3.6)$$

For a closed diagram which represents a $3nj$ symbol, reversal of the directions of the j -lines produces the factor $(-1)^{2J}$ in the associated expression, where J is the algebraic sum of all the momenta of the $3nj$ symbol (Table 11.6).

The reversal of the directions of three internal j -lines ($j_1 j_2 j_3$) coupled together in a node does not produce any additional phase factor, since $(-1)^{2(j_1+j_2+j_3)} = 1$. For example,




For a closed diagram associated with a $3nj$ symbol the reversal of directions of all the j -lines and simultaneous reversal of signs of all the nodes do not change the meaning of the expression.

11.3.5 Linking Subdiagrams

In this Section the graphical rules for the multiplication of subdiagrams are formulated. These are the rules for linking the subdiagrams (blocks) into one united diagram.


- (a) Two subdiagrams representing factors $\mathfrak{M}$ and $\mathfrak{N}$, which have at least one identical (for both subdiagrams) j -line, may be linked together into the combined diagram which displays the product of these factors:



The necessary condition for such a linking is that at least one of these subdiagrams should have no external line. The graphical rule is equivalent to the relation

$$\left(\sum_m \langle jm | \mathfrak{M} | jm \rangle \right) \times \left(\sum_{m'} \langle jm' | \mathfrak{N} | jm' \rangle \right) = (2j+1) \sum_{mm'} \langle jm | \mathfrak{M} | jm' \rangle \langle jm' | \mathfrak{N} | jm \rangle. \quad (11.3.7)$$


- (b) Two subdiagrams representing the factors $\mathfrak{M}$ and $\mathfrak{N}$ which have at least one identical (for both subdiagrams) node, i.e., three coupled j -lines ($j_1 j_2 j_3$) may be linked together into the combined diagram which corresponds to the product of these factors:




This is equivalent to the relation


$$\left[\sum_{m_1 m_2 m_3} \langle 00 | \mathfrak{M} | j_1 m_1 j_2 m_2 j_3 m_3 \rangle \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \right] \times \left[\sum_{m'_1 m'_2 m'_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix} \langle j_1 m'_1 j_2 m'_2 j_3 m'_3 | \mathfrak{N} | \rangle \right] \\ = \{j_1 j_2 j_3\} \sum_{m_1 m_2 m_3} \langle 00 | \mathfrak{M} | j_1 m_1 j_2 m_2 j_3 m_3 \rangle \langle j_1 m_1 j_2 m_2 j_3 m_3 | \mathfrak{N} | \rangle \quad (11.3.8)$$

For example,



As in Scheme (a), at least one of these subdiagrams should have no external lines. However, in the other subdiagrams the joined j -lines may be external. For instance,





Note that Scheme (a) is a special case of Scheme (b) when one of the momenta j_1, j_2, j_3 equals zero.

11.3.6 Cutting Diagram into Subdiagrams

The scalar product of two n -fold irreducible tensors of rank $j_1, j_2, \dots, j_n$, i.e., the n -fold invariant sum over projections

$$(\mathfrak{M}_{j_1 j_2 \dots j_n} \cdot \mathfrak{N}_{j_1 j_2 \dots j_n}) = \sum_{m_1 m_2 \dots m_n} \langle 00 | \mathfrak{M} | j_1 m_1 j_2 m_2 \dots j_n m_n \rangle \langle j_1 m_1 j_2 m_2 \dots j_n m_n | \mathfrak{N} | \rangle \quad (11.3.9)$$

may be transformed into a k -fold complete sum (over momenta $X_1, X_2, \dots, X_n$ with $k = n - 3$ and $n \geq 4$) of the product of factors:

$$(\mathfrak{M}_{j_1 j_2 \dots j_n} \cdot \mathfrak{N}_{j_1 j_2 \dots j_n}) = \sum_{X_1 X_2 \dots X_k}^{k=n-3} (2X_1 + 1)(2X_2 + 1) \dots (2X_k + 1) \mathfrak{M}_{j_1 j_2 \dots j_n}(X_1 X_2 \dots X_k) \mathfrak{N}_{j_1 j_2 \dots j_n}(X_1 X_2 \dots X_k). \quad (11.3.10)$$

These factors represent contractions of the original tensors and $3jm$ symbols:

$$\begin{aligned} & \mathfrak{M}_{j_1 j_2 \dots j_n}(X_1 X_2 \dots X_k) \\ \equiv & \sum_{m_i M_i} \mathfrak{M}_{j_1 m_1 j_2 m_2 \dots j_n m_n} \begin{pmatrix} j_1 & j_2 & X_1 \\ m_1 & m_2 & -M_1 \end{pmatrix} (-1)^{X_1 - M_1} \begin{pmatrix} X_1 & j_3 & X_2 \\ M_1 & m_3 & -M_2 \end{pmatrix} (-1)^{X_2 - M_2} \dots \begin{pmatrix} X_k & j_{n-1} & j_n \\ M_k & m_{n-1} & m_n \end{pmatrix} \\ \equiv & \sum_{m_i M_i} \mathfrak{N}_{j_1 j_2 \dots j_n}^{m_1 m_2 \dots m_n} \begin{pmatrix} j_1 & j_2 & X_1 \\ m_1 & m_2 & -M_1 \end{pmatrix} (-1)^{\mathfrak{N}_{j_1 j_2 \dots j_n}(X_1 X_2 \dots X_k)} \\ & \begin{pmatrix} X_1 & j_3 & X_2 \\ M_1 & m_3 & -M_2 \end{pmatrix} (-1)^{X_2 - M_2} \dots \begin{pmatrix} X_k & j_{n-1} & j_n \\ M_k & m_{n-1} & m_n \end{pmatrix} \end{aligned} \quad (11.3.11)$$


Again, at least one of these contracted quantities must be a true invariant, i.e., should not depend on any additional momentum projection or direction.

For $n \leq 3$ the sum over X_i may be reduced to the direct product of the contracted tensors. For example,

$$(\mathfrak{M}_{j_1 j_2 \dots j_n} \cdot \mathfrak{N}_{j_1 j_2 \dots j_n}) = \mathfrak{M}_{j_1 j_2 \dots j_n} \cdot \mathfrak{N}_{j_1 j_2 \dots j_n}. \quad (11.3.12)$$


This means that the original sum becomes factorized. Therefore, the cases $n \leq 3$ are the most important ones for practical applications.

The analytic transformations under study correspond to cutting the associated diagram into two subdiagrams which represent two factors. The necessary condition is that at least one of these subdiagrams should have no more external lines (apart from those j -lines which link the subdiagram together)



The most important cases of cutting of one, two, three, as well as four, five, and six j -lines are presented in Table 11.9.

In many cases, the successive use of these rules allows one to divide a diagram into several (three or more) subdiagrams representing invariant factors. For example,




The most important cases for the application of the cutting rules are in the separation of external lines of a diagram, i.e., in extracting the dependence on all the external variables in the form of the standard functions of the theory ($C_{j_1 m_1 j_2 m_2}^j, D_{mm'}^j(R), C_{jm}(\Omega)$, etc.). Such a separation is the essence of the Wigner-Eckart theorem which will be discussed in Sec. 11.4.2.

11.3.7 Graphical Method of Summation

According to Sec. 11.2.3, a complete invariant sum over angular momentum j and projection m is graphically represented by a thick j -line. Such a summation may be easily performed by the diagram technique, if this thick j -line connects two identical nodes.

Rule $\sum$: Any thick j -line which connects identical pairs of thin j -lines, may be removed provided the ends of the thin lines with equal momenta are linked together. This is displayed below:



Such a graphical operation does not change the meaning of a diagram and corresponds to the equation

$$\sum_{jm} (2j+1) \langle j_1 j_2 j m | \mathfrak{M} | j_1 j_2 j m \rangle = \sum_{m_1 m_2} \langle j_1 m_1 j_2 m_2 | \mathfrak{M} | j_1 m_1 j_2 m_2 \rangle \quad (11.3.13)$$


where

$$\langle j_1 m_1 j_2 m_2 | \mathfrak{M} | j'_1 m'_1 j'_2 m'_2 \rangle = \sum_{jm} (2j+1) \begin{pmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{pmatrix} \langle j_1 j_2 j m | \mathfrak{M} | j'_1 j'_2 j m \rangle \begin{pmatrix} j'_1 & j'_2 & j \\ m'_1 & m'_2 & m \end{pmatrix}. \quad (11.3.14)$$

For example,

$$\sum_{c\gamma} (2c+1) (-1)^{a-\alpha+b-\beta+c-\gamma} \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} \begin{pmatrix} a & b & c \\ \alpha' & \beta' & \gamma' \end{pmatrix} = \sum_{c\gamma} (2c+1) \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \begin{pmatrix} a & b & c \\ \alpha' & \beta' & \gamma' \end{pmatrix} = \delta_{\alpha\alpha'} \delta_{\beta\beta'}, \quad (11.3.15)$$

$$\sum_X (2X+1) \left\{ \begin{matrix} a & b & c \\ b' & d & e \\ c & e & X \end{matrix} \right\} = \{abc\} \{bde\} \frac{\delta_{bb'}}{(2b+1)} \quad (11.3.16)$$



When a common thick j -line is included into two or more unlinked subdiagrams, one should preliminarily link these subdiagrams with the aid of the rules of subdiagram multiplication (Sec. 11.3.5) in order to arrive at one thick j -line. Only after this operation can one perform graphical summation over j and m according to Rule $\sum$. These combined graphical operations are displayed in Table 11.9.

The complete invariant summation over j and m may also be performed graphically in the cases when the lines adjacent to the thick j -line are dashed Ω -lines. For instance,

$$\sum_{lm} Y_{lm}(\Omega) Y_{lm}^*(\Omega') = \delta(\Omega - \Omega') \quad (11.3.17)$$

$$\begin{array}{c} \Omega \\ \text{---} \end{array} \begin{array}{c} l \\ \text{---} \end{array} \begin{array}{c} \Omega' \\ \text{---} \end{array} = \begin{array}{c} \Omega \\ \text{---} \end{array} \begin{array}{c} \Omega' \\ \text{---} \end{array}$$

$$\sum_{JMM'} (2J+1) D_{MM'}^J(R_1) D_{M'M}^J(R_2) = 16\pi^2 \delta(R_1 - R_2), \quad (11.3.18)$$

In fact, all the rules for complete invariant summation formulated in this section follow from the completeness of $|jm\rangle$ -functions.

$$\sum_{jm} |jm\rangle \langle jm| = \sum_{jm} |jm\rangle (2j+1) \langle jm| = 1, \quad (11.3.19)$$

11.3.8 Replacing An Internal Ω -Line by a Thick j -Line

Each internal Ω -line which represents invariant integration over direction angles $\Omega\{\vartheta, \varphi\}$ may be replaced by a thick j -line representing the complete invariant sum over j and m , according to

$$\int d\Omega \langle \Omega | \mathfrak{M} | \Omega \rangle = \sum_{jm} \langle jm | \mathfrak{M} | jm \rangle, \quad (11.3.20)$$

where

$$\langle jm | \mathfrak{M} | j'm' \rangle \equiv \iint d\Omega d\Omega' \langle jm | \Omega' \rangle \langle \Omega' | \mathfrak{M} | \Omega \rangle \langle \Omega | j'm' \rangle. \quad (11.3.21)$$

Thus, all the Ω -lines in the inner part of a diagram may be converted into the j -lines. Hence, it is sufficient to formulate the rules of graphical operations only for diagrams in which all the internal lines are j -lines.


11.3.9 Elimination of $j = 0$ Line

In the particular case of $j = 0$ the corresponding expression is independent of the angular variables $\Omega\{\vartheta, \varphi\}$. Hence, the corresponding j -line may be removed from a diagram without changing its meaning. However, there remain the nodes which were initially connected with this $j = 0$ line. These nodes may also be extracted from the diagram in the form of a separated subdiagram corresponding to the factor

$$K = (\pm 1)^{2j_1} \frac{\delta_{j_1 j_2}}{\sqrt{2j_1 + 1}}. \quad (11.3.22)$$

The phase of K depends on the node sign and line direction. K is positive if the node sign corresponds to the cyclic order of the coupling momenta (j_1, j_1, j_2) , the j_1 -line is directed toward the node and the j_2 -line is directed outward from the node. In this case the resultant $j_1 = j_2$ -line has the same direction as that before the node's removal. In other cases the phase of K may be determined by means of the reduction of the diagram under study to the above case, using the rules of Sec. 11.3.2.


Thus elimination of the $j = 0$ line may be illustrated by



Some special cases of the $3jm$ and $3nj$ symbols with $j = 0$ are presented in Table 11.7.

11.3.10 Elimination of the $R = 1$ Line

In the particular case $R = 1$, a double dashed R -line corresponds to no rotation, i.e., it represents a fictitious dependence. Hence, an $R = 1$ line may be removed and the lines adjacent to the R -line have to be closed. This operation does not change the meaning of the diagram.



Some examples of the expressions and diagrams containing $R = 1$ are given in Table 11.8.

A summary of the rules of graphical operations is presented in Table 11.9.

11.4 Summary of the graphical technique

The preceding sections present the principles of the graphical methods of angular momentum theory, i.e., graphical representations of the basis functions and operations (Secs. 11.1 and 11.2) as well as the rules of handling them (Sec. 11.3). This section resumes the discussion of the general properties of diagrams, the Wigner-Eckart theorem in diagrammatic form and the scheme for the application of the graphical technique.

11.4.1 General Properties of Diagrams

Let us summarize the main properties of diagrams which are consequences of the graphical relationships discussed above.

1. Any diagram may be arbitrarily turned and deformed, bent or stretched. Along with these continuous transformations, discrete transformations such as reflection and inversion are also allowed.

A transformed diagram corresponds to the same analytic expression provided node signs are reversed for those nodes in which the cyclic order of the coupled momenta becomes opposite to the initial one.

2. Each j -line holds a fixed value of angular momentum j in spite of any deformation of the line. For an external j -line, the projection m is fixed as well as the j -value. Moreover, j and m remain unchanged along the line even if any block (subdiagram) is inserted into the line, provided this block has no additional external j -lines.
3. The total angular momentum of any diagram or subdiagram, i.e., the vector sum of angular momenta of all external lines, is equal to zero.

$$\sum_i \mathbf{j}_i = 0, \quad \sum_i m_i = 0. \quad (11.4.1)$$

Hence, for a diagram with one external j -line j has to be zero; for a diagram which has two external j -lines associated with j_1 and j_2 one has $j_1 = j_2$.

4. Any diagram may be cut into two subdiagrams in many different ways. The total angular momenta of both these subdiagrams are identical. Moreover, the angular momenta will be the same for all the diagram cross-sections with fixed sets of external lines for each subdiagram.

11.4.2 Generalized Wigner-Eckart Theorem in Diagrammatic Form

Standard functions of angular momentum theory may be used as a basis for the expansion of any expression. However, sometimes all the dependence of the expression on external variables may be factorized in the form of some standard function. Graphically such a function may be separated according to the rules of Sec. 11.3.6. The most important special cases are the following.

- (a) A diagram without external lines (i.e. a closed diagram) is invariant with respect to coordinate-system rotations.
- (b) A diagram with one external line also is invariant. This is a consequence of the fact that, in principle, one cannot distinguish any preferred direction in isotropic space. Therefore, any dependence on one direction is fictitious, so that such an external j -line has to correspond to $j = 0$ and may be eliminated in accordance with the rule of Sec. 11.3.9.

$$\langle 00 | \mathfrak{M} | jm \rangle = \delta_{j0} \delta_{m0} \langle 00 | \mathfrak{M} | 00 \rangle \quad (11.4.2)$$




Table 11.7. Zero Angular Momentum ($J = 0$).

Analytical Expression	Graphical Representation
<i>jm</i> Symbols	
$\begin{pmatrix} a & 0 & b \\ \alpha & 0 & \beta \end{pmatrix} = \frac{\delta_{ab}}{\sqrt{2a+1}} \begin{pmatrix} a \\ \alpha\beta \end{pmatrix} = (-1)^{a+\alpha} \frac{\delta_{ab}\delta_{\alpha,-\beta}}{\sqrt{2a+1}}$	
$\begin{pmatrix} a & b & 0 \\ \alpha & \beta & 0 \end{pmatrix} = \frac{\delta_{ab}}{\sqrt{2a+1}} \begin{pmatrix} a \\ \beta\alpha \end{pmatrix} = (-1)^{a-\alpha} \frac{\delta_{ab}\delta_{\alpha,-\beta}}{\sqrt{2a+1}}$	
$(-1)^{a-\alpha} \begin{pmatrix} a & 0 & b \\ -\alpha & 0 & \beta \end{pmatrix} = \frac{\delta_{ab}\delta_{\alpha\beta}}{\sqrt{2a+1}}$	
$(-1)^{a-\alpha} \begin{pmatrix} a & b & 0 \\ -\alpha & \beta & 0 \end{pmatrix} = (-1)^{2a} \frac{\delta_{ab}\delta_{\alpha\beta}}{\sqrt{2a+1}}$	
<i>nj</i> Symbols	
$\{a b 0\} = \delta_{ab}$	
$\begin{Bmatrix} 0 & 0 & 0 \\ a & b & c \end{Bmatrix} = (-1)^{2a} \frac{\delta_{ab}\delta_{bc}}{\sqrt{2a+1}}$	
$\begin{Bmatrix} a & a' & 0 \\ b & b' & 0 \end{Bmatrix} = (-1)^{2a} \frac{\delta_{aa'}\delta_{bb'}\delta_{ab}}{2a+1}$	
$\begin{Bmatrix} a & a' & 0 \\ b & b' & c \end{Bmatrix} = (-1)^{a+b+c} \frac{\delta_{aa'}\delta_{bb'}}{\sqrt{(2a+1)(2b+1)}} \{a b c\}$	
$\begin{Bmatrix} a & b & c \\ e & 0 & e' \\ f & b' & d \end{Bmatrix} = (-1)^{b+c+e+f} \frac{\delta_{bb'}\delta_{ee'}}{\sqrt{(2b+1)(2e+1)}} \begin{Bmatrix} a & b & c \\ d & e & f \end{Bmatrix}$	

Table 11.8. Zero Rotation ($R = 0$).

Analytical Expression	Graphical Representation
(a) For $R = 1$: $\delta(\Omega' - \hat{R}\Omega) = \delta(\Omega' - \Omega),$ i.e., $\langle \Omega' R \Omega \rangle = \langle \Omega' \Omega \rangle.$	
(b) For $R = 1$: $\delta_{jj'} D_{mm'}^j(000) = \delta_{jj'} \delta_{mm'},$ i.e., $\langle jm D(1) j'm' \rangle = \langle jm j'm' \rangle.$	
(c) For $R^{-1}R = 1$: $\int \delta(\Omega'' - R^{-1}\Omega') d\Omega' \delta(\Omega' - R\Omega) = \delta(\Omega'' - \Omega),$ i.e., $\langle \Omega'' R^{-1} \Omega' \rangle \langle \Omega' R \Omega \rangle = \langle \Omega'' \Omega \rangle.$	
(d) For $R^{-1}R = 1$: $\sum_{m'} D_{m''m'}^j(R^{-1}) D_{m'm}^j(R) = \delta_{m''m},$ i.e., $\langle jm'' D(R^{-1}) j'm' \rangle \langle j'm' D(R) jm \rangle = \langle jm'' jm \rangle.$	

Table 11.9. Rules for the Transformation of Diagrams

Analytical Expression	Graphical Representation
Invariant integration over directions $\int d\Omega \langle \Omega \mathfrak{M} \Omega \rangle = \sum_{JM} \langle JM \mathfrak{M} JM \rangle$ where $\langle JM \mathfrak{M} J'M' \rangle \equiv \iint \langle JM \Omega \rangle d\Omega \langle \Omega \mathfrak{M} \Omega' \rangle d\Omega' \langle \Omega' J'M' \rangle.$ Completeness Condition $\int \Omega \rangle d\Omega \langle \Omega = \sum_{JM} JM \rangle \langle JM = 1$	Replacing dashed Ω-line by thick solid j-line

Table 11.9. Cont'd

Analytical Expression	Graphical Representation
Complete summation over J and M $\sum_{JM} (2J+1) \langle j_1 j_2; JM \mathfrak{M} j_1 j_2; JM \rangle$ $= \sum_{m_1 m_2} \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_1 m_1 j_2 m_2 \rangle,$ $\sum_{JMM'} (2J+1) \langle j_1 j_2; JM \mathfrak{M} j_3 j_4; JM \rangle$ $\times \langle j_3 j_4; JM' \mathfrak{N} j_1 j_2; JM' \rangle$ $= \sum_{\substack{m_1 m_2 \\ m_3 m_4}} \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle$ $\times \langle j_3 m_3 j_4 m_4 \mathfrak{N} j_1 m_1 j_2 m_2 \rangle.$ $\sum_{JMM'M''} (2J+1) \langle j_1 j_2; JM \mathfrak{M} j_3 j_4; JM \rangle$ $\times \langle j_3 j_4; JM' \mathfrak{N} j_5 j_6; JM' \rangle \langle j_5 j_6; JM'' \mathfrak{L} j_1 j_2; JM'' \rangle$ $= \sum_{\substack{m_1 m_2 \\ m_3 m_4 \\ m_5 m_6}} \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle \times$ $\times \langle j_3 m_3 j_4 m_4 \mathfrak{N} j_5 m_5 j_6 m_6 \rangle \langle j_5 m_5 j_6 m_6 \mathfrak{L} j_1 m_1 j_2 m_2 \rangle.$ where $\langle j_1 j_2; JM \mathfrak{M} j_3 j_4; JM \rangle$ $\equiv \sum_{m_i} \begin{pmatrix} j_1 & j_2 & J \\ m_1 & m_2 & M \end{pmatrix} \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle \begin{pmatrix} j_3 & j_4 & J \\ m_3 & m_4 & M \end{pmatrix}.$ Transformation of product of invariant sums into invariant sum of products (at least one of two factors must be a perfect invariant). $\left(\sum_m \langle jm \mathfrak{M} jm \rangle \right) \left(\sum_{m'} \langle jm' \mathfrak{N} jm' \rangle \right)$ $= (2j+1) \sum_{mm'} \langle jm \mathfrak{M} jm' \rangle \langle jm' \mathfrak{N} jm \rangle$ $\left[\sum_{m_1 m_2 m_3} \langle \mathfrak{M} j_1 m_1 j_2 m_2 j_3 m_3 \rangle \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \right]$ $\times \left[\sum_{m_1 m_2 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \langle j_1 m_1 j_2 m_2 j_3 m_3 \mathfrak{N} \rangle \right]$ $= \sum_{m_1 m_2 m_3} \langle \mathfrak{M} j_1 m_1 j_2 m_2 j_3 m_3 \rangle \langle j_1 m_1 j_2 m_2 j_3 m_3 \mathfrak{N} \rangle.$ Expression reduction for $J = 0$ $\sum_{m'} \langle jm \mathfrak{M} j' m' \rangle (-1)^{j'-m'} \begin{pmatrix} j' & 0 & j \\ -m' & 0 & m \end{pmatrix}$ $= \frac{\delta_{jj'}}{\sqrt{2j+1}} \sum_m \langle jm \mathfrak{M} jm \rangle$	Elimination of thick solid j-line Linking subdiagrams containing identical j-line (at least one of subdiagrams should have no other external lines) Elimination of a $J = 0$ line in a diagram

Table 11.9. Cont'd





Analytical Expression	Graphical Representation
Expression reduction for $R = 1$ $\sum_{mm'} \langle jm \mathfrak{M} j'm' \rangle \langle j'm' D(000) jm \rangle$ $= \delta_{jj'} \sum_m \langle jm \mathfrak{M} jm \rangle$	Elimination of the $R = 1$ line 
Transformation of single, double, or triple invariant sum into direct product of two factors (at least one of the factors must be perfect invariant) $\sum_m \langle 00 \mathfrak{M} jm \rangle \langle jm \mathfrak{N} p \rangle = \delta_{j0} \delta_{m0} \langle 00 \mathfrak{M} 00 \rangle$ $\times \langle 00 \mathfrak{N} p \rangle.$	Cutting of a diagram into subdiagrams linked by one, two, or three j-lines (at least one of subdiagrams should have no additional external lines) 
$\sum_{m_1 m_2} \langle j_1 m_1 \mathfrak{M} j_2 m_2 \rangle \langle j_2 m_2 \mathfrak{N} j_1 m_1 p \rangle$ $= \frac{\delta_{j_1 j_2}}{(2j_1 + 1)} \left(\sum_{m_1} \langle j_1 m_1 \mathfrak{M} j_1 m_1 \rangle \right)$ $\times \left(\sum_{m_2} \langle j_2 m_2 \mathfrak{N} j_2 m_2 p \rangle \right)$	
$\sum_{m_1 m_2 m_3} \langle 00 \mathfrak{M} j_1 m_1 j_2 m_2 j_3 m_3 \rangle \langle j_1 m_1 j_2 m_2 j_3 m_3 \mathfrak{N} p \rangle$ $= \langle 0 \mathfrak{M} j_1 j_2 j_3 \rangle \langle j_1 j_2 j_3 \mathfrak{N} p \rangle$	
Here the reduced matrix elements are $\langle 0 \mathfrak{M} j_1 j_2 j_3 \rangle \equiv \sum_{m_1 m_2 m_3} \langle 00 \mathfrak{M} j_1 m_1 j_2 m_2 j_3 m_3 \rangle$ $\times \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}, \quad \langle j_1 j_2 j_3 \mathfrak{N} p \rangle \equiv \sum_{m_1 m_2 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$ $\times \langle j_1 m_1 j_2 m_2 j_3 m_3 \mathfrak{N} p \rangle.$	
Transformation of n-fold invariant sum ($n > 4$) over projections into a k-fold sum over momenta ($k = n - 3$) (at least one of factors under summation must be perfect invariant) $\sum_{\substack{m_1 m_2 \\ m_3 m_4}} \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle \langle j_3 m_3 j_4 m_4 \mathfrak{N} j_1 m_1 j_2 m_2 p \rangle$ $= \sum_J (2J + 1) \langle j_1 j_2 J \mathfrak{M} j_3 j_4 J \rangle \langle j_3 j_4 J \mathfrak{N} j_1 j_2 J p \rangle$	Cutting of a diagram into two subdiagrams linked by n_j-lines ($n > 4$) (at least one of subdiagrams should have no additional external lines) 
where $\langle j_1 j_2 J \mathfrak{M} j_3 j_4 J \rangle \equiv \sum_{\substack{m_1 m_2 M \\ m_3 m_4}} \begin{pmatrix} j_1 & j_2 & J \\ m_1 & m_2 & M \end{pmatrix}$ $\times \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle \begin{pmatrix} j_3 & j_4 & J \\ m_3 & m_4 & M \end{pmatrix}$ and similarly for $\langle j_3 j_4 J \mathfrak{N} j_1 j_2 J p \rangle$.	

Table 11.9. Cont'd

Analytical Expression	Graphical Representation
$\sum_{m_1 \dots m_5} \langle 00 \mathfrak{M} j_1 m_1 \dots j_5 m_5 \rangle \langle j_1 m_1 \dots j_5 m_5 q \mathfrak{N} p \rangle$ $= \sum_{J_{12} J_{45}} (2J_{12} + 1) (2J_{45} + 1) \langle 0 \mathfrak{M} J_{12} j_3 J_{45} \rangle$ $\times \langle J_{12} j_3 J_{45} q \mathfrak{N} p \rangle$	
where $\langle 0 \mathfrak{M} J_{12} j_3 J_{45} \rangle = \sum_{m_i M_{ik}} \langle 00 \mathfrak{M} j_1 m_1 \dots j_5 m_5 \rangle$ $\times \begin{pmatrix} j_1 & j_2 & J_{12} \\ m_1 & m_2 & M_{12} \end{pmatrix} (-1)^{j_3 - m_3} \begin{pmatrix} J_{12} & j_3 & J_{45} \\ M_{12} & -m_3 & M_{45} \end{pmatrix} \begin{pmatrix} j_4 & j_5 & J_{45} \\ m_4 & m_5 & M_{45} \end{pmatrix},$ and similarly for $\langle J_{12} j_3 J_{45} q \mathfrak{N} p \rangle$.	
$\sum_{m_1 \dots m_6} \langle 00 \mathfrak{M} j_1 m_1 \dots j_6 m_6 \rangle \langle j_1 m_1 \dots j_6 m_6 q \mathfrak{N} p \rangle$ $= \sum_{J_{ik}} (2J_{12} + 1) (2J_{34} + 1) (2J_{56} + 1) \langle 0 \mathfrak{M} J_{12} J_{34} J_{56} \rangle$ $\times \langle J_{12} J_{34} J_{56} q \mathfrak{N} p \rangle$	
where $\langle 0 \mathfrak{M} J_{12} J_{34} J_{56} \rangle = \sum_{m_i M_{ik}} \langle 00 \mathfrak{M} j_1 m_1 \dots j_6 m_6 \rangle$ $\times \begin{pmatrix} j_1 & j_2 & J_{12} \\ m_1 & m_2 & M_{12} \end{pmatrix} \begin{pmatrix} j_3 & j_4 & J_{34} \\ m_3 & m_4 & M_{34} \end{pmatrix} \begin{pmatrix} j_5 & j_6 & J_{56} \\ m_5 & m_6 & M_{56} \end{pmatrix} \begin{pmatrix} J_{12} & J_{34} & J_{56} \\ M_{12} & M_{34} & M_{56} \end{pmatrix},$ and similarly for $\langle J_{12} J_{34} J_{56} q \mathfrak{N} p \rangle$.	

Table 11.10. $3jm$ symbols

Analytical Expression	Graphical Representation
Reversal of Cyclic Order of Momentum Coupling	
$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = (-1)^{j_1 + j_2 + j_3} \begin{pmatrix} j_3 & j_2 & j_1 \\ m_3 & m_2 & m_1 \end{pmatrix}$	
Hermitian Conjugation	
$\sum_{m'_1} \begin{pmatrix} j_1 \\ m'_1 m_1 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m_2 & m_3 \end{pmatrix} = (-1)^{j_1 - m_1} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix}$ $= \sum_{m'_2 m'_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m'_2 & m'_3 \end{pmatrix} \begin{pmatrix} j_2 \\ m_2 m'_2 \end{pmatrix} \begin{pmatrix} j_3 \\ m_3 m'_3 \end{pmatrix} = (-1)^{j_2 + m_2 + j_3 + m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & -m_2 & -m_3 \end{pmatrix}$ $\sum_{m'_1 m'_2 m'_3} \begin{pmatrix} j_1 \\ m'_1 m_1 \end{pmatrix} \begin{pmatrix} j_2 \\ m'_2 m_2 \end{pmatrix} \begin{pmatrix} j_3 \\ m'_3 m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$ $= (-1)^{j_1 - m_1 + j_2 - m_2 + j_3 - m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & -m_2 & -m_3 \end{pmatrix} = \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$	

Table 11.10. Cont'd



Analytical Expression	Graphical Representation
Orthonormality	
$\sum_{m_1 m_2} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j'_3 \\ m_1 & m_2 & m'_3 \end{pmatrix}$ $= \sum_{m_1 m_2} (-1)^{j_1 - m_1 + j_2 - m_2 + j_3 - m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & -m_2 & -m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j'_3 \\ m_1 & m_2 & m'_3 \end{pmatrix}$ $= \frac{\delta_{j_3 j'_3} \delta_{m_3 m'_3}}{(2j_3 + 1)} \{j_1 j_2 j_3\}$	
Completeness	
$\sum_{j_3 m_3} (2j_3 + 1) \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m_3 \end{pmatrix}$ $= \sum_{j_3 m_3} (2j_3 + 1) (-1)^{j_1 - m_1 + j_2 - m_2 + j_3 - m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & -m_2 & -m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m_3 \end{pmatrix}$ $= \delta_{m_1 m'_1} \delta_{m_2 m'_2}$	

Table 11.10a.

Relation of Clebsch-Gordan Coefficients to 3jm Symbols.


Analytical Expression	Graphical Representation
$C_{j_1 m_1 j_2 m_2}^{J M} = \langle j_1 m_1 j_2 m_2 j_1 j_2 J M \rangle = \langle j_1 j_2 J M j_1 m_1 j_2 m_2 \rangle$ $= (-1)^{2j_1} \sum_{j_3 m_3} (2j_3 + 1) \begin{pmatrix} J & 0 & j_3 \\ M & 0 & m_3 \end{pmatrix} \begin{pmatrix} j_3 & j_2 & j_1 \\ m_3 & m_2 & m_1 \end{pmatrix}$ $= (-1)^{2j_1} \sqrt{2J+1} (-1)^{J+M} \begin{pmatrix} J & j_2 & j_1 \\ -M & m_2 & m_1 \end{pmatrix}$ $= (-1)^{2j_2} \sum_{j_3 m_3} (2j_3 + 1) \begin{pmatrix} 0 & J & j_3 \\ 0 & M & m_3 \end{pmatrix} \begin{pmatrix} j_3 & j_2 & j_1 \\ m_3 & m_2 & m_1 \end{pmatrix}$ $= (-1)^{2j_2} \sqrt{2J+1} (-1)^{J-M} \begin{pmatrix} J & j_2 & j_1 \\ -M & m_2 & m_1 \end{pmatrix}$	

Table 11.12. Metric Tensor for $(2j+1)$ -Dimensional Space of Functions $|jm\rangle$.

Analytical Expression	Graphical Representation
Definition and Properties	
$ jm\ jm'\rangle = \langle jm\ jm' = \begin{pmatrix} j \\ m\ m' \end{pmatrix} = (-1)^{j+m} \delta_{m,-m'}$	$\begin{array}{c} jm \quad jm' \\ \longleftrightarrow \end{array} = \begin{array}{c} jm \quad jm' \\ \rightarrow \leftarrow \end{array}$
$\langle jm' jm = jm' jm\rangle = \begin{pmatrix} j \\ m' m \end{pmatrix} = (-1)^{j-m} \delta_{m',-m}$	$\begin{array}{c} jm' \quad jm \\ \rightarrow \leftarrow \end{array} = \begin{array}{c} jm' \quad jm \\ \longleftrightarrow \end{array}$
$ jm' jm\rangle = (-1)^{2j} jm\ jm'\rangle$ $\langle jm' jm = (-1)^{2j} \langle jm\ jm' $	$\begin{array}{c} jm' \quad jm \\ \longleftrightarrow \end{array} = (-1)^{2j} \begin{array}{c} jm \quad jm' \\ \longleftrightarrow \end{array} =$ $\begin{array}{c} jm' \quad jm \\ \rightarrow \leftarrow \end{array} = (-1)^{2j} \begin{array}{c} jm \quad jm' \\ \rightarrow \leftarrow \end{array}$
$\sum_{m'} jm\ jm'\rangle \langle jm'' jm' = \langle jm'' jm $	$\sum_{m'} \begin{array}{c} jm \quad jm' \\ \longleftrightarrow \end{array} \begin{array}{c} jm' \quad jm'' \\ \rightarrow \leftarrow \end{array} = \begin{array}{c} jm \quad jm'' \\ \rightarrow \leftarrow \end{array}$
Covariant $\rightleftharpoons$ Contravariant. Transformation of the Irreducible Tensor	
$\sum_{m'} jm\ jm'\rangle \langle jm' \mathfrak{M} \rangle = \langle \mathfrak{M} jm \rangle$	$\sum_{m'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm' \quad jm \\ \longleftrightarrow \end{array} = \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \\ \rightarrow \leftarrow \end{array}$
$\sum_{m'} \langle \mathfrak{M} jm' \rangle \langle jm' jm = \langle jm \mathfrak{M} \rangle$	$\sum_{m'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm' \quad jm \\ \rightarrow \leftarrow \end{array} = \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \\ \rightarrow \leftarrow \end{array}$
Scalar Product of Irreducible Tensors	
$\sum_m \langle \mathfrak{M} jm \rangle \langle jm \mathfrak{N} \rangle$ $= \sum_{mm'} \langle \mathfrak{M} jm \rangle \langle \mathfrak{N} jm' \rangle \langle jm' jm $ $= \sum_{mm'} jm\ jm'\rangle \langle jm' \mathfrak{M} \rangle \langle jm \mathfrak{N} \rangle.$	$\begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} j \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array} = \sum_m \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$ $= \sum_{mm'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm' \quad jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$ $= \sum_{mm'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm' \quad jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$
$\sum_m \langle jm \mathfrak{M} \rangle \langle \mathfrak{N} jm \rangle$ $= \sum_{mm'} \langle \mathfrak{M} jm' \rangle \langle \mathfrak{N} jm \rangle \langle jm' jm $ $= \sum_{mm'} jm\ jm'\rangle \langle jm \mathfrak{M} \rangle \langle jm' \mathfrak{N} \rangle.$	$\begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} j \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array} = \sum_m \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$ $= \sum_{mm'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm' \quad jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$ $= \sum_{mm'} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} jm \quad jm \\ \rightarrow \leftarrow \end{array} \begin{array}{c} jm' \quad jm' \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$
$\sum_m \langle \mathfrak{M} jm \rangle \langle jm \mathfrak{N} \rangle = (-1)^{2j} \sum_m \langle jm \mathfrak{M} \rangle \langle \mathfrak{N} jm \rangle$	$\begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} j \\ \rightarrow \leftarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array} = (-1)^{2j} \begin{array}{c} \mathfrak{M} \end{array} \begin{array}{c} j \\ \leftarrow \rightarrow \end{array} \begin{array}{c} \mathfrak{N} \end{array}$

Table 11.13. Spherical Harmonics.









Analytical Expression	Graphical Representation
Definition	
$C_{lm}(\Omega) \equiv \sqrt{\frac{4\pi}{2l+1}} Y_{lm}(\Omega) = (\Omega lm),$	
$C_{lm}^*(\Omega) = (-1)^m C_{l-m}(\Omega) = (lm \Omega).$	
Addition Theorem	
$\sum_m C_{lm}(\Omega_1) C_{lm}^*(\Omega_2) = P_l(\cos \omega_{12}),$	
where ω_{12} is the angle between Ω_1 and Ω_2 .	
Completeness	
$\sum_{lm} (2l+1) C_{lm}(\Omega_1) C_{lm}^*(\Omega_2) = 4\pi \delta(\Omega_1 - \Omega_2).$	
Orthonormality	
$\int C_{l_1 m_1}^*(\Omega) C_{l_2 m_2}(\Omega) \frac{d\Omega}{4\pi} = \frac{\delta_{l_1 l_2} \delta_{m_1 m_2}}{(2l_1 + 1)}.$	
Integral of a Product of Three Harmonics ³	
$\int C_{l_1 m_1}(\Omega) C_{l_2 m_2}(\Omega) C_{l_3 m_3}(\Omega) \frac{d\Omega}{4\pi} = \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix}.$	
Clebsch-Gordan Expansion	
$C_{l_1 m_1}(\Omega) C_{l_2 m_2}(\Omega) = \sum_{l_3 m_3} \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} C_{l_3 m_3}^*(\Omega)$	
Sum of Scalar Products of Three Harmonics	
$\sum_{m_1 m_2 m_3} (2l_1 + 1) \begin{pmatrix} l_1 & l_2 & l_3 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & l_2 & l_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \times C_{l_1 m_1}(\Omega_1) C_{l_2 m_2}(\Omega_2) C_{l_3 m_3}(\Omega_3) = P_{l_2}(\cos \omega_{21}) P_{l_3}(\cos \omega_{31}).$	

Table 11.14. Wigner D -Functions.

Analytical Expression	Graphical Representation
Adjoint Matrix	
$\left(D_{mm'}^j(R)\right)^* = \sum_{\mu\mu'} \begin{pmatrix} j \\ m \mu \end{pmatrix} D_{\mu\mu'}^j(R) \begin{pmatrix} j \\ \mu' m' \end{pmatrix}$ $= (-1)^{m-m'} D_{-m, -m'}^j(R).$	
Inverse Matrix	
$\left(D_{mm'}^j(R^{-1})\right) = \left(D_{m'm}^j(R)\right)^*.$	
Unitarity	
$\sum_m \left(D_{mm'}^j(R)\right)^* D_{m\bar{m}'}^j(R) = \sum_m D_{m'm}^j(R^{-1}) D_{m\bar{m}'}^j(R) = \delta_{m'\bar{m}'}$	
$\sum_{m'} D_{mm'}^j(R) \left(D_{\bar{m}m'}^j(R)\right)^* = \sum_{m'} D_{mm'}^j(R) D_{m'\bar{m}}^j(R^{-1}) = \delta_{m\bar{m}}$	
Orthonormality	
$\int D_{m_1 m_1'}^j(R) D_{m_2 m_2'}^{j*}(R) \frac{dR}{16\pi^2} = \frac{\delta_{j_1 j_2} \delta_{m_1 m_2} \delta_{m_1' m_2'}}{(2j_1 + 1)}$	
Completeness	
$\sum_{j=0,1/2,1}^{\infty} (2j+1) \sum_{mm'} D_{mm'}^j(R_1) D_{mm'}^{j*}(R_2) = 16\pi^2 \delta(R_1 - R_2)$	
Rotation Matrix in Ω -Representation	
$\delta_{ll'} D_{mm'}^l(R) = \frac{2l+1}{4\pi} \iint d\Omega d\Omega' C_{lm}^*(\Omega) \delta(\Omega - R^{-1}\Omega') C_{l'm'}(\Omega'),$ $\langle lm R l' m' \rangle = \langle lm \Omega \rangle \langle \Omega R^{-1} \Omega' \rangle \langle \Omega' l' m' \rangle.$	
$\delta(\Omega' - R\Omega) = \sum_{lm m'} \frac{2l+1}{4\pi} C_{lm'}(\Omega') D_{m'm}^l(R^{-1}) C_{lm}^*(\Omega),$ $\langle \Omega' R \Omega \rangle = \langle \Omega' l' m' \rangle \langle l' m' R^{-1} lm \rangle \langle lm \Omega \rangle.$	

Table 11.14
Cnt'ed.

Analytical Expression	Graphical Representation
Clebsch-Gordan Expansion and Associated Relations	
$D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R)$ $= \sum_{j_3 m_3 m'_3} (2j_3 + 1) \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} D_{m_3 m'_3}^{j_3*}(R) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$	
$\sum_{\substack{m_1 m_2 \\ m'_1 m'_2}} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$ $= \frac{\delta_{j_3 j'_3}}{(2j_3 + 1)} \{j_1 j_2 j_3\} D_{m_3 m'_3}^{j_3*}(R)$	
$\sum_{m_1 m_2 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R) D_{m_3 m'_3}^{j_3}(R)$ $= \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$	
$\sum_{\substack{m_1 m_2 \\ m'_1 m'_2 \\ m''_3}} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R) D_{m_3 m''_3}^{j_3}(R) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$ $= \frac{\delta_{j_3 j'_3} \delta_{m_3 m'_3}}{(2j_3 + 1)} \{j_1 j_2 j_3\}$	
$D_{m_3 m'_3}^{j_3*}(R) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$ $= \sum_{m_1 m_2} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R)$	
Integral of Three D-Functions	
$\int D_{m_1 m'_1}^{j_1}(R) D_{m_2 m'_2}^{j_2}(R) D_{m_3 m'_3}^{j_3}(R) \frac{dR}{16\pi^2}$ $= \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix}$	

Table 11.14
Cnt'ed.

Analytical Expression	Graphical Representation
Addition Theorems	
$\sum_{m'} D_{mm'}^j(R_2) D_{m'm''}^j(R_1) = D_{mm''}^j(R_2 R_1)$	
$\sum_{\substack{j_3 m_3 \\ m'_1 m'_2 m'_3}} (2j_3 + 1) \begin{pmatrix} j_1 & j_2 & j_3 \\ \tilde{m}_1 & \tilde{m}_2 & m_3 \end{pmatrix} \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix} \\ \times D_{m_1 m'_1}^{j_1}(R_1) D_{m_2 m'_2}^{j_2}(R_2) D_{m_3 m'_3}^{j_3}(R_3) \\ = D_{m_1 \tilde{m}_1}^{j_1}(R_1 R_3^{-1}) D_{m_2 \tilde{m}_2}^{j_2}(R_2 R_3^{-1})$	
$\sum_{m'_1 m'_2 m'_3} D_{m_1 m'_1}^{j_1}(R_1) D_{m_2 m'_2}^{j_2}(R_2) D_{m_3 m'_3}^{j_3}(R_3) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix} \\ = \sum_{m'_1 m'_2} D_{m_1 m'_1}^{j_1}(R_1 R_3^{-1}) D_{m_2 m'_2}^{j_2}(R_2 R_3^{-1}) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m_3 \end{pmatrix}$	
$\left\{ \begin{matrix} j_1 & j_2 & j_3 \\ J_{23} & J_{31} & J_{12} \end{matrix} \right\} \sum_{m_1 m_2 m_3} \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} \\ \times D_{m_1 m'_1}^{j_1}(R_1) D_{m_2 m'_2}^{j_2}(R_2) D_{m_3 m'_3}^{j_3}(R_3) \\ = \sum_{\substack{M_{12} M_{23} M_{31} \\ M'_{12} M'_{23} M'_{31}}} (-1)^{J_{12} - M_{12} + J_{23} - M_{23} + J_{31} - M_{31}} \\ \times \begin{pmatrix} J_{12} & j_1 & J_{31} \\ M_{12} & m'_1 & -M'_{31} \end{pmatrix} \begin{pmatrix} J_{23} & j_2 & J_{12} \\ M_{23} & m'_2 & -M'_{12} \end{pmatrix} \begin{pmatrix} J_{31} & j_3 & J_{23} \\ M_{31} & m'_3 & -M'_{23} \end{pmatrix} \\ \times D_{M_{31} M'_{31}}^{J_{31}}(R_3^{-1} R_1) D_{M_{12} M'_{12}}^{J_{12}}(R_1^{-1} R_2) D_{M_{23} M'_{23}}^{J_{23}}(R_2^{-1} R_3)$	
$\sum_{j_1 j_2 j_3} (2j_1 + 1)(2j_2 + 1)(2j_3 + 1) \left\{ \begin{matrix} j_1 & j_2 & j_3 \\ J_{23} & J_{31} & J_{12} \end{matrix} \right\} \\ \times \sum_{\substack{m_1 m_2 m_3 \\ m'_1 m'_2 m'_3}} (-1)^{J_{12} - M_{12} + J_{23} - M_{23} + J_{31} - M_{31}} \\ \times \begin{pmatrix} J_{31} & j_1 & J_{12} \\ M'_{31} & m_1 & -M_{12} \end{pmatrix} \begin{pmatrix} J_{12} & j_2 & J_{23} \\ M'_{12} & m_2 & -M_{23} \end{pmatrix} \begin{pmatrix} J_{23} & j_3 & J_{31} \\ M'_{23} & m_3 & -M_{31} \end{pmatrix} \\ \times D_{m_1 m'_1}^{j_1}(R_1) D_{m_2 m'_2}^{j_2}(R_2) D_{m_3 m'_3}^{j_3}(R_3) \begin{pmatrix} j_1 & j_2 & j_3 \\ m'_1 & m'_2 & m'_3 \end{pmatrix} \\ = D_{M_{12} M'_{12}}^{J_{12}}(R_1 R_2^{-1}) D_{M_{23} M'_{23}}^{J_{23}}(R_2 R_3^{-1}) D_{M_{31} M'_{31}}^{J_{31}}(R_3 R_1^{-1})$	

Table 11.15. Matrix Elements of Irreducible Tensor Operators.


Analytical Expression	Graphical Representation
Matrix element of an irreducible tensor operator of rank k in the JM-representation (Wigner-Eckart theorem)	
$\langle \gamma JM \mathfrak{M}_{kq} \gamma' J' M' \rangle = (-1)^{J-M} \begin{pmatrix} J & k & J' \\ -M & q & M' \end{pmatrix} (\gamma J \ \mathfrak{M}_k \ \gamma' J').$ where $(\gamma J \ \mathfrak{M}_k \ \gamma' J')$ is the reduced matrix element with the renormalised states	
Reduced Matrix elements of irreducible tensor product of irreducible tensor operators.	
$(\gamma J \ \{ \mathfrak{M}_{k_1} \otimes \mathfrak{N}_{k_2} \}_k \ \gamma' J')$ $= (-1)^{J+k+J'} \sqrt{2k+1} \sum_{\gamma'' J''} \begin{Bmatrix} J & k & J' \\ k_2 & J'' & k_1 \end{Bmatrix}$ $\times (\gamma J \ \mathfrak{M}_{k_1} \ \gamma'' J'') (\gamma'' J'' \ \mathfrak{N}_{k_2} \ \gamma' J').$	
Two coupled subsystems in the $j_1 j_2; JM$ -representation. Reduced matrix element of the irreducible tensor product of two irreducible tensor operators, one of which depends only on variables of the first subsystem and the other depends only on variables of the second subsystem.	
$(\gamma j_1 j_2 J \ \{ \mathfrak{M}_{k_1}(1) \otimes \mathfrak{N}_{k_2}(2) \}_k \ \gamma' j'_1 j'_2 J')$ $= \sqrt{(2J+1)(2k+1)(2J'+1)} \begin{Bmatrix} j_1 & j'_1 & k_1 \\ j_2 & j'_2 & k_2 \\ J & J' & k \end{Bmatrix}$ $\times \sum_{\gamma''} (\gamma j_1 \ \mathfrak{M}_{k_1} \ \gamma'' j'_1) (\gamma'' j_2 \ \mathfrak{N}_{k_2} \ \gamma' j'_2)$	

Table 11.15
Cnt'ed.

Analytical Expression	Graphical Representation
Matrix element of the scalar product of two irreducible tensor operators which depend on variables of the first and the second subsystems, respectively.	
$\begin{aligned} & \langle \gamma j_1 j_2 J M (\mathfrak{M}_k(1) \cdot \mathfrak{N}_k(2)) \gamma' j'_1 j'_2 J' M' \rangle \\ &= (-1)^{j'_1 + j_2 + J} \delta_{JJ'} \delta_{MM'} \begin{Bmatrix} j_1 & j_2 & J \\ j'_2 & j'_1 & k \end{Bmatrix} \\ & \times \sum_{\gamma''} (\gamma j_1 \ \mathfrak{M}_k \ \gamma'' j'_1) (\gamma'' j_2 \ \mathfrak{N}_k \ \gamma' j'_2) \end{aligned}$	
Reduced matrix element of an irreducible tensor operator which depends only on variables of the first subsystem.	
$\begin{aligned} & (\gamma j_1 j_2 J \ \mathfrak{M}_k(1) \ \gamma' j'_1 j'_2 J') \\ &= (-1)^{j_1 + j_2 + J' + k} \delta_{j_2 j'_2} \sqrt{(2J+1)(2J'+1)} \\ & \times \begin{Bmatrix} J & k & J' \\ j'_1 & j_2 & j_1 \end{Bmatrix} (\gamma j_1 \ \mathfrak{M}_k \ \gamma' j'_1). \end{aligned}$	
$\begin{aligned} & (\gamma j_1 j_2 J \ \mathfrak{N}_k(2) \ \gamma' j'_1 j'_2 J') \\ &= (-1)^{j'_1 + j'_2 + J + k} \delta_{j_1 j'_1} \sqrt{(2J+1)(2J'+1)} \\ & \times \begin{Bmatrix} J & k & J' \\ j'_2 & j_1 & j_2 \end{Bmatrix} (\gamma j_2 \ \mathfrak{N}_k \ \gamma' j'_2). \end{aligned}$	

- (c) A diagram with two external lines corresponding to $\langle j_1 m_1 |$ and $| j_2 m_2 \rangle$ is non-zero only if $j_1 = j_2$ and $m_1 = m_2$

$$\langle j_1 m_1 | \mathfrak{M} | j_2 m_2 \rangle = \frac{\delta_{j_1 j_2} \delta_{m_1 m_2}}{2j_1 + 1} \sum_m \langle j_1 m | \mathfrak{M} | j_1 m \rangle \quad (11.4.3)$$




This follows from the fact that any dependence on two directions is, actually, the dependence on the angle between these directions, i.e. it may be represented by the scalar product of two functions $|jm\rangle$ and $\langle jm|$.

- (d) A diagram with three external j -lines which represent $|j_1 m_1\rangle$, $|j_2 m_2\rangle$ and $|j_3 m_3\rangle$ is proportional to the $3jm$ symbol $\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}$

$$\langle 00 | \mathfrak{M} | j_1 m_1 j_2 m_2 j_3 m_3 \rangle = \langle 00 | \mathfrak{M} | j_1 j_2 j_3 \rangle \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}, \quad (11.4.4)$$

where $\langle 00 | \mathfrak{M} | j_1 j_2 j_3 \rangle$ is the reduced matrix element defined by

$$\langle 00 | \mathfrak{M} | j_1 j_2 j_3 \rangle = \sum_{m_1 m_2 m_3} \langle 00 | \mathfrak{M} | j_1 m_1 j_2 m_2 j_3 m_3 \rangle \begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix}, \quad (11.4.5)$$




- (e) A diagram with four external j -lines corresponding $j_1 m_1, j_2 m_2, j_3 m_3$ and $j_4 m_4$ may be presented as a sum (over j) of the product of two $3jm$ symbols

$$\langle j_1 m_1 j_2 m_2 | \mathfrak{M} | j_3 m_3 j_4 m_4 \rangle = \sum_{jm} (2j+1) \begin{pmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{pmatrix} \langle j_1 j_2 jm | \mathfrak{M} | j_3 j_4 jm \rangle \begin{pmatrix} j_3 & j_4 & j \\ m_3 & m_4 & m \end{pmatrix}, \quad (11.4.6)$$

where

$$\langle j_1 j_2 jm | \mathfrak{M} | j_3 j_4 jm \rangle \equiv \sum_{m_1 m_2 m_3 m_4} \begin{pmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{pmatrix} \langle j_1 m_1 j_2 m_2 | \mathfrak{M} | j_3 m_3 j_4 m_4 \rangle \begin{pmatrix} j_3 & j_4 & j' \\ m_3 & m_4 & m' \end{pmatrix} \quad (11.4.7)$$



11.4.3 Scheme for the Application of the Graphical Technique

The aim of calculations inherent in the angular momentum theory is to transform an original formula in such a way as to express all the dependence on the orientation of the coordinate system in the form of a simple standard function $\left(C_{j_1 m_1 j_2 m_2}^{j m}, Y_{lm}(\Omega), D_{MM'}^J(R), \text{ or some related combination} \right)$. The remaining factor (or the expansion coefficient) is invariant and can be represented as a product (or sum of products) of the standard $3j, 6j, 9j$ symbols.

To carry out these calculations by the graphical method one has to perform the following operations:

- To represent the original analytic expression in diagrammatic form using the graphical representation of functions and operations in accordance with the rules formulated in Sections 11.1 and 11.2.
- To eliminate all fictitious $j = 0$ lines and $R = 1$ lines according to the rules of Sections 11.3.9 and 11.3.10.
- To link step-by-step, all the subdiagrams which have a common thick j -line, using the rules for graphical multiplication of subdiagrams (Sec. 11.3.5).

Table 11.16. Generalized Wigner-Eckart Theorem.

Analytical Expression	Graphical Representation
$\langle 00 \mathfrak{M} lm\rangle = \langle 00 \mathfrak{M} 00\rangle \delta_{l0} \delta_{m0}$	
$\langle jm \mathfrak{M} j'm'\rangle = \frac{\delta_{jj'} \delta_{mm'}}{(2j+1)} \langle j \mathfrak{M} j\rangle$	
where $\langle j \mathfrak{M} j\rangle \equiv \sum_m \langle jm \mathfrak{M} jm\rangle$	
$\langle \Omega \mathfrak{M} jm\rangle = \langle \Omega jm\rangle \frac{\langle j \mathfrak{M} j\rangle}{(2j+1)}$	
$\langle \Omega_1 \mathfrak{M} \Omega_2\rangle = \frac{1}{4\pi} \sum_l P_l(\cos \omega_{12}) \langle l \mathfrak{M} l\rangle$	
where ω_{12} is the angle $\widehat{\Omega_1 \Omega_2}$ and	
$P_l(\cos \omega_{12}) = \sum_m \langle \Omega_1 lm\rangle \langle lm \Omega_2\rangle$	
$\langle j_1 m_1 \mathfrak{M} j_2 m_2 j_3 m_3\rangle$ $= (-1)^{j_1 - m_1} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix} \langle j_1 \mathfrak{M}_{j_2} j_3\rangle$	
where $\langle j_1 \mathfrak{M}_{j_2} j_3\rangle \equiv \sum_{m_1 m_2 m_3} (-1)^{j_1 - m_1}$ $\times \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix} \langle j_1 m_1 \mathfrak{M} j_2 m_2 j_3 m_3\rangle$	
$\langle \Omega \mathfrak{M} j_2 m_2 j_3 m_3\rangle = \sum_{j_1 m_1} \langle \Omega j_1 m_1\rangle$ $\times (-1)^{j_1 - m_1} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix} \langle j_1 \mathfrak{M}_{j_2} j_3\rangle$	
$\langle \Omega_1 \mathfrak{M} j_2 m_2 \Omega_3\rangle = \sum_{j_1 m_1 j_3 m_3} \langle \Omega_1 j_1 m_1\rangle (-1)^{j_1 - m_1}$ $\times \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix} \langle j_3 m_3 \Omega_3\rangle \langle j_1 \mathfrak{M}_{j_2} j_3\rangle$	

Table 11.16. (Cont'd)

Analytical Expression	Graphical Representation
$\begin{aligned} & \langle j_1 m_1 j_2 m_2 \mathfrak{M} j_3 m_3 j_4 m_4 \rangle \\ &= \sum_{JM} (2J+1) \begin{pmatrix} j_1 & j_2 & J \\ m_1 & m_2 & M \end{pmatrix} \langle j_1 j_2; JM \mathfrak{M} j_3 j_4; JM \rangle \\ & \quad \times \begin{pmatrix} j_3 & j_4 & J \\ m_3 & m_4 & M \end{pmatrix} \end{aligned}$	
$\begin{aligned} & \langle \Omega_1 \mathfrak{M} \Omega_2 \Omega_3 \rangle = \sum_{j_1 j_2 j_3} \langle j_1 \mathfrak{M}_{j_2} j_3 \rangle \\ & \times \sum_{m_1 m_2 m_3} \langle \Omega_1 j_1 m_1 \rangle (-1)^{j_1 - m_1} \begin{pmatrix} j_1 & j_2 & j_3 \\ -m_1 & m_2 & m_3 \end{pmatrix} \\ & \quad \times \langle j_2 m_2 \Omega_2 \rangle \langle j_3 m_3 \Omega_3 \rangle \end{aligned}$	

hline

- (d) To replace each internal Ω -line by the corresponding thick j -line, using the rule of graphical integration (Sec. 11.3.8).
- (e) To remove thick j -lines with identical nodes $(j_1 j_2 j_3)$ and reconnect the remaining thin j_1 - and j_2 -lines in accordance with the rule of graphical summation (Sec. 11.3.7).
- (f) To cut the resultant diagram (according to the rules of Sec. 11.3.6) into subdiagrams which are topologically similar to diagrams of the standard functions. In particular, one should separate all the external lines in the form of the simplest possible subdiagram.
- (g) To determine the phase factor of a diagram by reducing each subdiagram to some standard diagram (Tables 11.1 and 11.2) by means of reversal (when necessary) of node signs and line directions according to the rules from Section 11.3.2 and 11.3.3.
- (h) To represent the final diagram in analytic form.

One can use a simplified version of the graphical method without taking the signs of the nodes and directions of the lines into account, if one is interested only in the absolute value of an expression but not in the phase. Such an approach is taken in the next chapter.

Chapter 12

Sums involving vector addition and recoupling coefficients

Many quantum-mechanical problems are concerned with calculations of sums containing products of the Clebsch-Gordan coefficients and $3nj$ symbols whose summation indices represent quantum numbers of angular momenta and momentum projections. Summation over different indices may be performed independently and successively. It is more convenient to start with the summation of $3jm$ symbol products over projections. This summation may be carried out with the aid of the equations presented in Sec. 12.1. Summation over projections yields invariant $3nj$ symbols. The next step is to sum the products of $3nj$ symbols over momenta; this may be done with the aid of the equations of Sec. 12.2.

12.1 Summation of products of $3jm$ symbols

This section contains a collection of sums involving the $3jm$ symbols. Each sum is illustrated by a diagram which shows the coupling scheme of angular momenta involved in the sum. The equations are grouped in accordance with the number of $3jm$ symbols in the products. Within each group the equations are arranged according to the number of fixed indices, i.e., the number of external lines of the corresponding diagram. The summations are of two kinds:

- (a) Summation index j represents an angular momentum and runs over all possible values consistent with the triangle condition. Summations of this kind are shown by thick internal j -lines in the diagrams.
- (b) Summation index m_j is a projection of j and runs over all allowed values, $j, j-1, j-2, \dots, -j$. Summations of such kind are shown by thin internal j -lines in the diagrams.

The sums are written in such a form that each summation index m_j is included in two $3jm$ symbols, although with the opposite signs. In addition, the sums contain the phase factors $(-1)^{j-m_j}$ which ensure that the results of summation are invariant under coordinate rotations (Sec. 11.2). Any sum over projections, which is invariant with respect to rotations, may be converted into such a form (Sec. 8.4).

Hereafter, we use the notation

$$\Pi_{ab\dots f} = \sqrt{(2a+1)(2b+1)\dots(2f+1)}. \quad (12.1.1)$$

12.1.1 Sum Involving One $3jm$ Symbol

$$\sum_{\alpha} (-1)^{a-\alpha} \begin{pmatrix} a & a & c \\ \alpha & -\alpha & \gamma \end{pmatrix} = \Pi_a \delta_{c0} \delta_{\gamma 0}. \quad (12.1.2)$$

12.1.2 Sums Involving Products of Two $3jm$ Symbols

$$\sum_{\psi\kappa} (-1)^{p-\psi+q-\kappa} \begin{pmatrix} a & p & q \\ -\alpha & \psi & \kappa \end{pmatrix} \begin{pmatrix} p & q & a' \\ -\psi & -\kappa & \alpha' \end{pmatrix} = \frac{(-1)^{a+\alpha}}{\Pi_a^2} \{apq\} \delta_{aa'} \delta_{\alpha\alpha'}, \quad (12.1.3)$$

$$\begin{array}{c} p \\ \text{---} \bigcirc \text{---} \\ q \end{array} \begin{array}{c} a\alpha \\ \text{---} \end{array} \begin{array}{c} a'\alpha' \\ \text{---} \end{array} = \begin{array}{c} a\alpha \\ \text{---} \end{array} \begin{array}{c} a'\alpha' \\ \text{---} \end{array} \begin{array}{c} p \\ \text{---} \bigcirc \text{---} \\ q \end{array}$$

$$\sum_{q\kappa} (-1)^{q-\kappa} \Pi_q^2 \begin{pmatrix} a & b & q \\ -\alpha & -\beta & \kappa \end{pmatrix} \begin{pmatrix} q & a & b \\ -\kappa & \alpha' & \beta' \end{pmatrix} = (-1)^{a+\alpha+b+\beta} \delta_{\alpha\alpha'} \delta_{\beta\beta'}, \quad (12.1.4)$$

$$\sum_{\kappa} (-1)^{q-\kappa} \begin{pmatrix} a & b & q \\ \alpha & \beta & -\kappa \end{pmatrix} \begin{pmatrix} q & d & c \\ \kappa & \delta & \gamma \end{pmatrix} = (-1)^{2a} \sum_{x\xi} (-1)^{x-\xi} \Pi_x^2 \begin{pmatrix} a & c & x \\ \alpha & \gamma & -\xi \end{pmatrix} \begin{pmatrix} x & d & b \\ \xi & \delta & \beta \end{pmatrix} \begin{Bmatrix} b & d & x \\ c & a & q \end{Bmatrix}, \quad (12.1.5)$$


12.1.3 Sums Involving Products of Three $3jm$ Symbols

$$\sum_{\kappa\psi\rho} (-1)^{p-\psi+q-\kappa+r-\rho} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & b & r \\ \kappa & \beta & -\rho \end{pmatrix} \begin{pmatrix} r & c & p \\ \rho & \gamma & -\psi \end{pmatrix} = \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} \begin{Bmatrix} a & b & c \\ r & p & q \end{Bmatrix}, \quad (12.1.6)$$


$$\sum_{\psi\kappa} \begin{pmatrix} a & b & p \\ \alpha & \beta & \psi \end{pmatrix} \begin{pmatrix} p & c & q \\ -\psi & -\gamma & -\kappa \end{pmatrix} \begin{pmatrix} q & d & e \\ \kappa & \delta & \varepsilon \end{pmatrix} = \sum_{\substack{xy \\ \varepsilon\eta}} \Pi_{xy}^2 \begin{pmatrix} a & d & x \\ \alpha & \delta & \xi \end{pmatrix} \begin{pmatrix} x & c & y \\ -\xi & -\gamma & -\eta \end{pmatrix} \begin{pmatrix} y & b & e \\ \eta & \beta & \varepsilon \end{pmatrix} \begin{Bmatrix} a & b & p \\ d & e & q \\ x & y & c \end{Bmatrix}, \quad (12.1.7)$$

12.1.4 Sums Involving Products of Four $3jm$ Symbols


$$\begin{aligned}
\sum_{\kappa\psi\rho\sigma\tau} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-\tau} & \begin{pmatrix} p & a & q \\ \psi & -\alpha & \kappa \end{pmatrix} \begin{pmatrix} q & r & t \\ -\kappa & \rho & \tau \end{pmatrix} \begin{pmatrix} r & a' & s \\ -\rho & \alpha' & \sigma \end{pmatrix} \begin{pmatrix} s & p & t \\ -\sigma & -\psi & -\tau \end{pmatrix} \\
&= \frac{(-1)^{a-\alpha}}{\Pi_a^2} \left\{ \begin{matrix} q & p & a \\ s & r & t \end{matrix} \right\} \delta_{aa'} \delta_{\alpha\alpha'}, \tag{12.1.8}
\end{aligned}$$




$$\begin{aligned}
\sum_{\psi\kappa\rho\sigma} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma} & \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & b & r \\ \kappa & \beta & -\rho \end{pmatrix} \begin{pmatrix} r & s & p \\ \rho & \sigma & -\psi \end{pmatrix} \begin{pmatrix} s & c & d \\ -\sigma & \gamma & \delta \end{pmatrix} \\
&= \left\{ \begin{matrix} a & b & s \\ r & p & q \end{matrix} \right\} \sum_{\sigma} (-1)^{s-\sigma} \begin{pmatrix} a & s & b \\ \alpha & \sigma & \beta \end{pmatrix} \begin{pmatrix} d & s & c \\ \delta & -\sigma & \gamma \end{pmatrix}, \tag{12.1.9}
\end{aligned}$$



$$\begin{aligned}
\sum_{\psi\kappa\rho\sigma} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma} & \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & b & r \\ \kappa & \beta & -\rho \end{pmatrix} \begin{pmatrix} r & c & s \\ \rho & \gamma & -\sigma \end{pmatrix} \begin{pmatrix} s & d & p \\ \sigma & \delta & -\psi \end{pmatrix} \\
&= (-1)^{s-a-d-q} \sum_{x\xi} (-1)^{x-\xi} \Pi_x^2 \begin{pmatrix} a & x & d \\ \alpha & -\xi & \delta \end{pmatrix} \begin{pmatrix} b & x & c \\ \beta & \xi & \gamma \end{pmatrix} \left\{ \begin{matrix} a & x & d \\ s & p & q \end{matrix} \right\} \left\{ \begin{matrix} b & x & c \\ s & r & q \end{matrix} \right\} \tag{12.1.10}
\end{aligned}$$




$$= (-1)^{2r-s-d-a-q} \sum_{x\xi} (-1)^{x-\xi} \Pi_x^2 \begin{pmatrix} a & x & c \\ \alpha & -\xi & \gamma \end{pmatrix} \begin{pmatrix} b & x & d \\ \beta & \xi & \delta \end{pmatrix} \left\{ \begin{matrix} a & p & q \\ x & d & b \\ c & s & r \end{matrix} \right\}, \tag{12.1.11}$$




12.1.5 Sums Involving Products of Five 3jm Symbols


$$\begin{aligned}
& \sum_{\substack{\psi \kappa \rho \sigma \\ \tau \nu \mu}} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-\tau+u-\nu+v-\mu} \begin{pmatrix} r & t & q \\ \rho & \tau & -\kappa \end{pmatrix} \begin{pmatrix} q & p & u \\ \kappa & \psi & -\nu \end{pmatrix} \begin{pmatrix} u & a & v \\ \nu & \alpha & -\mu \end{pmatrix} \begin{pmatrix} v & s & r \\ \mu & \sigma & -\rho \end{pmatrix} \begin{pmatrix} s & p & t \\ -\sigma & -\psi & -\tau \end{pmatrix} \\
& = (-1)^{2u} \frac{1}{\Pi_u} \delta_{uv} \delta_{a0} \delta_{\alpha 0} \left\{ \begin{matrix} q & p & u \\ s & r & t \end{matrix} \right\}, \tag{12.1.12}
\end{aligned}$$




$$\begin{aligned}
& \sum_{\substack{\psi \kappa \rho \sigma \\ \tau \nu}} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-\tau+u-\nu} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\chi \end{pmatrix} \begin{pmatrix} q & r & t \\ \kappa & \rho & -\tau \end{pmatrix} \begin{pmatrix} t & s & p \\ \tau & -\sigma & -\psi \end{pmatrix} \begin{pmatrix} s & c & u \\ \sigma & \gamma & -\nu \end{pmatrix} \begin{pmatrix} u & b & r \\ \nu & \beta & -\rho \end{pmatrix} \\
& = \begin{pmatrix} a & b & c \\ \alpha & \beta & \gamma \end{pmatrix} \left\{ \begin{matrix} a & b & c \\ u & s & r \end{matrix} \right\} \left\{ \begin{matrix} a & p & q \\ t & r & s \end{matrix} \right\}, \tag{12.1.13}
\end{aligned}$$




$$\begin{aligned}
& \sum_{\psi \kappa \rho \tau} (-1)^{p-\psi+q-\kappa+r-\rho+\rho-\sigma+t-r+u-\nu} \begin{pmatrix} p & a & q \\ \psi & \alpha & \kappa \end{pmatrix} \begin{pmatrix} q & t & r \\ -\kappa & -\tau & -\rho \end{pmatrix} \begin{pmatrix} r & b & s \\ \rho & \beta & \sigma \end{pmatrix} \begin{pmatrix} s & p & u \\ -\sigma & -\psi & -\nu \end{pmatrix} \begin{pmatrix} u & c & t \\ \nu & \gamma & r \end{pmatrix} \\
& = (-1)^{r+b+\theta} \begin{pmatrix} a & b & c \\ -\alpha & -\beta & -\gamma \end{pmatrix} \left\{ \begin{matrix} a & b & c \\ p & s & u \\ q & r & t \end{matrix} \right\}, \tag{12.1.14}
\end{aligned}$$




$$\begin{aligned}
& \sum_{\psi \kappa \rho} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-\tau} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & b & r \\ \kappa & \beta & -\rho \end{pmatrix} \begin{pmatrix} r & c & s \\ \rho & \gamma & -\sigma \end{pmatrix} \begin{pmatrix} s & d & t \\ \sigma & \delta & -\tau \end{pmatrix} \begin{pmatrix} t & e & p \\ \tau & \varepsilon & -\psi \end{pmatrix} \\
& = (-1)^{t-p-b-a-d-c} \sum_{x\xi y\eta} (-1)^{x-\xi+y-\eta} \Pi_{xy}^2 \begin{pmatrix} a & b & x \\ \alpha & \beta & \xi \end{pmatrix} \begin{pmatrix} x & e & y \\ -\xi & \varepsilon & -\eta \end{pmatrix} \begin{pmatrix} y & c & d \\ \eta & \gamma & \delta \end{pmatrix} \left\{ \begin{matrix} a & b & x \\ r & p & q \end{matrix} \right\} \left\{ \begin{matrix} x & e & y \\ t & r & p \end{matrix} \right\} \left\{ \begin{matrix} y & c & d \\ s & t & r \end{matrix} \right\} \tag{12.1.15}
\end{aligned}$$



$$= (-1)^{p-t-b-a-d-c} \sum_{x\xi y\eta} (-1)^{x-\xi+y-\eta} \Pi_{xy}^2 \begin{pmatrix} a & b & x \\ \alpha & \beta & \xi \end{pmatrix} \begin{pmatrix} x & d & y \\ -\xi & \delta & -\eta \end{pmatrix} \begin{pmatrix} y & c & e \\ \eta & \gamma & \varepsilon \end{pmatrix} \left\{ \begin{matrix} a & b & x \\ r & p & q \end{matrix} \right\} \left\{ \begin{matrix} c & e & y \\ r & p & x \\ s & t & d \end{matrix} \right\}, \quad (12.1.16)$$



$$= (-1)^{r-t+p+a+d} \sum_{x\xi y\eta z} (-1)^{x-\xi+y-\eta+x} \Pi_{xyz}^2 \begin{pmatrix} a & c & x \\ \alpha & \gamma & \xi \end{pmatrix} \begin{pmatrix} x & e & y \\ -\xi & \varepsilon & -\eta \end{pmatrix} \begin{pmatrix} y & b & d \\ \eta & \beta & \delta \end{pmatrix} \left\{ \begin{matrix} c & q & z \\ b & s & r \end{matrix} \right\} \left\{ \begin{matrix} a & c & x \\ z & p & q \end{matrix} \right\} \left\{ \begin{matrix} x & e & y \\ t & z & p \end{matrix} \right\} \left\{ \begin{matrix} y & b & d \\ s & t & z \end{matrix} \right\}, \quad (12.1.17)$$



12.1.6 Sums Involving Products of Six $3jm$ Symbols

$$\sum_{\substack{\psi \kappa \rho \sigma \\ \tau \nu \mu \lambda}} (-1)^{p-\psi+q-\kappa+r-\rho+\vartheta-\sigma+t-\tau+u-\nu+v-\mu+t'-\lambda} \begin{pmatrix} p & a & q \\ \psi & -\alpha & \kappa \end{pmatrix} \begin{pmatrix} q & r & t' \\ -\kappa & \rho & \lambda \end{pmatrix} \begin{pmatrix} t' & u & v \\ -\lambda & \nu & \mu \end{pmatrix} \begin{pmatrix} u & v & t \\ -\nu & -\mu & \tau \end{pmatrix} \\ \times \begin{pmatrix} r & a' & s \\ -\rho & \alpha' & \sigma \end{pmatrix} \begin{pmatrix} s & p & t \\ -\sigma & -\psi & -r \end{pmatrix} = \frac{(-1)^{2t+a-\alpha}}{\Pi_{at}^2} \left\{ \begin{matrix} a & p & q \\ t & r & s \end{matrix} \right\} t u v \delta + t' t \delta_{aa'} \delta_{\alpha\alpha'} \quad (12.1.18)$$

$$\sum_{\substack{\psi \kappa \rho \sigma \\ \nu \mu \tau \lambda}} (-1)^{p-\psi+q-\kappa+q'-\rho+p'-\sigma+u-\nu+v-\mu+t-\tau+\rho-\lambda} \begin{pmatrix} u & v & q \\ \nu & \mu & \kappa \end{pmatrix} \begin{pmatrix} u & v & q' \\ -\nu & -\mu & -\rho \end{pmatrix} \begin{pmatrix} t & s & p \\ \tau & \lambda & \psi \end{pmatrix} \begin{pmatrix} t & s & p' \\ -\tau & -\lambda & -\sigma \end{pmatrix} \\ \times \begin{pmatrix} q & p & a \\ \kappa & \psi & \alpha \end{pmatrix} \begin{pmatrix} q' & p' & a' \\ -\rho & -\sigma & -\alpha' \end{pmatrix} = \frac{(-1)^{\alpha-a}}{\Pi_{apq}^2} \delta_{qq'} \delta_{pp'} \delta_{aa'} \delta_{\alpha\alpha'} \{uvq\} \{pts\} \{apq\}, \quad (12.1.19)$$

$$\sum_{\psi \kappa \rho \sigma} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-\tau+v-\mu+u-\nu+g-\lambda} \begin{pmatrix} a & q & p \\ -\alpha & \kappa & \psi \end{pmatrix} \begin{pmatrix} p & t & g \\ -\psi & r & -\lambda \end{pmatrix} \begin{pmatrix} g & u & v \\ \lambda & -\nu & \mu \end{pmatrix} \begin{pmatrix} v & r & q \\ -\mu & \rho & -\kappa \end{pmatrix} \\ \times \begin{pmatrix} r & u & s \\ -\rho & \nu & -\sigma \end{pmatrix} \begin{pmatrix} s & t & a' \\ \sigma & -\tau & \alpha' \end{pmatrix} = \frac{(-1)^{s-p-\alpha-g}}{\Pi_a^2} \left\{ \begin{matrix} q & g & s \\ u & r & v \end{matrix} \right\} \left\{ \begin{matrix} a & p & q \\ g & s & t \end{matrix} \right\} \delta_{aa'} \delta_{\alpha\alpha'} \quad (12.1.20)$$

$$\sum_{\substack{\psi \kappa \rho \sigma \\ \nu \tau \mu \lambda}} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+u-\nu+t-r+v-\mu+g-\lambda} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & r & v \\ \kappa & \rho & -\mu \end{pmatrix} \begin{pmatrix} v & s & p \\ \mu & -\sigma & -\psi \end{pmatrix} \begin{pmatrix} u & r & g \\ \nu & -\rho & -\lambda \end{pmatrix} \\ \times \begin{pmatrix} t & a' & u \\ -\tau & -\alpha' & -\nu \end{pmatrix} \begin{pmatrix} s & t & a' \\ \sigma & -\tau & -\alpha' \end{pmatrix} = \frac{(-1)^{2g+a-\alpha}}{\Pi_a^2} \left\{ \begin{matrix} a & r & s \\ g & t & u \end{matrix} \right\} \left\{ \begin{matrix} a & r & s \\ v & p & q \end{matrix} \right\} \delta_{aa'} \delta_{\alpha\alpha'}, \quad (12.1.21)$$

$$\sum_{\substack{\psi \kappa \rho \sigma \\ \nu \tau \mu \lambda}} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+u-\nu+t-r+v-\mu+g-\lambda} \begin{pmatrix} a & p & q \\ \alpha & \psi & \kappa \end{pmatrix} \begin{pmatrix} q & g & s \\ -\kappa & -\lambda & -\sigma \end{pmatrix} \begin{pmatrix} s & t & v \\ \sigma & \tau & \mu \end{pmatrix} \begin{pmatrix} v & p & r \\ -\mu & -\psi & -\rho \end{pmatrix} \\ \times \begin{pmatrix} r & g & u \\ \rho & \lambda & \nu \end{pmatrix} \begin{pmatrix} u & t & a' \\ -\nu & -\tau & -\alpha' \end{pmatrix} = \frac{(-1)^{a-\alpha}}{\Pi_a^2} \left\{ \begin{matrix} a & p & q \\ u & r & g \\ t & v & s \end{matrix} \right\} \delta_{aa'} \delta_{\alpha\alpha'}, \quad (12.1.22)$$


$$\begin{aligned}
& \begin{array}{c} r \\ p \quad u \\ a\alpha \quad a'\alpha' \\ q \quad t \\ g \quad v \\ s \end{array} = \frac{a\alpha \quad a'\alpha'}{\quad} \begin{array}{c} r \\ p \quad u \\ a \\ q \quad t \\ g \quad v \\ s \end{array} \\
& \sum_{\substack{\psi\kappa\nu\rho \\ \mu\sigma\lambda}} (-1)^{p-\psi+q-\kappa+u-\nu+r-\rho+v-\mu+s-\sigma+g-\lambda} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & u & g \\ -\kappa & \nu & -\lambda \end{pmatrix} \begin{pmatrix} g & v & s \\ \lambda & \mu & -\sigma \end{pmatrix} \begin{pmatrix} s & d & p \\ \sigma & \delta & -\psi \end{pmatrix} \\
& \quad \times \begin{pmatrix} u & b & r \\ -\nu & \beta & \rho \end{pmatrix} \begin{pmatrix} r & c & v \\ -\rho & \gamma & -\mu \end{pmatrix} \\
& = (-1)^{s+b+u+v-q-a-d-c} \sum_{x\xi} \Pi_x^2 \begin{pmatrix} a & x & d \\ \alpha & -\xi & \beta \end{pmatrix} \begin{pmatrix} c & x & b \\ \gamma & \xi & \beta \end{pmatrix} \left\{ \begin{matrix} a & x & d \\ s & p & q \end{matrix} \right\} \left\{ \begin{matrix} c & x & b \\ u & r & v \end{matrix} \right\} \left\{ \begin{matrix} u & x & v \\ s & g & q \end{matrix} \right\}. \quad (12.1.23)
\end{aligned}$$

$$\begin{aligned}
& \begin{array}{c} a\alpha \quad b\beta \\ q \quad u \\ p \quad r \\ s \quad v \\ d\delta \quad c\gamma \end{array} = \begin{array}{c} a\alpha \quad b\beta \\ x \\ d\delta \quad c\gamma \end{array} \begin{array}{c} q \\ a \quad u \\ p \quad r \\ d \quad s \end{array} \begin{array}{c} q \quad u \\ g \\ s \quad v \end{array} \begin{array}{c} u \quad b \\ x \quad c \\ v \quad r \end{array} \\
& = (-1)^{v+u+r+p-g-a-c} \sum_{xy\eta} (-1)^{y-\eta} \Pi_{xy}^2 \begin{pmatrix} a & y & b \\ \alpha & -\eta & \beta \end{pmatrix} \begin{pmatrix} d & y & c \\ \delta & \eta & \gamma \end{pmatrix} \left\{ \begin{matrix} a & y & b \\ r & u & x \end{matrix} \right\} \left\{ \begin{matrix} d & y & c \\ r & v & x \end{matrix} \right\} \left\{ \begin{matrix} a & p & q \\ g & u & x \end{matrix} \right\} \left\{ \begin{matrix} v & g & s \\ p & d & x \end{matrix} \right\}, \quad (12.1.24)
\end{aligned}$$


$$\begin{aligned}
& \begin{array}{c} a\alpha \quad b\beta \\ q \quad u \\ p \quad r \\ s \quad v \\ d\delta \quad c\gamma \end{array} = \begin{array}{c} q \quad u \\ a \quad b \\ p \quad r \\ g \quad x \end{array} \begin{array}{c} a \quad b \\ u \quad r \\ x \quad y \end{array} \begin{array}{c} g \quad x \\ p \quad r \\ s \quad d \end{array} \begin{array}{c} x \quad r \\ v \quad c \\ d \quad y \end{array} \begin{array}{c} a\alpha \quad b\beta \\ y \\ d\delta \quad c\gamma \end{array} \\
& = (-1)^{r+s+d-q-c} \sum_{xy\eta} (-1)^{y-\eta+x} \Pi_{xy}^2 \begin{pmatrix} d & y & b \\ \delta & -\eta & \beta \end{pmatrix} \begin{pmatrix} a & y & c \\ \alpha & \eta & \gamma \end{pmatrix} \left\{ \begin{matrix} d & y & b \\ r & u & x \end{matrix} \right\} \left\{ \begin{matrix} a & y & c \\ r & v & x \end{matrix} \right\} \left\{ \begin{matrix} a & p & q \\ x & d & u \\ v & s & g \end{matrix} \right\}, \quad (12.1.25)
\end{aligned}$$

$$\begin{aligned}
& \begin{array}{c} a\alpha \quad b\beta \\ q \quad u \\ p \quad r \\ s \quad v \\ d\delta \quad c\gamma \end{array} = \begin{array}{c} q \quad u \\ a \quad d \\ p \quad r \\ s \quad v \end{array} \begin{array}{c} u \quad b \\ r \\ x \quad y \end{array} \begin{array}{c} x \quad y \\ r \\ v \quad c \end{array} \begin{array}{c} b\beta \quad d\delta \\ y \\ c\gamma \quad a\alpha \end{array}
\end{aligned}$$


$$\begin{aligned}
& \sum_{\psi \kappa \mu \rho} (-1)^{p-\psi+q-\kappa+v-\mu+r-\rho+s-\sigma+u-\nu+g-\lambda} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & v & g \\ \kappa & -\mu & -\lambda \end{pmatrix} \begin{pmatrix} g & u \\ \lambda & \nu \\ \nu & -\psi \end{pmatrix} \begin{pmatrix} v & b & -\rho \\ \mu & \beta & -\rho \end{pmatrix} \\
& \quad \times \begin{pmatrix} r & c & s \\ \rho & \gamma & -\sigma \end{pmatrix} \begin{pmatrix} s & d & u \\ \sigma & \delta & -\nu \end{pmatrix} \\
& = (-1)^{a+b+c+u+s} \left\{ \begin{matrix} a & p & q \\ g & v & u \end{matrix} \right\} \sum_{x\xi} (-1)^{x-\xi} \Pi_x^2 \begin{pmatrix} a & d & x \\ \alpha & \delta & \xi \end{pmatrix} \begin{pmatrix} x & b & c \\ -\xi & \beta & \gamma \end{pmatrix} \left\{ \begin{matrix} a & d & x \\ s & v & u \end{matrix} \right\} \left\{ \begin{matrix} x & b & c \\ r & s & v \end{matrix} \right\}, \quad (12.1.26)
\end{aligned}$$




$$\begin{aligned}
& \sum_{\psi \kappa \rho \sigma} (-1)^{p-\psi+q-\kappa+r-\rho+s-\sigma+t-r+u-\nu+v-\mu} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & u & r \\ \kappa & \nu & -\rho \end{pmatrix} \begin{pmatrix} r & c & s \\ \rho & \gamma & -\sigma \end{pmatrix} \begin{pmatrix} s & v & p \\ \sigma & \mu & -\psi \end{pmatrix} \\
& \quad \times \begin{pmatrix} u & b & t \\ -\nu & \beta & -\tau \end{pmatrix} \begin{pmatrix} t & d & v \\ \tau & \delta & -\mu \end{pmatrix} \\
& = (-1)^{q+v+t-o-d+2u} \sum_{xy\eta} (-1)^{x+y-\eta} \Pi_{xy}^2 \begin{pmatrix} a & b & y \\ \alpha & \beta & -\eta \end{pmatrix} \begin{pmatrix} y & c & d \\ \eta & \gamma & \delta \end{pmatrix} \left\{ \begin{matrix} p & r & x \\ c & v & s \end{matrix} \right\} \left\{ \begin{matrix} p & r & x \\ u & a & q \end{matrix} \right\} \left\{ \begin{matrix} a & b & y \\ t & x & u \end{matrix} \right\} \left\{ \begin{matrix} y & c & d \\ v & t & x \end{matrix} \right\} = \quad (12.1.27)
\end{aligned}$$



$$= (-1)^{q+u-a-s-b-d} \sum_{x\xi} (-1)^{x-\xi} \Pi_x^2 \begin{pmatrix} a & x & c \\ \alpha & -\xi & \gamma \end{pmatrix} \begin{pmatrix} d & x & b \\ \delta & \xi & \beta \end{pmatrix} \left\{ \begin{matrix} d & x & b \\ u & t & v \end{matrix} \right\} \left\{ \begin{matrix} a & x & c \\ p & v & s \\ q & u & r \end{matrix} \right\}, \quad (12.1.28)$$




$$\begin{aligned}
& \sum_{\substack{\psi \kappa \rho \\ \sigma \tau \nu}} (-1)^{p-\psi+q-\kappa+r-\rho+\rho-\sigma+t-\tau+u-\nu} \begin{pmatrix} p & a & q \\ \psi & \alpha & -\kappa \end{pmatrix} \begin{pmatrix} q & b & r \\ \kappa & \beta & -\rho \end{pmatrix} \begin{pmatrix} r & c & s \\ \rho & \gamma & -\sigma \end{pmatrix} \begin{pmatrix} s & d & t \\ \sigma & \delta & -\tau \end{pmatrix} \begin{pmatrix} t & e & u \\ \tau & \varepsilon & -\nu \end{pmatrix} \begin{pmatrix} u & f & p \\ \nu & \varphi & -\psi \end{pmatrix} \\
&= \sum_{\substack{xyz \\ \xi \eta s}} (-1)^{x-\xi+y-\eta+x-s} \Pi_{xyz}^2 \begin{pmatrix} a & x & b \\ \alpha & \xi & \beta \end{pmatrix} \begin{pmatrix} c & y & d \\ \gamma & \eta & \delta \end{pmatrix} \begin{pmatrix} e & z & f \\ \varepsilon & \zeta & \varphi \end{pmatrix} \begin{pmatrix} x & y & z \\ -\xi & -\eta & -s \end{pmatrix} \begin{Bmatrix} a & x & b \\ r & q & p \end{Bmatrix} \begin{Bmatrix} c & y & d \\ t & s & r \end{Bmatrix} \quad (12.1.29)
\end{aligned}$$




$$= (-1)^{2p-u-r+a+b+d+e-c} \sum_{\substack{xyz \\ \xi \eta \xi}} (-1)^x \Pi_{xyz}^2 (-1)^{x-\xi+y-\eta+x-\xi}$$

$$\times \begin{pmatrix} a & b & x \\ \alpha & \beta & -\xi \end{pmatrix} \begin{pmatrix} x & c & y \\ \xi & \gamma & -\eta \end{pmatrix} \begin{pmatrix} y & f & z \\ \eta & \varphi & -\zeta \end{pmatrix} \begin{pmatrix} z & d & e \\ \zeta & \delta & e \end{pmatrix} \begin{Bmatrix} a & b & x \\ r & p & q \end{Bmatrix} \begin{Bmatrix} x & c & y \\ s & p & r \end{Bmatrix} \begin{Bmatrix} y & f & z \\ u & s & p \end{Bmatrix} \begin{Bmatrix} z & d & e \\ t & u & s \end{Bmatrix} = \quad (12.1.30)$$



$$\begin{aligned}
&= (-1)^{u-d-c-r} \sum_{\substack{xyz \\ \xi \eta s}} (-1)^{x-\xi+y-\eta+x-s} \Pi_{xyz}^2 : \begin{pmatrix} a & x & b \\ \alpha & \xi & \beta \end{pmatrix} \begin{pmatrix} e & y & d \\ e & \eta & s \end{pmatrix} \begin{pmatrix} c & z & f \\ \gamma & \varsigma & \varphi \end{pmatrix} \begin{pmatrix} x & y & z \\ -\xi & -\eta & -\varsigma \end{pmatrix} \\
&\quad \times \begin{Bmatrix} a & x & b \\ r & q & p \end{Bmatrix} \begin{Bmatrix} e & y & d \\ s & t & u \end{Bmatrix} \begin{Bmatrix} x & y & z \\ p & u & f \end{Bmatrix} \begin{Bmatrix} r & s & c \end{Bmatrix} \quad (12.1.31)
\end{aligned}$$



12.2 Summation of products of $6j$ and $9j$ symbols

This section presents a collection of sums involving $6j$ and $9j$ symbols. All the sums are illustrated by diagrams which display the coupling scheme for angular momenta involved in the equation. The sums are grouped in accordance with the number of $3nj$ symbols in the products. Within each group the sums are arranged according to the number of summation indices. The summation indices are printed by capital letters in equations and marked by thick j -lines. In Sec. 12.2, just as in Sec. 12.1, we use the notation

$$\Pi_{ab\dots f} = [(2a+1)(2b+1)\dots(2f+1)]^{\frac{1}{2}}.$$

Some of the sums presented are taken from Refs. [21] and [34].

12.2.1 Sums Involving One $3nj$ Symbol

$$\sum_X \Pi_X^2 \{abX\} = \Pi_{ab}^2 \quad (12.2.1)$$

$$\begin{array}{c} | \\ a \text{---} X \text{---} b \\ | \end{array} = \begin{array}{c} \bigcirc \\ a \end{array} \quad \begin{array}{c} \bigcirc \\ b \end{array}$$

$$\sum_x (-1)^{2a+x} \Pi_x^2 \{aaX\} = \Pi_a^2 \quad (12.2.2)$$

$$\begin{array}{c} | \\ a \text{---} X \text{---} a \\ | \end{array} = \begin{array}{c} \bigcirc \\ a \end{array}$$

$$\sum_x \Pi_x^2 \left\{ \begin{array}{ccc} a & b & X \\ a & b & c \end{array} \right\} = (-1)^{2c} \{abc\}, \quad (12.2.3)$$

$$\begin{array}{c} a \quad b \\ | \quad | \\ \diagdown \quad \diagup \\ X \\ \diagup \quad \diagdown \\ b \quad a \end{array} = \begin{array}{c} \bigcirc \\ a \end{array} \begin{array}{c} \bigcirc \\ b \end{array}$$

$$\sum_X (-1)^{a+b+X} \Pi_X^2 \left\{ \begin{array}{ccc} a & b & x \\ b & a & c \end{array} \right\} = \Pi_{ab} \delta_{c0} \quad (12.2.4)$$

$$\begin{array}{c} a \quad b \\ | \quad | \\ \diagdown \quad \diagup \\ X \\ \diagup \quad \diagdown \\ a \quad b \end{array} = \begin{array}{c} \bigcirc \\ a \end{array} \text{---} c \text{---} \begin{array}{c} \bigcirc \\ b \end{array}$$

$$\sum_X \Pi_X^2 \left\{ \begin{array}{ccc} a & b & e \\ c & d & f \\ e & f & X \end{array} \right\} = \frac{\delta_{bc}}{\Pi_b^2} \{abe\} \{bdf\} \quad (12.2.5)$$

$$\sum_X (-1)^{a+b+c+d-X} \Pi_X^2 \begin{Bmatrix} a & b & e \\ c & d & f \\ f & e & X \end{Bmatrix} = \frac{\delta_{ad}}{\Pi_a^2} \{dbe\} \{acf\} \quad (12.2.6)$$

12.2.2 Sums Involving Products of Two $3nj$ Symbols

(a) (i) *Single summation*

$$\sum_X \Pi_X^2 \begin{Bmatrix} a & b & X \\ c & d & p \end{Bmatrix} \begin{Bmatrix} c & d & X \\ a & b & q \end{Bmatrix} = \frac{\delta_{pq}}{\Pi_p^2} \{adp\} \{bcp\} \quad (12.2.7)$$


$$\sum_X (-1)^{p+q+X} \Pi_X^2 \begin{Bmatrix} a & b & X \\ c & d & p \end{Bmatrix} \begin{Bmatrix} c & d & X \\ b & a & q \end{Bmatrix} = \begin{Bmatrix} c & a & q \\ d & b & p \end{Bmatrix} \quad (12.2.8)$$

$$\sum_X \Pi_X^2 \begin{Bmatrix} a & f & X \\ d & q & e \\ p & c & b \end{Bmatrix} \begin{Bmatrix} a & f & X \\ e & b & s \end{Bmatrix} = (-1)^{2s} \begin{Bmatrix} a & b & s \\ c & d & p \end{Bmatrix} \begin{Bmatrix} c & d & s \\ e & f & q \end{Bmatrix}, \quad (12.2.9)$$


$$\sum_X (-1)^{R+X} \Pi_X^2 \begin{Bmatrix} a & f & X \\ d & q & e \\ p & c & b \end{Bmatrix} \begin{Bmatrix} a & f & X \\ b & e & s \end{Bmatrix} = (-1)^{2s} \begin{Bmatrix} p & q & s \\ e & a & d \end{Bmatrix} \begin{Bmatrix} p & q & s \\ f & b & c \end{Bmatrix}, \quad (12.2.10)$$

where $R = a + b + c + d + e + f + p + q$

$$\sum_X \Pi_X^2 \begin{Bmatrix} a & b & p \\ c & d & q \\ r & s & X \end{Bmatrix} \begin{Bmatrix} e & f & p \\ g & h & q \\ r & s & X \end{Bmatrix} = (-1)^{b-c-f+g} \begin{Bmatrix} - & r & a & c \\ s & - & b & d \\ h & g & - & q \\ f & e & p & - \end{Bmatrix} \quad (12.2.11)$$




$$(-1)^{p+q-r-s} \sum_X \Pi_X^2 \begin{Bmatrix} a & b & p \\ c & d & q \\ r & s & X \end{Bmatrix} \begin{Bmatrix} e & f & p \\ g & h & q \\ r & s & X \end{Bmatrix} = (-1)^{b+g-c-f} \sum_Y \Pi_Y^2 \begin{Bmatrix} a & p & b \\ r & e & g \\ c & f & Y \end{Bmatrix} \begin{Bmatrix} d & s & b \\ q & h & g \\ c & f & Y \end{Bmatrix}, \quad (12.2.12)$$




(b) (ii) Double summation

$$\sum_{XY} \Pi_{XY}^2 \begin{Bmatrix} a & b & X \\ c & d & Y \\ e & f & j \end{Bmatrix} \begin{Bmatrix} a & b & X \\ c & d & Y \\ g & h & j \end{Bmatrix} = \frac{\delta_{eg} \delta_{fh}}{\Pi_{ef}^2} \{ace\} \{bdh\} \{gfj\}, \quad (12.2.13)$$



$$\sum_{XY} (-1)^Y \Pi_{XY}^2 \begin{Bmatrix} a & b & X \\ c & d & Y \\ e & f & j \end{Bmatrix} \begin{Bmatrix} a & b & X \\ d & c & Y \\ g & h & j \end{Bmatrix} = (-1)^{2b+f+h} \begin{Bmatrix} a & d & g \\ c & b & h \\ e & f & j \end{Bmatrix}, \quad (12.2.14)$$



(c) (iii) Triple summation

$$\sum_{XYZ} \Pi_{XYZ}^2 \begin{Bmatrix} X & Y & Z \\ a & b & c \end{Bmatrix}^2 = \Pi_{abc}^2 \quad (12.2.15)$$

$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} X & Y & Z \\ a & b & c \\ b & c & a \end{matrix} \right\} \left\{ \begin{matrix} X & Y & Z \\ c & a & b \end{matrix} \right\} = \{abc\} \quad (12.2.16)$$

$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} X & Y & Z \\ a & b & c \\ d & e & f \end{matrix} \right\}^2 = \{abc\}\{def\} \quad (12.2.17)$$

12.2.3 Sums Involving Products of Three $3nj$ Symbols


(a) *Single summation*

$$\sum_X (-1)^{R+X} \Pi_X^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ b & a & r \end{matrix} \right\} = \left\{ \begin{matrix} p & q & r \\ e & a & d \end{matrix} \right\} \left\{ \begin{matrix} p & q & r \\ f & b & c \end{matrix} \right\}, \quad (12.2.18)$$


where $R = a + b + c + d + e + f + p + q + r$,

$$\sum_X (-1)^{2X} \Pi_X^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ a & b & r \end{matrix} \right\} = \left\{ \begin{matrix} a & f & r \\ d & q & e \\ p & c & b \end{matrix} \right\}, \quad (12.2.19)$$

$$(-1)^{p+q-e-f} \sum_X \Pi_X^2 \left\{ \begin{matrix} a & b & p \\ c & d & q \\ e & f & X \end{matrix} \right\} \left\{ \begin{matrix} p & q & X \\ k & l & g \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ k & l & h \end{matrix} \right\} = \left\{ \begin{matrix} f & b & p & g \\ d & a & l & k \\ q & c & e & h \end{matrix} \right\} \quad (12.2.20)$$




$$\begin{aligned} & (-1)^{p+q-e-f} \sum_X \Pi_X^2 \left\{ \begin{matrix} a & b & p \\ c & d & q \\ e & f & X \end{matrix} \right\} \left\{ \begin{matrix} p & q & X \\ k & l & g \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ k & l & h \end{matrix} \right\} \\ & = (-1)^{b+g-c-h} \sum_Y \Pi_Y^2 \left\{ \begin{matrix} d & f & b \\ q & k & g \\ c & h & Y \end{matrix} \right\} \left\{ \begin{matrix} b & g & Y \\ l & a & p \end{matrix} \right\} \left\{ \begin{matrix} c & h & Y \\ l & a & e \end{matrix} \right\}. \end{aligned} \quad (12.2.21)$$




(b) Double summation


$$\sum_{XY} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ a & b & Y \end{matrix} \right\} \left\{ \begin{matrix} a & b & q \\ c & d & Y \end{matrix} \right\} = \left\{ \begin{matrix} a & b & q \\ c & d & p \end{matrix} \right\}, \quad (12.2.22)$$




$$\sum_{XY} (-1)^{X+Y+p} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} a & c & Y \\ d & b & X \end{matrix} \right\} \left\{ \begin{matrix} a & d & q \\ b & c & Y \end{matrix} \right\} = \frac{\delta_{pq}}{\Pi_p^2} \{adp\} \{bcp\} \quad (12.2.23)$$




$$\sum_{XY} (-1)^{X+Y+2a-s-t} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & p \\ c & d & X \\ q & Y & g \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ p & g & s \end{matrix} \right\} \left\{ \begin{matrix} b & d & Y \\ q & g & t \end{matrix} \right\} = \left\{ \begin{matrix} a & b & p \\ c & g & s \\ q & t & d \end{matrix} \right\}, \quad (12.2.24)$$




$$\sum_{XY} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & p \\ c & d & X \\ q & Y & g \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ g & p & s \end{matrix} \right\} \left\{ \begin{matrix} b & d & Y \\ g & q & t \end{matrix} \right\} = (-1)^{2s} \frac{\delta_{st}}{\Pi_s^2} \{dgs\} \left\{ \begin{matrix} a & b & p \\ s & c & q \end{matrix} \right\}, \quad (12.2.25)$$



$$\sum_{XY} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ e & f & g \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ Y & g & h \end{matrix} \right\} \left\{ \begin{matrix} c & d & Y \\ b & h & j \end{matrix} \right\} = (-1)^{2h} \frac{\delta_{fj}}{\Pi_f^2} \{bdf\} \left\{ \begin{matrix} e & j & g \\ h & a & c \end{matrix} \right\}, \quad (12.2.26)$$




$$\sum_{XY} \Pi_{XY}^2 \left\{ \begin{matrix} a & f & X \\ d & q & e \\ p & c & b \end{matrix} \right\} \left\{ \begin{matrix} a & f & X \\ h & r & e \\ Y & g & b \end{matrix} \right\} \left\{ \begin{matrix} a & h & Y \\ g & b & s \end{matrix} \right\} = \left\{ \begin{matrix} a & b & s \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & s \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & s \\ g & h & r \end{matrix} \right\}, \quad (12.2.27)$$




(c) (iii) Triple summation


$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & p \\ c & d & X \\ q & Y & Z \end{matrix} \right\} \left\{ \begin{matrix} p & X & Z \\ d & e & c \end{matrix} \right\} \left\{ \begin{matrix} q & Y & Z \\ d & e & b \end{matrix} \right\} = (-1)^{2e} \Pi_d^2 \left\{ \begin{matrix} a & b & p \\ e & c & q \end{matrix} \right\}, \quad (12.2.28)$$




$$\sum_{XYZ} (-1)^{X+Y-e-h} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Z \\ e & f & g \end{matrix} \right\} \left\{ \begin{matrix} g & b & Y \\ c & d & Z \\ h & j & a \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ g & Z & Y \end{matrix} \right\} = \frac{\delta_{fj}}{\Pi_d^2} \{bdf\} \left\{ \begin{matrix} e & f & g \\ h & c & a \end{matrix} \right\}, \quad (12.2.29)$$



$$\sum_{XYZ} (-1)^{X+f-a-l} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & p \\ c & d & X \\ q & Y & Z \end{matrix} \right\} \left\{ \begin{matrix} b & d & Y \\ c & f & Z \\ l & k & q \end{matrix} \right\} \left\{ \begin{matrix} p & X & Z \\ c & f & d \end{matrix} \right\} = \frac{\delta_{pk}}{\Pi_k^2} \{dfp\} \left\{ \begin{matrix} a & b & p \\ l & q & c \end{matrix} \right\}, \quad (12.2.30)$$




$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & p \\ c & d & X \\ q & Y & Z \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & Z \\ g & h & p \end{matrix} \right\} \left\{ \begin{matrix} b & d & Y \\ e & f & Z \\ j & k & q \end{matrix} \right\} = \frac{\delta_{hk}}{\Pi_h^2} \{dfh\} \left\{ \begin{matrix} a & b & p \\ c & e & g \\ q & j & h \end{matrix} \right\}, \quad (12.2.31)$$




12.2.4 Sums Involving Products of Four $3nj$ Symbols

(a) (i) Single summation

$$\begin{aligned}
& \sum_X (-1)^{R-X} \Pi_X^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ g & h & r \end{matrix} \right\} \left\{ \begin{matrix} g & h & X \\ b & a & s \end{matrix} \right\} \\
&= \sum_Y (-1)^{2Y+a+b+e+\rho} \Pi_Y^2 \left\{ \begin{matrix} s & h & b \\ g & r & f \\ a & e & Y \end{matrix} \right\} \left\{ \begin{matrix} b & f & Y \\ q & p & c \end{matrix} \right\} \left\{ \begin{matrix} a & e & Y \\ q & p & d \end{matrix} \right\} = \left\{ \begin{matrix} e & h & b & c \\ f & r & s & p & q & d \end{matrix} \right\}, \quad (12.2.32)
\end{aligned}$$


where $R = a + b + c + d + e + f + g + h + p + q + r + s$.

$$\begin{aligned}
& \sum_X \Pi_X^2 \left\{ \begin{matrix} a & b & X \\ c & d & p \end{matrix} \right\} \left\{ \begin{matrix} c & d & X \\ e & f & q \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ g & h & r \end{matrix} \right\} \left\{ \begin{matrix} g & h & X \\ a & b & s \end{matrix} \right\} \\
&= \sum_X \Pi_X^2 \left\{ \begin{matrix} a & f & X \\ d & q & e \\ p & c & b \end{matrix} \right\} \left\{ \begin{matrix} a & f & X \\ h & r & e \\ s & g & b \end{matrix} \right\} = (-1)^{-p+q-r+s} \left\{ \begin{matrix} - & a & d & p \\ f & - & q & a \\ g & s & - & b \\ r & h & e & - \end{matrix} \right\}, \quad (12.2.33)
\end{aligned}$$




(b) (ii) Double summation


$$\sum_{XY} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & c \\ d & X & Y \end{matrix} \right\} \left\{ \begin{matrix} a & e & f \\ g & X & Y \end{matrix} \right\} \left\{ \begin{matrix} c & g & p \\ f & d & X \end{matrix} \right\} \left\{ \begin{matrix} b & g & q \\ e & d & Y \end{matrix} \right\} = (-1)^{2(c+e)} \left\{ \begin{matrix} a & b & c \\ g & p & q \end{matrix} \right\} \left\{ \begin{matrix} a & e & f \\ d & p & q \end{matrix} \right\}, \quad (12.2.34)$$




$$\begin{aligned}
& \sum_{XY} (-1)^{X+Y} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & c \\ d & X & Y \end{matrix} \right\} \left\{ \begin{matrix} a & e & f \\ g & X & Y \end{matrix} \right\} \left\{ \begin{matrix} c & f & p \\ g & d & X \end{matrix} \right\} \left\{ \begin{matrix} b & e & q \\ g & d & Y \end{matrix} \right\} \\
&= (-1)^{-a+b+yc-d+e+f-e-p} \frac{\delta_{pq}}{\Pi_p^2} \{dgp\} \left\{ \begin{matrix} a & b & c \\ p & f & e \end{matrix} \right\} \quad (12.2.35)
\end{aligned}$$



$$\sum_{XY} (-1)^Y \Pi_{XY}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ p & q & r \end{matrix} \right\} \left\{ \begin{matrix} X & Y & r \\ j & h & g \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ h & g & e \end{matrix} \right\} \left\{ \begin{matrix} c & d & Y \\ j & g & f \end{matrix} \right\} = (-1)^{a-b+d+e-j-p+r} \left\{ \begin{matrix} e & b & h \\ f & d & j \\ p & q & r \end{matrix} \right\} \left\{ \begin{matrix} a & c & p \\ f & e & g \end{matrix} \right\}, \quad (12.2.36)$$




$$\begin{aligned} & \sum_{XY} (-1)^{X+Y} \Pi_{XY}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ p & q & s \end{matrix} \right\} \left\{ \begin{matrix} e & f & X \\ g & h & Y \\ r & t & s \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ f & e & k \end{matrix} \right\} \left\{ \begin{matrix} c & d & Y \\ h & g & l \end{matrix} \right\} \\ &= (-1)^{k+l+a+c+p-s-h-t-f} \sum_Z (-1)^Z \Pi_Z^2 \left\{ \begin{matrix} c & g & l \\ a & e & k \\ p & r & Z \end{matrix} \right\} \left\{ \begin{matrix} f & b & k \\ h & d & l \\ t & q & Z \end{matrix} \right\} \left\{ \begin{matrix} p & r & Z \\ t & q & s \end{matrix} \right\}, \end{aligned} \quad (12.2.37)$$




(c) (iii) Triple summation


$$\begin{aligned} & \sum_{X_1 X_2 X_3} (-1)^{X_1+X_2+X_3+a_1+a_2+a_3+b_1+b_2+b_3+c} \Pi_{X_1 X_2 X_3}^2 \left\{ \begin{matrix} a_1 & a_2 & a_3 \\ X_1 & X_2 & X_3 \end{matrix} \right\} \left\{ \begin{matrix} a_1 & b_2 & b_3 \\ c & X_3 & X_2 \end{matrix} \right\} \\ & \times \left\{ \begin{matrix} a_2 & b_3 & b_1 \\ c & X_1 & X_3 \end{matrix} \right\} \left\{ \begin{matrix} a_3 & b_1 & b_2 \\ c & X_2 & X_1 \end{matrix} \right\} = \Pi_c^2 \left\{ \begin{matrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{matrix} \right\}, \end{aligned} \quad (12.2.38)$$




$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} X & Y & Z \\ d & e & f \\ a & b & c \end{matrix} \right\} \left\{ \begin{matrix} X & Y & Z \\ f & c & g \end{matrix} \right\} \left\{ \begin{matrix} X & a & d \\ b' & g & c \end{matrix} \right\} \left\{ \begin{matrix} Y & b & e \\ d' & f & g \end{matrix} \right\} = (-1)^{2(b+d)} \frac{\delta_{bb'} \delta_{dd'}}{\Pi_{bd}^2} \{abc\} \{def\} \{bdg\}, \quad (12.2.39)$$




$$\sum_{XYZ} (-1)^{2Z} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ e' & f' & g \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ f & Z & d \end{matrix} \right\} \left\{ \begin{matrix} c & d & Y \\ Z & e & a \end{matrix} \right\} \left\{ \begin{matrix} e & f & g \\ X & Y & Z \end{matrix} \right\} = \frac{\delta_{ee'} \delta_{ff'}}{\Pi_{ef}^2} \{ace\} \{bdf\} \{egf\}, \quad (12.2.40)$$




$$\begin{aligned} & \sum_{XYZ} (-1)^Z \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & c' \\ d & e & f \\ X & Y & g \end{matrix} \right\} \left\{ \begin{matrix} l & e & j \\ a & b & c \\ Z & Y & h \end{matrix} \right\} \left\{ \begin{matrix} a & d & X \\ k & Z & l \end{matrix} \right\} \left\{ \begin{matrix} X & Y & g \\ h & k & Z \end{matrix} \right\} \\ & = (-1)^{2c+b-d+g+j} \frac{\delta_{cc'}}{\Pi_c^2} \{abc\} \left\{ \begin{matrix} k & j & f \\ c & g & h \end{matrix} \right\} \left\{ \begin{matrix} k & j & f \\ e & d & l \end{matrix} \right\}, \end{aligned} \quad (12.2.41)$$



$$\sum_{XYZ} (-1)^{2Y-Z} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ e & f & Z \end{matrix} \right\} \left\{ \begin{matrix} g & h & X \\ k & l & Y \\ f & e & Z \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ g & h & j \end{matrix} \right\} \left\{ \begin{matrix} c & d & Y \\ k & l & j' \end{matrix} \right\} = (-1)^{-b+c-h+k} \frac{\delta_{jj'}}{\Pi_j^2} \left\{ \begin{matrix} a & c & e \\ l & h & j \end{matrix} \right\} \left\{ \begin{matrix} b & d & f \\ k & g & j \end{matrix} \right\}, \quad (12.2.42)$$



$$\sum_{XYZ} (-1)^{Y-a-b-c-f-h-p+q} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & X \\ g & c & q \\ p & Z & Y \end{matrix} \right\} \left\{ \begin{matrix} b & d & f' \\ a & h & j \\ Z & Y & p \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ d & e & f \end{matrix} \right\} \left\{ \begin{matrix} d & h & Y \\ q & X & e \end{matrix} \right\} \\ = (-1)^{2j} \frac{\delta_{ff'}}{\Pi_f^2} \{bdf\} \left\{ \begin{matrix} e & g & j \\ p & f & a \end{matrix} \right\} \left\{ \begin{matrix} e & g & j \\ c & h & q \end{matrix} \right\}, \quad (12.2.43)$$



$$\sum_{XYZ} \Pi_{XYZ}^2 \left\{ \begin{matrix} a & b & X \\ c & d & Y \\ t & s & r \end{matrix} \right\} \left\{ \begin{matrix} a & b & X \\ h & j & q \\ s & f & Z \end{matrix} \right\} \left\{ \begin{matrix} k & l & p \\ c & d & Y \\ e & f & Z \end{matrix} \right\} \left\{ \begin{matrix} p & q & r \\ X & Y & Z \end{matrix} \right\} = \left\{ \begin{matrix} k & l & p \\ h & j & q \\ t & s & r \end{matrix} \right\} \left\{ \begin{matrix} k & h & t \\ a & c & e \end{matrix} \right\} \left\{ \begin{matrix} l & j & s \\ b & d & f \end{matrix} \right\}, \quad (12.2.44)$$

The diagram illustrates an identity involving the summation of products of $6j$ and $9j$ symbols. The left-hand side (LHS) features four 3D-like diagrams (cubes) with various labels (a, b, c, d, e, f, h, j, k, l, p, q, r, s, t, X, Y, Z) and internal lines. The right-hand side (RHS) features three 2D-like diagrams (squares and a cube) with similar labels and internal lines. An equals sign is placed between the two groups of diagrams.

Chapter 13

Matrix elements of irreducible tensor operators

We have added explicit expressions for reduced matrix elements each time the book gave an expression for a matrix element of an irreducible tensor operator. These equations are as yet unnumbered.

13.1 The Wigner-Eckart theorem and the evaluation of matrix elements

In this section we consider the evaluation of matrix elements of irreducible tensor operators in a general form. Let a quantum mechanical system be characterized by a set of the quantum numbers (njm) , where j and m are the angular momentum and its projection on the quantization axis, respectively. The symbol n stands for all other quantum numbers required to define the state of the system. Let Ψ_{njm} be the wave function of this state. The matrix element of any irreducible tensor operator $\widehat{\mathfrak{M}}_k$ of rank k between the states $|n'j'm'\rangle$ and $|njm\rangle$ is defined as

$$\langle n'j'm'|\widehat{\mathfrak{M}}_{k\kappa}|njm\rangle = \int \Psi_{n'j'm'}^* \widehat{\mathfrak{M}}_{k\kappa} \Psi_{njm} d\tau, \quad (13.1.1)$$

where $\widehat{\mathfrak{M}}_{k\kappa}$ is a tensor component.

13.1.1 Wigner-Eckart Theorem

The Wigner-Eckart theorem is widely used for evaluation of matrix elements. According to this theorem, the dependence of any matrix element on the orientation of the coordinate system, i.e., on the projections m , m' and κ , is entirely included in the $3jm$ symbol or the Clebsch-Gordan coefficient,

$$\langle n'j'm'|\widehat{\mathfrak{M}}_{k\kappa}|njm\rangle = (-1)^{j'-m'} \begin{pmatrix} j' & k & j \\ -m' & \kappa & m \end{pmatrix} \langle n'j'|\widehat{\mathfrak{M}}_k|nj\rangle = (-1)^{2k} C_{j \ m \ k \ \kappa}^{j' \ m'} \frac{\langle n'j'|\widehat{\mathfrak{M}}_k|nj\rangle}{\Pi_{j'}}. \quad (13.1.2)$$

The invariant factor $\langle n'j'|\widehat{\mathfrak{M}}_k|nj\rangle$ is called the reduced matrix element of the set of tensor operators $\widehat{\mathfrak{M}}_{k\kappa}$. As follows from the definition (13.1.2), the reduced matrix element of the unit operator is given by

$$\langle n'j'|\widehat{\mathbf{I}}|nj\rangle = \sqrt{2j+1} \delta_{n'n} \delta_{j'j}. \quad (13.1.3)$$

In this section, as well as in Chap. 12 we shall again use the notation

$$\Pi_{ab\dots c} \equiv [(2a+1)(2b+1)\dots(2c+1)]^{\frac{1}{2}}.$$

13.1.2 Sum Rules

Making use of the definition (13.1.2), one can obtain the following rules for sums of matrix elements of irreducible tensor operators:

$$\sum_{m\kappa} \left| \langle n'j'm' | \widehat{\mathfrak{M}}_{k\kappa} | njm \rangle \right|^2 = \frac{\left| \langle n'j' | \widehat{\mathfrak{M}}_k | nj \rangle \right|^2}{2j' + 1}, \quad (13.1.4)$$

$$\sum_{m'\kappa} \left| \langle n'j'm' | \widehat{\mathfrak{M}}_{k\kappa} | njm \rangle \right|^2 = \frac{\left| \langle n'j' | \widehat{\mathfrak{M}}_k | nj \rangle \right|^2}{2j + 1}, \quad (13.1.5)$$

$$\sum_{mm'} \left| \langle n'j'm' | \widehat{\mathfrak{M}}_{k\kappa} | njm \rangle \right|^2 = \frac{\left| \langle n'j' | \widehat{\mathfrak{M}}_k | nj \rangle \right|^2}{2k + 1}, \quad (13.1.6)$$

$$\sum_{mm'\kappa} \left| \langle n'j'm' | \widehat{\mathfrak{M}}_{k\kappa} | njm \rangle \right|^2 = \left| \langle n'j' | \widehat{\mathfrak{M}}_k | nj \rangle \right|^2. \quad (13.1.7)$$

13.1.3 Matrix Elements of Products of Irreducible Tensor Operators

From two irreducible tensor operators $\widehat{P}_{a\alpha}$ and $\widehat{Q}_{b\beta}$ one can compose the direct or reducible product $\widehat{P}_{a\alpha} \widehat{Q}_{b\beta}$ and the irreducible product $\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma}$ of rank c (see Chap. 3). A matrix element of the direct tensor product may be written in the form

$$\langle n'j'm' | \widehat{\mathbf{P}}_{a\alpha} \widehat{\mathbf{Q}}_{b\beta} | njm \rangle = \frac{(-1)^{2a+2b}}{\Pi_{j'}} \sum_{n_1j_1m_1} C_{j_1m_1a\alpha}^{j'm'} C_{j_1m_1b\beta}^{j_1m_1} \frac{1}{\Pi_{j_1}} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | nj \rangle. \quad (13.1.8)$$

Another frequently used relationship for the same matrix element is as follows:

$$\langle n'j'm' | \widehat{\mathbf{P}}_{a\alpha} \widehat{\mathbf{Q}}_{b\beta} | njm \rangle = \frac{(-1)^{j'+j}}{\Pi_{j'}} \sum_{c\gamma} (-1)^{-c} \Pi_c C_{j_1m_1c\gamma}^{j'm'} C_{a\alpha b\beta}^c \sum_{n_1j_1} \left\{ \begin{matrix} a & b & c \\ j & j' & j_1 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | nj \rangle. \quad (13.1.9)$$

Matrix elements of irreducible tensor products may be evaluated from the equation

$$\begin{aligned} \langle n'j'm' | \{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma} | njm \rangle &= (-1)^{j'+j-c} \frac{\Pi_c}{\Pi_{j'}} C_{j_1m_1c\gamma}^{j'm'} \sum_{n_1j_1} \left\{ \begin{matrix} a & b & c \\ j & j' & j_1 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | nj \rangle, \\ \langle n'j' | \{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_c | nj \rangle &= (-1)^{j'+j+c} \Pi_c \sum_{n_1j_1} \left\{ \begin{matrix} a & b & c \\ j & j' & j_1 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | nj \rangle. \end{aligned} \quad (13.1.10)$$

Putting $c = 0$ and $b = a$ in Eq. (13.1.10) and multiplying the resulting expression by $(-1)^{-a} \Pi_a$ (in accordance with Eq. (3.1.34)) we get a matrix element of scalar product of two tensors of rank a

$$\begin{aligned} \langle n'j'm' | \widehat{\mathbf{P}}_a \cdot \widehat{\mathbf{Q}}_a | njm \rangle &= \delta_{jj'} \delta_{mm'} \frac{1}{\Pi_j^2} \sum_{n_1j_1} (-1)^{-j+j_1} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_a | nj \rangle, \\ \langle n'j' | \widehat{\mathbf{P}}_a \cdot \widehat{\mathbf{Q}}_a | nj \rangle &= \delta_{jj'} \frac{1}{\Pi_j} \sum_{n_1j_1} (-1)^{-j+j_1} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_a | nj \rangle, \end{aligned} \quad (13.1.11)$$

From the three irreducible tensor operators $\widehat{P}_{a\alpha}$, $\widehat{Q}_{b\beta}$ and $\widehat{R}_{d\delta}$, one can compose the products

$$\begin{aligned} &\widehat{P}_{a\alpha} \widehat{Q}_{b\beta} \widehat{R}_{d\delta}, \quad \{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma} \widehat{R}_{d\delta}, \quad \widehat{P}_{a\alpha} \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_{f\varphi}, \\ &\{\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_c \otimes \widehat{\mathbf{R}}_d\}_{f\varphi}, \quad \{\widehat{\mathbf{P}}_a \otimes \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_c\}_{f\varphi}. \end{aligned}$$

Making permutations of operators one can construct other products which differ from the given above only by commutators of operators. The matrix element of the direct product of three tensor operators is given by

$$\begin{aligned} \langle n'j'm' | \widehat{P}_{a\alpha} \widehat{Q}_{b\beta} \widehat{R}_{d\delta} | njm \rangle &= \frac{(-1)^{2a+2b+2d}}{\Pi_{j'}} \sum_{n_1j_1n_2j_2} \frac{1}{\Pi_{j_1j_2}} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \\ &\quad \times \langle n_2j_2 | \widehat{\mathbf{R}}_d | nj \rangle \sum_{m_1m_2} C_{j_1m_1a\alpha}^{j'm'} C_{j_2m_2b\beta}^{j_1m_1} C_{j_2m_2d\delta}^{j_1m_1}. \end{aligned} \quad (13.1.12)$$

The evaluation of matrix elements of the products $\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma} \widehat{R}_{d\delta}$ and $\widehat{P}_{a\alpha} \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_{c\gamma}$ is reduced to the successive use of Eqs. (13.1.8)-(13.1.10).

The matrix element of the irreducible tensor product of three tensors may be represented in the form

$$\begin{aligned} \langle n'j'm' | \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f | njm \rangle &= (-1)^{f+c-j} \frac{\Pi_{fc}}{\Pi_{j'}} C_{j'm'f}^{j'c} \\ &\times \sum_{n_1j_1n_2j_2} (-1)^{j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j' & j & j_2 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_d | nj \rangle. \end{aligned} \quad (13.1.13)$$

$$\begin{aligned} \langle n'j' | \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f | nj \rangle &= (-1)^{f+c-j} \Pi_{fc} \sum_{n_1j_1n_2j_2} (-1)^{j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j' & j & j_2 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_d | nj \rangle. \end{aligned}$$

By the use of Eq. (3.3.1) the evaluation of the matrix elements of $\{\widehat{\mathbf{P}}_a \otimes \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_c\}_{f\varphi}$ is reduced to applying Eq. (13.1.13).

The matrix element of a scalar product may be obtained from Eq. (13.1.13). Taking into account the factor $(-1)^{-c}\Pi_c$, one gets

$$\begin{aligned} \langle n'j'm' | (\{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \cdot \widehat{\mathbf{R}}_c) | njm \rangle &= \delta_{jj'} \delta_{mm'} (-1)^{2j-c} \frac{\Pi_c}{\Pi_j^2} \\ &\times \sum_{n_1j_1n_2j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_c | nj \rangle. \end{aligned} \quad (13.1.14)$$

$$\begin{aligned} \langle n'j' | \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \cdot \widehat{\mathbf{R}}_c \} | nj \rangle &= \delta_{jj'} (-1)^{2j-c} \frac{\Pi_c}{\Pi_j} \sum_{n_1j_1n_2j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_c | nj \rangle. \end{aligned}$$

Of the four irreducible tensors $\widehat{P}_{a\alpha}$, $\widehat{Q}_{b\beta}$, $\widehat{R}_{d\delta}$ and $\widehat{S}_{e\epsilon}$ one can compose the following types of tensor products:

- (a) $\widehat{P}_{a\alpha} \widehat{Q}_{b\beta} \widehat{R}_{d\delta} \widehat{S}_{e\epsilon}$;
- (b) $\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma} \widehat{R}_{d\delta} \widehat{S}_{e\epsilon}$, $\widehat{P}_{a\alpha} \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_{k\kappa} \widehat{S}_{e\epsilon}$, $\widehat{P}_{a\alpha} \widehat{Q}_{b\beta} \{\widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e\}_{h\sigma}$;
- (c) $\{\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_c \otimes \widehat{\mathbf{R}}_d\}_f \widehat{S}_{e\epsilon}$, $\{\widehat{\mathbf{P}}_a \otimes \{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_h\}_f \widehat{S}_{e\epsilon}$, $\widehat{P}_{a\alpha} \{\{\widehat{\mathbf{Q}}_b \otimes \widehat{\mathbf{R}}_d\}_h \otimes \widehat{\mathbf{S}}_e\}_{f\varphi}$, $\widehat{P}_{a\alpha} \{\widehat{\mathbf{Q}}_b \otimes \{\widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e\}_h\}_{f\varphi}$, $\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_{c\gamma} \{\widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e\}_{h\sigma}$;
- (d) $\{\{\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_c \otimes \widehat{\mathbf{R}}_d\}_f \otimes \widehat{\mathbf{S}}_e\}_{k\kappa}$, $\{\widehat{\mathbf{P}}_a \otimes \{\widehat{\mathbf{Q}}_b \otimes \{\widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e\}_h\}_f\}_k$, $\{\{\widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b\}_c \otimes \{\widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e\}_f\}_{k\kappa}$.

Products with other orders of the operators can be reduced to the products listed here using the commutation rules.

A matrix element of a direct product may be evaluated as follows:

$$\begin{aligned} \langle n'j'm' | \widehat{P}_{a\alpha} \widehat{Q}_{b\beta} \widehat{R}_{d\delta} \widehat{S}_{e\epsilon} | njm \rangle &= (-1)^{2a+2b+2d+2e} \frac{1}{\Pi_{j'}} \sum_{\substack{n_1n_2n_3 \\ j_1j_2j_3}} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_d | n_3j_3 \rangle \\ &\times \langle n_3j_3 | \widehat{\mathbf{S}}_e | nj \rangle \frac{1}{\Pi_{j_1j_2j_3}} \sum_{m_1m_2m_3} C_{j_1m_1a\alpha}^{j'm'} C_{j_2m_2b\beta}^{j'm'} C_{j_3m_3d\delta}^{j'm'} C_{jm_e\epsilon}^{j'm'}. \end{aligned} \quad (13.1.15)$$

Matrix elements of products of (b)- and (c)-type products can be evaluated by using Eqs. (13.1.8)-(13.1.14).

The evaluation of matrix elements of irreducible tensor products yields

$$\begin{aligned} \langle n'j'm' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f \otimes \widehat{\mathbf{S}}_e \}_{k\kappa} | njm \rangle &= (-1)^{j-j'+f+c-k} \frac{\Pi_{fck}}{\Pi_{j'}} C_{j'm'k\kappa}^{j'm'} \sum_{\substack{n_1n_2n_3 \\ j_1j_2j_3}} (-1)^{j_2+j_3} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \\ &\times \left\{ \begin{matrix} d & c & f \\ j' & j_3 & j_2 \end{matrix} \right\} \left\{ \begin{matrix} e & f & k \\ j' & j & j_3 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_d | n_3j_3 \rangle \langle n_3j_3 | \widehat{\mathbf{S}}_e | nj \rangle, \end{aligned} \quad (13.1.16)$$

$$\begin{aligned} \langle n'j'm' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f \widehat{\mathbf{S}}_f \} | njm \rangle &= (-1)^{c-f-j} \frac{\Pi_{fc}}{\Pi_j^2} \delta_{j'j} \delta_{m'm} \sum_{\substack{n_1n_2n_3 \\ j_1j_2j_3}} (-1)^{j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j & j_3 & j_2 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1j_1 \rangle \langle n_1j_1 | \widehat{\mathbf{Q}}_b | n_2j_2 \rangle \langle n_2j_2 | \widehat{\mathbf{R}}_d | n_3j_3 \rangle \langle n_3j_3 | \widehat{\mathbf{S}}_f | nj \rangle. \end{aligned} \quad (13.1.17)$$

$$\begin{aligned} \langle n'j' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f \otimes \widehat{\mathbf{S}}_e \}_k | nj \rangle &= (-1)^{j-j'+f+c-k} \Pi_{fck} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} (-1)^{j_2+j_3} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j' & j_3 & j_2 \end{matrix} \right\} \left\{ \begin{matrix} e & f & k \\ j' & j & j_3 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_e | nj \rangle, \end{aligned}$$

$$\begin{aligned} \langle n'j' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \widehat{\mathbf{R}}_d \}_f \cdot \widehat{\mathbf{S}}_f \} | nj \rangle &= (-1)^{c-f-j} \frac{\Pi_{fck}}{\Pi_j} \delta_{j'j} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} (-1)^{j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j & j_3 & j_2 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_f | nj \rangle. \end{aligned}$$

Finally Eq. (3.3.5) allows one to simplify the calculation of matrix elements of the tensor product $\{ \widehat{\mathbf{P}}_a \otimes \{ \widehat{\mathbf{Q}}_b \otimes \{ \widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e \}_h \}_f \}_k$ to Eq. (13.1.16). Thus

$$\begin{aligned} \langle n'j'm' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \{ \widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e \}_f \}_k | njm \rangle &= (-1)^{2j'+c-k-f} \frac{\Pi_{cfk}}{\Pi_{j'}} C_{j'm'k}^{j' m' \kappa} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & e & f \\ j & j_2 & j_3 \end{matrix} \right\} \\ &\times \left\{ \begin{matrix} f & c & k \\ j' & j & j_2 \end{matrix} \right\} \langle n'j' | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_e | nj \rangle, \end{aligned} \quad (13.1.18)$$

$$\begin{aligned} \langle n'j' | \{ \{ \{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \otimes \{ \widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e \}_f \}_k | nj \rangle &= (-1)^{2j'+c-k-f} \Pi_{cfk} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} \left\{ \begin{matrix} a & b & c \\ j_2 & j' & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & e & f \\ j & j_2 & j_3 \end{matrix} \right\} \left\{ \begin{matrix} f & c & k \\ j' & j & j_2 \end{matrix} \right\} \\ &\times \langle n'j' | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_e | nj \rangle, \end{aligned}$$

$$\begin{aligned} \langle n'j'm' | (\{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \cdot \{ \widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e \}_e) | njm \rangle &= \delta_{j'j} \delta_{m'm} \frac{\Pi_c^2}{\Pi_j^2} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} (-1)^{-j+j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & e & c \\ j & j_2 & j_3 \end{matrix} \right\} \\ &\times \langle n'j | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_e | nj \rangle. \end{aligned} \quad (13.1.19)$$

$$\begin{aligned} \langle n'j' | (\{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_c \cdot \{ \widehat{\mathbf{R}}_d \otimes \widehat{\mathbf{S}}_e \}_c) | nj \rangle &= \delta_{j'j} \frac{\Pi_c^2}{\Pi_j} \sum_{\substack{n_1 n_2 n_3 \\ j_1 j_2 j_3}} (-1)^{-j+j_2} \left\{ \begin{matrix} a & b & c \\ j_2 & j & j_1 \end{matrix} \right\} \left\{ \begin{matrix} d & e & c \\ j & j_2 & j_3 \end{matrix} \right\} \\ &\times \langle n'j | \widehat{\mathbf{P}}_a | n_1 j_1 \rangle \langle n_1 j_1 | \widehat{\mathbf{Q}}_b | n_2 j_2 \rangle \langle n_2 j_2 | \widehat{\mathbf{R}}_d | n_3 j_3 \rangle \langle n_3 j_3 | \widehat{\mathbf{S}}_e | nj \rangle. \end{aligned}$$

13.1.4 Matrix Elements of Operators Which Depend on Variables of Two Subsystems

If a quantum-mechanical system consists of two subsystems, 1 and 2, and eigenstates of these subsystems are characterized by the sets of quantum numbers $(n_1 j_1 m_1)$ and $(n_2 j_2 m_2)$, it is often convenient to introduce the two following representations of matrix elements.

(a) The angular momenta of both subsystems are not coupled. Then

$$\langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \widehat{\mathbf{P}}(1, 2) | n_1 j_1 m_1; n_2 j_2 m_2 \rangle = \int \Psi_{n'_1 j'_1 m'_1}^*(1) \Psi_{n'_2 j'_2 m'_2}^*(2) \widehat{\mathbf{P}}(1, 2) \Psi_{n_1 j_1 m_1}(1) \Psi_{n_2 j_2 m_2}(2) d\tau_1 d\tau_2, \quad (13.1.20)$$

where $\Psi_{n_1 j_1 m_1}$ and $\Psi_{n_2 j_2 m_2}$ are wave functions that describe the eigenstates of each subsystem. Such a representation of matrix elements is called the $(n_1 j_1 m_1; n_2 j_2 m_2)$ -representation.

(b) If angular momenta $\mathbf{j}_1$ and $\mathbf{j}_2$ are coupled to the resultant angular momentum $\mathbf{j}$ with projection $m = m_1 + m_2$, then

$$\langle n'_1 j'_1 n'_2 j'_2 j' m' | \widehat{\mathbf{P}}(1, 2) | n_1 j_1 n_2 j_2 j m \rangle = \int \Psi_{n'_1 j'_1 n'_2 j'_2 j' m'}^*(1, 2) \widehat{\mathbf{P}}(1, 2) \Psi_{n_1 j_1 n_2 j_2 j m}(1, 2) d\tau_1 d\tau_2. \quad (13.1.21)$$

Here

$$\Psi_{n_1 j_1 n_2 j_2 j m}(1, 2) = \sum_{m_1 m_2} C_{j_1 m_1 j_2 m_2}^{j m} \Psi_{n_1 j_1 m_1}(1) \Psi_{n_2 j_2 m_2}(2). \quad (13.1.22)$$

This representation of matrix elements is called the $(n_1 j_1 n_2 j_2 j m)$ -representation.

In general case, we have

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \hat{\mathbf{P}}_{a\alpha}(1, 2) | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= (-1)^{2a} \sum_{j m j' m'} \frac{1}{\Pi_{j'}} C_{j_1 m_1 j_2 m_2}^{j m} C_{j'_1 m'_1 j'_2 m'_2}^{j' m'} C_{j m a \alpha}^{j' m'} \langle n'_1 j'_1 n'_2 j'_2 j' | \hat{\mathbf{P}}_a(1, 2) | n_1 j_1 n_2 j_2 j \rangle. \end{aligned} \quad (13.1.23)$$

Any operator $\hat{A}(1, 2)$ which depends on variables of two subsystems can frequently be expressed as a direct or irreducible product of two operators $\hat{P}_{a\alpha}(1)$ and $\hat{Q}_{b\beta}(2)$ which depend only on variables of the first and second subsystem, respectively:

$$\hat{A}(1, 2) = \hat{P}_{a\alpha}(1) \hat{Q}_{b\beta}(2) \quad \text{or} \quad \hat{A}(1, 2) = \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_{c\gamma}.$$

The matrix element for the direct-product case is given by

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \hat{P}_{a\alpha}(1) \hat{Q}_{b\beta}(2) | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= \frac{(-1)^{2a+2b}}{\Pi_{j'_1 j'_2}} C_{j_1 m_1 a \alpha}^{j'_1 m'_1} C_{j_2 m_2 b \beta}^{j'_2 m'_2} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \end{aligned} \quad (13.1.24)$$

whereas for the irreducible tensor product case it may be expressed in the form

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_{c\gamma} | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= \frac{(-1)^{2c}}{\Pi_{j'_1 j'_2}} \sum_{\alpha\beta} C_{j_1 m_1 a \alpha}^{j'_1 m'_1} C_{a \alpha b \beta}^{c \gamma} C_{j_2 m_2 b \beta}^{j'_2 m'_2} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle. \end{aligned} \quad (13.1.25)$$

The matrix element of a scalar product is given as follows:

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | (\hat{\mathbf{P}}_a(1) \cdot \hat{\mathbf{Q}}_a(2)) | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= \frac{1}{\Pi_{j'_1 j'_2}} \sum_{\alpha} (-1)^{-\alpha} C_{j_1 m_1 a \alpha}^{j'_1 m'_1} C_{j_2 m_2 a -\alpha}^{j'_2 m'_2} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_a(2) | n_2 j_2 \rangle. \end{aligned} \quad (13.1.26)$$

Matrix elements of the same operators can be evaluated in the $n_1 j_1 n_2 j_2 j m$ representation. This yields

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | \hat{P}_{a\alpha}(1) \hat{Q}_{b\beta}(2) | n_1 j_1 n_2 j_2 j m \rangle \\ &= (-1)^{2a+2b} \Pi_j \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle \sum_{c\gamma} \Pi_c C_{j m c \gamma}^{j' m'} C_{a \alpha b \beta}^{c \gamma} \left\{ \begin{matrix} a & b & c \\ j'_1 & j'_2 & j' \\ j_1 & j_2 & j \end{matrix} \right\}, \end{aligned} \quad (13.1.27)$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_{c\gamma} | n_1 j_1 n_2 j_2 j m \rangle \\ &= (-1)^{2c} \Pi_{c\gamma} C_{j m c \gamma}^{j' m'} \left\{ \begin{matrix} a & b & c \\ j'_1 & j'_2 & j' \\ j_1 & j_2 & j \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \end{aligned} \quad (13.1.28)$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c | n_1 j_1 n_2 j_2 j \rangle = \Pi_{c j j'} \left\{ \begin{matrix} a & b & c \\ j'_1 & j'_2 & j' \\ j_1 & j_2 & j \end{matrix} \right\} \\ & \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\hat{\mathbf{P}}_a(1) \cdot \hat{\mathbf{Q}}_a(2)) | n_1 j_1 n_2 j_2 j m \rangle \\ &= \delta_{j' j} \delta_{m' m} (-1)^{j+j_1+j'_2} \left\{ \begin{matrix} j'_1 & j_1 & a \\ j_2 & j'_2 & j \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_a(2) | n_2 j_2 \rangle. \end{aligned} \quad (13.1.29)$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' | (\hat{\mathbf{P}}_a(1) \cdot \hat{\mathbf{Q}}_a(2)) | n_1 j_1 n_2 j_2 j \rangle = \delta_{j' j} (-1)^{j+j_1+j'_2} \Pi_j \left\{ \begin{matrix} j'_1 & j_1 & a \\ j_2 & j'_2 & j \end{matrix} \right\} \\ & \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_a(2) | n_2 j_2 \rangle. \end{aligned}$$

Let us consider the products of three operators which depend only on variables of one of the subsystems. In this case the matrix elements of direct products of these operators can be evaluated by the successive use of Eqs. (13.1.24), (13.1.27) and (13.1.8). If two of three operators form an irreducible tensor product, Eqs. (13.1.25), (13.1.28), (13.1.9) and (13.1.10) must additionally be applied.

Matrix elements of irreducible tensor products of three tensor operators may be represented in the forms

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_{e\epsilon} | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= (-1)^{b-c-d-j_1-j'_1} \frac{\Pi_c}{\Pi_{j'_1 j'_2}} \sum_{fNJ} \Pi_f^2 C_{f\varphi b\beta}^{e\epsilon} C_{j_1 m_1 f\varphi}^{j'_1 m'_1} C_{j_2 m_2 b\beta}^{j'_2 m'_2} \left\{ \begin{matrix} a & b & c \\ e & d & f \end{matrix} \right\} \left\{ \begin{matrix} d & a & f \\ j'_1 & j_1 & J \end{matrix} \right\} \\ & \quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \quad (13.1.30) \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_{e\epsilon} | n_1 j_1 n_2 j_2 j m \rangle = (-1)^{b-c-d-j_1-j'_1} \Pi_{cje} C_{j m e\epsilon}^{j' m'} \sum_{fNJ} \Pi_f^2 \\ & \quad \left\{ \begin{matrix} a & b & c \\ e & d & f \end{matrix} \right\} \left\{ \begin{matrix} d & a & f \\ j'_1 & j_1 & J \end{matrix} \right\} \left\{ \begin{matrix} j_1 & f & j'_1 \\ j_2 & b & j'_2 \\ j & e & j' \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \quad (13.1.31) \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_e | n_1 j_1 n_2 j_2 j \rangle = (-1)^{b-c-d-j_1-j'_1} \Pi_{cjej'} \sum_{fNJ} \Pi_f^2 \\ & \quad \left\{ \begin{matrix} a & b & c \\ e & d & f \end{matrix} \right\} \left\{ \begin{matrix} d & a & f \\ j'_1 & j_1 & J \end{matrix} \right\} \left\{ \begin{matrix} j_1 & f & j'_1 \\ j_2 & b & j'_2 \\ j & e & j' \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle, \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \cdot \hat{\mathbf{R}}_c(1)) | n_1 j_1 n_2 j_2 j m \rangle \\ &= \delta_{j'j} \delta_{m'm} (-1)^{-a+b+j+j'_2-j'_1} \Pi_c \left\{ \begin{matrix} j'_1 & j_1 & b \\ j_2 & j'_2 & j \end{matrix} \right\} \sum_{NJ} \left\{ \begin{matrix} a & b & c \\ j_1 & J & j'_1 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{R}}_c(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle. \quad (13.1.32) \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \cdot \hat{\mathbf{R}}_c(1)) | n_1 j_1 n_2 j_2 j \rangle = \delta_{j'j} (-1)^{-a+b+j+j'_2-j'_1} \Pi_{cj} \left\{ \begin{matrix} j'_1 & j_1 & b \\ j_2 & j'_2 & j \end{matrix} \right\} \sum_{NJ} \left\{ \begin{matrix} a & b & c \\ j_1 & J & j'_1 \end{matrix} \right\} \\ & \quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{R}}_c(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | n_2 j_2 \rangle. \end{aligned}$$

The evaluation of matrix elements of the product $\{ \hat{\mathbf{P}}_a(1) \otimes \{ \hat{\mathbf{Q}}_b(2) \otimes \hat{\mathbf{R}}_d(1) \}_f \}_k$ can be performed using Eq. (3.3.1), while, for the product $\{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{R}}_d(1) \}_f \otimes \hat{\mathbf{Q}}_b(2) \}_k$, one can use Eqs. (13.1.25), (13.1.28), (13.1.10).

Matrix elements of direct products of four tensor operators can be obtained by the successive use of the foregoing relations.

We give below expressions for matrix elements of irreducible tensor products of four tensor operators:

$$\begin{aligned} & \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \otimes \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(2) \}_f \}_{k\kappa} | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\ &= (-1)^{2j'_1+2j'_2+2k} \frac{\Pi_{cj}}{\Pi_{j'_1 j'_2}} \sum_{gh\sigma\eta} \Pi_{gh}^2 C_{h\sigma g\eta}^{k\kappa} C_{h\sigma j_1 m_1}^{j'_1 m'_1} C_{g\eta j_2 m_2}^{j'_2 m'_2} \left\{ \begin{matrix} a & b & c \\ d & e & f \\ h & g & k \end{matrix} \right\} \sum_{N_1 J_1 N_2 J_2} \left\{ \begin{matrix} a & d & h \\ j_1 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} b & e & g \\ j_2 & j'_2 & J_2 \end{matrix} \right\} \\ & \quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \quad (13.1.33) \end{aligned}$$

$$\begin{aligned} & \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(2) \}_c \otimes \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(2) \}_f \}_{k\kappa} | n_1 j_1 n_2 j_2 j m \rangle \\ &= (-1)^{j+j'-k} \Pi_{cfjk} C_{j m k\kappa}^{j' m'} \sum_{gh} \Pi_{gh}^2 \left\{ \begin{matrix} a & b & c \\ d & e & f \\ h & g & k \end{matrix} \right\} \left\{ \begin{matrix} j_1 & j'_1 & h \\ j_2 & j'_2 & g \\ j & j' & k \end{matrix} \right\} \sum_{N_1 J_1 N_2 J_2} \left\{ \begin{matrix} d & a & h \\ j'_1 & j_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} e & b & g \\ j'_2 & j_2 & J_2 \end{matrix} \right\} \\ & \quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \hat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \quad (13.1.34) \end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \otimes \{ \widehat{\mathbf{R}}_d(1) \otimes \widehat{\mathbf{S}}_e(2) \}_f \}_k | n_1 j_1 n_2 j_2 j \rangle \\
&= (-1)^{j+j'-k} \Pi_{cfjkj'} \sum_{gh} \Pi_{gh}^2 \begin{Bmatrix} a & b & c \\ d & e & f \\ h & g & k \end{Bmatrix} \begin{Bmatrix} j_1 & j'_1 & h \\ j_2 & j'_2 & g \\ j & j' & k \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & h \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \\
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \cdot \{ \widehat{\mathbf{R}}_d(1) \otimes \widehat{\mathbf{S}}_e(2) \}_c) | n_1 j_1 n_2 j_2 j m \rangle = \delta_{j'j} \delta_{m'm} (-1)^{2c+j'_1+j_2-j-b-d} \Pi_c^2 \\
&\quad \times \sum_g \Pi_g^2 \begin{Bmatrix} a & b & c \\ e & d & g \end{Bmatrix} \begin{Bmatrix} j'_1 & j_1 & g \\ j_2 & j'_2 & j \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & g \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \quad (13.1.35)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \cdot \{ \widehat{\mathbf{R}}_d(1) \otimes \widehat{\mathbf{S}}_e(2) \}_c) | n_1 j_1 n_2 j_2 j \rangle = \delta_{j'j} (-1)^{2c+j'_1+j_2-j-b-d} \Pi_j \Pi_c^2 \\
&\quad \times \sum_g \Pi_g^2 \begin{Bmatrix} a & b & c \\ e & d & g \end{Bmatrix} \begin{Bmatrix} j'_1 & j_1 & g \\ j_2 & j'_2 & j \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & g \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle,
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \{ \{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \widehat{\mathbf{R}}_d(1) \}_f \widehat{\mathbf{S}}_e(2) \}_{k\kappa} | n_1 j_1 m_1; n_2 j_2 m_2 \rangle \\
&= (-1)^{c+d+k-e+2j'_1+2j'_2} \frac{\Pi_{cf}}{\Pi_{j'_1 j'_2}} \sum_{gh\sigma\eta} \Pi_{gh}^2 C_{h\sigma g\eta}^{k\kappa} C_{h\sigma j_1 m_1}^{j'_1 m'_1} C_{g\eta j_2 m_2}^{j'_2 m'_2} \begin{Bmatrix} a & d & h \\ f & b & c \end{Bmatrix} \begin{Bmatrix} b & e & g \\ k & h & f \end{Bmatrix} \\
&\quad \times \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & h \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \\
&\quad \times \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \quad (13.1.36)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \widehat{\mathbf{R}}_d(1) \}_f \widehat{\mathbf{S}}_e(2) \}_{k\kappa} | n_1 j_1 n_2 j_2 j m \rangle \\
&= (-1)^{c+d-e-j-j'} \Pi_{cfjk} C_{j m k \kappa}^{j'_1 m'_1} \sum_{gh} \Pi_{gh}^2 \begin{Bmatrix} a & d & h \\ f & b & c \end{Bmatrix} \begin{Bmatrix} b & e & g \\ k & h & f \end{Bmatrix} \begin{Bmatrix} j_1 & j'_1 & h \\ j_2 & j'_2 & g \\ j & j' & k \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & h \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \\
&\quad \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle, \quad (13.1.37)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \widehat{\mathbf{R}}_d(1) \}_f \widehat{\mathbf{S}}_e(2) \}_k | n_1 j_1 n_2 j_2 j \rangle \\
&= (-1)^{c+d-e-j-j'} \Pi_{cfjkj'} \sum_{gh} \Pi_{gh}^2 \begin{Bmatrix} a & d & h \\ f & b & c \end{Bmatrix} \begin{Bmatrix} b & e & g \\ k & h & f \end{Bmatrix} \begin{Bmatrix} j_1 & j'_1 & h \\ j_2 & j'_2 & g \\ j & j' & k \end{Bmatrix} \\
&\quad \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} d & a & h \\ j'_1 & j_1 & J_1 \end{Bmatrix} \begin{Bmatrix} e & b & g \\ j'_2 & j_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_e(2) | n_2 j_2 \rangle,
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \widehat{\mathbf{R}}_d(1) \}_f \cdot \widehat{\mathbf{S}}_f(2)) | n_1 j_1 n_2 j_2 j m \rangle \\
&= \delta_{j'j} \delta_{m'm} (-1)^{c+d-b+f-j+j'_1+j_2} \Pi_{cf} \sum_g \Pi_g^2 \begin{Bmatrix} a & b & c \\ f & d & g \end{Bmatrix} \begin{Bmatrix} j'_1 & j_1 & g \\ j_2 & j'_2 & j \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} a & d & g \\ j_1 & j'_1 & J_1 \end{Bmatrix} \begin{Bmatrix} b & f & g \\ j_2 & j'_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_f(2) | n_2 j_2 \rangle. \quad (13.1.38)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \{ \widehat{\mathbf{P}}_a(1) \otimes \widehat{\mathbf{Q}}_b(2) \}_c \widehat{\mathbf{R}}_d(1) \}_f \cdot \widehat{\mathbf{S}}_f(2)) | n_1 j_1 n_2 j_2 j \rangle \\
&= \delta_{j'j} (-1)^{c+d-b+f-j+j'_1+j_2} \Pi_{cfj} \sum_g \Pi_g^2 \begin{Bmatrix} a & b & c \\ f & d & g \end{Bmatrix} \begin{Bmatrix} j'_1 & j_1 & g \\ j_2 & j'_2 & j \end{Bmatrix} \sum_{N_1 J_1 N_2 J_2} \begin{Bmatrix} a & d & g \\ j_1 & j'_1 & J_1 \end{Bmatrix} \begin{Bmatrix} b & f & g \\ j_2 & j'_2 & J_2 \end{Bmatrix} \\
&\quad \times \langle n'_1 j'_1 | \widehat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \widehat{\mathbf{R}}_d(1) | n_1 j_1 \rangle \langle n'_2 j'_2 | \widehat{\mathbf{Q}}_b(2) | N_2 J_2 \rangle \langle N_2 J_2 | \widehat{\mathbf{S}}_f(2) | n_2 j_2 \rangle.
\end{aligned}$$

Matrix elements of irreducible tensor products with other coupling schemes of operators may be evaluated using the formulas for the recoupling of four irreducible tensors (see Eq. (3.3.3)-(3.3.5)).

If the wave functions and operators depend on variables of three or more subsystems, the evaluation of matrix elements is reduced to the successive use of the above equations.

13.1.5 Matrix Elements of Operators Which Depend on Variables of One of the Subsystems

Let $\hat{P}_{a\alpha}(1)$ be an operator which depends on variables of subsystem 1. Matrix elements of this operator are diagonal with respect to quantum numbers which characterize eigenstates of subsystem 2.

$$\langle n'_1 j'_1 m'_1; n'_2 j'_2 m'_2 | \hat{P}_{a\alpha}(1) | n_1 j_1 m_1; n_2 j_2 m_2 \rangle = \delta_{n'_2 n_2} \delta_{j'_2 j_2} \delta_{m'_2 m_2} \langle n'_1 j'_1 m'_1 | \hat{P}_{a\alpha}(1) | n_1 j_1 m_1 \rangle. \quad (13.1.39)$$

Further evaluation is reduced to the use of the foregoing relations.

If an eigenstate corresponds to some fixed resultant angular momentum $\mathbf{j} = \mathbf{j}_1 + \mathbf{j}_2$ and its projection $m = m_1 + m_2$, one has

$$\langle n'_1 j'_1 n'_2 j'_2 j' m' | \hat{P}_{a\alpha}(1) | n_1 j_1 n_2 j_2 j m \rangle = \delta_{j'_2 j_2} \delta_{n'_2 n_2} (-1)^{j+j'_1+j_2-a} \Pi_j C_{j m a \alpha}^{j'_1 m' j} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & a & j_1 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle. \quad (13.1.40)$$

$$\langle n'_1 j'_1 n'_2 j'_2 j' | \hat{\mathbf{P}}_a(1) | n_1 j_1 n_2 j_2 j \rangle = \delta_{j'_2 j_2} \delta_{n'_2 n_2} (-1)^{j+j'_1+j_2-a} \Pi_{jj'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & a & j_1 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | n_1 j_1 \rangle.$$

A matrix element of the direct tensor product of two operators may be represented in the form

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' m' | \hat{P}_{a\alpha}(1) \hat{Q}_{b\beta}(1) | n_1 j_1 n_2 j_2 j m \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j_2+j-j_1+2a+2b} \Pi_j \sum_{c\gamma} \Pi_c C_{j m c \gamma}^{j'_1 m' j} C_{a \alpha b \beta}^{c \gamma} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & c & j_1 \end{matrix} \right\} \\ &\times \sum_{NJ} \left\{ \begin{matrix} a & b & c \\ j_1 & j'_1 & J \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{Q}}_b(1) | n_1 j_1 \rangle. \end{aligned} \quad (13.1.41)$$

For an irreducible tensor product one gets the expressions

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_{c\gamma} | n_1 j_1 n_2 j_2 j m \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j_2+j-j_1+2c} \Pi_{jc} C_{j m c \gamma}^{j'_1 m' j} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & c & j_1 \end{matrix} \right\} \\ &\times \sum_{NJ} \left\{ \begin{matrix} a & b & c \\ j_1 & j'_1 & J \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{Q}}_b(1) | n_1 j_1 \rangle, \end{aligned} \quad (13.1.42)$$

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c | n_1 j_1 n_2 j_2 j \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j_2+j-j_1} \Pi_{jcj'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & c & j_1 \end{matrix} \right\} \\ &\times \sum_{NJ} \left\{ \begin{matrix} a & b & c \\ j_1 & j'_1 & J \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{Q}}_b(1) | n_1 j_1 \rangle, \end{aligned}$$

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\hat{\mathbf{P}}_a(1) \cdot \hat{\mathbf{Q}}_a(1)) | n_1 j_1 n_2 j_2 j m \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} \delta_{j'_1 j_1} \delta_{j' j} \delta_{m' m} \\ &\times \frac{1}{\Pi_{j_1}^2} (-1)^{-j_1} \sum_{JN} (-1)^J \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{Q}}_a(1) | n_1 j_1 \rangle. \end{aligned} \quad (13.1.43)$$

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' | (\hat{\mathbf{P}}_a(1) \cdot \hat{\mathbf{Q}}_a(1)) | n_1 j_1 n_2 j_2 j \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} \delta_{j'_1 j_1} \delta_{j' j} \frac{\Pi_j}{\Pi_{j_1}^2} (-1)^{-j_1} \\ &\times \sum_{JN} (-1)^J \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | NJ \rangle \langle NJ | \hat{\mathbf{Q}}_a(1) | n_1 j_1 \rangle. \end{aligned}$$

In the case of the irreducible tensor product of three operators we obtain

$$\begin{aligned} \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_{f\varphi} | n_1 j_1 n_2 j_2 j m \rangle &= \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j+j_1-j'_1+j_2+c} \Pi_{jfc} C_{j m f \varphi}^{j'_1 m' j} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & f & j_1 \end{matrix} \right\} \times \sum_{N_1 J_1 N_2 J_2} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j'_1 & j_1 & J_2 \end{matrix} \right\} \\ &\times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle, \end{aligned} \quad (13.1.44)$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_f | n_1 j_1 n_2 j_2 j \rangle = \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j+j_1-j'_1+j_2+c} \Pi_{jfcj'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & f & j'_1 \end{matrix} \right\} \\
& \times \sum_{N_1 J_1 N_2 J_2} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j'_1 & j_1 & J_2 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | n_1 j_1 \rangle, \\
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \cdot \hat{\mathbf{R}}_c(1)) | n_1 j_1 n_2 j_2 j m \rangle = \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} \delta_{m' m} (-1)^{-c+2j_1} \\
& \times \frac{\Pi_c}{\Pi_{j_1}^2} \sum_{\substack{N_1 N_2 \\ J_1 J_2}} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_c(1) | n_1 j_1 \rangle. \quad (13.1.45)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \cdot \hat{\mathbf{R}}_c(1)) | n_1 j_1 n_2 j_2 j \rangle = \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} (-1)^{-c+2j_1} \frac{\Pi_{cj}}{\Pi_{j_1}^2} \\
& \times \sum_{\substack{N_1 N_2 \\ J_1 J_2}} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_c(1) | n_1 j_1 \rangle.
\end{aligned}$$

The matrix elements of irreducible tensor products of four operators may be evaluated using the expressions given below:

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(1) \}_f \}_{k\kappa} | n_1 j_1 n_2 j_2 j m \rangle \\
& = \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j+j'_1+j_2-c-f} \Pi_{cfkj} C_{j m k \kappa}^{j' m'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & k & j'_1 \end{matrix} \right\} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{2J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} e & d & f \\ J_2 & j_1 & J_3 \end{matrix} \right\} \left\{ \begin{matrix} f & c & k \\ j'_1 & j_1 & J_2 \end{matrix} \right\} \\
& \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle, \quad (13.1.46)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(1) \}_f \}_{k\kappa} | n_1 j_1 n_2 j_2 j \rangle \\
& = \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j+j'_1+j_2-c-f} \Pi_{cfkj} C_{j m k \kappa}^{j' m'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & k & j'_1 \end{matrix} \right\} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{2J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} e & d & f \\ J_2 & j_1 & J_3 \end{matrix} \right\} \left\{ \begin{matrix} f & c & k \\ j'_1 & j_1 & J_2 \end{matrix} \right\} \\
& \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle,
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \cdot \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(1) \}_c) | n_1 j_1 n_2 j_2 j m \rangle \\
& = \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} \delta_{m' m} \frac{(-1)^{-j_1} \Pi_j}{\Pi_{j_1}^2} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} e & d & c \\ J_2 & j_1 & J_3 \end{matrix} \right\} \\
& \times \Pi_c^2 \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle, \quad (13.1.47)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \cdot \{ \hat{\mathbf{R}}_d(1) \otimes \hat{\mathbf{S}}_e(1) \}_c) | n_1 j_1 n_2 j_2 j \rangle \\
& = \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} \frac{(-1)^{-j_1} \Pi_j}{\Pi_{j_1}^2} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} e & d & c \\ J_2 & j_1 & J_3 \end{matrix} \right\} \\
& \times \Pi_c^2 \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle,
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | \{ \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_f \otimes \hat{\mathbf{S}}_e(1) \}_{k\kappa} | n_1 j_1 n_2 j_2 j m \rangle \\
& = \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j_1+j_2+j+j'+c} \Pi_{fcjk} C_{j m k \kappa}^{j' m'} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & k & j'_1 \end{matrix} \right\} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2+J_3} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j'_1 & J_3 & J_2 \end{matrix} \right\} \left\{ \begin{matrix} e & f & k \\ j'_1 & j_1 & J_3 \end{matrix} \right\} \\
& \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle, \quad (13.1.48)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | \{ \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_f \otimes \hat{\mathbf{S}}_e(1) \}_k | n_1 j_1 n_2 j_2 j \rangle \\
&= \delta_{n'_2 n_2} \delta_{j'_2 j_2} (-1)^{j_1 + j_2 + j + j' + c} \Pi_{fckj'j} \left\{ \begin{matrix} j_1 & j_2 & j \\ j' & k & j_1' \end{matrix} \right\} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2 + J_3} \left\{ \begin{matrix} a & b & c \\ J_2 & j'_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j'_1 & J_3 & J_2 \end{matrix} \right\} \left\{ \begin{matrix} e & f & k \\ j'_1 & j_1 & J_3 \end{matrix} \right\} \\
&\quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_e(1) | n_1 j_1 \rangle, \\
& \langle n'_1 j'_1 n'_2 j'_2 j' m' | (\{ \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_f \cdot \hat{\mathbf{S}}_f(1)) | n_1 j_1 n_2 j_2 j m \rangle \\
&= \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} \delta_{m' m} (-1)^{c-f-j_1} \frac{\Pi_{fjc}}{\Pi_{j_1}} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j_1 & J_3 & J_2 \end{matrix} \right\} \\
&\quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_f(1) | n_1 j_1 \rangle. \quad (13.1.49)
\end{aligned}$$

$$\begin{aligned}
& \langle n'_1 j'_1 n'_2 j'_2 j' | (\{ \{ \{ \hat{\mathbf{P}}_a(1) \otimes \hat{\mathbf{Q}}_b(1) \}_c \otimes \hat{\mathbf{R}}_d(1) \}_f \cdot \hat{\mathbf{S}}_f(1)) | n_1 j_1 n_2 j_2 j \rangle \\
&= \delta_{n'_2 n_2} \delta_{j'_1 j_1} \delta_{j'_2 j_2} \delta_{j' j} (-1)^{c-f-j_1} \frac{\Pi_{fcj}}{\Pi_{j_1}} \sum_{\substack{N_1 N_2 N_3 \\ J_1 J_2 J_3}} (-1)^{J_2} \left\{ \begin{matrix} a & b & c \\ J_2 & j_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} d & c & f \\ j_1 & J_3 & J_2 \end{matrix} \right\} \\
&\quad \times \langle n'_1 j'_1 | \hat{\mathbf{P}}_a(1) | N_1 J_1 \rangle \langle N_1 J_1 | \hat{\mathbf{Q}}_b(1) | N_2 J_2 \rangle \langle N_2 J_2 | \hat{\mathbf{R}}_d(1) | N_3 J_3 \rangle \langle N_3 J_3 | \hat{\mathbf{S}}_f(1) | n_1 j_1 \rangle.
\end{aligned}$$

Matrix elements for tensor products with other coupling schemes of operators may be obtained from the foregoing equations by the use of the recoupling rules and the commutation relations for irreducible tensors (Chap. 3).

13.2 Matrix elements of basic tensor operators

13.2.1 Some Introductory Remarks

In this section we shall consider matrix elements of the following operators and their tensor products:

- (a) the unit operator $\hat{\mathbf{I}}$;
- (b) the unit vector operator $\hat{\mathbf{n}} \equiv \hat{\mathbf{n}}_1$;
- (c) the gradient operators $\hat{\nabla} \equiv \hat{\nabla}_1$ and $\hat{\nabla}_\Omega \equiv (\hat{\nabla}_\Omega)_1$;
- (d) the total angular momentum operator $\hat{\mathbf{J}} \equiv \hat{\mathbf{J}}_1$;
- (e) the orbital angular momentum operator $\hat{\mathbf{L}} \equiv \hat{\mathbf{L}}_1$;
- (f) the spin angular momentum operator $\hat{\mathbf{S}} \equiv \hat{\mathbf{S}}_1$;
- (g) the spherical harmonic operator $Y_{lm}(\vartheta, \varphi)$.

All matrix elements may be evaluated from the equations of Sec. 13.1.

A wave function which describes an eigenstate of some quantum system will generally depend on the position (r, ϑ, φ) as well as on the spin variables (ξ) . It may be defined by

$$\Phi(r, \vartheta, \varphi; \xi) = \Psi_n(r) Y_{lm}(\vartheta, \varphi) \chi_{sm_s}(\xi) = \langle r, \vartheta, \varphi; \xi | n, l, m; sm_s \rangle. \quad (13.2.1)$$

where l and m are the orbital momentum and its projection; s and m_s are the spin momentum and its projection. The symbol n has been defined previously in Sec. 13.1.1.

The wave functions (13.2.1) determine the $nlmsm_s$ -representation of operators.

If the angular momenta $\hat{\mathbf{L}}$ and $\hat{\mathbf{S}}$ are coupled into the total angular momentum $\hat{\mathbf{J}}$ with its projection M , the wave function which describes an eigenstate of the quantum system is defined as

$$\Phi(r, \vartheta, \varphi; \xi) = \Psi_n(r) \sum_{mm_s} C_{lmsm_s}^{JM} Y_{lm}(\vartheta, \varphi) \chi_{sm_s}(\xi) = \Psi_n(r) Y_{JM}^{ls}(\vartheta, \varphi) = \langle r, \vartheta, \varphi; \xi | n, l, s, J, M \rangle. \quad (13.2.2)$$

These wave functions determine the $nl sJM$ -representation of operators.

If an operator acts only on position variables, the evaluation of matrix elements in the $nlmsm_s$ -representation is reduced to calculating matrix elements in the nlm -representation.

$$\langle n'l'm'; s'm'_s | \widehat{\mathfrak{M}}_{k\kappa}(r, \vartheta, \varphi) | nlm; sm_s \rangle = \delta_{ss'} \delta_{m_s m'_s} \langle n'l'm' | \widehat{\mathfrak{M}}_{k\kappa}(r, \vartheta, \varphi) | nlm \rangle. \quad (13.2.3)$$

If an operator depends only on spin variables, we get a similar relationship

$$\langle n'l'm'; s'm'_s | \widehat{\mathfrak{N}}_{k\kappa}(\xi) | nlm; sm_s \rangle = \delta_{ll'} \delta_{mm'} \langle n'l'm' | \widehat{\mathfrak{N}}_{k\kappa}(\xi) | nsm_s \rangle. \quad (13.2.4)$$

Hereafter matrix elements of operators which depend only on position variables or only on spin variables will be evaluated in the nlm or the nsm_s representation, respectively.

To determine matrix elements of operators $\widehat{\mathfrak{M}}_{k\kappa}(r, \vartheta, \varphi)$ and $\widehat{\mathfrak{N}}_{k\kappa}(\xi)$ in the $nl sJM$ representation one should use the equations

$$\langle n'l's'J' | \widehat{\mathfrak{M}}_k | nl sJ \rangle = \delta_{ss'} (-1)^{J+l'+s+k} \Pi_{JJ'} \left\{ \begin{matrix} l & s & J \\ J' & k & l' \end{matrix} \right\} \langle n'l' | \mathfrak{M}_k | nl \rangle, \quad (13.2.5)$$

$$\langle n'l's'J' | \widehat{\mathfrak{N}}_k | nl sJ \rangle = \delta_{ll'} (-1)^{J'+l+s+k} \Pi_{JJ'} \left\{ \begin{matrix} s & l & J \\ J' & k & s' \end{matrix} \right\} \langle n's' | \mathfrak{N}_k | ns \rangle. \quad (13.2.6)$$

If some operator $\widehat{R}_{a\alpha}(r, \vartheta, \varphi; \xi)$ acts on position as well as on spin variables, its matrix elements will be given in the $nl sJM$ -representation. These matrix elements may be evaluated in the $nlmsm_s$ representation with aid of

$$\langle n'l'm'; s'm'_s | \widehat{R}_{a\alpha}(r, \vartheta, \varphi; \xi) | nlm; sm_s \rangle = \sum_{\substack{J'M' \\ J'M'}} C_{l'm's'm_s}^{JM} C_{l'm's'm'_s}^{J'M'} \langle n'l's'J'M' | \widehat{R}_{a\alpha}(r, \vartheta, \varphi; \xi) | nl sJM \rangle. \quad (13.2.7)$$

or from the corresponding equations of the section above.

From now on, we will omit the arguments of operators in the expressions for matrix elements to simplify the equations. In addition, the index n will also be dropped in all the cases for which it is not essential.

13.2.2 Matrix Elements of the Unit Operator $\widehat{\mathbf{I}}$

The unit operator $\widehat{\mathbf{I}}$ is an irreducible tensor operator of rank zero. It does not act on the variables $r, \vartheta, \varphi, \xi$. Hence, in any representation its diagonal matrix elements are determined by

$$\langle \lambda' | \widehat{\mathbf{I}} | \lambda \rangle = \langle \lambda' | \lambda \rangle = \delta_{\lambda\lambda'}. \quad (13.2.8)$$

All non-diagonal matrix elements vanish. A reduced matrix element of the operator $\widehat{\mathbf{I}}$ may be represented in the form

$$\langle l | \widehat{\mathbf{I}} | l \rangle = \Pi_l, \quad \langle s | \widehat{\mathbf{I}} | s \rangle = \Pi_s, \quad \langle l's'J | \widehat{\mathbf{I}} | lsJ \rangle = \Pi_J \delta_{ll'} \delta_{ss'}. \quad (13.2.9)$$

13.2.3 Matrix Elements of the Unit Vector $\widehat{\mathbf{n}}(\vartheta, \varphi) \equiv \widehat{\mathbf{n}}_1(\vartheta, \varphi)$

The unit vector operator $\widehat{\mathbf{n}}_1(\vartheta, \varphi)$ acts only on variables ϑ, φ . In the lm -representation matrix elements of spherical components of this operator are given by¹

$$\langle l'm' | \widehat{n}_{1\mu} | lm \rangle = \frac{\langle l' | \widehat{\mathbf{n}}_1 | l \rangle}{\sqrt{2l'+1}} C_{l'm'1\mu}^{l'0}, \quad (13.2.10)$$

where

$$\langle l' | \widehat{\mathbf{n}}_1 | l \rangle = \Pi_l C_{l'010}^{l'0}. \quad (13.2.11)$$

Taking into account Eq. (13.2.5), we obtain a reduced matrix element in the lsJ -representation

$$\langle l's'J' | \widehat{\mathbf{n}}_1 | lsJ \rangle = \delta_{ss'} (-1)^{s+J+l'+1} \Pi_{JJ'} \sqrt{(2J'+1)} C_{l'010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}. \quad (13.2.12)$$

¹the notation $\mathbf{n}$ inside a reduced matrix element is slightly confusing, but correct. The result is a scalar, not a vector!

The matrix element (13.2.10) vanishes unless $l' = l \pm 1$. Using Eq. (13.2.11) one can rewrite Eq. (13.2.10) in component form as

$$\begin{aligned}\langle l+1m'|\hat{n}_{1\pm 1}|lm\rangle &= \sqrt{\frac{(l\pm m+1)(l\pm m+2)}{2(2l+1)(2l+3)}}\delta_{m'm\pm 1}, \\ \langle l+1m'|\hat{n}_{10}|lm\rangle &= \sqrt{\frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)}}\delta_{m'm}, \\ \langle l-1m'|\hat{n}_{1\pm 1}|lm\rangle &= -\sqrt{\frac{(l\mp m-1)(l\mp m)}{2(2l+1)(2l-1)}}\delta_{m'm\pm 1}, \\ \langle l-1m'|\hat{n}_{10}|lm\rangle &= \sqrt{\frac{(l-m)(l+m)}{(2l+1)(2l-1)}}\delta_{m'm}.\end{aligned}\tag{13.2.13}$$

All other matrix elements of the spherical components of the operator in question are equal to zero. The matrix elements of the cartesian components of the operator $\hat{\mathbf{n}}_1(\vartheta, \varphi)$ are given by

$$\langle l'm \pm 1|\hat{n}_x|lm\rangle = \mp \frac{1}{2}\sqrt{\frac{(l\pm m+1)(l\pm m+2)}{(2l+1)(2l+3)}}\delta_{l'l+1} \pm \frac{1}{2}\sqrt{\frac{(l\mp m-1)(l\mp m)}{(2l+1)(2l-1)}}\delta_{l'l-1},\tag{13.2.14}$$

$$\langle l'm \pm 1|\hat{n}_y|lm\rangle = \frac{i}{2}\sqrt{\frac{(l\pm m+1)(l\pm m+2)}{(2l+1)(2l+3)}}\delta_{l'l+1} - \frac{i}{2}\sqrt{\frac{(l\mp m-1)(l\mp m)}{(2l+1)(2l-1)}}\delta_{l'l-1},\tag{13.2.15}$$

$$\langle l'm|\hat{n}_z|lm\rangle = \sqrt{\frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)}}\delta_{l'l+1} + \sqrt{\frac{(l-m)(l+m)}{(2l+1)(2l-1)}}\delta_{l'l-1}.\tag{13.2.16}$$

All other matrix elements of the cartesian components of $\hat{\mathbf{n}}_1(\vartheta, \varphi)$ vanish.

It is easy to obtain the matrix elements of direct products of two operators involving $\hat{\mathbf{n}}_1(\vartheta, \varphi)$:

$$\langle l'm'|\hat{n}_{1\mu}\hat{n}_{1\nu}|lm\rangle = \frac{(-1)^\mu}{3}\delta_{ll'}\delta_{mm'}\delta_{\mu,-\nu} + \sqrt{\frac{2}{3}}\frac{\Pi_l}{\Pi_{l'}}C_{l020}^{l'0}C_{1\mu1\nu}^{2\kappa}C_{lm2\kappa}^{l'm'}.\tag{13.2.17}$$

These matrix elements vanish unless $l' = l, l \pm 2$. To derive the matrix elements of direct products of cartesian components of $\hat{\mathbf{n}}_1(\vartheta, \varphi)$ one may express these products in terms of the products of spherical components. In this way we get

$$\begin{aligned}\langle l'm \pm 2|\hat{n}_x\hat{n}_x|lm\rangle &= \frac{1}{2}\langle l'm \pm 2|\hat{n}_{1\pm 1}\hat{n}_{1\pm 1}|lm\rangle, \\ \langle l'm|\hat{n}_x\hat{n}_x|lm\rangle &= -\langle l'm|\hat{n}_{1\pm 1}\hat{n}_{1\mp 1}|lm\rangle, \\ \langle l'm \pm 2|\hat{n}_y\hat{n}_y|lm\rangle &= -\langle l'm \pm 2|\hat{n}_{1\pm 1}\hat{n}_{1\pm 1}|lm\rangle, \\ \langle l'm|\hat{n}_y\hat{n}_y|lm\rangle &= -\langle l'm|\hat{n}_{1\pm 1}\hat{n}_{1\mp 1}|lm\rangle, \\ \langle l'm|\hat{n}_x\hat{n}_z|lm\rangle &= \langle l'm|\hat{n}_{10}\hat{n}_{10}|lm\rangle, \\ \langle l'm \pm 2|\hat{n}_x\hat{n}_y|lm\rangle &= \mp \frac{i}{2}\langle l'm \pm 2|\hat{n}_{1\pm 1}\hat{n}_{1\pm 1}|lm\rangle, \\ \langle l'm \pm 1|\hat{n}_x\hat{n}_z|lm\rangle &= \mp \frac{1}{\sqrt{2}}\langle l'm \pm 1|\hat{n}_{1\pm 1}\hat{n}_{10}|lm\rangle, \\ \langle l'm \pm 1|\hat{n}_y\hat{n}_z|lm\rangle &= \frac{i}{\sqrt{2}}\langle l'm \pm 1|\hat{n}_{1\pm 1}\hat{n}_{10}|lm\rangle.\end{aligned}\tag{13.2.18}$$

The matrix elements of the irreducible tensor product of k unit vector operators $\hat{\mathbf{n}}_1(\vartheta, \varphi)$ may be represented in the form

$$\begin{aligned}\langle l'm'|\{\dots\{\{\hat{\mathbf{n}}_1 \otimes \hat{\mathbf{n}}_1\}_{q_2} \otimes \hat{\mathbf{n}}_1\}_{q_3} \otimes \dots \otimes \hat{\mathbf{n}}_1\}_{q_k \kappa_k}|lm\rangle &= \frac{\Pi_l}{\Pi_{l'}}C_{l0q_k0}^{l'0} \prod_{i=2}^k C_{q_{i-1}010}^{q_i0} C_{lmq_k\kappa_k}^{l'm'}, \\ \langle l' || \{\dots\{\{\hat{\mathbf{n}}_1 \otimes \hat{\mathbf{n}}_1\}_{q_2} \otimes \hat{\mathbf{n}}_1\}_{q_3} \otimes \dots \otimes \hat{\mathbf{n}}_1\}_{q_k} || l\rangle &= \Pi_l C_{l0q_k0}^{l'0} \prod_{i=2}^k C_{q_{i-1}010}^{q_i0},\end{aligned}\tag{13.2.19}$$

where $q_1 = 1$. In particular, if $q_2 = 2, q_3 = 3, q_4 = 4, \dots$ one has

$$\langle l'm'|\{\dots\{\{\hat{\mathbf{n}}_1 \otimes \hat{\mathbf{n}}_1\}_2 \otimes \hat{\mathbf{n}}_1\}_3 \otimes \dots \otimes \hat{\mathbf{n}}_1\}_{k\kappa}|lm\rangle = \sqrt{\frac{k!}{(2k-1)!!}} \frac{\Pi_l}{\Pi_{l'}} C_{l0k0}^{l'0} C_{lmk\kappa}^{l'm'}.\tag{13.2.20}$$

$$\langle l' || \{ \dots \{ \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{n}}_1 \}_2 \otimes \hat{\mathbf{n}}_1 \}_3 \otimes \dots \otimes \hat{\mathbf{n}}_1 \}_k || l \rangle = \sqrt{\frac{k!}{(2k-1)!!}} \Pi_l C_{l0k}^{l'0}.$$

From Eq. (13.2.19) one obtains two obvious equalities,

$$\begin{aligned} \langle l'm' | (\hat{\mathbf{n}} \cdot \hat{\mathbf{n}}) | lm \rangle &= \delta_{ll'} \delta_{mm'}, \\ \langle l'm' | [\hat{\mathbf{n}} \times \hat{\mathbf{n}}] | lm \rangle &= 0. \end{aligned}$$

13.2.4 Matrix Elements of the Operator $\hat{\nabla}(r, \vartheta, \varphi) \equiv \hat{\nabla}_1(r, \vartheta, \varphi)$

The operator $\hat{\nabla}_1(r, \vartheta, \varphi)$ acts only on variables r, ϑ, φ . The matrix elements of its spherical components are determined by

$$\langle n'l'm' | \hat{\nabla}_{1\mu} | nlm \rangle = \frac{\langle n'l' || \hat{\nabla}_1 || nl \rangle}{\sqrt{2l'+1}} C_{l'm'1\mu}^{l'0}, \quad (13.2.21)$$

where

$$\langle n'l' || \hat{\nabla}_1 || nl \rangle = \sqrt{l+1} A_{n'l'nl} \delta_{l'l+1} - \sqrt{l} B_{n'l'nl} \delta_{l'l-1}. \quad (13.2.22)$$

The quantities $A_{n'l'nl}$ and $B_{n'l'nl}$ in Eq. (13.2.22) may be represented as the integrals

$$\begin{aligned} A_{n'l'nl} &= \int_0^\infty \Psi_{n'l'}^*(r) \left(\frac{\partial}{\partial r} - \frac{l}{r} \right) \Psi_{nl}(r) r^2 dr, \\ B_{n'l'nl} &= \int_0^\infty \Psi_{n'l'}^*(r) \left(\frac{\partial}{\partial r} + \frac{l+1}{r} \right) \Psi_{nl}(r) r^2 dr. \end{aligned} \quad (13.2.23)$$

Here $\Psi_{nl}(r)$ is the radial part of the corresponding wave function.

Reduced matrix elements of the operator $\hat{\nabla}_1(r, \vartheta, \varphi)$ in the $nl sJM$ representation may be obtained from Eq. (13.2.5) by replacing $\hat{\mathfrak{M}}_k \rightarrow \hat{\nabla}_1$.

The matrix elements (13.2.21) vanish unless $l' = l \pm 1$. In detailed form Eq. (13.2.21) reads

$$\begin{aligned} \langle n'l+1m' | \hat{\nabla}_{1\pm 1} | nlm \rangle &= \sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{2(2l+1)(2l+3)}} A_{n'l+1, nl} \delta_{m'm \pm 1}, \\ \langle n'l+1m' | \hat{\nabla}_{10} | nlm \rangle &= \sqrt{\frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)}} A_{n'l+1, nl} \delta_{m'm}, \\ \langle n'l-1m' | \hat{\nabla}_{1\pm 1} | nlm \rangle &= -\sqrt{\frac{(l \mp m - 1)(l \mp m)}{2(2l+1)(2l-1)}} B_{n'l-1, nl} \delta_{m'm \pm 1}, \\ \langle n'l-1m' | \hat{\nabla}_{10} | nlm \rangle &= \sqrt{\frac{(l-m)(l+m)}{(2l+1)(2l-1)}} B_{n'l-1, nl} \delta_{m'm}. \end{aligned} \quad (13.2.24)$$

All other matrix elements equal zero. The matrix elements of the cartesian components of $\hat{\nabla}_1(r, \vartheta, \varphi)$ are as follows:

$$\begin{aligned} \langle n'l'm \pm 1 | \hat{\nabla}_x | nlm \rangle &= \mp \frac{1}{2} \sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{(2l+1)(2l+3)}} A_{n'l+1, nl} \delta_{l'l+1}, \\ &\quad \pm \frac{1}{2} \sqrt{\frac{(l \mp m - 1)(l \mp m)}{(2l+1)(2l-1)}} B_{n'l-1, nl} \delta_{l'l-1}, \\ \langle n'l'm \pm 1 | \hat{\nabla}_y | nlm \rangle &= \frac{i}{2} \sqrt{\frac{(l \pm m + 1)(l \pm m + 2)}{(2l+1)(2l+3)}} A_{n'l+1, nl} \delta_{l'l+1}, \\ &\quad - \frac{i}{2} \sqrt{\frac{(l \mp m - 1)(l \mp m)}{(2l+1)(2l-1)}} B_{n'l-1, nl} \delta_{l'l-1}, \\ \langle n'l'm | \hat{\nabla}_z | nlm \rangle &= \sqrt{\frac{(l-m+1)(l+m+1)}{(2l+1)(2l+3)}} A_{n'l+1, nl} \delta_{l'l+1}, \\ &\quad + \sqrt{\frac{(l-m)(l+m)}{(2l+1)(2l-1)}} B_{n'l-1, nl} \delta_{l'l-1}. \end{aligned} \quad (13.2.25)$$

All other matrix elements are equal zero.

The matrix elements of the angular part of $\widehat{\nabla}(r, \vartheta, \varphi)$ may be presented in a simpler form. According to Eq. (1.3.10),

$$\widehat{\nabla}_\Omega = -i[\widehat{\mathbf{n}} \times \widehat{\mathbf{L}}].$$

This operator is independent of r . The use of the Wigner-Eckart theorem yields the following expression for the matrix elements of the spherical components of $(\widehat{\nabla}_\Omega)_1$,

$$\langle l'm' | (\widehat{\nabla}_\Omega)_{1\mu} | lm \rangle = \frac{\langle l' || (\widehat{\nabla}_\Omega)_1 || l \rangle}{\sqrt{2l'+1}} C_{l'm'1\mu}^{l'm'} \quad (13.2.26)$$

where

$$\langle l' || (\widehat{\nabla}_\Omega)_1 || l \rangle = -\{l\sqrt{l+1}\delta_{l'l+1} + (l+1)\sqrt{l}\delta_{l'l-1}\}. \quad (13.2.27)$$

Equation (13.2.27) may be obtained from (13.2.22) by the formal substitution

$$A_{n'l',nl} \rightarrow -l, \quad B_{n'l',nl} \rightarrow l+1. \quad (13.2.28)$$

Similarly, Eqs. (13.2.24) and (13.2.25) become valid for $(\widehat{\nabla}_\Omega)_1$ after the same substitution (13.2.28).

Let us consider the matrix elements of products of two operators $(\widehat{\nabla}_\Omega)_1$. In the case of the direct product we obtain

$$\begin{aligned} \langle l'm' | (\widehat{\nabla}_\Omega)_{1\mu} (\widehat{\nabla}_\Omega)_{1\nu} | lm \rangle &= \frac{(-1)^{1-\mu}}{3} l(l+1) \delta_{l'l} \delta_{m'm'} \delta_{\mu,-\nu} + C_{1\mu 1\nu}^{2\kappa} \left\{ \frac{l(l+1)\sqrt{(l+1)(l+2)}}{\sqrt{(2l+3)(2l+5)}} C_{l'm'2\kappa}^{l+2m+\kappa} \delta_{l'l+2} \right. \\ &\quad \left. + \frac{1}{2l+1} \sqrt{\frac{l(l+1)}{6(2l-1)(2l+3)}} [4l^3 + 6l^2 - 4l - 3] C_{l'm'2\kappa}^{l+m+\kappa} \delta_{l'l'} + \frac{l(l+1)\sqrt{l(l-1)}}{\sqrt{(2l-1)(2l-3)}} C_{l'm'2\kappa}^{l-2m+\kappa} \delta_{l'l-2} \right\}. \end{aligned} \quad (13.2.29)$$

The matrix elements of the product of cartesian components $(\widehat{\nabla}_\Omega)_i (\widehat{\nabla}_\Omega)_k$ ($i, k = x, y, z$) are related to the matrix elements (13.2.29) by means of Eqs. (13.2.18) where one should make the replacement $\widehat{n}_i \widehat{n}_k \rightarrow (\widehat{\nabla}_\Omega)_i (\widehat{\nabla}_\Omega)_k$ ($i, k = x, y, z$). In the case of the irreducible tensor product of two operators $(\widehat{\nabla}_\Omega)_1$ one has

$$\begin{aligned} \langle l'm' | \{ (\widehat{\nabla}_\Omega)_1 \otimes (\widehat{\nabla}_\Omega)_1 \}_{k\kappa} | lm \rangle \\ = (-1)^{k+l+l'} \Pi_k / \Pi_{l'} C_{l'm'k\kappa}^{l'm'} \sum_{l''} \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l'' \end{matrix} \right\} \langle l' || (\widehat{\nabla}_\Omega)_1 || l'' \rangle \langle l'' || (\widehat{\nabla}_\Omega)_1 || l \rangle. \end{aligned} \quad (13.2.30)$$

$$\langle l' || \{ (\widehat{\nabla}_\Omega)_1 \otimes (\widehat{\nabla}_\Omega)_1 \}_k || l \rangle = (-1)^{k+l+l'} \Pi_k \sum_{l''} \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l'' \end{matrix} \right\} \langle l' || (\widehat{\nabla}_\Omega)_1 || l'' \rangle \langle l'' || (\widehat{\nabla}_\Omega)_1 || l \rangle.$$

In particular, Eq. (13.2.30) yields the well known expression for the matrix elements of scalar product of two operators $(\widehat{\nabla}_\Omega)_1$

$$\langle l'm' | ((\widehat{\nabla}_\Omega)_1 \cdot (\widehat{\nabla}_\Omega)_1) | lm \rangle = -\delta_{l'l} \delta_{m'm} l(l+1).$$

Matrix elements of products of the operators $\widehat{\mathbf{n}}_1$ and $(\widehat{\nabla}_\Omega)_1$ are of interest in applications. For the matrix elements of the direct product of these operators one obtains

$$\begin{aligned} \langle l'm' | \widehat{n}_{1\mu} (\widehat{\nabla}_\Omega)_{1\nu} | lm \rangle &= -\delta_{l'l} \sqrt{\frac{l(l+1)}{2}} \left\{ C_{1\mu 1\nu}^{2\kappa} C_{l'm'1\kappa}^{l'm'} + \sqrt{\frac{3}{(2l-1)(2l+3)}} C_{1\mu 1\nu}^{2\kappa} C_{l'm'2\kappa}^{l'm'} \right\} \\ &\quad - \delta_{l'l+2} \sqrt{\frac{(l+1)(l+2)}{(2l+3)(2l+5)}} C_{1\mu 1\nu}^{2\kappa} C_{l'm'2\kappa}^{l'm'} + \delta_{l'l-2} (l+1) \sqrt{\frac{l(l-1)}{(2l-1)(2l-3)}} C_{1\mu 1\nu}^{2\kappa} C_{l'm'2\kappa}^{l'm'}. \end{aligned} \quad (13.2.31)$$

The matrix element of the direct product $(\widehat{\nabla}_\Omega)_{1\mu} \widehat{n}_{1\nu}$ differs from the foregoing one by the factor

$$\langle l'm' | \mathfrak{R}_{1\nu 1\mu} (\widehat{n}_{1\nu}, (\nabla_\Omega)_{1\mu}) | lm \rangle,$$

where $\mathfrak{R}_{1\nu 1\mu}$ is the commutator of the operator components in question (see Sec. 3.1.7)

$$\mathfrak{R}_{1\nu 1\mu} (\widehat{n}_{1\nu}, (\widehat{\nabla}_\Omega)_{1\mu}) = -(\widehat{\nabla}_\Omega)_{1\mu} \widehat{n}_{1\nu} = (-1)^{1+\nu} \frac{2}{3} \delta_{\nu,-\mu} + 2\sqrt{\frac{2\pi}{15}} C_{1\mu 1\nu}^{2\mu+\nu} Y_{2\mu+\nu}(\vartheta, \varphi) \quad (13.2.32)$$

and

$$\langle l'm' | \mathfrak{R}_{1\mu 1\nu}(\hat{n}_{1\nu}, (\hat{\nabla}\Omega)_{1\mu}) | lm \rangle = (-1)^{1+\nu} \frac{2}{3} \delta_{\nu, -\mu} \delta_{ll'} \delta_{mm'} + \sqrt{\frac{2}{3}} \Pi_l / \Pi_{l'} C_{l020}^{l'0} C_{1\mu 1\nu}^{2\mu+\nu} C_{lm 2\mu+\nu}^{l'm'} \quad (13.2.32a)$$

To determine the matrix elements of products of cartesian components of the operators $\hat{\mathbf{n}}_1$ and $(\hat{\nabla}\Omega)_1$ one can transform these products to the products of spherical components (see Eq. (1.2.11)) and use Eq. (13.2.31).

For the irreducible tensor product of $\hat{\mathbf{n}}_1(\vartheta, \varphi)$ and $(\hat{\nabla}\Omega)_1$ one obtains

$$\langle l'm' | \{ \hat{\mathbf{n}}_1 \otimes (\hat{\nabla}\Omega)_1 \}_{k\kappa} | lm \rangle = (-1)^{l'+l+k} C_{lm k \kappa}^{l' m'} \Pi_k / \Pi_{l'} \sum_{l''} \Pi_{l''} \left\{ \begin{matrix} 1 & l & l'' \\ l' & 1 & k \end{matrix} \right\} C_{l''010}^{l'0} \langle l'' | (\hat{\nabla}\Omega)_1 | l \rangle. \quad (13.2.33)$$

$$\langle l' | \{ \hat{\mathbf{n}}_1 \otimes (\hat{\nabla}\Omega)_1 \}_k | l \rangle = (-1)^{l'+l+k} \Pi_k \sum_{l''} \Pi_{l''} \left\{ \begin{matrix} 1 & l & l'' \\ l' & 1 & k \end{matrix} \right\} C_{l''010}^{l'0} \langle l'' | (\hat{\nabla}\Omega)_1 | l \rangle.$$

The reduced matrix element of $(\hat{\nabla}\Omega)_1$ in (13.2.33) is given by Eq. (13.2.27).

$$\langle l'm' | \{ (\hat{\nabla}\Omega)_1 \otimes \hat{\mathbf{n}}_1 \}_{k\kappa} | lm \rangle = (-1)^k \langle l'm' | \{ \hat{\mathbf{n}}_1 \otimes (\nabla\Omega)_1 \}_{k\kappa} | lm \rangle - (-1)^k \langle l'm' | \mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, (\hat{\nabla}\Omega)_1) | lm \rangle. \quad (13.2.34)$$

The second term in Eq. (13.2.34) is the matrix element of the commutator of the irreducible tensor product in question,

$$\mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, (\hat{\nabla}\Omega)_1) = \frac{2}{\sqrt{3}} \delta_{k0} + 2\sqrt{\frac{2\pi}{15}} Y_{2\kappa}(\vartheta, \varphi) \delta_{k2}. \quad (13.2.35)$$

Using Eqs. (13.2.104) and (13.2.105), we find

$$\langle l'm' | \mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, (\hat{\nabla}\Omega)_1) | lm \rangle = \frac{2}{\sqrt{3}} \delta_{k0} \delta_{ll'} \delta_{mm'} + \sqrt{\frac{2}{3}} \Pi_l / \Pi_{l'} C_{l020}^{l'0} C_{lm 2\kappa}^{l'm'} \delta_{k2}. \quad (13.2.36)$$

13.2.5 Matrix Elements of the Total Angular Momentum Operator $\hat{\mathbf{J}} \equiv \hat{\mathbf{J}}_1$

The operator of the total angular momentum $\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$ acts on position as well as spin variables. Making use of the Wigner-Eckart theorem, we get

$$\langle l's'J'M' | \hat{\mathbf{J}}_{1\mu} | lsJM \rangle = \frac{\langle l's'J' || \hat{\mathbf{J}}_1 || lsJ \rangle}{\sqrt{2J'+1}} C_{JM 1\mu}^{J'M'} \quad (13.2.37)$$

where

$$\langle l's'J' || \hat{\mathbf{J}}_1 || lsJ \rangle = \delta_{ll'} \delta_{ss'} \delta_{JJ'} \Pi_J \sqrt{J(J+1)}. \quad (13.2.38)$$

For simplicity in equations (13.2.41)-(13.2.47) we shall omit the quantum numbers l and s in notation for matrix elements. Thus,

$$\langle J'M' | \hat{\mathbf{J}}_{1\mu} | JM \rangle = \frac{\langle J' || \hat{\mathbf{J}}_1 || J \rangle}{\sqrt{2J'+1}} C_{JM 1\mu}^{J'M'} \quad (13.2.39)$$

where

$$\langle J' || \hat{\mathbf{J}}_1 || J \rangle = \delta_{JJ'} \Pi_J \sqrt{J(J+1)}. \quad (13.2.40)$$

In expanded form

$$\begin{aligned} \langle JM \pm 1 | \hat{J}_{1\pm 1} | JM \rangle &= \mp \sqrt{\frac{(J \pm M + 1)(J \mp M)}{2}}, \\ \langle JM | \hat{J}_{10} | JM \rangle &= M. \end{aligned} \quad (13.2.41)$$

All other matrix elements are equal to zero. Let us present the non-vanishing matrix elements of the cartesian components of $\hat{\mathbf{J}}$:

$$\begin{aligned} \langle JM \pm 1 | \hat{J}_x | JM \rangle &= \frac{1}{2} \sqrt{(J \pm M + 1)(J \mp M)}, \\ \langle JM \pm 1 | \hat{J}_y | JM \rangle &= \mp \frac{i}{2} \sqrt{(J \pm M + 1)(J \mp M)}, \\ \langle JM | \hat{J}_z | JM \rangle &= M. \end{aligned} \quad (13.2.42)$$

The matrix element of the operator $\hat{J}^2$ is as follows:

$$\langle J' M' | \hat{J}^2 | J M \rangle = \delta_{JJ'} \delta_{MM'} J(J+1). \quad (13.2.43)$$

Let us consider matrix elements of products of the $\hat{\mathbf{J}}_1$ components. Matrix elements of direct product of spherical components may be represented in the form

$$\langle J' M' | \hat{J}_{1\mu} \hat{J}_{1\nu} | J M \rangle = J(J+1) C_{J M' -\mu 1 \mu}^{J M'} C_{J M 1 \nu}^{J M'} \delta_{JJ'}. \quad (13.2.44)$$

In component form

$$\begin{aligned} \langle J M \pm 2 | \hat{J}_{1\pm 1} \hat{J}_{1\pm 1} | J M \rangle &= \frac{1}{2} \sqrt{(J \pm M + 2)(J \pm M + 1)(J \mp M - 1)(J \mp M)}, \\ \langle J M \pm 1 | \hat{J}_{1\pm 1} \hat{J}_{10} | J M \rangle &= \mp M \sqrt{\frac{(J \pm M + 1)(J \mp M)}{2}}, \\ \langle J M \pm 1 | \hat{J}_{10} \hat{J}_{1\pm 1} | J M \rangle &= \mp (M \pm 1) \sqrt{\frac{(J \pm M + 1)(J \mp M)}{2}}, \\ \langle J M | \hat{J}_{1\pm 1} \hat{J}_{1\mp 1} | J M \rangle &= -\frac{1}{2} (J \pm M)(J \mp M + 1), \\ \langle J M | \hat{J}_{10} \hat{J}_{10} | J M \rangle &= M^2. \end{aligned} \quad (13.2.45)$$

The non-vanishing matrix elements of product of the cartesian components are as follows:

$$\begin{aligned} \langle J M \pm 2 | \hat{J}_x \hat{J}_x | J M \rangle &= \frac{1}{4} \sqrt{(J \pm M + 2)(J \pm M + 1)(J \mp M - 1)(J \mp M)}, \\ \langle J M | \hat{J}_x \hat{J}_x | J M \rangle &= \frac{1}{2} [J(J+1) - M^2], \\ \langle J M \pm 2 | \hat{J}_y \hat{J}_y | J M \rangle &= -\frac{1}{4} \sqrt{(J \pm M + 2)(J \pm M + 1)(J \mp M - 1)(J \mp M)}, \\ \langle J M | \hat{J}_y \hat{J}_y | J M \rangle &= \frac{1}{2} [J(J+1) - M^2], \\ \langle J M | \hat{J}_z \hat{J}_z | J M \rangle &= M^2, \\ \langle J M \pm 2 | \hat{J}_x \hat{J}_y | J M \rangle &= \mp \frac{i}{4} \sqrt{(J \pm M + 2)(J \pm M + 1)(J \mp M - 1)(J \mp M)}, \\ \langle J M | \hat{J}_x \hat{J}_y | J M \rangle &= \frac{i}{2} M, \\ \langle J M \pm 2 | \hat{J}_y \hat{J}_x | J M \rangle &= \mp \frac{i}{4} \sqrt{(J \pm M + 2)(J \pm M + 1)(J \mp M - 1)(J \mp M)}, \\ \langle J M | \hat{J}_y \hat{J}_x | J M \rangle &= -\frac{i}{2} M, \\ \langle J M \pm 1 | \hat{J}_x \hat{J}_z | J M \rangle &= \frac{M}{2} \sqrt{(J \pm M + 1)(J \mp M)}, \\ \langle J M \pm 1 | \hat{J}_z \hat{J}_x | J M \rangle &= \frac{M \pm 1}{2} \sqrt{(J \pm M + 1)(J \mp M)}, \\ \langle J M \pm 1 | \hat{J}_y \hat{J}_z | J M \rangle &= \mp \frac{i}{2} M \sqrt{(J \pm M + 1)(J \mp M)}, \\ \langle J M \pm 1 | \hat{J}_z \hat{J}_y | J M \rangle &= \mp \frac{i}{2} (M \pm 1) \sqrt{(J \pm M + 1)(J \mp M)}. \end{aligned} \quad (13.2.46)$$

In the case of the irreducible tensor product of two operators $\hat{\mathbf{J}}$ one obtains

$$\langle J' M' | \{ \hat{\mathbf{J}}_1 \otimes \hat{\mathbf{J}}_1 \}_{k\kappa} | J M \rangle = \delta_{JJ'} (-1)^{2J+k} J(J+1) \Pi_{kJ} \left\{ \begin{matrix} 1 & 1 & k \\ J & J & J \end{matrix} \right\} C_{J M k \kappa}^{J M'}. \quad (13.2.47)$$

$$\langle J' || \{ \hat{\mathbf{J}}_1 \otimes \hat{\mathbf{J}}_1 \}_k || J \rangle = \delta_{JJ'} (-1)^{2J+k} J(J+1) \Pi_{kJJ'} \left\{ \begin{matrix} 1 & 1 & k \\ J & J & J \end{matrix} \right\}.$$

In particular, from this equation one has two obvious relations

$$\begin{aligned} \langle J' M' | (\hat{\mathbf{J}} \cdot \hat{\mathbf{J}}) | J M \rangle &= \delta_{JJ'} \delta_{MM'} J(J+1), \\ \langle J' M' | [\hat{\mathbf{J}} \times \hat{\mathbf{J}}]_{\kappa} | J M \rangle &= \delta_{JJ'} i \sqrt{J(J+1)} C_{J M 1 \kappa}^{J M'}. \end{aligned}$$

In applications, the expressions for matrix elements of products of the operator $\widehat{\mathbf{J}}$ and the operators $\widehat{\mathbf{n}}_1$ or $(\widehat{\nabla}_\Omega)_1$ are of particular interest. Some relations for the matrix elements in questions will be considered below. The matrix elements of the direct tensor product of operators $\widehat{\mathbf{n}}_1$ and $\widehat{\mathbf{J}}_1$ are given by

$$\langle l' s' J' M' | \widehat{n}_{1\mu} \widehat{J}_{1\nu} | l s J M \rangle = \delta_{ss'} (-1)^{s+J+l'+1} \sqrt{J(J+1)} \Pi_{lJ} C_{l010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M + \nu}^{J' M'} C_{J M 1 \nu}^{J M + \nu}, \quad (13.2.48)$$

$$\langle l' s' J' M' | \widehat{J}_{1\nu} \widehat{n}_{1\mu} | l s J M \rangle = \langle l' s' J' M' | \widehat{n}_{1\mu} \widehat{J}_{1\nu} | l s J M \rangle - \langle l' s' J' M' | \Re_{1\mu 1\nu}(n_{1\mu}, J_{1\nu}) | l s J M \rangle. \quad (13.2.49)$$

The commutator of components of these operators is as follows:

$$\Re_{1\mu 1\nu}(\widehat{n}_{1\mu}, \widehat{J}_{1\nu}) = -\sqrt{\frac{8\pi}{3}} C_{1\mu 1\nu}^{1\mu+\nu} Y_{1\mu+\nu}(\vartheta, \varphi). \quad (13.2.50)$$

Using Eqs. (13.2.104) and (13.2.105), one finds

$$\langle l' s' J' M' | \Re_{1\mu 1\nu}(\widehat{n}_{1\mu}, \widehat{J}_{1\nu}) | l s J M \rangle = (-1)^{J+l'+s} \delta_{ss'} \sqrt{2} \Pi_{lJ} C_{l010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{1\mu 1\nu}^{1\mu+\nu} C_{J M 1 \mu+\nu}^{J' M'}. \quad (13.2.51)$$

In the case of the irreducible tensor product of the operators $\widehat{\mathbf{n}}_1$ and $\widehat{\mathbf{J}}_1$ one obtains

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} | l s J M \rangle &= (-1)^{s+l'-J'+k+1} \delta_{ss'} \sqrt{J(J+1)} \Pi_{kl} \\ &\times \Pi_{JJ} C_{l010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M k \kappa}^{J' M'}, \end{aligned} \quad (13.2.52)$$

$$\langle l' s' J' | \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k | l s J \rangle = (-1)^{s+l'-J'+k+1} \delta_{ss'} \sqrt{J(J+1)} \Pi_{klJ'} \Pi_{JJ} C_{l010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}.$$

The commutator of the irreducible tensor product may be expressed as

$$\Re_{k\kappa}^{11}(\widehat{\mathbf{n}}_1, \widehat{\mathbf{J}}_1) = -\sqrt{\frac{8\pi}{3}} Y_{1\kappa}(\vartheta, \varphi) \delta_{k1}. \quad (13.2.53)$$

Taking Eqs. (13.2.104) and (13.2.105) into account, we get

$$\langle l' s' J' M' | \Re_{k\kappa}^{11}(\widehat{\mathbf{n}}_1, \widehat{\mathbf{J}}_1) | l s J M \rangle = (-1)^{J+l'+s} \delta_{ss'} \sqrt{2} \Pi_{Jl} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{l010}^{l'0} C_{J M 1 \kappa}^{J' M'} \delta_{k1}. \quad (13.2.54)$$

$$\langle l' s' J' | \Re_k^{11}(\widehat{\mathbf{n}}_1, \widehat{\mathbf{J}}_1) | l s J \rangle = (-1)^{J+l'+s} \delta_{ss'} \sqrt{2} \Pi_{Jl'} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{l010}^{l'0} \delta_{k1}.$$

The matrix elements of $\{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{n}}_1 \}_{k\kappa}$ are determined by

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{n}}_1 \}_{k\kappa} | l s J M \rangle \\ = (-1)^k \langle l' s' J' M' | \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} | l s J M \rangle + (-1)^{k+1} \langle l' s' J' M' | \Re_{k\kappa}^{11}(\widehat{\mathbf{n}}_1, \widehat{\mathbf{J}}_1) | l s J M \rangle. \end{aligned} \quad (13.2.55)$$

This expression may also be rewritten in the form

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{n}}_1 \}_{k\kappa} | l s J M \rangle &= (-1)^{s+l'-J'+k+1} \delta_{ss'} \Pi_{lkJ'J} \\ &\times \sqrt{J'(J'+1)} C_{l010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M k \kappa}^{J' M'}, \end{aligned} \quad (13.2.56)$$

$$\langle l' s' J' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{n}}_1 \}_k | l s J \rangle = (-1)^{s+l'-J'+k+1} \delta_{ss'} \Pi_{lkJ}(2J'+1) \sqrt{J'(J'+1)} C_{l010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}.$$

Let us consider the matrix elements of products of the operators $(\widehat{\nabla}_\Omega)_1$ and $\widehat{\mathbf{J}}_1$. In the case of the direct product of these operators we have

$$\langle l' s' J' M' | (\widehat{\nabla}_\Omega)_{1\mu} \widehat{J}_{1\nu} | l s J M \rangle = (-1)^{J+l'+s+1} \sqrt{J(J+1)} \Pi_J \left\{ \begin{matrix} J & J' & 1 \\ l' & l & s \end{matrix} \right\} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle C_{J M + \nu}^{J' M'} C_{J M 1 \nu}^{J M + \nu}. \quad (13.2.57)$$

As in the foregoing case,

$$\langle l' s' J' M' | \hat{J}_{1\nu} (\hat{\nabla}_\Omega)_{1\mu} | l s J M \rangle = \langle l' s' J' M' | (\hat{\nabla}_\Omega)_{1\mu} \hat{J}_{1\nu} | l s J M \rangle - \langle l' s' J' M' | \Re_{1\mu 1\nu} ((\hat{\nabla}_\Omega)_{1\mu}, J_{1\nu}) | l s J M \rangle. \quad (13.2.58)$$

The commutator of components of the operators $(\hat{\nabla}_\Omega)_1$ and $\hat{J}_1$ is

$$\Re_{1\mu 1\nu} ((\hat{\nabla}_\Omega)_{1\mu}, \hat{J}_{1\nu}) = -\sqrt{2} C_{1\mu 1\nu}^{1\lambda} (\hat{\nabla}_\Omega)_\lambda. \quad (13.2.59)$$

Using Eq. (13.2.26), we find

$$\langle l' s' J' M' | \Re_{1\mu 1\nu} ((\hat{\nabla}_\Omega)_{1\mu}, J_{1\nu}) | l s J M \rangle = (-1)^{J+l'-s} \delta_{ss'} \sqrt{2} \Pi_J \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{1\mu 1\nu}^{1\mu+\nu} C_{J M 1\mu+\nu}^{J' M'} \langle l' || (\nabla_\Omega)_1 || l \rangle. \quad (13.2.60)$$

In the case of the irreducible tensor product of these operators, $\{(\nabla_\Omega)_1 \otimes \hat{J}_1\}_{k\kappa}$ one has

$$\begin{aligned} \langle l' s' J' M' | \{(\hat{\nabla}_\Omega)_1 \otimes \hat{J}_1\}_{k\kappa} | l s J M \rangle &= (-1)^{s+l'-J'+k+1} \delta_{ss'} \sqrt{J(J+1)} \Pi_k \\ &\times \Pi_{JJ} \left\{ \begin{matrix} J & J' & 1 \\ l' & l & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \langle l' || (\nabla_\Omega)_1 || l \rangle C_{J M k\kappa}^{J' M'}, \end{aligned} \quad (13.2.61)$$

$$\langle l' s' J' || \{(\hat{\nabla}_\Omega)_1 \otimes \hat{J}_1\}_k || l s J \rangle = (-1)^{s+l'-J'+k+1} \delta_{ss'} \sqrt{J(J+1)} \Pi_{kJ'} \Pi_{JJ} \left\{ \begin{matrix} J & J' & 1 \\ l' & l & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \langle l' || (\nabla_\Omega)_1 || l \rangle,$$

while for the irreducible tensor product $\{\hat{J}_1 \otimes (\hat{\nabla}_\Omega)_1\}_{k\kappa}$ we have

$$\begin{aligned} \langle l' s' J' M' | \{\hat{J}_1 \otimes (\hat{\nabla}_\Omega)_1\}_{k\kappa} | l s J M \rangle &= (-1)^k \langle l' s' J' M' | \{(\hat{\nabla}_\Omega)_1 \otimes \hat{J}_1\}_{k\kappa} | l s J M \rangle \\ &+ (-1)^{k+1} \langle l' s' J' M' | \Re_{k\kappa}^{11} ((\hat{\nabla}_\Omega)_1, \hat{J}_1) | l s J M \rangle. \end{aligned} \quad (13.2.62)$$

The commutator of $\{(\hat{\nabla}_\Omega)_1 \otimes \hat{J}_1\}_{k\kappa}$ is determined by

$$\Re_{k\kappa}^{11} ((\hat{\nabla}_\Omega)_1, \hat{J}_1) = -\sqrt{2} (\hat{\nabla}_\Omega)_{1\kappa} \delta_{k1}, \quad (13.2.63)$$

i.e.,

$$\langle l' s' J' M' | \Re_{k\kappa}^{11} ((\hat{\nabla}_\Omega)_1, \hat{J}_1) | l s J M \rangle = (-1)^{J+l'+s} \delta_{ss'} \delta_{k1} \sqrt{2} \Pi_J \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} \langle l' || (\hat{\nabla}_\Omega)_1 || l \rangle C_{J M 1\kappa}^{J' M'}, \quad (13.2.64)$$

$$\langle l' s' J' || \Re_k^{11} ((\hat{\nabla}_\Omega)_1, \hat{J}_1) || l s J \rangle = (-1)^{J+l'+s} \delta_{ss'} \delta_{k1} \sqrt{2} \Pi_{JJ'} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} \langle l' || (\hat{\nabla}_\Omega)_1 || l \rangle C_{J M 1\kappa}^{J' M'}.$$

Equation (13.2.62) may also be represented by

$$\begin{aligned} \langle l' s' J' M' | \{\hat{J}_1 \otimes (\hat{\nabla}_\Omega)_1\}_{k\kappa} | l s J M \rangle &= (-1)^{s+l'-J'+k+1} \delta_{ss'} \Pi_{klJ'} \\ &\times \sqrt{J'(J'+1)} \left\{ \begin{matrix} J & J' & 1 \\ l' & l & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \langle l' || (\nabla_\Omega)_1 || l \rangle C_{J M k\kappa}^{J' M'}, \end{aligned} \quad (13.2.65)$$

$$\langle l' s' J' || \{\hat{J}_1 \otimes (\hat{\nabla}_\Omega)_1\}_k || l s J \rangle = (-1)^{s+l'-J'+k+1} \delta_{ss'} \Pi_{klJ'J'} \sqrt{J'(J'+1)} \left\{ \begin{matrix} J & J' & 1 \\ l' & l & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \langle l' || (\nabla_\Omega)_1 || l \rangle.$$

13.2.6 Matrix Elements of the Orbital Angular Momentum Operator $\hat{\mathbf{L}} \equiv \hat{\mathbf{L}}_1$

This operator acts only on the position variables θ, φ . In accordance with the Wigner-Eckart theorem, we get

$$\langle l' m' | \hat{L}_{1\mu} | l m \rangle = \frac{\langle l' || \hat{\mathbf{L}}_1 || l \rangle}{\Pi_{l'}} C_{l' m' 1\mu}^{l m} \quad (13.2.66)$$

where

$$\langle l' || \hat{\mathbf{L}}_1 || l \rangle = \delta_{ll'} \sqrt{l(l+1)} \Pi_l. \quad (13.2.67)$$

Equations (13.2.39)-(13.2.47) are obtained for the matrix elements of the total angular momentum operator $\hat{\mathbf{J}}$ and remain valid for the operator $\hat{\mathbf{L}}$ after the replacement,

$$J \rightarrow l, \quad M \rightarrow m, \quad \hat{J}_{1\mu} \rightarrow \hat{L}_{1\mu}.$$

A reduced matrix element in the lsJ -representation is given by

$$\langle l' s' J' | \hat{\mathbf{L}}_1 | l s J \rangle = (-1)^{s+J+l+1} \delta_{ss'} \delta_{ll'} \Pi_{JJ'l} \sqrt{l(l+1)} \begin{Bmatrix} l & s & J \\ J' & 1 & l \end{Bmatrix}. \quad (13.2.68)$$

We present matrix elements of products of the operators $\hat{\mathbf{L}}_1$ and $\mathbf{n}_1$. For the direct product of these operators we have

$$\langle l' m' | \hat{n}_{1\mu} \hat{L}_{1\nu} | l m \rangle = \sqrt{l(l+1)} \Pi_l / \Pi_{l'} C_{l'010}^{l'0} C_{l'm+\nu 1\mu}^{l'm'} C_{lm1\nu}^{l'm+\nu}. \quad (13.2.69)$$

The commutator of the components $\hat{n}_{1\mu}$ and $\hat{L}_{1\nu}$ may be written in the form

$$\mathfrak{R}_{1\mu 1\nu}(\hat{n}_{1\mu}, \hat{L}_{1\nu}) = -\sqrt{\frac{8\pi}{3}} C_{1\mu 1\nu}^{1\mu+\nu} Y_{1\mu+\nu}(\vartheta, \varphi). \quad (13.2.70)$$

Taking Eqs. (13.2.104) and (13.2.105) in account one finds

$$\langle l' m' | \mathfrak{R}_{1\mu 1\nu}(\hat{n}_{1\mu}, \hat{L}_{1\nu}) | l m \rangle = -\sqrt{2} \Pi_l / \Pi_{l'} C_{l'010}^{l'0} C_{1\mu 1\nu}^{1\mu+\nu} C_{lm1\mu+\nu}^{l'm'}. \quad (13.2.71)$$

The matrix element of the direct product $\hat{L}_{1\nu} \hat{n}_{1\mu}$ is

$$\langle l' m' | \hat{L}_{1\nu} \hat{n}_{1\mu} | l m \rangle = \Pi_l / \Pi_{l'} \sqrt{l'(l'+1)} C_{l'010}^{l'0} C_{l'm+\mu 1\nu}^{l'm'} C_{lm1\mu}^{l'm+\mu}. \quad (13.2.72)$$

In the case of irreducible tensor products of the operators in question one gets

$$\langle l' m' | \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_{k\kappa} | l m \rangle = (-1)^{l'+l+k} C_{l'm k \kappa}^{l' m'} \Pi_{kl} / \Pi_{l'} \sqrt{l(l+1)} C_{l'010}^{l'0} \begin{Bmatrix} 1 & 1 & k \\ l' & l & l \end{Bmatrix}, \quad (13.2.73)$$

$$\langle l' m' | \{ \hat{\mathbf{L}}_1 \otimes \hat{\mathbf{n}}_1 \}_{k\kappa} | l m \rangle = (-1)^{l'+l+k} C_{l'm k \kappa}^{l' m'} \Pi_{kl} \sqrt{l'(l'+1)} C_{l'010}^{l'0} \begin{Bmatrix} 1 & 1 & k \\ l' & l & l \end{Bmatrix}, \quad (13.2.74)$$

$$\langle l' | \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_k | l \rangle = (-1)^{l'+l+k} \Pi_{kl} \sqrt{l(l+1)} C_{l'010}^{l'0} \begin{Bmatrix} 1 & 1 & k \\ l' & l & l \end{Bmatrix},$$

$$\langle l' | \{ \hat{\mathbf{L}}_1 \otimes \hat{\mathbf{n}}_1 \}_k | l \rangle = (-1)^{l'+l+k} \Pi_{kl} \sqrt{l'(l'+1)} C_{l'010}^{l'0} \begin{Bmatrix} 1 & 1 & k \\ l' & l & l \end{Bmatrix}.$$

Both equations are related by the expression

$$\langle l' m' | \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_{k\kappa} | l m \rangle = (-1)^k \langle l' m' | \{ \hat{\mathbf{L}}_1 \otimes \hat{\mathbf{n}}_1 \}_{k\kappa} | l m \rangle + \langle l' m' | \mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, \hat{\mathbf{L}}_1) | l m \rangle, \quad (13.2.75)$$

where

$$\mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, \hat{\mathbf{L}}_1) = -\sqrt{\frac{8\pi}{3}} Y_{1\kappa} \delta_{k1}, \quad (13.2.76)$$

$$\langle l' m' | \mathfrak{R}_{k\kappa}^{11}(\hat{\mathbf{n}}_1, \hat{\mathbf{L}}_1) | l m \rangle = -\frac{\Pi_l}{\Pi_{l'}} C_{l'010}^{l'0} C_{l'm 1\kappa}^{l'm'}, \quad (13.2.77)$$

$$\langle l' | \mathfrak{R}_k^{11}(\hat{\mathbf{n}}_1, \hat{\mathbf{L}}_1) | l \rangle = -\Pi_l C_{l'010}^{l'0} \delta_{k1}.$$

Let us now consider matrix elements of products of the operators $(\hat{\nabla}_\Omega)_1$ and $\hat{\mathbf{L}}_1$. The matrix elements of the direct product of these operators are

$$\langle l' m' | (\hat{\nabla}_\Omega)_{1\mu} \hat{L}_{1\nu} | l m \rangle = \frac{\sqrt{l(l+1)}}{\Pi_{l'}} \langle l' | (\hat{\nabla}_\Omega)_1 | l \rangle C_{l'm+\nu 1\mu}^{l'm'} C_{lm1\nu}^{l'm+\nu}, \quad (13.2.78)$$

$$\langle l' m' | \hat{L}_{1\nu} (\hat{\nabla}_\Omega)_{1\mu} | l m \rangle = \frac{\sqrt{l(l+1)}}{\Pi_{l'}} \langle l' | (\hat{\nabla}_\Omega)_1 | l \rangle C_{l'm+\mu 1\nu}^{l'm'} C_{lm1\mu}^{l'm+\mu}. \quad (13.2.79)$$

Equations (13.2.78) and (13.2.79) differ by a quantity which is the matrix element of the commutator of $\hat{L}_{1\nu}$ and $(\hat{\nabla}_\Omega)_{1\mu}$,

$$\mathfrak{R}_{1\mu 1\nu}((\hat{\nabla}_\Omega)_{1\mu}, \hat{L}_{1\nu}) = -\sqrt{2} C_{1\mu 1\nu}^{1\mu+\nu} (\hat{\nabla}_\Omega)_{1\mu+\nu}, \quad (13.2.80)$$

$$\langle l' m' | \mathfrak{R}_{1\mu 1\nu}((\hat{\nabla}_\Omega)_{1\mu}, \hat{L}_{1\nu}) | l m \rangle = -\frac{\sqrt{2}}{\Pi_{l'}} \langle l' | (\hat{\nabla}_\Omega)_1 | l \rangle C_{1\mu 1\nu}^{1\mu+\nu} C_{lm1\mu+\nu}^{l'm'}. \quad (13.2.81)$$

In the case of the irreducible tensor product of the operators $(\widehat{\nabla}_\Omega)_1$ and $\widehat{\mathbf{L}}_1$ we find

$$\langle l'm' | \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{L}}_1 \}_{k\kappa} | lm \rangle = (-1)^{l'+l+k} \Pi_{kl} / \Pi_{l'} \sqrt{l(l+1)} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l \end{matrix} \right\} C_{l'm'k\kappa}^{l' m'}, \quad (13.2.82)$$

$$\begin{aligned} \langle l' | \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{L}}_1 \}_k | l \rangle &= (-1)^{l'+l+k} \Pi_{kl} \sqrt{l(l+1)} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l \end{matrix} \right\}, \\ \langle l'm' | \{ \widehat{\mathbf{L}}_1 \otimes (\widehat{\nabla}_\Omega)_1 \}_{k\kappa} | lm \rangle &= (-1)^{l'+l+k} \Pi_k \sqrt{l'(l'+1)} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l' \end{matrix} \right\} C_{l'm'k\kappa}^{l' m'}, \\ \langle l' | \{ \widehat{\mathbf{L}}_1 \otimes (\widehat{\nabla}_\Omega)_1 \}_k | l \rangle &= (-1)^{l'+l+k} \Pi_{kl'} \sqrt{l'(l'+1)} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle \left\{ \begin{matrix} 1 & 1 & k \\ l' & l & l' \end{matrix} \right\}. \end{aligned} \quad (13.2.83)$$

The commutator of the product $\{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{L}}_1 \}_{k\kappa}$ may be represented in the form

$$\mathfrak{R}_{k\kappa}^{11}((\widehat{\nabla}_\Omega)_1, \widehat{\mathbf{L}}_1) = -\sqrt{2}(\widehat{\nabla}_\Omega)_{1\kappa} \delta_{k1}, \quad (13.2.84)$$

Using Eq. (13.2.26), one obtains

$$\begin{aligned} \langle l'm' | \mathfrak{R}_{k\kappa}^{11}((\widehat{\nabla}_\Omega)_1, \widehat{\mathbf{L}}_1) | lm \rangle &= -\frac{\sqrt{2}}{\Pi_{l'}} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle C_{l'm'1\kappa}^{l' m'}, \\ \langle l' | \mathfrak{R}_{k\kappa}^{11}((\widehat{\nabla}_\Omega)_1, \widehat{\mathbf{L}}_1) | l \rangle &= -\sqrt{2} \langle l' | (\widehat{\nabla}_\Omega)_1 | l \rangle. \end{aligned} \quad (13.2.85)$$

The matrix elements of the irreducible tensor product of the operators $\widehat{\mathbf{L}}_1$ and $\widehat{\mathbf{J}}_1$ are

$$\begin{aligned} \langle l's'J'M' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} | lsJM \rangle &= (-1)^{s+l-J'+k+1} \delta_{ll'} \delta_{ss'} C_{J'M'k\kappa}^{J' M'} \Pi_{JJ} \\ &\quad \times \Pi_{kl} \sqrt{l(l+1)J(J+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\}, \end{aligned} \quad (13.2.86)$$

$$\langle l's'J' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k | lsJ \rangle = (-1)^{s+l-J'+k+1} \delta_{ll'} \delta_{ss'} \Pi_{JJ} \Pi_{J'} \Pi_{kl} \sqrt{l(l+1)J(J+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\},$$

$$\begin{aligned} \langle l's'J'M' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{L}}_1 \}_{k\kappa} | lsJM \rangle &= (-1)^{s+l-J'+k+1} \delta_{l'l} \delta_{ss'} C_{J'M'k\kappa}^{J' M'} \\ &\quad \times \Pi_{klJJ'} \sqrt{l(l+1)J'(J'+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\}. \end{aligned} \quad (13.2.87)$$

$$\langle l's'J' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{L}}_1 \}_k | lsJ \rangle = (-1)^{s+l-J'+k+1} \delta_{l'l} \delta_{ss'} \Pi_{klJJ'} \Pi_{J'} \sqrt{l(l+1)J'(J'+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\}.$$

The commutator of the irreducible tensor product $\{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa}$ is defined as

$$\mathfrak{R}_{k\kappa}^{11}(\widehat{\mathbf{L}}_1, \widehat{\mathbf{J}}_1) = -\sqrt{2} \widehat{L}_{1\kappa} \delta_{k1}, \quad (13.2.88)$$

To evaluate its matrix elements one can use Eqs. (13.2.66) and (13.2.67). Then

$$\langle l's'J'M' | \mathfrak{R}_{k\kappa}^{11}(\widehat{\mathbf{L}}_1, \widehat{\mathbf{J}}_1) | lsJM \rangle = (-1)^{s+J+l} \delta_{ll'} \delta_{ss'} \delta_{k1} \Pi_{Jl} \sqrt{2l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\} C_{J'M'1\kappa}^{J' M'}. \quad (13.2.89)$$

To determine the matrix elements of the direct tensor product of the operators $\widehat{\mathbf{L}}_1$ and $\widehat{\mathbf{J}}_1$ one can decompose this direct product into irreducible products. In this way one obtains

$$\langle l's'J'M' | \widehat{L}_{1\nu} \widehat{J}_{1\mu} | lsJM \rangle = \sum_{k\kappa} C_{1\mu 1\nu}^{k\kappa} \langle l's'J'M' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} | lsJM \rangle. \quad (13.2.90)$$

$$\langle l's'J'M' | \widehat{J}_{1\mu} \widehat{L}_{1\nu} | lsJM \rangle = \sum_{k\kappa} C_{1\mu 1\nu}^{k\kappa} \langle l's'J'M' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{L}}_1 \}_{k\kappa} | lsJM \rangle. \quad (13.2.91)$$

The commutator of the operator components $\widehat{L}_{1\mu}$ and $\widehat{J}_{1\nu}$ may be presented in the form

$$\mathfrak{R}_{1\mu 1\nu}(\widehat{L}_{1\mu}, \widehat{J}_{1\nu}) = -\sqrt{2} C_{1\mu 1\nu}^{1\mu+\nu} \widehat{L}_{1\mu+\nu}. \quad (13.2.92)$$

From Eqs. (13.2.92), (13.2.66) and (13.2.67) we find

$$\langle l's'J'M' | \mathfrak{R}_{1\mu 1\nu}(\widehat{L}_{1\mu}, \widehat{J}_{1\nu}) | lsJM \rangle = (-1)^{s+J+l} \delta_{ll'} \delta_{ss'} \Pi_{Jl} \sqrt{2l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\} C_{1\mu 1\nu}^{1\mu+\nu} C_{J'M'1\mu+\nu}^{J' M'}. \quad (13.2.93)$$

13.2.7 Matrix Elements of the Spin Angular Momentum Operator $\widehat{\mathbf{S}} \equiv \widehat{\mathbf{S}}_1$

This operator acts only on spin variables. Using the Wigner-Eckart theorem, we get

$$\langle s' m'_s | \widehat{S}_{1\mu} | s m_s \rangle = \frac{\langle s' || \widehat{\mathbf{S}}_1 || s \rangle}{\Pi_{s'}} C_{s m_s 1 \mu}^{s' m'_s}, \quad (13.2.94)$$

$$\langle s' || \widehat{\mathbf{S}}_1 || s \rangle = \delta_{ss'} \Pi_s \sqrt{s(s+1)}. \quad (13.2.95)$$

Equations (13.2.39)-(13.2.47) were obtained for the matrix elements of the operator $\widehat{\mathbf{J}}_1$ and they remain valid for the spin operator $\widehat{\mathbf{S}}_1$ after the replacement

$$J \rightarrow s, \quad M \rightarrow m_s, \quad \widehat{\mathbf{J}}_1 \rightarrow \widehat{\mathbf{S}}_1.$$

The use of Eqs. (13.2.6) and (13.2.95) yields an expression for the reduced matrix element of the spin operator in the (lsJ) -representation

$$\langle l' s' J' || \widehat{\mathbf{S}}_1 || lsJ \rangle = (-1)^{l+s+J'+1} \delta_{ll'} \delta_{ss'} \Pi_{sJJ'} \sqrt{s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\}. \quad (13.2.96)$$

The operators $\widehat{\mathbf{n}}_1, (\widehat{\nabla}_\Omega)_1, \widehat{\mathbf{L}}_1$ commute with the spin operator $\widehat{\mathbf{S}}_1$. Matrix elements of the irreducible tensor products of these operators and the spin operator are given by

$$\langle l' s' J' M' | \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} | lsJM \rangle = \delta_{ss'} \Pi_{kJls} \sqrt{s(s+1)} C_{l'010}^{l'0} C_{JMk\kappa}^{J'M'} \left\{ \begin{matrix} l & 1 & l' \\ s & 1 & s \\ J & k & J' \end{matrix} \right\}, \quad (13.2.97)$$

$$\langle l' s' J' M' | \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} | lsJM \rangle = \delta_{ss'} \Pi_{kJs} \sqrt{s(s+1)} \langle l' || (\widehat{\nabla}_\Omega)_1 || l \rangle C_{JMk\kappa}^{J'M'} \left\{ \begin{matrix} l & 1 & l' \\ s & 1 & s \\ J & k & J' \end{matrix} \right\}, \quad (13.2.98)$$

$$\langle l' s' J' M' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} | lsJM \rangle = \delta_{ll'} \delta_{ss'} \Pi_{kJls} \sqrt{l(l+1)s(s+1)} C_{JMk\kappa}^{J'M'} \left\{ \begin{matrix} l & 1 & l \\ s & 1 & s \\ J & k & J' \end{matrix} \right\}, \quad (13.2.99)$$

$$\langle l' s' J' || \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{S}}_1 \}_k || lsJ \rangle = \delta_{ss'} \Pi_{kJlsJ'} \sqrt{s(s+1)} C_{l'010}^{l'0} \left\{ \begin{matrix} l & 1 & l' \\ s & 1 & s \\ J & k & J' \end{matrix} \right\},$$

$$\langle l' s' J' || \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{S}}_1 \}_k || lsJ \rangle = \delta_{ss'} \Pi_{kJsJ'} \sqrt{s(s+1)} \langle l' || (\widehat{\nabla}_\Omega)_1 || l \rangle \left\{ \begin{matrix} l & 1 & l' \\ s & 1 & s \\ J & k & J' \end{matrix} \right\},$$

$$\langle l' s' J' || \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{S}}_1 \}_k || lsJ \rangle = \delta_{ll'} \delta_{ss'} \Pi_{kJlsJ'} \sqrt{l(l+1)s(s+1)} \left\{ \begin{matrix} l & 1 & l \\ s & 1 & s \\ J & k & J' \end{matrix} \right\}.$$

The operators $\widehat{\mathbf{S}}_1$ and $\widehat{\mathbf{J}}_1$ do not commute. Matrix elements of irreducible tensor products of these operators depend on the order of the operators.

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} | lsJM \rangle &= (-1)^{J+k-l-s+1} \delta_{s's} \delta_{l'l} \Pi_{JJks} \sqrt{s(s+1)J(J+1)} \\ &\times \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\} C_{JMk\kappa}^{J'M'}, \end{aligned} \quad (13.2.100)$$

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} | lsJM \rangle &= (-1)^{J+k-l-s+1} \delta_{s's} \delta_{l'l} \Pi_{kJJ's} \sqrt{s(s+1)J'(J'+1)} \\ &\times \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\} C_{JMk\kappa}^{J'M'}, \end{aligned} \quad (13.2.101)$$

$$\langle l' s' J' || \{ \widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k || lsJ \rangle = (-1)^{J+k-l-s+1} \delta_{s's} \delta_{l'l} \Pi_{JJksJ'} \sqrt{s(s+1)J(J+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J \end{matrix} \right\} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\},$$

$$\langle l' s' J' || \{ \widehat{\mathbf{J}}_1 \otimes \widehat{\mathbf{S}}_1 \}_k || lsJ \rangle = (-1)^{J+k-l-s+1} \delta_{s's} \delta_{l'l} \Pi_{JJksJ'} \sqrt{s(s+1)J'(J'+1)} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J & J' \end{matrix} \right\} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\}.$$

The commutator of the product $\{\widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1\}_{k\kappa}$ is as follows:

$$\mathfrak{R}_{k\kappa}^{11}(\widehat{\mathbf{S}}_1, \widehat{\mathbf{J}}_1) = -\sqrt{2}\widehat{S}_{1\kappa}\delta_{k1}, \quad (13.2.102)$$

Using Eqs. (13.2.94) and (13.2.95), we find

$$\langle l' s' J' M' | \mathfrak{R}_{k\kappa}^{11}(\widehat{\mathbf{S}}_1, \widehat{\mathbf{J}}_1) | l s J M \rangle = (-1)^{l+s+J'} \delta_{ll'} \delta_{ss'} \delta_{k1} \Pi_{sJ} \sqrt{2s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\} C_{JMk\kappa}^{J'M'}, \quad (13.2.103)$$

$$\langle l' s' J' | \mathfrak{R}_k^{11}(\widehat{\mathbf{S}}_1, \widehat{\mathbf{J}}_1) | l s J \rangle = (-1)^{l+s+J'} \delta_{ll'} \delta_{ss'} \delta_{k1} \Pi_{sJJ'} \sqrt{2s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\}.$$

To obtain matrix elements of direct products of the operator $\widehat{\mathbf{S}}_1$ and the operators $\widehat{\mathbf{n}}_1$, $(\widehat{\nabla}_\Omega)_1$, $\widehat{\mathbf{L}}_1$ or $\widehat{\mathbf{J}}_1$ one should decompose the direct products into irreducible products in accordance with Eq. (3.1.22) and use Eqs. (13.2.97)-(13.2.101).

13.2.8 Matrix Elements of the Spherical Harmonic Operator $\widehat{Y}_{L\nu} \equiv Y_{L\nu}(\vartheta, \varphi)$

A spherical harmonic operator, $Y_{L\nu}$, depends only on position variables ϑ, φ . From the Wigner-Eckart theorem it follows that

$$\langle l' m' | \widehat{Y}_{L\nu} | l m \rangle = \frac{\langle l' || \widehat{\mathbf{Y}}_L || l \rangle}{\Pi_{l'}} C_{l m L \nu}^{l' m'}, \quad (13.2.104)$$

$$\langle l' || \widehat{\mathbf{Y}}_L || l \rangle = \Pi_{Ll} \sqrt{\frac{1}{4\pi}} C_{l 0 L 0}^{l' 0}. \quad (13.2.105)$$

Similar relationships for an operator $\widehat{C}_{L\nu}(\vartheta, \varphi)$ (see Eq. (5.1.7)) may be represented in the form

$$\langle l' m' | \widehat{C}_{L\nu} | l m \rangle = \frac{\langle l' || \widehat{\mathbf{C}}_L || l \rangle}{\Pi_{l'}} C_{l m L \nu}^{l' m'}, \quad (13.2.106)$$

$$\langle l' || \widehat{\mathbf{C}}_L || l \rangle = \Pi_l C_{l 0 L 0}^{l' 0}. \quad (13.2.107)$$

Let us consider the matrix elements of some tensor products involving $\widehat{Y}_{L\nu}(\vartheta, \varphi)$. For products of the commuting operators $\widehat{\mathbf{n}}_1$ and $\widehat{\mathbf{Y}}_L$, we get

$$\begin{aligned} \langle l' m' | \widehat{\mathbf{n}}_{1\mu} \widehat{Y}_{L\nu} | l m \rangle &= -\frac{\Pi_{Ll}}{\Pi_{l'}} \sqrt{\frac{1}{4\pi}} \left\{ \sqrt{\frac{l'+1}{2l'+3}} C_{l 0 L 0}^{l'+1 0} C_{l m L \nu}^{l'+1 m+\nu} C_{l'+1 m+\nu 1 \mu}^{l' m'} \right. \\ &\quad \left. - \sqrt{\frac{l'}{2l'-1}} C_{l 0 L 0}^{l'-1 0} C_{l m L \nu}^{l'-1 m+\nu} C_{l'-1 m+\nu 1 \mu}^{l' m'} \right\}, \end{aligned} \quad (13.2.108)$$

$$\langle l' m' | \{\widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{Y}}_L\}_{L'\nu'} | l m \rangle = (-1)^{L'+l'+l} \frac{\Pi_{LL'l}}{\Pi_{l'}} \sqrt{\frac{1}{4\pi}} C_{l m L' \nu'}^{l' m'} \times \sum_k \Pi_k C_{k 0 1 0}^{l' 0} C_{l 0 L 0}^{k 0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & k \end{matrix} \right\}, \quad (13.2.109)$$

$$\langle l' || \{\widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{Y}}_L\}_{L'} || l \rangle = (-1)^{L'+l'+l} \Pi_{LL'l} \sqrt{\frac{1}{4\pi}} \sum_k \Pi_k C_{k 0 1 0}^{l' 0} C_{l 0 L 0}^{k 0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & k \end{matrix} \right\}.$$

The matrix elements of products of the non-commuting operators $(\widehat{\nabla}_\Omega)_1$ and $\widehat{\Psi}_L$ are determined by

$$\begin{aligned} \langle l' m' | (\widehat{\nabla}_\Omega)_1 \widehat{Y}_{L\nu} | l m \rangle &= -\Pi_{Ll} / \Pi_{l'} \sqrt{\frac{1}{4\pi}} \left\{ (l'-1) \sqrt{\frac{l'}{2l'-1}} C_{l 0 L 0}^{l'-1 0} C_{l m L \nu}^{l'-1 m+\nu} C_{l'-1 m+\nu 1 \mu}^{l' m'} \right. \\ &\quad \left. + (l'+2) \sqrt{\frac{l'+1}{2l'+3}} C_{l 0 L 0}^{l'+1 0} C_{l m L \nu}^{l'+1 m+\nu} C_{l'+1 m+\nu 1 \mu}^{l' m'} \right\}, \end{aligned} \quad (13.2.110)$$

$$\langle l' m' | \{(\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{Y}}_L\}_{L'\nu'} | l m \rangle = (-1)^{L'+l'+1} \Pi_{LL'l} / \Pi_{l'} \sqrt{\frac{1}{4\pi}} \sum_k \langle l' || (\widehat{\nabla}_\Omega)_1 || k \rangle C_{l 0 L 0}^{k 0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & k \end{matrix} \right\} C_{l m L' \nu'}^{l' m'}, \quad (13.2.111)$$

$$\langle l' || \{(\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{Y}}_L\}_{L'} || l \rangle = (-1)^{L'+l'+1} \Pi_{LL'l} \sqrt{\frac{1}{4\pi}} \sum_k \langle l' || (\widehat{\nabla}_\Omega)_1 || k \rangle C_{l 0 L 0}^{k 0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & k \end{matrix} \right\}.$$

The commutator of components of these operators is as follows

$$\Re_{1\mu L\nu}((\hat{\nabla}_\Omega)_{1\mu}, \hat{Y}_{L\nu}) = -\sqrt{6L(L+1)}\Pi_{LL} \sum_{L'} \frac{1}{\sqrt{2L'+1}} C_{L'010}^{L'0} C_{1\mu L\nu}^{L'\mu+\nu} \left\{ \begin{matrix} 1 & 1 & 1 \\ L' & L & L \end{matrix} \right\} \hat{Y}_{L'\mu+\nu}(\vartheta, \varphi). \quad (13.2.112)$$

The commutator of the irreducible product of the same operators may be expressed as

$$\Re_{L'\nu'}^{1L}((\hat{\nabla}_\Omega)_1, \hat{Y}_L) = -\sqrt{6L(L+1)}\Pi_{LL}/\Pi_{L'} C_{L'010}^{L'0} \left\{ \begin{matrix} 1 & 1 & 1 \\ L' & L & L \end{matrix} \right\} \hat{Y}_{L'\nu'}(\vartheta, \varphi). \quad (13.2.113)$$

One can easily obtain the matrix elements of these commutators:

$$\langle l'm' | \Re_{1\mu L\nu}((\hat{\nabla}_\Omega)_{1\mu}, \hat{Y}_{L\nu}) | lm \rangle = -\Pi_{lLL}/\Pi_{l'} \sqrt{\frac{6L(L+1)}{4\pi}} C_{L'010}^{L'0} C_{1\mu L\nu}^{L'\mu+\nu} C_{lm L'\mu+\nu}^{l'm'} \left\{ \begin{matrix} 1 & 1 & 1 \\ L' & L & L \end{matrix} \right\}, \quad (13.2.114)$$

$$\langle l'm' | \Re_{L'\nu'}^{1L}((\hat{\nabla}_\Omega)_1, \hat{Y}_L) | lm \rangle = -\Pi_{lLL}/\Pi_{l'} \sqrt{\frac{6L(L+1)}{4\pi}} C_{lm L'\nu'}^{l'm'} C_{L'010}^{L'0} C_{l'0 L'}^{l'0} \left\{ \begin{matrix} 1 & 1 & 1 \\ L' & L & L \end{matrix} \right\}, \quad (13.2.115)$$

$$\langle l' | \Re_{L'\nu'}^{1L}((\hat{\nabla}_\Omega)_1, \hat{Y}_L) | l \rangle = -\Pi_{lLL} \sqrt{\frac{6L(L+1)}{4\pi}} C_{L'010}^{L'0} C_{l'0 L'}^{l'0} \left\{ \begin{matrix} 1 & 1 & 1 \\ L' & L & L \end{matrix} \right\}.$$

Of particular interest are the matrix elements of direct and irreducible products of the angular momentum operators and the spherical harmonic operator.

The matrix elements of products of the operators $\hat{\mathbf{L}}_1$ and $\hat{\mathbf{Y}}_L$ are given by

$$\langle l'm' | \hat{L}_{1\mu} \hat{Y}_{L\nu} | lm \rangle = \Pi_{Ll}/\Pi_{l'} \sqrt{\frac{l'(l'+1)}{4\pi}} C_{l'0 L 0}^{l'0} C_{lm L\nu}^{l'm+\nu} C_{l'm+\nu 1\mu}^{l'm'}, \quad (13.2.116)$$

$$\langle l'm' | \{\hat{\mathbf{L}}_1 \otimes \hat{\mathbf{Y}}_L\}_{L'\nu'} | lm \rangle = (-1)^{L'+l'+l} \Pi_{L'Ll} \sqrt{\frac{(l'+1)l'}{4\pi}} C_{lm L'\nu'}^{l'm'} C_{l'0 L 0}^{l'0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & l' \end{matrix} \right\}, \quad (13.2.117)$$

$$\langle l' | \{\hat{\mathbf{L}}_1 \otimes \hat{\mathbf{Y}}_L\}_{L'} | l \rangle = (-1)^{L'+l'+l} \Pi_{L'Lll'} \sqrt{\frac{(l'+1)l'}{4\pi}} C_{l'0 L 0}^{l'0} \left\{ \begin{matrix} L & 1 & L' \\ l' & l & l' \end{matrix} \right\}.$$

The commutators of these operators are

$$\Re_{1\mu L\nu}(\hat{L}_{1\mu}, \hat{Y}_{L\nu}) = \sqrt{L(L+1)} C_{L\nu 1\mu}^{L\mu+\nu} \hat{Y}_{L\mu+\nu}, \quad (13.2.118)$$

$$\Re_{L'\nu'}^{1L}(\hat{\mathbf{L}}_1, \hat{\mathbf{Y}}_L) = -\sqrt{L(L+1)} \hat{Y}_{L\mu+\nu} \delta_{L'L} \delta_{\nu'\mu+\nu}. \quad (13.2.119)$$

Using Eqs. (13.2.104) and (13.2.105), we find the expressions for the matrix elements of the commutators,

$$\langle l'm' | \Re_{1\mu L\nu}(\hat{L}_{1\mu}, \hat{Y}_{L\nu}) | lm \rangle = \Pi_{Ll}/\Pi_{l'} \sqrt{\frac{L(L+1)}{4\pi}} C_{lm L\mu+\nu}^{l'm'} C_{L\nu 1\mu}^{L\mu+\nu} C_{l'0 L 0}^{l'0}, \quad (13.2.120)$$

$$\langle l'm' | \Re_{L'\nu'}^{1L}(\hat{\mathbf{L}}_1, \hat{\mathbf{Y}}_L) | lm \rangle = -\Pi_{Ll}/\Pi_{l'} \sqrt{\frac{L(L+1)}{4\pi}} C_{l'0 L 0}^{l'0} C_{lm L\mu+\nu}^{l'm'} \delta_{L'L}. \quad (13.2.121)$$

In the case of products of the operators $\hat{\mathbf{S}}_1$ and $\hat{\mathbf{Y}}_L$ the matrix elements may be evaluated in the $lsJM$ representation:

$$\begin{aligned} \langle l's'J'M' | \hat{S}_{1\mu} \hat{Y}_{L\nu} | lsJM \rangle &= \delta_{ss'} \Pi_{JlLs} \sqrt{\frac{s(s+1)}{4\pi}} C_{l'0 L 0}^{l'0} \\ &\times \sum_k (-1)^{L+k+1} \Pi_k C_{JM k\mu+\nu}^{J'M'} C_{1\mu L\nu}^{k\mu+\nu} \left\{ \begin{matrix} l & L & l' \\ s & 1 & s \\ J & k & J' \end{matrix} \right\}; \end{aligned} \quad (13.2.122)$$

$$\langle l's'J'M' | \{\hat{\mathbf{Y}}_L \otimes \hat{\mathbf{S}}_1\}_{L'\nu'} | lsJM \rangle = \delta_{ss'} \Pi_{JL'Ll's} \sqrt{\frac{s(s+1)}{4\pi}} C_{l'0 L 0}^{l'0} C_{JM L'\nu'}^{J'M'} \left\{ \begin{matrix} l & L & l' \\ s & 1 & s \\ J & L' & J' \end{matrix} \right\}, \quad (13.2.123)$$

$$\langle l' s' J' | \{ \hat{\mathbf{Y}}_L \otimes \hat{\mathbf{S}}_1 \}_{L'} | l s J \rangle = \delta_{ss'} \Pi_{JL' L s J'} \sqrt{\frac{s(s+1)}{4\pi}} C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & L & l' \\ s & 1 & s \\ J & L' & J' \end{Bmatrix}.$$

The matrix elements of products of the operators $\hat{\mathbf{J}}_1$ and $\hat{\mathbf{Y}}_L$ in the same representation may be obtained by using the relations

$$\begin{aligned} \langle l' s' J' M' | \hat{J}_{1\mu} \hat{Y}_{L\nu} | l s J M \rangle &= (-1)^{l'+s+J+L} \delta_{ss'} \Pi_{LlJ} \sqrt{\frac{J'(J'+1)}{4\pi}} \\ &\times C_{J M L \nu}^{J' M' + \nu} C_{J' M' + \nu 1 \mu}^{J' M'} C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix}; \quad (13.2.124) \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | \{ \hat{\mathbf{J}}_1 \otimes \hat{\mathbf{Y}}_L \}_{L' \nu'} | l s J M \rangle &= (-1)^{l+s+L'-J'} \delta_{ss'} \Pi_{LL' l J J'} \sqrt{\frac{J'(J'+1)}{4\pi}} \\ &\times C_{J M L' \nu'}^{J' M'} \begin{Bmatrix} L & 1 & L' \\ J' & J & J' \end{Bmatrix} C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix}, \quad (13.2.125) \end{aligned}$$

$$\langle l' s' J' | \{ \hat{\mathbf{J}}_1 \otimes \hat{\mathbf{Y}}_L \}_{L'} | l s J \rangle = (-1)^{l+s+L'-J'} \delta_{ss'} \Pi_{LL' l J J'} \sqrt{\frac{J'(J'+1)}{4\pi}} \begin{Bmatrix} L & 1 & L' \\ J' & J & J' \end{Bmatrix} C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix}.$$

The commutators of these operators are reduced to the commutators of $\hat{\mathbf{L}}_1$ and $\hat{\mathbf{Y}}_L$. Namely,

$$\begin{aligned} \Re_{1\mu L\nu}(\hat{J}_{1\mu}, \hat{Y}_{L\nu}) &= \Re_{1\mu L\nu}(\hat{L}_{1\mu}, \hat{Y}_{L\nu}) \\ \Re_{L'\nu'}^L(\hat{J}_1, \hat{\mathbf{Y}}_L) &= \Re_{L'\nu'}^L(\hat{\mathbf{L}}_1, \hat{\mathbf{Y}}_L). \end{aligned}$$

The matrix elements of these commutators in the $lsJM$ -representation are given by

$$\begin{aligned} \langle l' s' J' M' | \Re_{1\mu L\nu}(\hat{J}_{1\mu}, \hat{Y}_{L\nu}) | l s J M \rangle &= (-1)^{J+l'+s+L} \delta_{ss'} \Pi_{LlJ} \sqrt{\frac{L(L+1)}{4\pi}} \\ &\times C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix} C_{L\nu 1 \mu}^{L\mu+\nu} C_{J M L \mu+\nu}^{J' M'}, \quad (13.2.126) \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | \Re_{L'\nu'}^{LL}(\hat{\mathbf{J}}_1, \hat{\mathbf{Y}}_L) | l s J M \rangle &= (-1)^{J+l'+s+L+1} \delta_{ss'} \delta_{LL'} \Pi_{LlJ} \sqrt{\frac{L(L+1)}{4\pi}} \\ &\times C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix} C_{J M L \nu'}^{J' M'}. \quad (13.2.127) \end{aligned}$$

$$\langle l' s' J' | \Re_{L'}^{LL}(\hat{\mathbf{J}}_1, \hat{\mathbf{Y}}_L) | l s J \rangle = (-1)^{J+l'+s+L+1} \delta_{ss'} \delta_{LL'} \Pi_{l' L l J} \sqrt{\frac{L(L+1)}{4\pi}} C_{l' 0 L 0}^{l' 0} \begin{Bmatrix} l & s & J \\ J' & L & l' \end{Bmatrix}.$$

Below we also present expressions for matrix elements of some more complex tensor products which include the spherical harmonic operator:

$$\begin{aligned} \langle l' m' | \{ \hat{\mathbf{n}}_1 \otimes (\hat{\nabla}_\Omega)_1 \}_{k\kappa} \hat{Y}_{L\nu} | l m \rangle &= \sum_{L'\nu'l''} (-1)^{l'+L'+k} \Pi_{Llkl''} / \Pi_{l'L'} \sqrt{\frac{1}{4\pi}} \\ &\times C_{l m L \nu}^{l' m'} C_{L'\nu' k \kappa}^{l' m'} C_{l' 0 L 0}^{l' 0} C_{l'' 0 1 0}^{l' 0} \begin{Bmatrix} 1 & L' & l'' \\ l' & 1 & k \end{Bmatrix} \langle l'' | (\hat{\nabla}_\Omega)_1 | L' \rangle, \quad (13.2.128) \end{aligned}$$

$$\begin{aligned} \langle l' m' | \{ \{ \hat{\mathbf{n}}_1 \otimes (\hat{\nabla}_\Omega)_1 \}_k \otimes \hat{\mathbf{Y}}_L \}_{F\varphi} | l m \rangle &= (-1)^{F+l+k} \Pi_{Ll k F} / \Pi_{l'} \sqrt{\frac{1}{4\pi}} \\ &\times C_{l m F \varphi}^{l' m'} \sum_{L'l''} (-1)^{L'} \Pi_{l''} C_{l' 0 L 0}^{L' 0} C_{l'' 0 1 0}^{l' 0} \begin{Bmatrix} 1 & L' & l'' \\ l' & 1 & k \end{Bmatrix} \begin{Bmatrix} L & k & F \\ l' & 1 & L' \end{Bmatrix} \langle l'' | (\hat{\nabla}_\Omega)_1 | L' \rangle, \quad (13.2.129) \end{aligned}$$

$$\begin{aligned} \langle l' | \{ \{ \hat{\mathbf{n}}_1 \otimes (\hat{\nabla}_\Omega)_1 \}_k \otimes \hat{\mathbf{Y}}_L \}_F | l \rangle &= (-1)^{F+l+k} \Pi_{Ll k F} \sqrt{\frac{1}{4\pi}} \\ &\times \sum_{L'l''} (-1)^{L'} \Pi_{l''} C_{l' 0 L 0}^{L' 0} C_{l'' 0 1 0}^{l' 0} \begin{Bmatrix} 1 & L' & l'' \\ l' & 1 & k \end{Bmatrix} \begin{Bmatrix} L & k & F \\ l' & 1 & L' \end{Bmatrix} \langle l'' | (\hat{\nabla}_\Omega)_1 | L' \rangle. \end{aligned}$$

$$\begin{aligned} \langle l'm' | \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_{k\kappa} \hat{Y}_{L\nu} | lm \rangle &= \sum_{L'\nu'} (-1)^{l'+L'+k} \Pi_{LlkL'} / \Pi_{l'} \sqrt{\frac{L'(L'+1)}{4\pi}} \\ &\times C_{lmL\nu}^{L'\nu'} C_{L'\nu'k\kappa}^{l'm'} C_{l0L0}^{L'0} C_{L'010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\}, \quad (13.2.130) \end{aligned}$$

$$\begin{aligned} \langle l'm' | \{ \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_k \otimes \mathbf{Y}_L \}_{F\varphi} | lm \rangle &= (-1)^{F+l+k} \Pi_{LlkF} / \Pi_{l'} \sqrt{\frac{1}{4\pi}} \\ &\times C_{lmF\varphi}^{l'm'} \sum_{L'} (-1)^{L'} \Pi_{L'} \sqrt{L'(L'+1)} C_{l0L0}^{L'0} C_{L'010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ l' & l & L' \end{matrix} \right\}, \quad (13.2.131) \end{aligned}$$

$$\begin{aligned} \langle l' || \{ \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{L}}_1 \}_k \otimes \mathbf{Y}_L \}_F || l \rangle &= (-1)^{F+l+k} \Pi_{LlkF} \sqrt{\frac{1}{4\pi}} \\ &\times \sum_{L'} (-1)^{L'} \Pi_{L'} \sqrt{L'(L'+1)} C_{l0L0}^{L'0} C_{L'010}^{l'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ l' & l & L' \end{matrix} \right\}. \end{aligned}$$

$$\begin{aligned} \langle l'm' | \{ (\hat{\nabla}_\Omega)_1 \otimes \hat{\mathbf{L}}_1 \}_{k\kappa} \hat{Y}_{L\nu} | lm \rangle &= \sum_{L'\nu'} (-1)^{l'+L'+k} \Pi_{kLl} / \Pi_{l'} \sqrt{\frac{L'(L'+1)}{4\pi}} \\ &\times C_{lmL\nu}^{L'\nu'} C_{L'\nu'k\kappa}^{l'm'} C_{l0L0}^{L'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\} \langle l' || (\hat{\nabla}_\Omega)_1 || L' \rangle, \quad (13.2.132) \end{aligned}$$

$$\begin{aligned} \langle l'm' | \{ \{ (\hat{\nabla}_\Omega)_1 \otimes \hat{\mathbf{L}}_1 \}_k \otimes \hat{\mathbf{Y}}_L \}_{F\varphi} | lm \rangle &= (-1)^{F+l+k} \Pi_{kLlF} / \Pi_{l'} \sqrt{\frac{1}{4\pi}} \\ &\times C_{lmF\varphi}^{l'm'} \sum_{L'} (-1)^{L'} \Pi_{L'} \sqrt{L'(L'+1)} C_{l0L0}^{L'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ l' & l & L' \end{matrix} \right\} \langle l' || (\hat{\nabla}_\Omega)_1 || L' \rangle, \quad (13.2.133) \end{aligned}$$

$$\begin{aligned} \langle l' || \{ \{ (\hat{\nabla}_\Omega)_1 \otimes \hat{\mathbf{L}}_1 \}_k \otimes \hat{\mathbf{Y}}_L \}_F || l \rangle &= (-1)^{F+l+k} \Pi_{kLlF} \sqrt{\frac{1}{4\pi}} \\ &\times \sum_{L'} (-1)^{L'} \Pi_{L'} \sqrt{L'(L'+1)} C_{l0L0}^{L'0} \left\{ \begin{matrix} 1 & 1 & k \\ l' & L' & L' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ l' & l & L' \end{matrix} \right\} \langle l' || (\hat{\nabla}_\Omega)_1 || L' \rangle. \end{aligned}$$

$$\begin{aligned} \langle l's'J'M' | \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{S}}_1 \}_{k\kappa} \hat{Y}_{L\nu} | lsJM \rangle &= (-1)^{l+s+J} \delta_{ss'} \Pi_{LlJks} \sqrt{\frac{s(s+1)}{4\pi}} \\ &\times \sum_{J_1L_1M_1} \Pi_{J_1L_1} C_{JM L\nu}^{J_1M_1} C_{J_1M_1k\kappa}^{J'M'} C_{L0l0}^{L_10} C_{L_1010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}, \quad (13.2.134) \end{aligned}$$

$$\begin{aligned} \langle l's'J'M' | \{ \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{S}}_1 \}_k \otimes \mathbf{Y}_L \}_{F\varphi} | lsJM \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} C_{JM F\varphi}^{J'M'} \\ &\times \Pi_{LJlFs} \sqrt{\frac{s(s+1)}{4\pi}} \sum_{J_1L_1} \Pi_{L_1J_1J_1} \\ &\times C_{L0l0}^{L_10} C_{L_1010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}, \quad (13.2.135) \end{aligned}$$

$$\begin{aligned} \langle l's'J' || \{ \{ \hat{\mathbf{n}}_1 \otimes \hat{\mathbf{S}}_1 \}_k \otimes \mathbf{Y}_L \}_F || lsJ \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} \Pi_{LJlFsJ'} \sqrt{\frac{s(s+1)}{4\pi}} \\ &\times \sum_{J_1L_1} \Pi_{L_1J_1J_1} C_{L0l0}^{L_10} C_{L_1010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}. \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{l+s+J} \delta_{ss'} \Pi_{LlJk} \sqrt{\frac{s(s+1)}{4\pi}} \\ &\times \sum_{J_1 L_1 M_1} \Pi_{J_1} C_{JM L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} C_{L 0 l 0}^{L_1 0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\} \langle l' | \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle, \end{aligned} \quad (13.2.136)$$

$$\begin{aligned} \langle l' s' J' M' | \{ \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{S}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} C_{JM F \varphi}^{J' M'} \\ &\times \Pi_{LlJkFs} \sqrt{\frac{s(s+1)}{4\pi}} \sum_{J_1 L_1} (2J_1 + 1) C_{L 0 l 0}^{L_1 0} \\ &\times \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\} \langle l' | \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle, \end{aligned} \quad (13.2.137)$$

$$\begin{aligned} \langle l' s' J' | \| \{ \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{S}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F \| l s J \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} \Pi_{LlJkFsJ'} \sqrt{\frac{s(s+1)}{4\pi}} \\ &\times \sum_{J_1 L_1} (2J_1 + 1) C_{L 0 l 0}^{L_1 0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\} \langle l' | \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle. \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{2s+J-J'+k+L} \delta_{ss'} \\ &\times \Pi_{LlJk} \sqrt{\frac{1}{4\pi}} \sum_{L_1 J_1 M_1} \sqrt{J_1(J_1+1)} \Pi_{L_1 J_1 J_1} \\ &\times C_{JM L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} C_{L 0 l 0}^{L_1 0} C_{L_1 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & s & J_1 \\ J' & 1 & l' \end{matrix} \right\}, \end{aligned} \quad (13.2.138)$$

$$\begin{aligned} \langle l' s' J' M' | \{ \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{F+k+L} \delta_{ss'} C_{JM F \varphi}^{J' M'} \\ &\times \Pi_{FLlJk} \sqrt{\frac{1}{4\pi}} \sum_{L_1 J_1} \sqrt{J_1(J_1+1)} \Pi_{L_1} \\ &\times (2J_1 + 1)^{\frac{3}{2}} C_{L 0 l 0}^{L_1 0} C_{L_1 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & s & J_1 \\ J' & 1 & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}, \end{aligned} \quad (13.2.139)$$

$$\begin{aligned} \langle l' s' J' | \| \{ \{ \widehat{\mathbf{n}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F \| l s J \rangle &= (-1)^{F+k+L} \delta_{ss'} \Pi_{FLlJkJ'} \sqrt{\frac{1}{4\pi}} \\ &\times \sum_{L_1 J_1} \sqrt{J_1(J_1+1)} \Pi_{L_1} (2J_1 + 1)^{\frac{3}{2}} C_{L 0 l 0}^{L_1 0} C_{L_1 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & s & J_1 \\ J' & 1 & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}. \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{l+l'-J-J'+k+1} \delta_{ss'} \Pi_{LlJk} \frac{1}{\sqrt{4\pi}} \\ &\times \sum_{L_1 J_1} \sqrt{J_1(J_1+1)} \Pi_{J_1}^2 C_{JM L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} C_{L 0 l 0}^{L_1 0} \langle l' | \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J' & 1 \\ l' & L_1 & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\}, \end{aligned} \quad (13.2.140)$$

$$\begin{aligned} \langle l' s' J' M' | \{ \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{l+l'+F+k+1} C_{JM F \varphi}^{J' M'} \\ &\times \Pi_{LlJkF} \sqrt{\frac{1}{4\pi}} \sum_{L_1 J_1} \sqrt{J_1(J_1+1)} (2J_1 + 1)^{\frac{3}{2}} C_{L 0 l 0}^{L_1 0} \\ &\times \langle l' | \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J' & 1 \\ l' & L_1 & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}, \end{aligned} \quad (13.2.141)$$

$$\begin{aligned}
\langle l' s' J' \| \{ \{ (\widehat{\nabla}_\Omega)_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F \| l s J \rangle &= (-1)^{l+l'+F+k+1} \Pi_{LlJkFJ'} \sqrt{\frac{1}{4\pi}} \\
&\times \sum_{L_1 J_1} \sqrt{J_1(J_1+1)(2J_1+1)^{\frac{3}{2}}} C_{L_0 l 0}^{L_1 0} \langle l' \| (\widehat{\nabla}_\Omega)_1 \| L_1 \rangle \left\{ \begin{matrix} l & s & J \\ J_1 & L & L_1 \end{matrix} \right\} \left\{ \begin{matrix} J_1 & J' & 1 \\ l' & L_1 & s \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}. \\
\langle l' s' J' M' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{S}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{l+s+J'} \delta_{ss'} C_{L_0 l 0}^{l' 0} \Pi_{Llksl'J} \sqrt{\frac{s(s+1)l'(l'+1)}{4\pi}} \\
&\times \sum_{J_1 M_1} \Pi_{J_1} C_{J_1 M_1 L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} l' & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}, \quad (13.2.142)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' M' | \{ \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{S}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} C_{J_1 M_1 F \varphi}^{J' M'} C_{L_0 l 0}^{l' 0} \\
&\times \Pi_{LlJkFsl'J} \sqrt{\frac{s(s+1)l'(l'+1)}{4\pi}} \\
&\times \sum_{J_1} (2J_1+1) \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l' & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}, \quad (13.2.143)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' \| \{ \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{S}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F \| l s J \rangle &= (-1)^{l+s-J'-F} \delta_{ss'} C_{L_0 l 0}^{l' 0} \Pi_{LlJkFsl'J} \sqrt{\frac{s(s+1)l'(l'+1)}{4\pi}} \\
&\times \sum_{J_1} (2J_1+1) \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l' & 1 & l' \\ s & 1 & s \\ J_1 & k & J' \end{matrix} \right\}.
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' M' | \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{l+l'+J+J'+k+1} \delta_{ss'} C_{L_0 l 0}^{l' 0} \\
&\times \Pi_{LlJkl'} \sqrt{\frac{l'(l'+1)}{4\pi}} \sum_{J_1 M_1} (2J_1+1) \sqrt{J_1(J_1+1)} C_{J_1 M_1 L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} \\
&\times \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l' & s & J_1 \\ J' & 1 & l' \end{matrix} \right\}, \quad (13.2.144)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' M' | \{ \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{l+l'-F+k+1} \delta_{ss'} C_{L_0 l 0}^{l' 0} C_{J_1 M_1 F \varphi}^{J' M'} \\
&\times \Pi_{LlJkFl'} \sqrt{\frac{l'(l'+1)}{4\pi}} \sum_{J_1} \sqrt{J_1(J_1+1)(2J_1+1)^{\frac{3}{2}}} \\
&\times \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l' & s & J_1 \\ J' & 1 & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}, \quad (13.2.145)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' \| \{ \{ \widehat{\mathbf{L}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F \| l s J \rangle &= (-1)^{l+l'-F+k+1} \delta_{ss'} C_{L_0 l 0}^{l' 0} \Pi_{LlJkFl'J'} \sqrt{\frac{l'(l'+1)}{4\pi}} \\
&\times \sum_{J_1} \sqrt{J_1(J_1+1)(2J_1+1)^{\frac{3}{2}}} \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l' & s & J_1 \\ J' & 1 & l' \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}.
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' M' | \{ \widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1 \}_{k\kappa} \widehat{Y}_{L\nu} | l s J M \rangle &= (-1)^{J+L+k+1} \delta_{ss'} C_{L_0 l 0}^{l' 0} \\
&\times \Pi_{LlJks} \sqrt{\frac{s(s+1)}{4\pi}} \sum_{J_1 M_1} C_{J_1 M_1 L \nu}^{J_1 M_1} C_{J_1 M_1 k \kappa}^{J' M'} \sqrt{J_1(J_1+1)(2J_1+1)} \\
&\times (-1)^{J_1} \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} s & l' & J_1 \\ J' & 1 & s \end{matrix} \right\}, \quad (13.2.146)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' M' | \{ \{ \widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_{F\varphi} | l s J M \rangle &= (-1)^{L+F-J'+k+1} \delta_{ss'} C_{L 0 1 0}^{l' 0} C_{J M F \varphi}^{J' M'} \\
&\times \Pi_{LlJkFs} \sqrt{\frac{s(s+1)}{4\pi}} \sum_{J_1} \sqrt{J_1(J_1+1)(2J_1+1)}^{\frac{3}{2}} \\
&\times (-1)^{J_1} \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} s & l' & J_1 \\ J' & 1 & s \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}, \quad (13.2.147)
\end{aligned}$$

$$\begin{aligned}
\langle l' s' J' | \{ \{ \widehat{\mathbf{S}}_1 \otimes \widehat{\mathbf{J}}_1 \}_k \otimes \widehat{\mathbf{Y}}_L \}_F | l s J \rangle &= (-1)^{L+F-J'+k+1} \delta_{ss'} C_{L 0 1 0}^{l' 0} \Pi_{LlJkFsJ'} \sqrt{\frac{s(s+1)}{4\pi}} \\
&\times \sum_{J_1} \sqrt{J_1(J_1+1)(2J_1+1)}^{\frac{3}{2}} (-1)^{J_1} \left\{ \begin{matrix} l & s & J \\ J_1 & L & l' \end{matrix} \right\} \left\{ \begin{matrix} 1 & 1 & k \\ J' & J_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} s & l' & J_1 \\ J' & 1 & s \end{matrix} \right\} \left\{ \begin{matrix} L & k & F \\ J' & J & J_1 \end{matrix} \right\}.
\end{aligned}$$

The commutators of the operators in Eqs. (13.2.128)-(13.2.147) may be represented in a more general form as

$$\Re_{k\kappa, L\nu}(\{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_{k\kappa}, \widehat{\mathbf{Y}}_{L\nu}) = \sum_{\alpha\beta} C_{a\alpha b\beta}^{k\kappa} \{ \Re_{a\alpha, L\nu}(\widehat{P}_{a\alpha}, \widehat{Y}_{L\nu}) \widehat{Q}_{b\beta} + \widehat{P}_{a\alpha} \Re_{b\beta, L\nu}(\widehat{Q}_{b\beta}, \widehat{Y}_{L\nu}) \}, \quad (13.2.148)$$

$$\Re_{F\varphi}^{kL}(\{ \widehat{\mathbf{P}}_a \otimes \widehat{\mathbf{Q}}_b \}_k, \widehat{\mathbf{Y}}_L) = \sum_{\alpha\beta\kappa\nu} C_{a\alpha b\beta}^{k\kappa} C_{k\kappa L\nu}^{F\varphi} \{ \Re_{a\alpha, L\nu}(\widehat{P}_{a\alpha}, \widehat{Y}_{L\nu}) \widehat{Q}_{b\beta} + \widehat{P}_{a\alpha} \Re_{b\beta, L\nu}(\widehat{Q}_{b\beta}, \widehat{Y}_{L\nu}) \}. \quad (13.2.149)$$

The evaluation of matrix elements of these commutators may be performed using the foregoing equations.

13.2.9 Matrix Elements of Some Scalar and Vector Products

Using Eq. (13.2.33), one can easily obtain matrix elements of scalar and vector products of the operators $\widehat{\mathbf{n}}$ and $\widehat{\nabla}_\Omega$

$$\langle l' m' | (\widehat{\mathbf{n}} \cdot \widehat{\nabla}_\Omega) | l m \rangle = 0, \quad (13.2.150)$$

$$\langle l' m' | (\widehat{\mathbf{n}} \times \widehat{\nabla}_\Omega)_\mu | l m \rangle = i\sqrt{l(l+1)} C_{l m 1 \mu}^{l' m'} \delta_{ll'}. \quad (13.2.151)$$

$$\langle l' | (\widehat{\mathbf{n}} \times \widehat{\nabla}_\Omega) | l \rangle = i\sqrt{l(l+1)} \Pi_{l'} \delta_{ll'}.$$

Equation (13.2.150) follows from the orthogonality of the vectors in question. Equation (13.2.151) may also be obtained by taking into account the fact that

$$[\widehat{\mathbf{n}} \times \widehat{\nabla}_\Omega] = i\widehat{\mathbf{L}}.$$

Matrix elements of scalar and vector products of the operator $\widehat{\mathbf{J}}$ and the operators $\widehat{\mathbf{n}}$ and $\widehat{\nabla}_\Omega$ may be derived from Eqs. (13.2.52)-(13.2.56) and (13.2.61)-(13.2.65)

$$\langle l' s' J' M' | (\widehat{\mathbf{n}} \cdot \widehat{\mathbf{J}}) | l s J M \rangle = \delta_{s's} \delta_{J'J} \delta_{M'M} (-1)^{s+l'+J+1} \Pi_{Jl} \sqrt{J(J+1)} C_{l 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J & 1 & l' \end{matrix} \right\}, \quad (13.2.152)$$

$$\langle l' s' J' | (\widehat{\mathbf{n}} \cdot \widehat{\mathbf{J}}) | l s J \rangle = \delta_{s's} \delta_{J'J} (-1)^{s+l'+J+1} \Pi_{Jl} \sqrt{J(J+1)} C_{l 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J & 1 & l' \end{matrix} \right\},$$

and

$$\begin{aligned}
\langle l' s' J' M' | [\widehat{\mathbf{n}} \times \widehat{\mathbf{J}}]_\kappa | l s J M \rangle &= \frac{i}{2} (-1)^{J+s+l'} \{ J'(J'+1) - J(J+1) - 2 \} \Pi_{Jl} \\
&\quad C_{l 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M 1 \kappa}^{J' M'}, \quad (13.2.153)
\end{aligned}$$

$$\langle l' s' J' | [\widehat{\mathbf{n}} \times \widehat{\mathbf{J}}] | l s J \rangle = \frac{i}{2} (-1)^{J+s+l'} \{ J'(J'+1) - J(J+1) - 2 \} \Pi_{JlJ'} C_{l 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}.$$

The scalar product of the operators $\widehat{\mathbf{n}}$ and $\widehat{\mathbf{J}}$ is commutative. At the same time, the vector product of these operators is not commutative, i.e.,

$$\begin{aligned}
&\langle l' s' J' M' | [\widehat{\mathbf{J}} \times \widehat{\mathbf{n}}]_\kappa | l s J M \rangle \\
&= \frac{i}{2} (-1)^{J+s+l'+1} \{ J'(J'+1) - J(J+1) + 2 \} \Pi_{Jl} C_{l 0 1 0}^{l' 0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M 1 \kappa}^{J' M'}, \quad (13.2.154)
\end{aligned}$$

$$\langle l' s' J' \| [\hat{\mathbf{J}} \times \hat{\mathbf{n}}] \| l s J \rangle = \frac{i}{2} (-1)^{J+s+l'+1} \{ J'(J'+1) - J(J+1) + 2 \} \Pi_{Jl'J'} C_{l'010}^{l'0} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}.$$

In a similar manner we get

$$\begin{aligned} \langle l' s' J' M' | (\hat{\mathbf{V}}_{\Omega} \cdot \hat{\mathbf{J}}) | l s J M \rangle &= \delta_{J'J} \delta_{s's} \delta_{M'M} (-1)^{s+l'+J} \Pi_{Jl'} \sqrt{J(J+1)} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J & 1 & l' \end{matrix} \right\}, \end{aligned} \quad (13.2.155)$$

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{V}}_{\Omega} \times \hat{\mathbf{J}}]_{\kappa} | l s J M \rangle &= \frac{i}{2} (-1)^{J+s+l'+1} \Pi_{Jl'} \{ J'(J'+1) - J(J+1) - 2 \} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.156)$$

$$\begin{aligned} \langle l' s' J' \| (\hat{\mathbf{V}}_{\Omega} \cdot \hat{\mathbf{J}}) \| l s J \rangle &= \delta_{J'J} \delta_{s's} (-1)^{s+l'+J} \Pi_{Jl'} \sqrt{J(J+1)} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J & 1 & l' \end{matrix} \right\}, \end{aligned}$$

$$\begin{aligned} \langle l' s' J' \| [\hat{\mathbf{V}}_{\Omega} \times \hat{\mathbf{J}}] \| l s J \rangle &= \frac{i}{2} (-1)^{J+s+l'+1} \Pi_{Jl'} \Pi_{J'} \{ J'(J'+1) - J(J+1) - 2 \} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}. \end{aligned}$$

As in the foregoing case, the scalar product of the vector operators $\hat{\mathbf{V}}_{\Omega}$ and $\hat{\mathbf{J}}$ is commutative, while the vector product is not commutative. Therefore,

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{J}} \times \hat{\mathbf{V}}_{\Omega}]_{\kappa} | l s J M \rangle &= \frac{i}{2} (-1)^{J+s+l'} \{ J'(J'+1) - J(J+1) + 2 \} \Pi_{Jl'} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J M 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.157)$$

$$\begin{aligned} \langle l' s' J' \| [\hat{\mathbf{J}} \times \hat{\mathbf{V}}_{\Omega}] \| l s J \rangle &= \frac{i}{2} (-1)^{J+s+l'} \{ J'(J'+1) - J(J+1) + 2 \} \Pi_{Jl'J'} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}. \end{aligned}$$

The matrix elements of scalar and vector products of the operator $\hat{\mathbf{L}}$ and the operators $\hat{\mathbf{n}}$ and $\hat{\mathbf{V}}_{\Omega}$ may be obtained from Eqs. (13.2.73)-(13.2.77) and (13.2.82)-(13.2.85). In the case of scalar products we get

$$\begin{aligned} \langle l' m' | (\hat{\mathbf{n}} \cdot \hat{\mathbf{L}}) | l m \rangle &= 0, \\ \langle l' m' | (\hat{\mathbf{V}}_{\Omega} \cdot \hat{\mathbf{L}}) | l m \rangle &= 0. \end{aligned} \quad (13.2.158)$$

These results follow from the orthogonality of the vectors $\hat{\mathbf{n}}$, $\hat{\mathbf{V}}_{\Omega}$ and $\hat{\mathbf{L}}$. The vector products $[\hat{\mathbf{n}} \times \hat{\mathbf{L}}]$ and $[\hat{\mathbf{V}}_{\Omega} \times \hat{\mathbf{L}}]$ are not commutative. Hence,

$$\langle l' m' | [\hat{\mathbf{n}} \times \hat{\mathbf{L}}]_{\kappa} | l m \rangle = -\frac{i}{2} [l'(l'+1) - l(l+1) - 2] \frac{\Pi_l}{\Pi_{l'}} C_{l'010}^{l'0} C_{l m 1 \kappa}^{l' m'} = i \langle l' m' | (\hat{\mathbf{V}}_{\Omega})_{1\kappa} | l m \rangle, \quad (13.2.159)$$

$$\langle l' m' | [\hat{\mathbf{L}} \times \hat{\mathbf{n}}]_{\kappa} | l m \rangle = \frac{i}{2} [l'(l'+1) - l(l+1) + 2] \frac{\Pi_l}{\Pi_{l'}} C_{l'010}^{l'0} C_{l m 1 \kappa}^{l' m'}, \quad (13.2.160)$$

$$\langle l' \| [\hat{\mathbf{n}} \times \hat{\mathbf{L}}] \| l \rangle = -\frac{i}{2} [l'(l'+1) - l(l+1) - 2] \Pi_l C_{l'010}^{l'0},$$

$$\langle l' \| [\hat{\mathbf{L}} \times \hat{\mathbf{n}}] \| l \rangle = \frac{i}{2} [l'(l'+1) - l(l+1) + 2] \Pi_l C_{l'010}^{l'0}$$

and also

$$\begin{aligned} \langle l'm' | \{ \widehat{\mathbf{V}}_{\Omega} \times \widehat{\mathbf{L}} \}_{\kappa} | lm \rangle \\ = \frac{i}{2} [l'(l'+1) - l(l+1) - 2] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] C_{l'm'1\kappa}^{l' m'} \end{aligned} \quad (13.2.161)$$

$$\begin{aligned} \langle l'm' | \{ \widehat{\mathbf{L}} \times \widehat{\mathbf{V}}_{\Omega} \}_{\kappa} | lm \rangle \\ = -\frac{i}{2} [l'(l'+1) - l(l+1) + 2] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] C_{l'm'1\kappa}^{l' m'} \end{aligned} \quad (13.2.162)$$

$$\langle l' || \{ \widehat{\mathbf{V}}_{\Omega} \times \widehat{\mathbf{L}} \} || l \rangle = \frac{i}{2} [l'(l'+1) - l(l+1) - 2] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \Pi_{l'},$$

$$\langle l' || \{ \widehat{\mathbf{L}} \times \widehat{\mathbf{V}}_{\Omega} \} || l \rangle = -\frac{i}{2} [l'(l'+1) - l(l+1) + 2] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \Pi_{l'}.$$

The matrix elements of scalar and vector products of the operators $\widehat{\mathbf{L}}$ and $\widehat{\mathbf{J}}$ may be derived from Eqs. (13.2.86)-(13.2.89). The scalar product is commutative:

$$\begin{aligned} \langle l's'J'M' | (\widehat{\mathbf{L}} \cdot \widehat{\mathbf{J}}) | lsJM \rangle &= \delta_{l'l} \delta_{s's} \delta_{J'J} \delta_{M'M} \frac{J(J+1) + l(l+1) - s(s+1)}{2}, \\ \langle l's'J' || (\widehat{\mathbf{L}} \cdot \widehat{\mathbf{J}}) || lsJ \rangle &= \delta_{l'l} \delta_{s's} \delta_{J'J} \Pi_J \frac{J(J+1) + l(l+1) - s(s+1)}{2}. \end{aligned} \quad (13.2.163)$$

The vector product is not commutative:

$$\begin{aligned} \langle l's'J'M' | \{ \widehat{\mathbf{L}} \times \widehat{\mathbf{J}} \}_{\kappa} | lsJM \rangle &= \frac{i}{2} (-1)^{s+l+J+1} \delta_{l'l} \delta_{s's} [J(J+1) - J'(J'+1) + 2] \\ &\times \Pi_{Jl} \sqrt{l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\} C_{J'M'1\kappa}^{J' M'}, \end{aligned} \quad (13.2.164)$$

$$\langle l's'J' || \{ \widehat{\mathbf{L}} \times \widehat{\mathbf{J}} \} || lsJ \rangle = \frac{i}{2} (-1)^{s+l+J+1} \delta_{l'l} \delta_{s's} [J(J+1) - J'(J'+1) + 2] \Pi_{JlJ'} \sqrt{l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l \end{matrix} \right\}$$

$$\begin{aligned} \langle l's'J'M' | [\widehat{\mathbf{J}} \times \widehat{\mathbf{L}}]_{\kappa} | lsJM \rangle &= \frac{i}{2} (-1)^{s+l+J+1} \delta_{l'l} \delta_{s's} [J'(J'+1) - J(J+1) + 2] \\ &\times \Pi_{Jl} \sqrt{l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\} C_{J'M'1\kappa}^{J' M'}, \end{aligned} \quad (13.2.165)$$

$$\langle l's'J' || [\widehat{\mathbf{J}} \times \widehat{\mathbf{L}}] || lsJ \rangle = \frac{i}{2} (-1)^{s+l+J+1} \delta_{l'l} \delta_{s's} [J'(J'+1) - J(J+1) + 2] \times \Pi_{JlJ'} \sqrt{l(l+1)} \left\{ \begin{matrix} l & s & J \\ J' & 1 & l' \end{matrix} \right\}.$$

Matrix elements of scalar and vector products of the spin operator $\widehat{\mathbf{S}}$ and the operators $\widehat{\mathbf{n}}$, $\widehat{\mathbf{V}}_{\Omega}$ and $\widehat{\mathbf{L}}$ (which commutative with $\widehat{\mathbf{S}}$) may be found from Eqs. (13.2.95)-(13.2.99):

$$\langle l's'J'M' | (\widehat{\mathbf{n}} \cdot \widehat{\mathbf{S}}) | lsJM \rangle = (-1)^{l+s+J} \delta_{JJ'} \delta_{MM'} \delta_{ss'} \Pi_{sl} \sqrt{s(s+1)} C_{l'010}^{l' 0} \left\{ \begin{matrix} 1 & s & s \\ J & l & l' \end{matrix} \right\}, \quad (13.2.166)$$

$$\langle l's'J' || (\widehat{\mathbf{n}} \cdot \widehat{\mathbf{S}}) || lsJ \rangle = (-1)^{l+s+J} \delta_{JJ'} \delta_{ss'} \Pi_{slJ} \sqrt{s(s+1)} C_{l'010}^{l' 0} \left\{ \begin{matrix} 1 & s & s \\ J & l & l' \end{matrix} \right\}.$$

$$\begin{aligned} \langle l's'J'M' | (\widehat{\mathbf{V}}_{\Omega} \cdot \widehat{\mathbf{S}}) | lsJM \rangle &= (-1)^{l+s+J+1} \delta_{JJ'} \delta_{MM'} \delta_{ss'} \Pi_{sl'} \sqrt{s(s+1)} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} 1 & s & s \\ J & l & l' \end{matrix} \right\}, \end{aligned} \quad (13.2.167)$$

$$\begin{aligned} \langle l's'J' || (\widehat{\mathbf{V}}_{\Omega} \cdot \widehat{\mathbf{S}}) || lsJ \rangle &= (-1)^{l+s+J+1} \delta_{JJ'} \delta_{ss'} \Pi_{sl'J} \sqrt{s(s+1)} \\ &\times \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] \left\{ \begin{matrix} 1 & s & s \\ J & l & l' \end{matrix} \right\} \end{aligned}$$

$$\langle l' s' J' M' | (\hat{\mathbf{L}} \cdot \hat{\mathbf{S}}) | l s J M \rangle = \delta_{ll'} \delta_{ss'} \delta_{JJ'} \delta_{MM'} \frac{1}{2} [J(J+1) - l(l+1) - s(s+1)], \quad (13.2.168)$$

$$\langle l' s' J' | (\hat{\mathbf{L}} \cdot \hat{\mathbf{S}}) | l s J \rangle = \delta_{ll'} \delta_{ss'} \delta_{JJ'} \Pi_J \frac{1}{2} [J(J+1) - l(l+1) - s(s+1)].$$

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{n}} \times \hat{\mathbf{S}}]_{\kappa} | l s J M \rangle &= i(-1)^{l'+J+s} \delta_{ss'} \Pi_{Jl} C_{l010}^{l'0} \\ &\times \frac{1}{2} [(J' - l')(J' + l' + 1) - (J - l)(J + l + 1)] \left\{ \begin{matrix} J' & l' & s \\ l & J & 1 \end{matrix} \right\} C_{J M' 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.169)$$

$$\begin{aligned} \langle l' s' J' | [\hat{\mathbf{n}} \times \hat{\mathbf{S}}] | l s J \rangle &= i(-1)^{l'+J+s} \delta_{ss'} \Pi_{Jl} C_{l010}^{l'0} \\ &\times \frac{1}{2} [(J' - l')(J' + l' + 1) - (J - l)(J + l + 1)] \left\{ \begin{matrix} J' & l' & s \\ l & J & 1 \end{matrix} \right\} \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | [(\hat{\nabla}_{\Omega}) \times \hat{\mathbf{S}}]_{\kappa} | l s J M \rangle &= i(-1)^{l'+J+s+1} \delta_{ss'} \Pi_{Jl'} \left\{ \begin{matrix} J' & l' & s \\ l & J & 1 \end{matrix} \right\} \\ &\times \frac{1}{2} [(J' - l')(J' + l' + 1) - (J - l)(J + l + 1)] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right] C_{J M' 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.170)$$

$$\begin{aligned} \langle l' s' J' | [(\hat{\nabla}_{\Omega}) \times \hat{\mathbf{S}}] | l s J \rangle &= i(-1)^{l'+J+s+1} \delta_{ss'} \Pi_{Jl' J'} \left\{ \begin{matrix} J' & l' & s \\ l & J & 1 \end{matrix} \right\} \\ &\times \frac{1}{2} [(J' - l')(J' + l' + 1) - (J - l)(J + l + 1)] \left[l \sqrt{\frac{l+1}{2l+3}} \delta_{l'l+1} + (l+1) \sqrt{\frac{l}{2l-1}} \delta_{l'l-1} \right]. \end{aligned}$$

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{L}}_1 \times \hat{\mathbf{S}}_1]_{\kappa} | l s J M \rangle &= i(-1)^{l'+J+s} \Pi_{Jl} \sqrt{l(l+1)} \delta_{ll'} \delta_{ss'} \\ &\times \frac{1}{2} [(J' - l)(J' + l + 1) - (J - l)(J + l + 1)] \left\{ \begin{matrix} J' & l & s \\ l & J & 1 \end{matrix} \right\} C_{J M' 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.171)$$

$$\begin{aligned} \langle l' s' J' | [\hat{\mathbf{L}}_1 \times \hat{\mathbf{S}}_1] | l s J \rangle &= i(-1)^{l'+J+s} \Pi_{Jl J'} \sqrt{l(l+1)} \delta_{ll'} \delta_{ss'} \\ &\times \frac{1}{2} [(J' - l)(J' + l + 1) - (J - l)(J + l + 1)] \left\{ \begin{matrix} J' & l & s \\ l & J & 1 \end{matrix} \right\}. \end{aligned}$$

The scalar product of the operators $\hat{\mathbf{S}}$ and $\hat{\mathbf{J}}$ is commutative and its matrix elements are determined by

$$\langle l' s' J' M' | (\hat{\mathbf{S}} \cdot \hat{\mathbf{J}}) | l s J M \rangle = \delta_{ll'} \delta_{ss'} \delta_{JJ'} \delta_{MM'} \frac{1}{2} [J(J+1) + s(s+1) - l(l+1)], \quad (13.2.172)$$

$$\langle l' s' J' | (\hat{\mathbf{S}} \cdot \hat{\mathbf{J}}) | l s J \rangle = \delta_{ll'} \delta_{ss'} \delta_{JJ'} \Pi_J \frac{1}{2} [J(J+1) + s(s+1) - l(l+1)].$$

The vector product of the same operators is not commutative. The matrix elements of this product are given by

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{S}} \times \hat{\mathbf{J}}]_{\kappa} | l s J M \rangle &= \frac{i}{2} (-1)^{l'+s+J'+1} \delta_{ss'} \delta_{ll'} [J(J+1) - J'(J'+1) + 2] \\ &\times \Pi_{Js} \sqrt{s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\} C_{J M' 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.173)$$

$$\begin{aligned} \langle l' s' J' M' | [\hat{\mathbf{J}} \times \hat{\mathbf{S}}]_{\kappa} | l s J M \rangle &= \frac{i}{2} (-1)^{l'+s+J'+1} \delta_{ll'} \delta_{ss'} [J'(J'+1) - J(J+1) + 2] \\ &\times \Pi_{Js} \sqrt{s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\} C_{J M' 1 \kappa}^{J' M'}, \end{aligned} \quad (13.2.174)$$

$$\langle l' s' J' | [\hat{\mathbf{S}} \times \hat{\mathbf{J}}] | l s J \rangle = \frac{i}{2} (-1)^{l'+s+J'+1} \delta_{ss'} \delta_{ll'} [J(J+1) - J'(J'+1) + 2] \Pi_{Js J'} \sqrt{s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\},$$

$$\langle l' s' J' | [\hat{\mathbf{J}} \times \hat{\mathbf{S}}] | l s J \rangle = \frac{i}{2} (-1)^{l'+s+J'+1} \delta_{ll'} \delta_{ss'} [J'(J'+1) - J(J+1) + 2] \Pi_{Js J'} \sqrt{s(s+1)} \left\{ \begin{matrix} s & l & J \\ J' & 1 & s \end{matrix} \right\}.$$

Chapter 14

Computer generation of numerical and algebraic expressions

The algebraic tables of Secs. 8.12, 9.11 and 10.11, and the numerical tables of Secs. 8.13 and 10.12, together with Tables 9.9–9.11, may be reproduced entry by entry by computer algebra. This chapter records the method used, so that any individual entry can be regenerated and checked rather than taken on trust. The programs are listed in Sec. 14.5.

14.1 Scope

Each algebraic table fixes one small angular momentum and leaves the remaining arguments symbolic: b in the Clebsch-Gordan tables, d in the $6j$ tables, and the triad $(\alpha\beta\gamma)$ in the $9j$ tables. The corresponding sum is then finite with a number of terms bounded by that fixed argument, and is therefore a finite symbolic computation. Nothing in the method is restricted to the ranges printed here; the same programs generate entries beyond them, at greater cost.

14.2 Algebraic generation

14.2.1 The Clebsch-Gordan coefficients

For fixed b and β , and with $c = a + k$, the summation index of the Van der Waerden-Racah sum of Sec. 8.2.1 runs over at most $2b + 1$ values, so the coefficient is a short symbolic sum. The difficulty is that its terms contain factorials of $a \pm \alpha$, which are not polynomial in a . It is removed by the change of variable

$$p = a + \alpha, \quad q = a - \alpha, \quad (14.2.1)$$

both of which are non-negative *integers* even when a and α are half-integral. Every factorial argument is then an explicit integer offset from p , q or $p + q = 2a$, and each such factorial may be replaced by a rational function through

$$\frac{(x+m)!}{x!} = \begin{cases} (x+1)(x+2)\cdots(x+m), & m \geq 0, \\ [x(x-1)\cdots(x+m+1)]^{-1}, & m < 0. \end{cases} \quad (14.2.2)$$

The factor $p!q!$ produced by the sum cancels exactly against the one under the square root, and the coefficient takes the form

$$C_{a\alpha b\beta}^{c\gamma} = T(p, q) [W(p, q)]^{1/2}, \quad (14.2.3)$$

with T and W explicit rational functions. No general-purpose simplification of factorials or gamma functions is required at any stage.

Eq. (14.2.3) does not fix the presentation, since any factor may be moved under the radical as its square. The tables of Sec. 8.12 absorb every factor that is of one sign throughout the physical domain $|\gamma| \leq c$, such as $(c + \gamma)$ or $(c - \gamma + 1)$, and leave outside only those that change sign, such as γ itself. Applying that rule reproduces the printed entries.

14.2.2 The 6j symbols

The analogous integer variables for the 6j symbols are the triangle deficits

$$u = s - 2a, \quad v = s - 2b, \quad w = s - 2c, \quad s = a + b + c = u + v + w. \quad (14.2.4)$$

Writing the summation index of Eq. (9.2.1) as $N = s + t$, and taking $e = c + n$ and $f = b + m$ as in the tables, the seven factorial arguments of that equation become

$$\begin{aligned} N - a - b - c &= t, & N - a - e - f &= t - n - m, \\ N - b - d - f &= v + t - d - m, & N - c - d - e &= w + t - d - n, \\ a + b + d + e - N &= d + n - t, & a + c + d + f - N &= d + m - t, \\ b + c + e + f - N &= u + n + m - t, \end{aligned} \quad (14.2.5)$$

each an explicit integer offset from t , u , v or w . The index t runs over the explicit finite range

$$\max(0, n + m) \leq t \leq \min(d + n, d + m), \quad (14.2.6)$$

so the sum has at most $2d + 1$ terms. Here the cancellation is complete: $\Delta(abc)$ contributes $[u! v! w! / (s + 1)!]^{1/2}$, the sum contributes $(s + 1)! / (u! v! w!)$, and $\Delta(aef)$ contributes the inverse, leaving

$$\left\{ \begin{matrix} a & b & c \\ d & e & f \end{matrix} \right\} = (-1)^s T(u, v, w) [W(u, v, w)]^{1/2} \quad (14.2.7)$$

with no symbolic factorial remaining. The phase arises because $(-1)^N = (-1)^s (-1)^t$, which is why every entry of Sec. 9.11 carries $(-1)^s$ or $(-1)^{s+1}$.

The factors of T and W are linear in a , b , c and are printed in the notation of the tables by reading off their coefficients: $(1, 1, 1)$ gives $s + k$, $(-1, 1, 1)$ gives $s - 2a + k$, $(0, 2, 0)$ gives $2b + k$, and so on. The factor left outside the radical is expressed as a polynomial in X by division, X being quadratic in a .

14.2.3 The 9j symbols

The 9j symbols of Sec. 10.11 are not given by a single sum of the above type. They are obtained from the sum over three 6j symbols, Eq. (10.2.20). For the tabulated symbols the triad (βax) restricts the intermediate momentum to

$$x = a + \xi, \quad -\beta \leq \xi \leq \beta, \quad (14.2.8)$$

an explicit sum of at most $2\beta + 1$ terms. Each of the three 6j symbols is carried by a permutation of its columns together with an interchange of the upper and lower arguments in two of them into the form

$$\left\{ \begin{matrix} A & B & C \\ d & C + n & B + m \end{matrix} \right\}$$

treated in Sec. 14.2.2, so the same machinery evaluates all three; only the deficits (14.2.4) need re-expressing in terms of those of (abc) .

The phases collapse. Summing the three factors $(-1)^{A+B+C}$ with the $(-1)^{2x}$ of Eq. (10.2.20) gives the exponent $4a + 4\xi + 2(a + b + c) + \beta + \mu$, and since $2a$, 2ξ and $a + b + c$ are integers every term is even except the last two:

$$\left\{ \begin{matrix} a + \lambda & b + \mu & c + \nu \\ a & b & c \\ \alpha & \beta & \gamma \end{matrix} \right\} = (-1)^{\beta + \mu} \sum_{\xi} (2a + 2\xi + 1) T_1 T_2 T_3 [W_1 W_2 W_3]^{1/2}, \quad (14.2.9)$$

independent of ξ and of a , b , c . The radicands of the individual terms differ by perfect squares of rational functions, so a common radical may be extracted and the whole reduces to a rational multiple of a single square root. That the ratios are perfect squares is tested, not assumed.

14.3 Numerical evaluation

Numerical values are obtained in exact rational arithmetic. Since every coefficient of Secs. 8.13 and 10.12, and of Tables 9.9–9.11, is a signed square root of a rational number, the value V is squared, V^2 is confirmed to be rational, and the entry is presented as

$$V = \pm \left(\frac{P}{Q} \right)^{1/2}, \quad (14.3.1)$$

the decimal being derived from the fraction rather than the reverse. The 12j symbols are evaluated from their defining single sums over four 6j symbols, Eqs. (10.13.6) and (10.13.26).

14.4 Verification

Each generator is checked against an independent implementation of the same coefficient at a large number of numerical points, at two levels: the sum itself, and the regrouping into the printed form, so that an error in a phase or in the choice of factors placed under the radical cannot pass unnoticed. Points are chosen asymmetrically as well as symmetrically, since a factor such as X or Z changes sign only when one momentum dominates the other two, and a symmetric sample would not detect a factor wrongly treated as of one sign. All entries of the tables cited in Sec. 14.1 agree. The relations of Sec. 10.13 are checked against each other in the same way.

14.5 The programs

The generators and their checks are written in Python using SymPy. Each generator produces a single entry, given the fixed argument and the row and column indices of the table:

```
gen_8_12_cg_tables.py --b 1 --beta=0 --k=0
gen_9_11_6j_tables.py --d 1 --m=0 --n=0
gen_10_11_9j_tables.py --alpha 3/2 --beta 3/2 --gamma 0 --lam=1/2 --mu=1/2 --nu=0
```

Omitting the row and column indices produces a whole table. The companion programs `check_8_12.py`, `check_9_11.py`, `check_10_11.py` and `check_10_13.py` perform the verification of Sec. 14.4. The same functions are available interactively in the two notebooks described in Sec. 14.6.

14.6 Use of the notebooks

Two Jupyter notebooks make the generators available without recourse to the command line. Both require only SymPy; if `ipywidgets` is also present the arguments are set from menus, and if it is not, each quantity is instead obtained by editing the arguments at the head of a cell and running it. Every result is displayed both as a typeset formula and as the corresponding SymPy expression, the latter being the form to copy if the entry is to be used in further computation.

14.6.1 Algebraic expressions

The notebook `vmk_algebraic_tables.ipynb` has one part for each of the three families of Sec. 14.2. In each, one selects the fixed argument and the row and column of the table:

```
clebsch(b=1, beta=0, k=0) # <a alpha, b beta / c gamma>, c = a+k
sixj(d=1, m=0, n=0) # {a b c; d e f}, e = c+n, f = b+m
ninej(1, 1, 0, lam=1, mu=0, nu=0) # alpha, beta, gamma, then lambda, mu, nu
```

A further cell in each part prints an entire table at once. Setting `show_source=True` adds the L^AT_EX source of the entry. The Clebsch-Gordan entries are recomputed as soon as an argument is changed; the $6j$ and $9j$ entries are not, since a single entry of the larger tables takes appreciably longer to construct, and are instead computed on request. A closing part runs the verification programs of Sec. 14.4 over a reduced range of arguments.

14.6.2 Numerical values

The notebook `vmk_numeric_tables.ipynb` evaluates the Clebsch-Gordan coefficients and the $6j$, $9j$ and $12j$ symbols at given numerical arguments, in the form (14.3.1):

```
clebsch(1, 0, 1, 0, 2) # gamma = alpha + beta is supplied
sixj(1, 1, 1, 1, 1, 1)
ninej(0.5, 0.5, 1, 0.5, 0.5, 1, 1, 1, 2)
twelvej1(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1) # eq (10.13.6)
twelvej2(1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1) # eq (10.13.26)
```

A coefficient that vanishes is reported together with the condition that fails, whether a triangular condition on a particular triad or a projection outside its permitted range, so that a zero is never returned without explanation. Further cells reproduce the numerical tables themselves under the conditions stated there, and a final part evaluates the algebraic expressions of Sec. 14.2 at numerical arguments and compares them with the values obtained directly, the two being independent routes to the same quantity.

REFERENCES

I. Books and Reviews

- [1] N. E. Cochín. *Vector Calculus and Principles of Tensor Calculus*. Moscow: USSR Academy of Sciences, 1951.
- [2] V. I. Smirnov. *Course of Higher Mathematics*. Vol. 2. Moscow: Fizmatgiz, 1958.
- [3] V. I. Smirnov. *Course of Higher Mathematics*. Vol. 3. Moscow: Fizmatgiz, 1958.
- [4] M. E. Rose. *Multiple Fields*. New York: John Wiley, 1955.
- [5] R. G. Newton. *Scattering Theory of Waves and Particles*. New York: McGraw-Hill, 1969.
- [6] M. E. Rose. *Elementary Theory of Angular Momentum*. New York: John Wiley, 1957.
- [7] D. M. Brink and G. R. Satchler. *Angular Momentum*. Oxford: Clarendon Press, 1968.
- [8] L. D. Landau and E. M. Lifshitz. *Quantum Mechanics*. New York: Pergamon, 1977. a) Messiah, A., *Quantum Mechanics*, North-Holland, Amsterdam (1960).
- [9] M. Tinkham. *Group Theory and Quantum Mechanics*. New York: McGraw-Hill, 1964.
- [10] A. M. Baldin et al. *Kinematics of Nuclear Reactions*. Moscow: Atomizdat, 1968.
- [11] A. De-Shalit and I. Talmi. *Nuclear Shell Theory*. New York: Academic Press, 1963.
- [12] A. Z. Dolginov. “Angular Correlations at Radiative Transitions of Nuclei.” In: *Gamma Rays*. Ed. by L. A. Sliv. Moscow-Leningrad: USSR Academy of Sciences, 1961.
- [13] A. S. Davydov. *Quantum Mechanics*. Moscow: Fizmatgiz, 1963.
- [14] A. Bohr and B. R. Mottelson. *Nuclear Structure*. Vol. 1. New York: W. A. Benjamin, 1969.
- [15] E. P. Wigner. *Group Theory*. New York: Academic Press, 1959.
- [16] U. Fano and G. Racah. *Irreducible Tensorial Sets*. New York: Academic Press, 1959.
- [17] V. B. Berestetskii, E. M. Lifshitz, and L. P. Pitaevskii. *Quantum Electrodynamics*. Oxford: Pergamon, 1982.
- [18] I. M. Gel’fand, R. A. Minlos, and Z. Ya. Shapiro. *Representations of the Rotation and Lorentz Groups and Their Applications*. London: Pergamon, 1963.
- [19] N. Ya. Vilenkin. *Special Functions and the Theory of Group Representations*. Vol. 22. Chap. VII, Sec. 1, Subsec. 3. Providence, R. I.: AMS Translations, 1968.
- [20] A. P. Yutsis and A. A. Bandsaitis. *The Theory of Angular Momentum in Quantum Mechanics*. Vilnius: Mintus, 1965.
- [21] B. R. Judd. *Operator Techniques in Atomic Spectroscopy*. New York: McGraw-Hill, 1963.
- [22] E. U. Condon and Q. W. Shortley. *The Theory of Atomic Spectra*. Cambridge: Cambridge Univ. Press, 1935.
- [23] H. Bateman and A. Erdelyi. *Higher Transcendental Functions*. Vol. 1–3. New York: McGraw-Hill, 1953.
- [24] E. W. Hobson. *The Theory of Spherical and Ellipsoidal Harmonics*. Cambridge: Cambridge University Press, 1931.
- [25] W. Magnus, F. Oberhettinger, and R. P. Soni. *Formulas and Theorems for the Special Functions of Mathematical Physics*. Berlin-Heidelberg: Springer, 1966. DOI: [10.1007/978-3-662-11761-3](https://doi.org/10.1007/978-3-662-11761-3).
- [26] A. I. Akhiezer and V. B. Berestetskii. *Quantum Electrodynamics*. New York: Interscience, 1965.
- [27] J. M. Blatt and V. F. Weisskopf. *Theoretical Nuclear Physics*. New York: McGraw-Hill, 1952.
- [28] J. D. Jackson. *Classical Electrodynamics*. New York: John Wiley, 1975.
- [29] B. L. van der Waerden. *Die Gruppentheoretische Methode in der Quantenmechanik*. Berlin: Springer, 1932.
- [30] Ya. A. Smorodinskii and L. A. Shelepin. “Clebsch-Gordan Coefficients Treated from Different Sides.” In: *Usp. Fiz. Nauk (Sov.)* 106 (1972). DOI: [10.3367/ufnr.0106.197201a.0003](https://doi.org/10.3367/ufnr.0106.197201a.0003).
- [31] C. Eckart. “The Application of Group Theory to the Quantum Dynamics of Monatomic Systems.” In: *Rev. Mod. Phys.* 2 (1930), pp. 305–380. DOI: [10.1103/RevModPhys.2.305](https://doi.org/10.1103/RevModPhys.2.305).
- [32] G. Ya. Lyubarskii. *The Application of Group Theory in Physics*. Oxford: Pergamon Press, 1960.
- [33] A. R. Edmonds. *Angular Momentum in Quantum Mechanics*. Princeton: Princeton Univ. Press, 1957.

- [34] A. P. Yutsis, I. B. Levinson, and V. V. Vanagas. *The Mathematical Apparatus of the Theory of Angular Momentum*. Jerusalem: Israel Program for Scientific Translation, 1962.
- [35] E. El-Baz and B. Castel. “Graphical Methods of Spin Algebras.” In: *Atomic, Nuclear and Particle Physics*. New York: Marcel Dekker, 1972.
- [36] K. Adler. “Beiträge zur Theorie der Richtungskorrelation.” In: *Helv. Phys. Acta* 25 (1952), pp. 235–258. URL: <https://www.e-periodica.ch/digbib/view?pid=hpa-001%3A1952%3A25%3A%3A237>.
- [37] B. F. Baymann. “Some Lectures on Groups and Their Applications to Spectroscopy.” Copenhagen; unpublished. 1957.
- [38] L. A. Sliv, ed. *Deformation of Atomic Nuclei*. Moscow: IL, 1958.
- [39] V. Fock. *Principles of Quantum Mechanics*. Leningrad: Kubuch, 1932.
- [40] H. Goldstein. *Classical Mechanics*. Reading, MA: Addison-Wesley, 1953.
- [41] L. C. Biedenharn and H. Van Dam, eds. *Quantum Theory of Angular Momentum*. New York: Academic Press, 1965.
- [42] J. C. Slater. *Quantum Theory of Atomic Structure*. Vol. 2. New York: McGraw-Hill, 1960.
- [43] I. I. Sobelman. *Atomic Spectra and Radiative Transitions*. Berlin: Springer-Verlag, 1979.
- [44] *Spectroscopic and Group Theoretical Methods in Physics, Racah Memorial Volume*. Amsterdam: North-Holland, 1968.
- [45] G. N. Watson. *A Treatise on the Theory of Bessel Functions*. Vol. 1. Cambridge: Cambridge Univ. Press, 1945.

II. Papers on Special Problems

- [46] G. Ponzano and T. Regge. “Semiclassical Limit of Racah Coefficients.” In: *Spectroscopic and Group Theoretical Methods in Physics*. Amsterdam: North-Holland, 1968.
- [47] A. Edmonds. *Angular Momenta in Quantum Mechanics*. CERN report 55-26. Geneva: CERN, 1955.
- [48] P. R. Fontana. “Symmetric Expansion of One- and Two-center Coulomb Potentials.” In: *J. Math. Phys.* 2 (1961), pp. 825–828. DOI: [10.1063/1.1724228](https://doi.org/10.1063/1.1724228).
- [49] B. Freidman and J. Russak. “Addition Theorems for Spherical Waves.” In: *Quart. of Appl. Math.* 12 (1954), pp. 13–23. DOI: [10.1090/qam/60649](https://doi.org/10.1090/qam/60649).
- [50] R. A. Sack. “Generalisation of Laplace’s Expansion to Arbitrary Charge Distributions.” In: *J. Math. & Phys.* 37 (1957), pp. 215–222.
- [51] R. A. Sack. “Three-dimensional Addition Theorem for Arbitrary Functions, involving Expansions in Spherical Harmonics.” In: *J. Math. Phys.* 5 (1964), pp. 252–259. DOI: [10.1063/1.1704115](https://doi.org/10.1063/1.1704115).
- [52] K. G. Kay, H. D. Todd, and H. J. Silverstone. “Dirac Delta Functions in the Laplace-type Expansion of $r^n Y_{lm}(\vartheta, \varphi)$.” In: *J. Chem. Phys.* 51 (1969), pp. 2359–2362. DOI: [10.1063/1.1672352](https://doi.org/10.1063/1.1672352).
- [53] V. B. Berestetskii, A. Z. Dolginov, and K. A. Ter-Martirosian. “Angular Functions of Particles with Spins.” In: *JETP* 20 (1950), pp. 527–535.
- [54] E. P. Wigner. “On the Matrices Which Reduce the Kronecker Products of Representations of Simply Reducible Groups.” In: *Quantum Theory of Angular Momentum*. Ed. by L. C. Biedenharn and H. van Dam. New York: Academic Press, 1965.
- [55] T. Regge. “Symmetry Properties of Clebsch-Gordan Coefficients.” In: *Nuovo Cimento* 10 (1958), pp. 544–545. DOI: [10.1007/BF02859841](https://doi.org/10.1007/BF02859841).
- [56] G. Racah. “Theory of Complex Spectra. II.” In: *Phys. Rev.* 62 (1942), pp. 438–462. DOI: [10.1103/PhysRev.62.438](https://doi.org/10.1103/PhysRev.62.438).
- [57] V. A. Fock. “New Deduction of the Vector Model.” In: *JETP* 10 (1940), pp. 383–393.
- [58] S. D. Majumdar. “The Clebsch-Gordan Coefficients.” In: *Prog. Theor. Phys.* 20 (1958), pp. 798–803. DOI: [10.1143/PTP.20.798](https://doi.org/10.1143/PTP.20.798).
- [59] A. Bandzaitis and A. Yutsis. “Once More on the Formulas for the Clebsch-Gordan Coefficients.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 45–49.
- [60] S. M. R. Ansari. “Quasi-binomial Representation of the Clebsch-Gordan Coefficients.” In: *Nuovo Cimento* 38 (1965), pp. 1883–1886. DOI: [10.1007/BF02750104](https://doi.org/10.1007/BF02750104).

- [61] A. M. Bincer. “Interpretation of the Symmetry of the Clebsch-Gordan Coefficients Discovered by Regge.” In: *J. Math. Phys.* 11 (1970), pp. 1835–1844. DOI: [10.1063/1.1665333](https://doi.org/10.1063/1.1665333).
- [62] E. L. Akim and A. A. Levin. “Generating Function for Clebsch-Gordan Coefficients.” In: *Dokl. Acad. Nauk USSR* 138 (1961), pp. 503–505.
- [63] A. P. Stone. “Expressions for Certain Wigner Coefficients.” In: *Proc. Phys. Soc.* 70A (1957), pp. 980–909. DOI: [10.1088/0370-1298/70/12/111](https://doi.org/10.1088/0370-1298/70/12/111).
- [64] L. A. Shelepin. “SU_n Symmetry in the Theory of Clebsch-Gordan Coefficients.” In: *JETP* 48 (1965), pp. 360–367.
- [65] J. Schwinger. “On Angular Momentum.” In: *Quantum Theory of Angular Momentum*. Ed. by L. C. Biedenharn and H. van Dam. New York: Academic Press, 1965.
- [66] P. J. Brussaard and H. A. Tolhoek. “Classical Limits of Clebsch-Gordan Coefficients. Racah Coefficients and $D_{mn}^l(\varphi, \vartheta, \psi)$ -functions.” In: *Physica* 23 (1957), pp. 955–971. DOI: [10.1016/S0031-8914\(57\)95547-7](https://doi.org/10.1016/S0031-8914(57)95547-7).
- [67] S. F. Boys. “Electronic Wave Functions IV.” In: *Proc. Roy. Soc., London* A207 (1951), pp. 181–197. DOI: [10.1098/rspa.1951.0110](https://doi.org/10.1098/rspa.1951.0110).
- [68] U. Fano. *Statistical Matrix Techniques and Their Application to the Directional Correlation of Radiation*. Report 1214. US National Bureau of Standards, 1951.
- [69] W. Sharp. *Racah Algebra and the Contraction of Groups*. AECL-1098. Ontario, 1960.
- [70] L. A. Shelepin. “On the Symmetry of the Clebsch-Gordan Coefficients.” In: *JETP* 46 (1964), pp. 1033–1038.
- [71] T. Regge. “Symmetry Properties of Racah’s Coefficients.” In: *Nuovo Cimento* 11 (1959), pp. 116–117. DOI: [10.1007/BF02724914](https://doi.org/10.1007/BF02724914).
- [72] L. C. Biedenharn, J. M. Blatt, and M. E. Rose. “Some Properties of the Racah and Associated Coefficients.” In: *Rev. Mod. Phys.* 24 (1952), pp. 249–257. DOI: [10.1103/RevModPhys.24.249](https://doi.org/10.1103/RevModPhys.24.249).
- [73] V. Bargmann. “On the Representations of the Rotation Group.” In: *Rev. Mod. Phys.* 34 (1962), pp. 829–845. DOI: [10.1103/RevModPhys.34.829](https://doi.org/10.1103/RevModPhys.34.829).
- [74] K. Alder et al. “Study of Nuclear Structure by Electromagnetic Excitation with Accelerated Ions.” In: *Rev. Mod. Phys.* 28 (1956), pp. 432–542. DOI: [10.1103/RevModPhys.28.432](https://doi.org/10.1103/RevModPhys.28.432).
- [75] G. Racah. “Directional Correlation of Successive Nuclear Radiations.” In: *Phys. Rev.* 84 (1951), pp. 910–912. DOI: [10.1103/PhysRev.84.910](https://doi.org/10.1103/PhysRev.84.910).
- [76] H. A. Jahn. “Theoretical Studies in Nuclear Structure. II.” In: *Proc. Roy. Soc. London* A205 (1951), pp. 192–237. DOI: [10.1098/rspa.1951.0026](https://doi.org/10.1098/rspa.1951.0026).
- [77] A. Wu. “Structure of the Wigner 9j Coefficients in the Bargmann Approach.” In: *J. Math. Phys.* 13 (1972), pp. 84–90. DOI: [10.1063/1.1665855](https://doi.org/10.1063/1.1665855).
- [78] A. Z. Dolginov and A. N. Moskalov. “Relativistic Spherical Harmonics. III.” In: *JETP* 37 (1959), pp. 1697–1707.
- [79] H. A. Jahn and J. Hope. “Symmetry Properties of the Wigner 9j Symbol.” In: *Phys. Rev.* 93 (1954), pp. 318–321. DOI: [10.1103/PhysRev.93.318](https://doi.org/10.1103/PhysRev.93.318).
- [80] A. Karosiene, S. Ališauskas, and A. Bandzaitis. “9j Coefficients with One Parameter Equal to Unity.” In: *Litov. Fiz. Sb.* 5 (1965), pp. 13–21.
- [81] S. Jang. “Relationship among the Wigner 9j Symbols.” In: *J. Math. Phys.* 9 (1968), pp. 397–402. DOI: [10.1063/1.1664592](https://doi.org/10.1063/1.1664592).
- [82] A. Arima, H. Horie, and Y. Tanabe. “Generalized Racah Coefficient and its Application.” In: *Prog. Theor. Phys.* 11 (1954), pp. 143–154. DOI: [10.1143/PTP.11.143](https://doi.org/10.1143/PTP.11.143).
- [83] R. T. Sharp. “Stretched χ -coefficients.” In: *Nucl. Phys.* A95 (1967), pp. 222–228. DOI: [10.1016/0375-9474\(67\)90163-7](https://doi.org/10.1016/0375-9474(67)90163-7).
- [84] J. M. Kennedy and M. J. Cliff. *Transformation Coefficients Between LS and JJ Couplings*. Chalk River Report CRT. Ontario, 1957.
- [85] I. B. Levinson. “Sum of Products of Wigner Coefficients and Their Graphical Representation.” In: *Liet. TSR Mokslu Akad. Darbai B Ser.* 2 (1956), pp. 17–30.
- [86] A. Bandzaitis et al. “On the Calculation of 6j Coefficients.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 35–43.
- [87] A. Bandzaitis, A. Karosiene, and A. Yutsis. “On the Calculation of 9j Coefficients.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 457–466.

- [88] V. B. Berestetskii. “Electromagnetic Multipole Field.” In: *JETP* 17 (1947), pp. 12–18.
- [89] L. C. Biedenharn. “An Identity Satisfied by the Racah Coefficients.” In: *J. Math. & Phys.* 31 (1953), pp. 287–293. DOI: [10.1002/sapm1952311287](https://doi.org/10.1002/sapm1952311287).
- [90] A. Z. Dolginov. “Relativistic Spherical Harmonics.” In: *JETP* 30 (1956), pp. 746–755.
- [91] A. Z. Dolginov and I. N. Toptygin. “Relativistic Spherical Harmonics. II.” In: *JETP* 37 (1959), pp. 1441–1451.
- [92] J. P. Elliot. “Theoretical Studies in Nuclear Structure. V.” In: *Proc. Roy. Soc., London* A218 (1953), pp. 345–370. DOI: [10.1098/rspa.1953.0109](https://doi.org/10.1098/rspa.1953.0109).
- [93] V. Fock. “Zur Theorie des Wasserstoffatoms.” In: *Z. Phys.* 98 (1936), pp. 145–154. DOI: [10.1007/BF01336904](https://doi.org/10.1007/BF01336904).
- [94] W. Happer. “A Partial-wave Expansion of the Finite Rotation Operator.” In: *Ann. Phys.* 48 (1968), pp. 579–591. DOI: [10.1016/0003-4916\(68\)90146-2](https://doi.org/10.1016/0003-4916(68)90146-2).
- [95] F. R. Innes and C. W. Ufford. “Microwave Zeeman Effect and Theory of Complex Spectra.” In: *Phys. Rev.* 111 (1958), pp. 194–202. DOI: [10.1103/PhysRev.111.194](https://doi.org/10.1103/PhysRev.111.194).
- [96] A. Karosiene et al. “6j Coefficients with Negative Parameters.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 187–196.
- [97] J. D. Louk. “New Recursion Relation for Clebsch-Gordan Coefficients.” In: *Phys. Rev.* 110 (1958), pp. 815–816. DOI: [10.1103/PhysRev.110.815](https://doi.org/10.1103/PhysRev.110.815).
- [98] S. D. Majumdar. “Coupling of Three Angular Momenta.” In: *Acta Phys. Acad. Scient. Hung.* 26 (1969), pp. 311–318. DOI: [10.1007/BF03157469](https://doi.org/10.1007/BF03157469).
- [99] M. S. Marinov. “Representation of the Finite Rotation Matrix as a Sum over Clebsch-Gordan Coefficients.” In: *Yad. Fiz.* 5 (1967), pp. 1321–1323.
- [100] M. A. Melvin and N. V. V. J. Swamy. “Evaluation of Certain Physically Interesting Integrals and Hypergeometric Sums.” In: *J. Math. & Phys.* 36 (1957), pp. 157–163. DOI: [10.1002/sapm1957361157](https://doi.org/10.1002/sapm1957361157).
- [101] M. Micu. “Recursion Relation for the 3j Symbols.” In: *Nucl. Phys.* A113 (1968), pp. 215–220. DOI: [10.1016/0375-9474\(68\)90895-6](https://doi.org/10.1016/0375-9474(68)90895-6).
- [102] H. E. Moses. “Irreducible Representation of the Rotation Group in Terms of the Axis and Angle of Rotation.” In: *Ann. Phys.* 37 (1966), pp. 224–226. DOI: [10.1016/0003-4916\(66\)90040-6](https://doi.org/10.1016/0003-4916(66)90040-6).
- [103] G. I. Petrashen. “Solution of the Vector Limit Problem of Mathematical Physics in the Case of a Sphere.” In: *Dokl. Acad. Nauk* 46 (1945), pp. 291–294.
- [104] G. Racah. “Theory of Complex Spectra. I.” In: *Phys. Rev.* 61 (1942), pp. 186–197. DOI: [10.1103/PhysRev.61.186](https://doi.org/10.1103/PhysRev.61.186).
- [105] G. Racah. “Theory of Complex Spectra. III.” In: *Phys. Rev.* 63 (1943), pp. 367–382. DOI: [10.1103/PhysRev.63.367](https://doi.org/10.1103/PhysRev.63.367).
- [106] M. E. Rose. “The Electrostatic Interaction of Two Arbitrary Charge Distributions.” In: *J. Math. & Phys.* 37 (1958), pp. 215–222. DOI: [10.1002/sapm1958371215](https://doi.org/10.1002/sapm1958371215).
- [107] M. Sato. “General Formula of the Racah Coefficient.” In: *Prog. Theor. Phys.* 13 (1955), pp. 405–414. DOI: [10.1143/PTP.13.405](https://doi.org/10.1143/PTP.13.405).
- [108] A. Savukynas et al. “The Symmetry of Mirror Reflection in the Theory of Angular Momentum.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 467–478.
- [109] R. T. Sharp. “A Generalised Regge Identity for Wigner Coefficients.” In: *Nuovo Cimento* 47 (1967), pp. 860–868. DOI: [10.1007/BF02771400](https://doi.org/10.1007/BF02771400).
- [110] J. D. Talman and W. W. True. “An Identity Involving 9j Coefficients.” In: *Can. J. Phys.* 42 (1964), pp. 1081–1086. DOI: [10.1139/p64-101](https://doi.org/10.1139/p64-101).
- [111] E. Wigner. “On Representations of Certain Finite Groups.” In: *Am. J. Math.* 63 (1941), pp. 57–63. DOI: [10.2307/2371276](https://doi.org/10.2307/2371276).
- [112] A. Yutsis et al. “Clebsch-Gordan Coefficients with Negative Angular Momentum Quantum Numbers.” In: *Litov. Fiz. Sb.* 4 (1964), pp. 173–185.

III. Tables

- [113] H. A. Buckmaster. “Tables of Angular Momentum Transformation Matrix Elements $d_{mm'}^j(\beta)$ ($j = 2, 4, 6$).” In: *Can. J. Phys.* 42 (1964), pp. 386–391. DOI: [10.1139/p64-036](https://doi.org/10.1139/p64-036).

- [114] G. F. Wolters. “Simple Method for the Explicit Calculation of d-Functions.” In: *Nucl. Phys.* B18 (1970), pp. 625–653. DOI: [10.1016/0550-3213\(70\)90140-9](https://doi.org/10.1016/0550-3213(70)90140-9).
- [115] A. N. Behkami. “Tables of Rotational Wavefunctions d_{MK}^J , $J \leq 13$ and $J \leq 25/2$.” In: *Nuclear Data Tables* 10 (1971), pp. 1–48. DOI: [10.1016/S0092-640X\(71\)80020-7](https://doi.org/10.1016/S0092-640X(71)80020-7).
- [116] T. Shimpuku. “General Theory and Numerical Tables of Clebsch-Gordan Coefficients.” In: *Prog. Theor. Phys. Suppl.* 13 (1960), pp. 1–136. DOI: [10.1143/PTPS.13.1](https://doi.org/10.1143/PTPS.13.1).
- [117] L. C. Biedenharn. *Tables of the Racah Coefficients*. ONRL-1098. 1952.
- [118] D. L. Falkoff, C. S. Calladay, and R. E. Sells. “Transformation Amplitudes for Vector Addition of Angular Momentum, $(j_3 m m' | j_3 J M)$.” In: *Can. J. Phys.* 30 (1952), pp. 253–256. DOI: [10.1139/p52-024](https://doi.org/10.1139/p52-024).
- [119] M. Yamada and M. Morita. “On the β -ray Angular Correlations.” In: *Prog. Theor. Phys.* 8 (1952), pp. 431–442. DOI: [10.1143/PTP.8.431](https://doi.org/10.1143/PTP.8.431).
- [120] A. Simon. *Numerical Tables of the Clebsch-Gordan Coefficients*. ORNL-1718. 1954.
- [121] H. Appel. *Numerical Tables for 3j, 6j, 9j Symbols*. Vol. 3. Landolt-Börnstein (Group 1). Berlin: Springer, 1968.
- [122] H. Matsunobu and H. Takebe. “Tables of U-Coefficients.” In: *Prog. Theor. Phys.* 14 (1955), pp. 589–605. DOI: [10.1143/PTP.14.589](https://doi.org/10.1143/PTP.14.589).
- [123] J. Vizbaraite et al. *The Tables of 9j Coefficients for Integer Values of Parameters with One Parameter Equal Unity*. Moscow: Computer Center of the USSR Academy of Sciences, 1968.
- [124] T. Ishidzu. *Tables of the Racah Coefficients*. Tokyo: Pan-Pacific Press, 1960.
- [125] K. Kumar. “Tables of Certain Clebsch-Gordan Coefficients and of Matrix Elements.” In: *Can. J. Phys.* 35 (1957). Errata 35 1401 (1957), pp. 341–345. DOI: [10.1139/p57-038](https://doi.org/10.1139/p57-038).
- [126] M. A. Melvin and N. V. V. J. Swamy. “Algebraic Table of Vector Addition Coefficients for $j = 5/2$.” In: *Phys. Rev.* 107 (1957), pp. 186–189. DOI: [10.1103/PhysRev.107.186](https://doi.org/10.1103/PhysRev.107.186).
- [127] M. Morita, R. Morita, and T. Tsukamoto. “Clebsch-Gordan Coefficients for $j = 5/2, 3$ and $7/2$.” In: *Prog. Theor. Phys. Suppl.* 26 (1963), pp. 64–74. DOI: [10.1143/PTPS.26.64](https://doi.org/10.1143/PTPS.26.64).
- [128] A. F. Nikiforov, V. B. Uvarov, and Yu. I. Levitan. *Tables of Racah Coefficients*. New York: Macmillan, 1965.
- [129] M. Rotenberg et al. *The 3j and 6j Symbols*. London: Crosby Lockwood, 1959.
- [130] R. Saito and M. Morita. “Clebsch-Gordan Coefficients for $j = 5/2$.” In: *Prog. Theor. Phys.* 13 (1955), pp. 540–542. DOI: [10.1143/PTP.13.540](https://doi.org/10.1143/PTP.13.540).

IV. References Added in This Edition

- [131] H. De Meyer and G. Vanden Berghe. “Traces of Products of Angular Momentum Operators in the Spherical Basis.” In: *J. Phys. A, Math. Gen.* 11 (1978), pp. 494–500. DOI: [10.1088/0305-4470/11/3/009](https://doi.org/10.1088/0305-4470/11/3/009).
- [132] P. R. Subramanian and V. Devanathan. “Trace Techniques for Angular Momentum Operators.” In: *J. Phys. A, Math. Gen.* 7 (1974), pp. 1995–2007. DOI: [10.1088/0305-4470/7/16/004](https://doi.org/10.1088/0305-4470/7/16/004).
- [133] J. Thakur. “General Expression for the Trace of the Product of an Arbitrary Number of Components of Angular Momentum.” In: *Ind. J. Pure and Appl. Phys.* 13 (1975), pp. 792–793.
- [134] Ya. A. Smorodinskii and S. K. Suslov. “Clebsch-Gordan Coefficients for SU(2) and Hahn Polynomials.” In: *Yad. Fiz.* 35 (1982), pp. 192–201.
- [135] Ya. A. Smorodinskii and S. K. Suslov. “6j Symbols and Orthogonal Polynomials.” In: *Yad. Fiz.* 36 (1982), pp. 1066–1071.
- [136] J. D. Morgan III. “Further Relations Involving 3j Symbols.” In: *J. Phys. A, Math. Gen.* 9 (1976), pp. 1231–1233. DOI: [10.1088/0305-4470/9/8/009](https://doi.org/10.1088/0305-4470/9/8/009).
- [137] A. M. Din. “A Simple Sum Formula for Clebsch-Gordan Coefficients.” In: *Letters in Math. Phys.* 5 (1981), pp. 207–211. DOI: [10.1007/BF00420700](https://doi.org/10.1007/BF00420700).
- [138] B. I. Dunlap and B. R. Judd. “Novel Identities for Simple nj-Symbols.” In: *J. Math. Phys.* 16 (1975), pp. 318–319. DOI: [10.1063/1.522543](https://doi.org/10.1063/1.522543).
- [139] K. Schulten and R. G. Gordon. “Semiclassical Approximations to 3j and 6j Coefficients for Quantum Mechanical Coupling of Angular Momenta.” In: *J. Math. Phys.* 16 (1975), pp. 1971–1988. DOI: [10.1063/1.522427](https://doi.org/10.1063/1.522427).

- [140] S. J. Ališauskas and A. P. Yutsis. “Weight Lowering Operators and the Multiplicity-free Isoscalar Factors for the Group R_5 .” In: *J. Math. Phys.* 12 (1971), pp. 594–605. DOI: [10.1063/1.1665626](https://doi.org/10.1063/1.1665626).
- [141] V. K. Khersonskii, A. N. Moskalev, and D. A. Varshalovich. *Tables of the 12j Symbols of the First Kind*. Preprint 499. Leningrad: Ioffe Phys. Tech. Institute, 1975.
- [142] W. T. Sharp. *Some Formal Properties of the 12j Symbol*. Chalk River Report TPJ-81. Ontario, 1955.
- [143] L. C. Biedenharn and J. D. Louck. *Angular Momentum in Quantum Physics. Theory and Application*. Encyclopedia of Mathematics and Its Applications. Reading, Massachusetts: Addison-Wesley, 1981.
- [144] V. K. Khersonskii, A. N. Moskalev, and D. A. Varshalovich. “On the Evaluation of the 12j Symbols of the First Kind.” In: *Izv. Acad. Nauk SSSR, Ser. Fiz.* 40 (1976), pp. 702–706.
- [145] V. K. Khersonskii, A. N. Moskalev, and D. A. Varshalovich. “On the Evaluation of the 12j Symbols of the Second Kind.” In: *Izv. Acad. Nauk SSSR, Ser. Fiz.* 40 (1976), pp. 706–711.

V. Additional Papers Since the Book Came Out

(none)

