

Example script for VAST for spatio-temporal analysis of single-species catch-rate data

James Thorson

October 10, 2016

Contents

1	Overview	2
2	Getting started	2
2.1	Further information	2
2.2	Related tools	2
2.3	How to cite VAST	3
3	Settings	4
3.1	Spatial settings	4
3.2	Model settings	4
3.3	Potential outputs	5
3.4	Stratification for results	5
3.5	Derived objects	6
3.6	Save settings	6
4	Prepare the data	6
4.1	Data-frame for catch-rate data	6
4.2	Extrapolation grid	7
4.3	Derived objects for spatio-temporal estimation	7
5	Build and run model	8
5.1	Build model	8
5.2	Estimate fixed effects and predict random effects	8
6	Diagnostic plots	8
6.1	Plot data	9
6.2	Convergence	11
6.3	Diagnostics for encounter-probability component	12
6.4	Diagnostics for positive-catch-rate component	12
6.5	Diagnostics for plotting residuals on a map	14
6.6	Model selection	17

7	Model output	17
7.1	Direction of “geometric anisotropy”	17
7.2	Density surface for each year	18
7.3	Index of abundance	18
7.4	Center of gravity and range expansion/contraction	19

```
## package 'pander' successfully unpacked and MD5 sums checked
##
## The downloaded binary packages are in
## C:\Users\James.Thorson\AppData\Local\Temp\RtmpWEm8vp\downloaded_packages
```

1 Overview

This tutorial will walk through a simple example of how to use VAST for estimating single-species abundance indices, distribution shifts, and range expansion.

2 Getting started

To install TMB on a windows machine, we need to first install [Rtools](#). During the installation, please select the option to have Rtools included in your system path. On other operating systems, it is not necessary to install Rtools. We then install VAST

```
devtools::install_github("james-thorson/VAST")
devtools::install_github("james-thorson/utilities")
```

Next load libraries.

```
library(TMB) # Can instead load library(TMBdebug)
```

```
## Warning: package 'TMB' was built under R version
## 3.3.2
```

```
library(VAST)
```

2.1 Further information

If you have further questions after reading this tutorial, please explore the [GitHub repo](#) mainpage, wiki, and glossary. Also please explore the R help files, e.g., `?Data_Fn` for explanation of data inputs, or `?Param_Fn` for explanation of parameters.

2.2 Related tools

Related tools for spatio-temporal fisheries analysis are currently housed at www.FishStats.org. These include [SpatialDeltaGLMM](#), a single-species antecedent of VAST, and www.FishViz.org, a tool for visualizing single-species results using worldwide. VAST and [SpatialDeltaGLMM](#) both use continuous integration to confirm that they give identical estimates when applied to single-species data.

2.3 How to cite VAST

VAST has involved many publications for developing individual features. If using VAST, please read and cite:

```
citation("VAST")
```

```
##
## Please cite 2016 (ICES J. Mar. Sci. J.
## Cons.) if using the package; 2016 (Glob.
## Ecol. Biogeogr) if exploring factor
## decomposition of spatio-temporal variation;
## 2015 (ICES J. Mar. Sci. J. Cons.) if
## calculating an index of abundance; 2016
## (Methods Ecol. Evol.) if using the
## center-of-gravity metric; 2016 (Fish. Res.)
## if using the bias-correction feature; 2016
## (Proc R Soc B) if using the
## effective-area-occupied metric.
##
## Thorson, J.T., and Barnett, L.A.K. In
## press. Comparing estimates of abundance
## trends and distribution shifts using
## single- and multispecies models of fishes
## and biogenic habitat. ICES J. Mar. Sci. J.
## Cons
##
## Thorson, J.T., Ianelli, J.N., Larsen, E.,
## Ries, L., Scheuerell, M.D., Szuwalski, C.,
## and Zipkin, E. 2016. Joint dynamic species
## distribution models: a tool for community
## ordination and spatiotemporal monitoring.
## Glob. Ecol. Biogeogr. 25(9): 1144-1158.
## doi:10.1111/geb.12464. url:
## http://onlinelibrary.wiley.com/doi/10.1111/geb.12464/abstract
##
## Thorson, J.T., Shelton, A.O., Ward, E.J.,
## Skaug, H.J., 2015. Geostatistical
## delta-generalized linear mixed models
## improve precision for estimated abundance
## indices for West Coast groundfishes. ICES
## J. Mar. Sci. J. Cons. 72(5), 1297-1310.
## doi:10.1093/icesjms/fsu243. URL:
## http://icesjms.oxfordjournals.org/content/72/5/1297
##
## Thorson, J.T., and Kristensen, K. 2016.
## Implementing a generic method for bias
## correction in statistical models using
## random effects, with spatial and
## population dynamics examples. Fish. Res.
## 175: 66-74.
## doi:10.1016/j.fishres.2015.11.016. url:
## http://www.sciencedirect.com/science/article/pii/S0165783615301399
##
## Thorson, J.T., Pinsky, M.L., Ward, E.J.,
```

```
## 2016. Model-based inference for estimating
## shifts in species distribution, area
## occupied, and center of gravity. Methods
## Ecol. Evol. 7(8), 990-1008.
## doi:10.1111/2041-210X.12567. URL:
## http://onlinelibrary.wiley.com/doi/10.1111/2041-210X.12567/full
##
## Thorson, J.T., Rindorf, A., Gao, J.,
## Hanselman, D.H., and Winker, H. 2016.
## Density-dependent changes in effective
## area occupied for sea-bottom-associated
## marine fishes. Proc R Soc B 283(1840):
## 20161853. doi:10.1098/rspb.2016.1853. URL:
## http://rspb.royalsocietypublishing.org/content/283/1840/20161853.
```

and also browse the [GitHub list](#) of papers

3 Settings

First chose an example data set for this script, as archived with package

```
Data_Set = c("Chatham_rise_hake", "Iceland_cod", "WCGBTS_canary",
             "GSL_american_plaice", "BC_pacific_cod", "EBS_pollock",
             "GOA_Pcod", "GOA_pollock", "GB_spring_haddock",
             "GB_fall_haddock", "SAWC_jacopever", "Aleutian_islands_POP")[6]
```

Next use latest version for CPP code

```
Version = "VAST_v2_0_0"
```

3.1 Spatial settings

The following settings define the spatial resolution for the model, and whether to use a grid or mesh approximation

```
Method = c("Grid", "Mesh")[2]
grid_size_km = 25
n_x = c(100, 250, 500, 1000, 2000)[1] # Number of stations
Kmeans_Config = list( "randomseed"=1, "nstart"=100, "iter.max"=1e3 )
```

3.2 Model settings

The following settings define whether to include spatial and spatio-temporal variation, whether its autocorrelated, and whether there's overdispersion

```
FieldConfig = c(Omega1 = 1, Epsilon1 = 1, Omega2 = 1,
                Epsilon2 = 1)
RhoConfig = c(Beta1 = 0, Beta2 = 0, Epsilon1 = 0, Epsilon2 = 0)
OverdispersionConfig = c(Vessel = 0, VesselYear = 0)
ObsModel = c(2, 0)
```

3.3 Potential outputs

The following settings define what types of output we want to calculate

```
Options = c(SD_site_density = 0, SD_site_logdensity = 0,  
  Calculate_Range = 1, Calculate_evenness = 0, Calculate_effective_area = 1,  
  Calculate_Cov_SE = 0, Calculate_Synchrony = 0,  
  Calculate_Coherence = 0)
```

3.4 Stratification for results

We also define any potential stratification of results, and settings specific to any case-study data set

```
# Default  
if (Data_Set %in% c("GSL_american_plaice", "BC_pacific_cod",  
  "EBS_pollock", "SAWC_jacopever", "Chatham_rise_hake",  
  "Aleutian_islands_POP")) {  
  strata.limits <- data.frame(STRATA = "All_areas")  
}  
# Specific (useful as examples)  
if (Data_Set %in% c("WGBTS_canary", "Sim")) {  
  # In this case, it will calculate a coastwide  
  # index, and also a separate index for each state  
  # (although the state lines are approximate)  
  strata.limits <- data.frame(STRATA = c("Coastwide",  
    "CA", "OR", "WA"), north_border = c(49, 42,  
    46, 49), south_border = c(32, 32, 42, 46),  
    shallow_border = c(55, 55, 55, 55), deep_border = c(1280,  
    1280, 1280, 1280))  
  # Override default settings for vessels  
  VesselConfig = c(Vessel = 0, VesselYear = 1)  
}  
if (Data_Set %in% c("GOA_Pcod", "GOA_pollock")) {  
  # In this case, will calculating an unrestricted  
  # index and a separate index restricted to west of  
  # -140W  
  strata.limits <- data.frame(STRATA = c("All_areas",  
    "west_of_140W"), west_border = c(-Inf, -Inf),  
    east_border = c(Inf, -140))  
}  
if (Data_Set %in% c("GB_spring_haddock", "GB_fall_haddock")) {  
  # For NEFSC indices, strata must be specified as a  
  # named list of area codes  
  strata.limits = list(Georges_Bank = c(1130, 1140,  
    1150, 1160, 1170, 1180, 1190, 1200, 1210, 1220,  
    1230, 1240, 1250, 1290, 1300))  
}  
if (Data_Set %in% c("Iceland_cod")) {  
  strata.limits = data.frame(STRATA = "All_areas")  
  # Turn off all spatial, temporal, and  
  # spatio-temporal variation in probability of  
  # occurrence, because they occur almost everywhere  
  FieldConfig = c(Omega1 = 0, Epsilon1 = 0, Omega2 = 1,
```

```

    Epsilon2 = 1)
  RhoConfig = c(Beta1 = 3, Beta2 = 0, Epsilon1 = 0,
    Epsilon2 = 0)
}

```

3.5 Derived objects

Depending on the case study, we define a **Region** used when extrapolating or plotting density estimates. If its a different data set, it will define **Region**="Other", and this is a recognized level for all uses of **Region** (which attempts to define reasonable settings based on the location of sampling). For example **Data_Set**="Iceland_cod" has no associated meta-data for the region, so it uses **Region**="Other" by default.

```

Region = switch( Data_Set, "Chatham_rise_hake"="New_Zealand",
  "WCGBTS_canary"="California_current",
  "GSL_american_plaice"="Gulf_of_St_Lawrence",
  "BC_pacific_cod"="British_Columbia",
  "EBS_pollock"="Eastern_Bering_Sea",
  "GOA_Pcod"="Gulf_of_Alaska",
  "GOA_pollock"="Gulf_of_Alaska",
  "GB_spring_haddock"="Northwest_Atlantic",
  "GB_fall_haddock"="Northwest_Atlantic",
  "SAWC_jacopever"="South_Africa",
  "Aleutian_islands_POP"="Aleutian_Islands",
  "Other")

```

3.6 Save settings

We then set the location for saving files.

```

DateFile = paste0(getwd(), '/VAST_output/')
dir.create(DateFile)

```

I also like to save all settings for later reference, although this is not necessary.

```

Record = ThorsonUtilities::bundlelist(c("Data_Set",
  "Version", "Method", "grid_size_km", "n_x", "FieldConfig",
  "RhoConfig", "OverdispersionConfig", "ObsModel",
  "Kmeans_Config"))
save(Record, file = file.path(DateFile, "Record.RData"))
capture.output(Record, file = paste0(DateFile, "Record.txt"))

```

4 Prepare the data

4.1 Data-frame for catch-rate data

Depending upon the **Data_Set** chosen, we load archived data sets that are distributed with the package. Each archived data set is then reformatted to create a data-frame **Data_Geostat** with a standardized set of columns. For a new data set, the user is responsible for formatting **Data_Geostat** appropriately to match this format. We show the first six rows of **Data_Geostat** given that **Data_Set** = **Data_Set**.

Catch_KG	Year	Vessel	AreaSwept_km2	Lat	Lon	Pass
22.2	1982	missing	0.01	55.3	-165	0
26.3	1982	missing	0.01	55.3	-167	0
132	1982	missing	0.01	55.3	-164	0
11.2	1982	missing	0.01	55.3	-166	0
52.3	1982	missing	0.01	55.3	-165	0
9.24	1982	missing	0.01	55.4	-163	0

4.2 Extrapolation grid

We also generate the extrapolation grid appropriate for a given region. For new regions, we use `Region="Other"`.

```
if (Region %in% c("California_current", "Eastern_Bering_Sea",
  "Gulf_of_Alaska", "Aleutian_Islands", "Northwest_Atlantic",
  "Gulf_of_St_Lawrence", "New_Zealand")) {
  Extrapolation_List = SpatialDeltaGLMM::Prepare_Extrapolation_Data_Fn(Region = Region,
    strata.limits = strata.limits)
}
if (Region == "British_Columbia") {
  Extrapolation_List = SpatialDeltaGLMM::Prepare_Extrapolation_Data_Fn(Region = Region,
    strata.limits = strata.limits, strata_to_use = c("HS",
    "QCS"))
}
if (Region == "South_Africa") {
  Extrapolation_List = SpatialDeltaGLMM::Prepare_Extrapolation_Data_Fn(Region = Region,
    strata.limits = strata.limits, region = "west_coast")
}
if (Region == "Other") {
  Extrapolation_List = SpatialDeltaGLMM::Prepare_Extrapolation_Data_Fn(Region = Region,
    strata.limits = strata.limits, observations_LL = Data_Geostat[,
    c("Lat", "Lon")], maximum_distance_from_sample = 15)
}
```

4.3 Derived objects for spatio-temporal estimation

And we finally generate the information used for conducting spatio-temporal parameter estimation, bundled in list `Spatial_List`

```
Spatial_List = SpatialDeltaGLMM::Spatial_Information_Fn(grid_size_km = grid_size_km,
  n_x = n_x, Method = Method, Lon = Data_Geostat[,
  "Lon"], Lat = Data_Geostat[, "Lat"], Extrapolation_List = Extrapolation_List,
  randomseed = Kmeans_Config[["randomseed"]], nstart = Kmeans_Config[["nstart"]],
  iter.max = Kmeans_Config[["iter.max"]], DirPath = DateFile,
  Save_Results = FALSE)
# Add knots to Data_Geostat
Data_Geostat = cbind(Data_Geostat, Spatial_List$loc_UTM,
  knot_i = Spatial_List$knot_i)
```

5 Build and run model

5.1 Build model

To estimate parameters, we first build a list of data-inputs used for parameter estimation. `Data_Fn` has some simple checks for buggy inputs, but also please read the help file `?Data_Fn`.

```
TmbData = Data_Fn(Version = Version, FieldConfig = FieldConfig,
  OverdispersionConfig = OverdispersionConfig, RhoConfig = RhoConfig,
  ObsModel = ObsModel, c_i = rep(0, nrow(Data_Geostat)),
  b_i = Data_Geostat[, "Catch_KG"], a_i = Data_Geostat[,
    "AreaSwept_km2"], v_i = as.numeric(Data_Geostat[,
    "Vessel"]) - 1, s_i = Data_Geostat[, "knot_i"] -
  1, t_i = Data_Geostat[, "Year"], a_xl = Spatial_List$a_xl,
  MeshList = Spatial_List$MeshList, GridList = Spatial_List$GridList,
  Method = Spatial_List$Method, Options = Options)
```

```
##   Omega1 Epsilon1   Omega2 Epsilon2
##       1           1         1         1
##   Vessel VesselYear
##       -1           -1
```

We then build the TMB object.

```
TmbList = Build_TMB_Fn(TmbData = TmbData, RunDir = DateFile,
  Version = Version, RhoConfig = RhoConfig, loc_x = Spatial_List$loc_x)
Obj = TmbList[["Obj"]]
```

5.2 Estimate fixed effects and predict random effects

Next, we use a gradient-based nonlinear minimizer to identify maximum likelihood estimates for fixed-effects

```
Opt = TMBhelper::Optimize(obj = Obj, lower = TmbList[["Lower"]],
  upper = TmbList[["Upper"]], getsd = TRUE, savedir = DateFile,
  bias.correct = FALSE)
```

Finally, we bundle and save output

```
Report = Obj$report()
Save = list("Opt"=Opt, "Report"=Report, "ParHat"=Obj$env$parList(Opt$par), "TmbData"=TmbData)
save(Save, file=paste0(DateFile,"Save.RData"))
```

6 Diagnostic plots

We first apply a set of standard model diagnostics to confirm that the model is reasonable and deserves further attention. If any of these do not look reasonable, the model output should not be interpreted or used.

6.1 Plot data

It is always good practice to conduct exploratory analysis of data. Here, I visualize the spatial distribution of data. Spatio-temporal models involve the assumption that the probability of sampling a given location is statistically independent of the probability distribution for the response at that location. So if sampling “follows” changes in density, then the model is probably not appropriate!

```
SpatialDeltaGLMM::Plot_data_and_knots(Extrapolation_List = Extrapolation_List,  
    Spatial_List = Spatial_List, Data_Geostat = Data_Geostat,  
    PlotDir = DateFile)
```




Figure 1: Spatial extent and location of knots




Figure 2: Spatial distribution of catch-rate data

6.2 Convergence

Here I print the diagnostics generated during parameter estimation, and I confirm that (1) no parameter is hitting an upper or lower bound and (2) the final gradient for each fixed-effect is close to zero. For explanation of parameters, please see ?Data_Fn.

```
pander::pandoc.table( Opt$diagnostics[,c('Param', 'Lower', 'MLE', 'Upper', 'final_gradient')] )
```

Param	Lower	MLE	Upper	final_gradient
ln_H_input	-50	0.2315	50	0.0008786
ln_H_input	-50	-0.9657	50	0.0009001
beta1_ct	-50	4.12	50	-0.0004914
beta1_ct	-50	4.229	50	-0.0001649
beta1_ct	-50	4.323	50	6.758e-05
beta1_ct	-50	5.093	50	0.0003083
beta1_ct	-50	5.428	50	0.0005221
beta1_ct	-50	4.105	50	0.0007122
beta1_ct	-50	5.056	50	7.428e-06
beta1_ct	-50	4.168	50	-0.0006779
beta1_ct	-50	4.333	50	-0.0003862
beta1_ct	-50	5.989	50	2.901e-05
beta1_ct	-50	4.524	50	1.19e-05
beta1_ct	-50	5.265	50	6.13e-05
beta1_ct	-50	5.647	50	0.0003327
beta1_ct	-50	4.886	50	-5.081e-05
beta1_ct	-50	5.073	50	-0.0004524
beta1_ct	-50	4.753	50	-9.86e-05
beta1_ct	-50	4.997	50	0.0001197
beta1_ct	-50	6.219	50	8.882e-05
beta1_ct	-50	5.125	50	0.0002557
beta1_ct	-50	5.707	50	0.0007568
beta1_ct	-50	4.809	50	-8.683e-05
beta1_ct	-50	4.535	50	0.0001293
beta1_ct	-50	5.454	50	-0.001194
beta1_ct	-50	4.747	50	-5.943e-05
beta1_ct	-50	4.572	50	0.0002242
beta1_ct	-50	4.198	50	0.000267
beta1_ct	-50	2.877	50	0.0002274
beta1_ct	-50	3.426	50	0.0002083
beta1_ct	-50	2.986	50	0.0001771
beta1_ct	-50	4.66	50	-0.0001344
beta1_ct	-50	4.657	50	-0.0002736
beta1_ct	-50	5.189	50	-0.0002172
beta1_ct	-50	6.231	50	-0.0004938
L_omega1_z	-50	1.946	50	9.472e-05
L_epsilon1_z	-50	0.9753	50	0.001679
logkappa1	-6.01	-4.12	-2.574	-0.000908
beta2_ct	-50	7.517	50	-5.443e-05
beta2_ct	-50	8.74	50	-3.202e-05
beta2_ct	-50	7.844	50	-5.82e-05
beta2_ct	-50	8.535	50	4.756e-05
beta2_ct	-50	8.097	50	-1.987e-05
beta2_ct	-50	8.459	50	1.503e-05

Param	Lower	MLE	Upper	final_gradient
beta2_ct	-50	8.287	50	-3.478e-05
beta2_ct	-50	8.243	50	2.595e-05
beta2_ct	-50	8.046	50	-3.521e-05
beta2_ct	-50	8.17	50	7.08e-06
beta2_ct	-50	8.063	50	-2.384e-05
beta2_ct	-50	8.212	50	-8.04e-05
beta2_ct	-50	8.008	50	-6.509e-05
beta2_ct	-50	7.516	50	-4.389e-05
beta2_ct	-50	7.73	50	3.487e-05
beta2_ct	-50	7.887	50	5.193e-05
beta2_ct	-50	7.663	50	0.0001318
beta2_ct	-50	7.405	50	-6.556e-05
beta2_ct	-50	8.198	50	6.193e-07
beta2_ct	-50	8.166	50	-3.368e-05
beta2_ct	-50	7.847	50	-0.0001007
beta2_ct	-50	8.542	50	-1.254e-05
beta2_ct	-50	7.983	50	1.183e-05
beta2_ct	-50	7.832	50	-6.843e-05
beta2_ct	-50	7.13	50	-0.0001259
beta2_ct	-50	6.996	50	-7.861e-05
beta2_ct	-50	6.544	50	7.068e-07
beta2_ct	-50	6.056	50	-1.221e-05
beta2_ct	-50	7.291	50	6.297e-05
beta2_ct	-50	7.546	50	5.785e-05
beta2_ct	-50	7.247	50	-0.0001146
beta2_ct	-50	7.513	50	-9.088e-05
beta2_ct	-50	8.565	50	-1.622e-05
L_omega2_z	-50	1.106	50	0.002481
L_epsilon2_z	-50	-1.123	50	-0.001173
logkappa2	-6.01	-4.535	-2.574	0.001304
logSigmaM	-50	0.1682	10	-0.003915

6.3 Diagnostics for encounter-probability component

Next, we check whether observed encounter frequencies for either low or high probability samples are within the 95% predictive interval for predicted encounter probability

```
Enc_prob = SpatialDeltaGLMM::Check_encounter_prob(Report = Report,
  Data_Geostat = Data_Geostat, DirName = DateFile)
```

6.4 Diagnostics for positive-catch-rate component

We can visualize fit to residuals of catch-rates given encounters using a Q-Q plot. A good Q-Q plot will have residuals along the one-to-one line.

```
Q = SpatialDeltaGLMM::QQ_Fn(TmbData = TmbData, Report = Report,
  FileName_PP = paste0(DateFile, "Posterior_Predictive.jpg"),
  FileName_Phist = paste0(DateFile, "Posterior_Predictive-Histogram.jpg"),
  FileName_QQ = paste0(DateFile, "Q-Q_plot.jpg"),
  FileName_Qhist = paste0(DateFile, "Q-Q_hist.jpg")) # SpatialDeltaGLMM::
```




Figure 3: Expectedated probability and observed frequency of encounter for “encounter probability” component




Figure 4: Quantile-quantile plot indicating residuals for “positive catch rate” component

6.5 Diagnostics for plotting residuals on a map

Finally, we visualize residuals on a map. To do so, we first define years to plot and generate plotting inputs. useful plots by first determining which years to plot (`Years2Include`), and labels for each plotted year (`Year_Set`)

```
# Get region-specific settings for plots
MapDetails_List = SpatialDeltaGLMM::MapDetails_Fn( "Region"=Region, "NN_Extrap"=Spatial_List$PolygonList
# Decide which years to plot
Year_Set = seq(min(Data_Geostat[, 'Year']), max(Data_Geostat[, 'Year']))
Years2Include = which( Year_Set %in% sort(unique(Data_Geostat[, 'Year'])))
```

We then plot Pearson residuals. If there are visible patterns (areas with consistently positive or negative residuals accross or within years) then this is an indication of the model “overshrinking” results towards the intercept, and model results should then be treated with caution.

```
SpatialDeltaGLMM::plot_residuals(Lat_i = Data_Geostat[,
  "Lat"], Lon_i = Data_Geostat[, "Lon"], TmbData = TmbData,
  Report = Report, Q = Q, savedir = DateFile, MappingDetails = MapDetails_List[["MappingDetails"]],
  PlotDF = MapDetails_List[["PlotDF"]], MapSizeRatio = MapDetails_List[["MapSizeRatio"]],
  Xlim = MapDetails_List[["Xlim"]], Ylim = MapDetails_List[["Ylim"]],
  FileName = DateFile, Year_Set = Year_Set, Years2Include = Years2Include,
  Rotate = MapDetails_List[["Rotate"]], Cex = MapDetails_List[["Cex"]],
  Legend = MapDetails_List[["Legend"]], zone = MapDetails_List[["Zone"]],
  mar = c(0, 0, 2, 0), oma = c(3.5, 3.5, 0, 0), cex = 1.8)
```




Figure 5: Pearson residuals for encounter-probability by knot




Figure 6: Pearson residuals for positive catch rates by knot

6.6 Model selection

To select among models, we recommend using the Akaike Information Criterion, AIC, via `Opt$AIC=1.149\times 10^{\{5\}}`.

7 Model output

Last but not least, we generate pre-defined plots for visualizing results

7.1 Direction of “geometric anisotropy”

We can visualize which direction has faster or slower decorrelation (termed “geometric anisotropy”)

```
SpatialDeltaGLMM::PlotAniso_Fn(FileName = paste0(DateFile,  
  "Aniso.png"), Report = Report, TmbData = TmbData)
```




Figure 7: Decorrelation distance for different directions

7.2 Density surface for each year

We can visualize many types of output from the model. Here I only show predicted density, but other options are obtained via other integers passed to `plot_set` as described in `?PlotResultsOnMap_Fn`

```
SpatialDeltaGLMM::PlotResultsOnMap_Fn(plot_set = c(3),
  MappingDetails = MapDetails_List[["MappingDetails"]],
  Report = Report, Sdreport = Opt$SD, PlotDF = MapDetails_List[["PlotDF"]],
  MapSizeRatio = MapDetails_List[["MapSizeRatio"]],
  Xlim = MapDetails_List[["Xlim"]], Ylim = MapDetails_List[["Ylim"]],
  FileName = DateFile, Year_Set = Year_Set, Years2Include = Years2Include,
  Rotate = MapDetails_List[["Rotate"]], Cex = MapDetails_List[["Cex"]],
  Legend = MapDetails_List[["Legend"]], zone = MapDetails_List[["Zone"]],
  mar = c(0, 0, 2, 0), oma = c(3.5, 3.5, 0, 0), cex = 1.8,
  plot_legend_fig = FALSE)
```

7.3 Index of abundance

The index of abundance is generally most useful for stock assessment models.

```
Index = SpatialDeltaGLMM::PlotIndex_Fn(DirName = DateFile,
  TmbData = TmbData, Sdreport = Opt[["SD"]], Year_Set = Year_Set,
  Years2Include = Years2Include, strata_names = strata.limits[,
    1], use_biascorr = TRUE)
pander::pandoc.table(Index$Table[, c("Year", "Fleet",
  "Estimate_metric_tons", "SD_log", "SD_mt")])
```

Year	Fleet	Estimate_metric_tons	SD_log	SD_mt
1982	All_areas	2391518	0.08975	214645
1983	All_areas	5743674	0.08742	502140
1984	All_areas	4037774	0.09583	386953
1985	All_areas	4903450	0.1109	543679
1986	All_areas	4323628	0.09065	391957
1987	All_areas	4848402	0.09324	452086
1988	All_areas	6494203	0.09803	636641
1989	All_areas	5765367	0.08935	515127
1990	All_areas	6362280	0.1095	696834
1991	All_areas	4530972	0.09176	415743
1992	All_areas	4203668	0.09303	391081
1993	All_areas	5014039	0.08535	427968
1994	All_areas	4662240	0.09004	419794
1995	All_areas	4099802	0.102	418012
1996	All_areas	2672989	0.07938	212187
1997	All_areas	3040908	0.08515	258930
1998	All_areas	2335759	0.08682	202789
1999	All_areas	3274280	0.09699	317587
2000	All_areas	4512363	0.08594	387783
2001	All_areas	3857100	0.08484	327254
2002	All_areas	4240923	0.07631	323610
2003	All_areas	6840989	0.08879	607392
2004	All_areas	3595980	0.07969	286556

Year	Fleet	Estimate_metric_tons	SD_log	SD_mt
2005	All_areas	4270664	0.08614	367858
2006	All_areas	2702639	0.08798	237790
2007	All_areas	3810938	0.1063	405098
2008	All_areas	2605632	0.1023	266616
2009	All_areas	1863568	0.1169	217879
2010	All_areas	3263269	0.1035	337901
2011	All_areas	2919661	0.09075	264960
2012	All_areas	3125148	0.08222	256949
2013	All_areas	4185026	0.09032	377980
2014	All_areas	7295635	0.07829	571191

7.4 Center of gravity and range expansion/contraction

We can detect shifts in distribution or range expansion/contraction.

```
SpatialDeltaGLMM::Plot_range_shifts(Report = Report,
  TmbData = TmbData, Sdreport = Opt[["SD"]], Znames = colnames(TmbData$Z_xm),
  PlotDir = DateFile, Year_Set = Year_Set)
```




Figure 8: Density maps for each year




Figure 9: Index of abundance plus/minus 1 standard error




Figure 10: Center of gravity (COG) indicating shifts in distribution plus/minus 1 standard error




Figure 11: Effective area occupied indicating range expansion/contraction plus/minus 1 standard error