Optimization of CUDA-based Monte Carlo Simulation for Radiation Therapy

GTC 2014

N. Henderson & K. Murakami

The collaboration





Special thanks to the CUDA Center of Excellence Program Makoto Asai, SLAC
Joseph Perl, SLAC
Andrea Dotti, SLAC
Koichi Murakami, KEK
Takashi Sasaki, KEK
Margot Gerritsen, ICME
Nick Henderson, ICME

Radiotherapy simulation methods

Analytic


- time: seconds to minutes
- accurate within 3-5%
- used in treatment planning

Monte Carlo

- time: several hours to days of CPU time
- accurate within 1-2%
- used to verify treatment plans in certain cases

Project overview


GPU based dose calculation for radiation therapy



Features

- GPU code based on Geant4
- Voxelized geometry
- DICOM interface
- Material is water with variable density
- Limited Geant4 EM physics: electron/positron/gamma
- Scoring of dose in each voxel
- Process secondary particles


Geant4




Images from: Geant4 gallery and gMocren

Basic idea

- Particles are transported through material in steps
- Physics processes act on particles and may generate secondaries
- Particles are removed when out of domain or out of energy




A single step




G4CU: CUDA implementation

- Each GPU thread processes a single particle until the particle exits the geometry
- Each thread has a stack for secondary particles
- Every thread takes a step in each iteration
- Algorithm is periodically interrupted for various actions



Take step

G4CU: parallel execution



- CUDA kernels are applied to all particles
- Threads must wait if physics process does not apply to particle
- Parallel execution is achieved:
 - maintenance operations
 - transport process
 - same process on same particle

Performance and validation

Hardware and software:


- GPU:
 - Tesla K20 Kepler
 - 2496 cores, 706 MHz, 5GB GDDR5 (ECC)
- CPU
 - Xeon X5680 (3.33GHz)
 - single thread operation
- SDK:
 - CUDA 5.5
 - Geant4 release 9.6 p2









Phantom geometry



Particle sources


- 20 MeV electron / 6 MeV photons (mono-energy)
- 6 MV & 18 MV spectrum photons







Depth dose distribution: water phantom

- 6MeV photon, 20 MeV electron
- dose along central axis






Comparisons for slab phantoms



6MV photon, Lung/Bone as inner slab material

Visualization with gMocren



- Image for demonstration purposes only!
- 50M 6MeV photons
- Pencil beam configuration

Challenges

- Geant4 is complex
 - Implement targeted subset
- Varied character of physics process
 - Bad occupancy
 - No clear optimization target
- Random simulation
 - Thread divergence

- Lookup (interpolation) tables
 - Thread divergence
 - Non-ideal memory access
- Energy deposition in global array
 - Possible race condition
 - Non-ideal memory access

Physics Processes


| Process | Lines of code | Initialization kernel registers | Step length kernel registers | Action kernel registers |
|-----------------------|---------------|---------------------------------|------------------------------|-------------------------|
| Compton scattering | 116 | 14 | 18 | 49 |
| Photo electric effect | 118 | 14 | 18 | 40 |
| Gamma conversion | 160 | 14 | 18 | 36 |
| Bremsstrahlung | 156 | 14 | 18 | 44 |
| Ionization | 394 | 14 | (18, <mark>18</mark>) | (60,36) |
| Scattering | 575 | | 12 | 28 |
| Positron annihilation | 156 | | 12 | 26 |
| Electron removal | 44 | | 12 | 8 |
| Transport | 491 | 20 | 22 | (20,22) |

Physics Profile (6 MeV gamma)

| Sum of Time(%) | step | along-step | after-step | other | init | at-rest | Grand Total |
|-----------------------|------|------------|------------|-------|------|---------|----------------|
| em-helper | 18.4 | | | | 4.2 | | 22.5 |
| ionization | 2.5 | 15.3 | 2.1 | | | | 19.9 |
| multiple-scattering | 11.6 | 7.7 | | | | | 19.3 |
| management | 3.7 | | | 8.7 | 0.5 | | 12.8 |
| transport | 2.4 | 1.7 | 3.3 | | 1.4 | | 8.9 |
| bremsstrahlung | | | 5.7 | | | | 5.7 |
| positron-annihilation | 2.1 | | 0.9 | | 0.6 | 0.7 | 4.3 |
| compton-scattering | | | 2.6 | | | | 2.6 |
| gamma-conversion | | | 1.4 | | | | 1.4 |
| photo-electric-effect | | | 1.3 | | | | 1.3 |
| electron-removal | | | | | | 1.2 | 1.2 |
| Grand Total | 40.7 | 24.6 | 17.4 | 8.7 | 6.6 | 2.0 | 100.0 |

Optimization 1: dose storage

- Old idea: use thrust to join and reduce dose stacks into global array
- Better idea: use atomicAdd
- Speedup: ~1.5 (at the time)



Optimization 2: device configuration

- 6 MeV gamma pencil beam
- 3,000,000 events
- Table shows run time / shortest time
- Voxels: 61 x 61 x 150
- 1 = 22 seconds
- ~ 136 events / ms

| | | threads per block | | | | |
|-----|-----|-------------------|------|------|------|------|
| blo | cks | 32 | 64 | 128 | 256 | 512 |
| 3 | 2 | 17.98 | 9.85 | 5.62 | 3.21 | 2.04 |
| 6 | 4 | 9.85 | 5.63 | 3.2 | 2 | 1.48 |
| 12 | 28 | 5.62 | 3.21 | 1.98 | 1.39 | 1.22 |
| 25 | 6 | 3.55 | 2.15 | 1.36 | 1.1 | 1.14 |
| 51 | .2 | 2.42 | 1.46 | 1.08 | 1.02 | 1.15 |
| 10 | 24 | 1.8 | 1.22 | 1 | 1.01 | 1.61 |

| | threads per block | | | |
|--------|-------------------|------|------|--|
| blocks | 32 | 64 | 128 | |
| 1000 | 2.2 | 1.33 | 1 | |
| 2000 | 1.82 | 1.19 | 1.03 | |
| 3000 | 1.8 | 1.19 | 1.04 | |
| 4000 | 1.74 | 1.27 | 1.15 | |

Other optimizations

- New hardware!
- Refactoring
- New features sometime slow us down
- Results from:
 - 512 x 512 x 256 geometry
 - 6 MeV gamma pencil beam

| Date | Hardware | Feature | Time (min) | Events / ms |
|---------|----------|-------------------------------|---------------|-------------|
| March | C2070 | | 72 | 23.1 |
| May | K20 | | 60 | 27.8 |
| June | K20 | atomicAdd | 41 | 40.7 |
| July | K20 | multiple scattering | 42.5 | 39.2 |
| August | K20 | world volume | 49 | 34.0 |
| Current | K20 | Refactoring, device config | 23 | 72.5 |

Comparison to Geant4 on CPU

- Geometry:61 x 61 x 150
- GPU: Tesla K20
- CPU: Xeon X5680
 - 6 core, 12 thread
- Our measurement was with single threaded G4
- New G4 is multi-threaded
- We multiply by 12 to get CPU events / ms

| | | 20 MeV electron (pencil) | 6 MeV photon (pencil) |
|--------------------|--------------------|--------------------------------|-----------------------------|
| | Time (h) | 29.44 | 12.42 |
| CPU | Events / ms / core | 0.94 | 2.24 |
| | Events / ms | 11.32 | 26.85 |
| CDII | Time (min) | 22.23 | 9.65 |
| GPU | Events / ms | 74.99 | 172.69 |
| speedup (GPU/core) | | 79.49 | 77.19 |
| speedup (GPU/CPU) | | 6.62 | 6.43 |

Other optimization experiments

- 6 MeV gamma pencil beam
- 1,000,000 events
- Voxels: 61 x 61
 x 150

| Experiment | Run time (s) | Events / ms |
|------------------------------------------------------------------|--------------|-------------|
| standard | 9.33 | 107.2 |
| prefer L1 cache | 9.28 | 107.8 |
| disable L1 cache | 9.27 | 107.9 |
| remove atomicAdd | 9.31 | 107.4 |
| limit to 32 registers | 9.29 | 107.6 |
| no thread divergence, no atomic add | 3.25 | 307.7 |
| no thread divergence, no atomic add, limit to 32 registers | 3.297 | 303.3 |
| add bounds checking | 17.91 | 55.8 |

Going forward

- Hope to get factor 1.5 speed up from splitting particles into separate CUDA streams
 - Reduce thread divergence
 - Allow concurrent kernel launches
- Some possible benefits from using texture memory and hardware interpolation for lookup tables
- Look for other opportunities in expensive kernels

Acknowledgements

- NVidia for support of the CUDA Center of Excellence
- Koichi Murakami
- Makoto Asai, Joseph Pearl, Andrea Dotti @ SLAC
- Takashi Sasaki @ KEK
- Akinori Kimura, Ashikaga Institute of Technology