

Contrast coding in linear (mixed-effects) models

Nora Wickelmaier

Last modified: April 25, 2025

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors
- We need to assign numeric values often called contrasts to our categorical variables to enter them into a regression model

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors
- We need to assign numeric values often called contrasts to our categorical variables to enter them into a regression model
- Contrasts allow for comparisons between one or more levels of a categorical variable

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors
- We need to assign numeric values often called contrasts to our categorical variables to enter them into a regression model
- Contrasts allow for comparisons between one or more levels of a categorical variable
- For any categorical variable with n levels, $n - 1$ contrasts (or dummy variables) can be defined

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors
- We need to assign numeric values often called contrasts to our categorical variables to enter them into a regression model
- Contrasts allow for comparisons between one or more levels of a categorical variable
- For any categorical variable with n levels, $n - 1$ contrasts (or dummy variables) can be defined
- The default contrast in R is the treatment contrast (or dummy coding)

Categorical predictors

- For all regression models (including mixed-effects model), we need to make a choice in how to treat categorical predictors
- We need to assign numeric values often called contrasts to our categorical variables to enter them into a regression model
- Contrasts allow for comparisons between one or more levels of a categorical variable
- For any categorical variable with n levels, $n - 1$ contrasts (or dummy variables) can be defined
- The default contrast in R is the treatment contrast (or dummy coding)
- For ANOVAs, so-called sum or effects coding is usually used

① 2 × 2 example for linear mixed-effects model

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*
 2. *Foods* with levels *Salad* and *Soup*

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*
 2. *Foods* with levels *Salad* and *Soup*
- The dependent measure is *RT* (speed of eating in minutes)

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*
 2. *Foods* with levels *Salad* and *Soup*
- The dependent measure is *RT* (speed of eating in minutes)
- We will simulate 20 subjects and 10 food ingredients (items)

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*
 2. *Foods* with levels *Salad* and *Soup*
- The dependent measure is *RT* (speed of eating in minutes)
- We will simulate 20 subjects and 10 food ingredients (items)
- The data is generated by the following model

$$RT = \beta_0 + \beta_1 \textit{Utensils} + \beta_2 \textit{Foods} + \beta_3 \textit{Utensils} \times \textit{Foods} + v_0 + \eta_0 + \varepsilon$$

with $v_0 \sim N(0, \sigma_v^2)$, $\eta_0 \sim N(0, \sigma_\eta^2)$, $\varepsilon \sim N(0, \sigma_\varepsilon^2)$, all i.i.d.

Example by Brehm and Alday (2022)

- We will look at the impact of different contrasts with a small simulated example
- We have two manipulated factors:
 1. *Utensils* with levels *Fork* and *Spoon*
 2. *Foods* with levels *Salad* and *Soup*
- The dependent measure is *RT* (speed of eating in minutes)
- We will simulate 20 subjects and 10 food ingredients (items)
- The data is generated by the following model

$$RT = \beta_0 + \beta_1 \text{UtensilsSpoon} + \beta_2 \text{FoodsSoup} + \beta_3 \text{UtensilsSpoon} \times \text{FoodsSoup} + v_0 + \eta_0 + \varepsilon$$

with $v_0 \sim N(0, \sigma_v^2)$, $\eta_0 \sim N(0, \sigma_\eta^2)$, $\varepsilon \sim N(0, \sigma_\varepsilon^2)$, all i.i.d.

Example by Brehm and Alday (2022)

```
library(lme4)
set.seed(1012)

# Average time to eat in min
SpoonSoup <- 5
ForkSoup <- 10
SpoonSalad <- 10
ForkSalad <- 5
# Standard deviation for all groups
Groupsd <- 2
# Number of subjects (ps) and ingredients (ii)
ps <- 20
ii <- 10
```

Example by Brehm and Alday (2022)

```
# Create data frame for given design
ds <- data.frame(
  Utensils      = factor(rep(c("Spoon", "Fork"), each = ps * ii, times = 2)),
  Foods         = factor(rep(c("Soup", "Salad"), each = ps * ii * 2)),
  Participant   = factor(rep(paste0("p", 1:ps), times = ii * 4)),
  Item          = factor(rep(paste0("i", 1:ii), each = ps, times = 4)))

# Simulate data
ds$RT <- c(rnorm(ps * ii, mean = SpoonSoup, sd = Groupsd),
          rnorm(ps * ii, mean = ForkSoup, sd = Groupsd),
          rnorm(ps * ii, mean = SpoonSalad, sd = Groupsd),
          rnorm(ps * ii, mean = ForkSalad, sd = Groupsd))
psre <- rnorm(ps, mean = 0, sd = Groupsd / 5)
iire <- rnorm(ii, mean = 0, sd = Groupsd / 5)
ds$RT <- ds$RT + psre[ds$Participant] + iire[ds$Item]
```

Treatment coding

- The default coding in R is treatment (or dummy) coding
- The first level of each factor is assigned as the reference category

```
contrasts(ds$Utensils)
#      Spoon
# Fork      0
# Spoon      1
contrasts(ds$Foods)
#      Soup
# Salad    0
# Soup      1
```

Effects coding


- In effects (or sum) coding, one level is set as negative and one positive, with zero as the mean of the two levels
- For more than two levels, there will also be a reference category, which is the *last* factor level in R

```
contrasts(ds$Utensils)
#           [,1]
# Fork         1
# Spoon        -1
contrasts(ds$Foods)
#           [,1]
# Salad         1
# Soup         -1
```

Exercise

- Fit the model on slide 4 to your simulated data
 1. with dummy coding
 2. with effects coding
- Interpret the fixed parameters and compare the variance components for the random effects for the two models
- What is the interpretation of the intercept for the two models? Why does it differ?

Treatment coding




- Means from simulated data

	Salad	Soup
Fork	4.81	9.94
Spoon	9.87	4.90

- Reference categories: Salad and Fork
- Estimated coefficients

	β	SE	t
(Intercept)	4.81	0.21	23.45
UtensilsSpoon	5.06	0.20	25.19
FoodsSoup	5.13	0.20	25.52
UtensilsSpoon:FoodsSoup	-10.10	0.28	-35.52

Effects coding



- Means from simulated data

	Salad	Soup
Fork	4.81	9.94
Spoon	9.87	4.90

- Estimated coefficients

	β	SE	t
(Intercept)	7.38	0.16	44.97
Utensils1	-0.01	0.07	-0.09
Foods1	-0.04	0.07	-0.57
Utensils1:Foods1	-2.52	0.07	-35.52

2 Defining contrasts

General linear model

$$\mathbf{y}_i = \mathbf{X}_i \boldsymbol{\beta} + \varepsilon_i$$

which corresponds to

$$\begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \\ y_N \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & x_{12} & \dots & x_{1p} \\ 1 & x_{21} & x_{22} & \dots & x_{2p} \\ 1 & x_{31} & x_{32} & \dots & x_{3p} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_{N1} & x_{N2} & \dots & x_{Np} \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \varepsilon_3 \\ \vdots \\ \varepsilon_N \end{pmatrix}$$

Example by Schad et al. (2020)

- In order to keep it reasonably complex, let us consider a design with one between-subjects factor with four levels

Example by Schad et al. (2020)

- In order to keep it reasonably complex, let us consider a design with one between-subjects factor with four levels
- The four levels F_1 to F_4 reflect levels of word frequency with levels low, medium-low, medium-high, and high frequency words, and the dependent variable reflects some response time

Example by Schad et al. (2020)

- In order to keep it reasonably complex, let us consider a design with one between-subjects factor with four levels
- The four levels F_1 to F_4 reflect levels of word frequency with levels low, medium-low, medium-high, and high frequency words, and the dependent variable reflects some response time
- From four factor levels, we can build three contrasts

Example by Schad et al. (2020)

- Let us consider 8 subjects for our 4-level between-subjects design

$$\mathbf{y}_i = \mathbf{X}_i \boldsymbol{\beta} + \boldsymbol{\varepsilon}_i$$

$$\begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \\ y_6 \\ y_7 \\ y_8 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \varepsilon_3 \\ \varepsilon_4 \\ \varepsilon_5 \\ \varepsilon_6 \\ \varepsilon_7 \\ \varepsilon_8 \end{pmatrix}$$

- For the four means of the factor levels, we get

$$\boldsymbol{\mu} = \mathbf{C} \boldsymbol{\beta}$$

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses
 1. Treatment contrasts

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses
 1. Treatment contrasts
 2. Sum contrasts

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses
 1. Treatment contrasts
 2. Sum contrasts
 3. Helmert contrasts


Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses
 1. Treatment contrasts
 2. Sum contrasts
 3. Helmert contrasts
 4. Sequential difference contrasts

Example by Schad et al. (2020)

- The contrast matrix \mathbf{C} defines how the model parameters are “combined” to estimate μ
- μ is our data and will always be the same
- \mathbf{C} depends on the contrast coding we choose and, therefore, so does β
- We will go through different possibilities, all testing different hypotheses
 1. Treatment contrasts
 2. Sum contrasts
 3. Helmert contrasts
 4. Sequential difference contrasts
 5. Custom contrasts

Example by Schad et al. (2020)



```
design <- data.frame(  
  F = factor(c("F1", "F2", "F3", "F4")),  
  mu = c(10, 20, 10, 40))  
  
beta <- lm(mu ~ F, design) |>  
  coef() |>  
  zapsmall()  
  
# Contrast matrix  
mm <- model.matrix(~ F, design)  
  
# Compare to mu  
mm %*% beta
```

Contrast vs. hypothesis matrix

- In the literature on contrasts, we sometimes find the distinction between contrast and hypothesis matrix (e. g., Schad et al., 2020)

Contrast vs. hypothesis matrix

- In the literature on contrasts, we sometimes find the distinction between contrast and hypothesis matrix (e. g., Schad et al., 2020)
- When the contrast matrix defines how we add up the parameters to obtain μ , then the hypothesis matrix defines how to add up the means to obtain the parameters

$$\mu = \mathbf{C} \beta \quad \Leftrightarrow \quad \beta = \mathbf{C}^{-1} \mu$$

Contrast vs. hypothesis matrix

- In the literature on contrasts, we sometimes find the distinction between contrast and hypothesis matrix (e. g., Schad et al., 2020)
- When the contrast matrix defines how we add up the parameters to obtain μ , then the hypothesis matrix defines how to add up the means to obtain the parameters

$$\mu = \mathbf{C} \beta \quad \Leftrightarrow \quad \beta = \mathbf{C}^{-1} \mu$$

- Hence, the hypothesis matrix is the inverse of the contrast matrix

Contrast vs. hypothesis matrix

- In the literature on contrasts, we sometimes find the distinction between contrast and hypothesis matrix (e. g., Schad et al., 2020)
- When the contrast matrix defines how we add up the parameters to obtain μ , then the hypothesis matrix defines how to add up the means to obtain the parameters

$$\mu = \mathbf{C} \beta \quad \Leftrightarrow \quad \beta = \mathbf{C}^{-1} \mu$$

- Hence, the hypothesis matrix is the inverse of the contrast matrix
- It is sometimes easier to define the hypothesis matrix and then obtain the contrast matrix based on it

Treatment contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

$$\mu_1 = \beta_0$$

$$\mu_2 = \beta_0 + \beta_1$$

$$\mu_3 = \beta_0 + \beta_2$$

$$\mu_4 = \beta_0 + \beta_3$$

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

$$\beta_0 = \mu_1$$

$$\beta_1 = \mu_1 - \mu_2$$

$$\beta_2 = \mu_1 - \mu_3$$

$$\beta_3 = \mu_1 - \mu_4$$

Hypotheses being tested: $H_{01}: \mu_1 = \mu_2$, $H_{02}: \mu_1 = \mu_3$, $H_{03}: \mu_1 = \mu_4$

Sum contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & -1 & -1 & -1 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ 3/4 & -1/4 & -1/4 & -1/4 \\ -1/4 & 3/4 & -1/4 & -1/4 \\ -1/4 & -1/4 & 3/4 & -1/4 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

$$\mu_1 = \beta_0 + \beta_1$$

$$\mu_2 = \beta_0 + \beta_2$$

$$\mu_3 = \beta_0 + \beta_3$$

$$\mu_4 = \beta_0 - \beta_1 - \beta_2 - \beta_3$$

$$\beta_0 = \frac{\mu_1 + \mu_2 + \mu_3 + \mu_4}{4}$$

$$\beta_1 = \frac{3}{4}\mu_1 - \frac{1}{4}(\mu_2 + \mu_3 + \mu_4)$$

$$\beta_2 = \frac{3}{4}\mu_2 - \frac{1}{4}(\mu_1 + \mu_3 + \mu_4)$$

$$\beta_3 = \frac{3}{4}\mu_3 - \frac{1}{4}(\mu_1 + \mu_2 + \mu_4)$$

Hypotheses being tested: $H_{01}: \frac{\mu_2 + \mu_3 + \mu_4}{3} = \mu_1$, $H_{02}: \frac{\mu_1 + \mu_3 + \mu_4}{3} = \mu_2$, $H_{03}: \frac{\mu_1 + \mu_2 + \mu_4}{3} = \mu_3$

Helmert contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & -1 & -1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & 0 & 2 & -1 \\ 1 & 0 & 0 & 3 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

$$\mu_1 = \beta_0 - \beta_1 - \beta_2 - \beta_3$$

$$\mu_2 = \beta_0 + \beta_1 - \beta_2 - \beta_3$$

$$\mu_3 = \beta_0 + 2 \cdot \beta_2 - \beta_3$$

$$\mu_4 = \beta_0 + 3 \cdot \beta_3$$

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ -1/2 & 1/2 & 0 & 0 \\ -1/6 & -1/6 & 1/3 & 0 \\ -1/12 & -1/12 & -1/12 & 1/4 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

$$\beta_0 = \frac{\mu_1 + \mu_2 + \mu_3 + \mu_4}{4}$$

$$\beta_1 = -\frac{1}{2}\mu_1 + \frac{1}{2}\mu_2$$

$$\beta_2 = -\frac{1}{6}(\mu_1 + \mu_2) + \frac{1}{3}\mu_3$$

$$\beta_3 = -\frac{1}{12}(\mu_1 + \mu_2 + \mu_3) + \frac{1}{4}\mu_4$$

Hypotheses being tested: $H_{01}: \mu_1 = \mu_2$, $H_{02}: \frac{\mu_1 + \mu_2}{2} = \mu_3$, $H_{03}: \frac{\mu_1 + \mu_2 + \mu_3}{3} = \mu_4$

Sequential difference contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & -3/4 & -1/2 & -1/4 \\ 1 & 1/4 & -1/2 & -1/4 \\ 1 & 1/4 & 1/2 & -1/4 \\ 1 & 1/4 & 1/2 & 3/4 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

$$\mu_1 = \beta_0 - \frac{3}{4}\beta_1 - \frac{1}{2}\beta_2 - \frac{1}{4}\beta_3$$

$$\mu_2 = \beta_0 + \frac{1}{4}\beta_1 - \frac{1}{2}\beta_2 - \frac{1}{4}\beta_3$$

$$\mu_3 = \beta_0 + \frac{1}{4}\beta_1 + \frac{1}{2}\beta_2 - \frac{1}{4}\beta_3$$

$$\mu_4 = \beta_0 + \frac{1}{4}\beta_1 + \frac{1}{2}\beta_2 + \frac{3}{4}\beta_3$$

$$\beta_0 = \frac{\mu_1 + \mu_2 + \mu_3 + \mu_4}{4}$$

$$\beta_1 = \mu_2 - \mu_1$$

$$\beta_2 = \mu_3 - \mu_2$$

$$\beta_3 = \mu_4 - \mu_3$$

Hypotheses being tested: $H_{01}: \mu_1 = \mu_2$, $H_{02}: \mu_2 = \mu_3$, $H_{03}: \mu_3 = \mu_4$

Custom contrasts

- All contrasts that we looked at so far are standard and directly available in R

Custom contrasts

- All contrasts that we looked at so far are standard and directly available in R
- It is also possible to construct contrasts testing custom hypotheses

Custom contrasts

- All contrasts that we looked at so far are standard and directly available in R
- It is also possible to construct contrasts testing custom hypotheses
- Let us assume for our example, we want to test if the first two means (F_1 and F_2) are identical, but that means for levels F_3 and F_4 increase linearly

Custom contrasts

- All contrasts that we looked at so far are standard and directly available in R
- It is also possible to construct contrasts testing custom hypotheses
- Let us assume for our example, we want to test if the first two means (F_1 and F_2) are identical, but that means for levels F_3 and F_4 increase linearly
- Hence, we assume a potential hypothetical outcome, such as $\mu_1 = 10$, $\mu_2 = 10$, $\mu_3 = 20$, and $\mu_4 = 30$

Custom contrasts

- All contrasts that we looked at so far are standard and directly available in R
- It is also possible to construct contrasts testing custom hypotheses
- Let us assume for our example, we want to test if the first two means (F_1 and F_2) are identical, but that means for levels F_3 and F_4 increase linearly
- Hence, we assume a potential hypothetical outcome, such as $\mu_1 = 10$, $\mu_2 = 10$, $\mu_3 = 20$, and $\mu_4 = 30$
- These means can easily be turned into a contrast by centering them and then maybe do some scaling

μ_1	μ_2	μ_3	μ_4
10	10	20	30
-7.5	-7.5	2.5	12.5
-3	-3	1	5

Custom contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & -3 \\ 1 & -3 \\ 1 & 1 \\ 1 & 5 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix}$$

$$\mu_1 = \beta_0 - 3\beta_1$$

$$\mu_2 = \beta_0 - 3\beta_1$$

$$\mu_3 = \beta_0 + \beta_1$$

$$\mu_4 = \beta_0 + 5\beta_1$$

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix} = \begin{pmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ -0.068 & -0.068 & 0.023 & 0.114 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

$$\beta_0 = \frac{\mu_1 + \mu_2 + \mu_3 + \mu_4}{4}$$

$$\beta_1 = -\frac{3}{44}\mu_1 - \frac{3}{44}\mu_2 + \frac{1}{44}\mu_3 + \frac{5}{44}\mu_4$$

Hypothesis being tested: $H_{01}: -3\mu_1 - 3\mu_2 + \mu_3 + 5\mu_4 = 0$

Custom contrasts

Contrast Matrix

$$\begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix} = \begin{pmatrix} 1 & -3 & -0.511 & -0.533 \\ 1 & -3 & 0.131 & 0.727 \\ 1 & 1 & 0.760 & -0.387 \\ 1 & 5 & -0.380 & 0.194 \end{pmatrix} \cdot \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

- It is often a good idea to add orthogonal contrasts, so any variance in your data that is independent of your contrast can be accounted for
- R will automatically add orthogonal contrasts if you provide less than $n - 1$ contrasts for a factor with n levels

Hypothesis Matrix

$$\begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 1/4 & 1/4 & 1/4 & 1/4 \\ -0.068 & -0.068 & 0.023 & 0.114 \\ -0.511 & 0.131 & 0.760 & -0.380 \\ -0.533 & 0.727 & -0.387 & 0.194 \end{pmatrix} \cdot \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}$$

Hypothesis being tested: $H_{01}: -3\mu_1 - 3\mu_2 + \mu_3 + 5\mu_4 = 0$

Simulate data

```
set.seed(1700)

n <- 20
dat <- data.frame(id = 1:n,
                  F = factor(rep(c("F1", "F2", "F3", "F4"),
                                times = n / 4)),
                  DV = rnorm(n, mean = c(10, 20, 10, 40), sd = 5))

datm <- aggregate(DV ~ F, dat, mean)
aggregate(DV ~ F, dat, sd)

lattice::xyplot(DV ~ F, dat, type = c("a", "p"))
```

Exercise

- Now fit linear models with all contrasts we looked at
 1. Treatment contrasts
 2. Sum contrasts
 3. Helmert contrasts
 4. Sequential difference contrasts
 5. Custom contrasts
- Interpret the fixed parameters
- Predict data with each model and compare the results
- Look again at the intercepts for all models and compare

References

- Brehm, L., & Alday, P. M. (2022). Contrast coding choices in a decade of mixed models. *Journal of Memory and Language*, 125, 104334.
<https://doi.org/10.1016/j.jml.2022.104334>
- Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2020). How to capitalize on a priori contrasts in linear (mixed) models: A tutorial. *Journal of Memory and Language*, 110, 104038. <https://doi.org/10.1016/j.jml.2019.104038>