OLLIS POULTRY ANALYTICS (OPA)

COMPREHENSIVE TECHNICAL PROSPECTUS & USER MANUAL

---

1. Executive Platform Overview

Ollis Poultry Analytics (OPA) is a cutting-edge, mathematically rigorous software ecosystem engineered specifically for the precision requirements of commercial poultry research, biostatistics, and flock management. The application replaces disjointed spreadsheets with a robust, centralized relational database, instantly calculating complex biological trajectories, nutritional efficiencies, and economic margins.

The Business Value of OPA

While OPA is a scientific instrument, its primary function is economic protection. 1. **Automated Error Prevention:** The system actively catches transcription errors and biological impossibilities (e.g., negative weight gain) *before* they ruin a feed trial. 2. **Instant Economic Forensics:** Instead of waiting weeks for a statistician to process data, OPA calculates the exact **Margin Over Feeding Cost (MOFC)** instantly, proving the true financial ROI of a dietary supplement over the control group. 3. **Paperless to Paper translation:** OPA handles the heavy lifting digitally, but allows management to print comprehensive, beautifully formatted PDF reports for physical boardroom review.

Designed to operate securely on a local environment or a private cloud architecture, OPA processes live data through an industrial-grade **Python/SciPy statistical engine**. This ensures that absolutely every metric presented—from a daily average body weight to a complex Gompertz sigmoidal growth projection—is scientifically verifiable, repeatable, and peer-review ready. **Crucially, all data remains structurally owned by the facility and can be bulk-exported back to Excel (CSV) at any time, ensuring zero platform lock-in.**

1.1 Core Architecture Paths

The system tracks the chronological sequence of a trial from conceptualization to final economic evaluation: 1. **Metadata & Protocol Setup**: Define interventions, genetic strains, and block randomization. 2. **Applied Nutrition Framework**: Map dietary formulations, linking phase-specific metabolizable energy and costs directly to specific pens. 3. **Daily Biological Input**: Mortality tracking, feed weigh-backs, and weight

measurements. 4. **Laboratory & Pathogen Tracking**: Direct ingestion of bacteriological plates, OPG (Oocysts Per Gram), and morphological lesion scoring. 5. **The Analytical Engine**: Instantaneous processing of raw data into ANOVA algorithms, visualizations, and economic forecasts.

---

2. Advanced Biological & Growth Mathematics

The core of OPA's value lies in its ability to transform discrete weight measurements into continuous vectors of understanding. The application utilizes universally accepted biometrical formulas to prevent misrepresentation of performance.

2.1 Mortality-Adjusted Feed Conversion Ratio (FCR_{adj})

Location in App: `Dashboard -> Key Performance Indicators`

The traditional FCR metric is dangerously flawed in trials with high late-stage mortality, as the feed consumed by the deceased birds drives up the feed ratio without yielding terminal weight. OPA corrects this using the strictly defined **Mass Balance Formula**.

$$FCR_{adj} = \frac{\sum Feed_Consumed_(kg)}{(\sum Live_Weight - \sum Start_Biomass) + \sum Mortality_Weight + \sum Culled_Weight}$$

> **Scientific Application:** By crediting the weight of dead or sampled birds back to the mass pool, the formulation is judged strictly on its *metabolic digestive efficiency*, isolated entirely from pathogenic mortality events.

2.2 European Production Efficiency Factor (EPEF)

Location in App: `Dashboard -> Key Performance Indicators`

A holistic performance index combining survival, mass velocity, and metabolic conversion into a single comparable integer. Industry targets often exceed 400.

$$EPEF = \frac{\left(\frac{Liveability\%}{100} \times Final_Live_Weight_(kg) \times 10,000\right)}{Trial_Duration_(Days) \times FCR_{adj}}$$

2.3 Nutrient Conversion Efficiencies

Location in App: `Analytics -> Analysis & Visualization -> ⚡ Nutrient Conversion Ratios`

Volume conversion is often misleading when comparing high-density vs. low-density diets. OPA calculates conversion down to the sub-macronutrient level. * **Caloric Conversion:** $ $ * **Specific Amino Acid Conversion:** Details strict utilization, e.g., $ $.

---

3. Hatchery, Parent Stock, & Egg Analytics

Location in App: `Dashboard -> Egg Production`

OPA models the entire layer lifecycle, tracking long-term persistency, peak-phase shifts, and egg quality metrics.

3.1 Core Laying Metrics

- **Hen-Day Production (HD%):** The daily velocity of lay. Target: ≥85%.
$$HD\% = \left(\frac{\text{Live Eggs Collected}}{\text{Live Hens Present in Pen Today}}\right) \times 100$$
- **Hen-Housed Production (HH%):** The lifetime economic potential of the initial placement, heavily penalizing early mortality.
$$HH\% = \left(\frac{\text{Total Lifetime Eggs}}{\text{Initial Hens Placed on Day 0}}\right) \times 100$$
- **Daily Egg Mass:** The true physical output volume combining rate with egg sizing.
$$Egg_Mass_(g/hen/day) = \frac{HD\%}{100} \times \text{Avg Egg Weight (g)}$$
- **Egg Quality & Uniformity:** OPA tracks Average Egg Weight and calculates the Coefficient of Variation (CV%) to ensure flock uniformity, a critical metric for grading and incubation success.
- **Egg-Specific Feed Efficiency:** Calculates biological efficiency through two distinct economic lenses: FCR_{dozen} (Kg of feed per dozen eggs) and FCR_{mass} (Kg of feed per Kg of egg mass produced).

3.2 Automated Persistency Modeling

OPA utilizes linear regression on post-peak production data to generate a **Persistency Slope**. By isolating data physically succeeding the peak, the system calculates the daily percentage rate of lay decline (e.g., -0.15%/day), providing statistically sound predictions for when the flock drops below economic viability ($< 80\%$ lay).

4. The SciPy Statistical Engine: Automated Analysis

Location in App: `Dashboard -> Statistical Analysis`

Scientists spend extensive hours exporting data to SAS, JMP, or R (traditional statistical software packages). OPA entirely removes this step, executing massive statistical pipelines automatically and rendering the models natively.

4.1 Automated Validation of Assumptions

Prior to running testing arrays, OPA mathematically assesses dataset structures: 1. **Shapiro-Wilk Test:** Evaluates Gaussian normality of the dataset points. (Target: $P > 0.05$) 2. **Levene's Test:** Assesses homogeneity of variance across multiple experimental blocks to ensure that no single group is skewing the dispersion envelope.

4.2 Dynamic Hypothesis Testing Pipeline

Based on the specific results of the validations above, OPA autonomously selects the mathematically sound operation: * **Strictly Normal & Equal Variance:** Performs **One-Way ANOVA** coupled with **Tukey's Honest Significant Difference (HSD)** post-hoc analysis. The system features a greedy set-cover algorithm to automatically construct **Compact Letter Displays (CLD)** (e.g., Groups 'A', 'AB', 'B') upon the UI. * **Normal but Unequal Variance:** Deploys **Welch's t-test**. * **Non-Parametric Datasets:** If normality assumptions are categorically destroyed, OPA routes calculations to the **Mann-Whitney U Test** or the non-parametric **Kruskal-Wallis H-test** (critical for ordinal categorical data like 0-4 point integer Lesion Scores).

4.3 Advanced Processing Analysis (ANCOVA vs ANOVA)

When analyzing processing data (such as Breast Yield, Fat Pad Yield, or Hot WOG), analyzing absolute organ weights using standard ANOVA is mathematically dangerous if the treatments resulted in birds of distinct total body weights. Conversely, analyzing sheer percentages (e.g. Breast Yield %) using ANOVA violates assumptions of normality unless heavily transformed.

To resolve this, OPA utilizes **Analysis of Covariance (ANCOVA)** as its default processing engine: * **The Covariate:** OPA automatically utilizes the bird's *Pre-Slaughter Live Weight* as a continuous covariate. * **The Mechanism:** By running a mixed-effects ANCOVA, the system mathematically "adjusts" the response variable (e.g., Absolute Breast Weight). It asks the statistical question: *"If all the birds in all the treatments had weighed the exact same amount on the day of processing, would this diet still have*

produced a larger breast?" * **The Output:** This eliminates the confounding variable of total body size, allowing for mathematically pure comparisons of physiological part-yields between dietary groups.

4.4 Predictive Mathematical Modeling

Rather than simply looking at historical data, OPA projects trial outcomes. * **Gompertz Sigmoidal Growth Analytics:** Poultry does NOT grow linearly; it grows in an "S" curve. OPA fits raw scatterplot weight data to the Gompertz biological formula:

$$W(t) = W_{inf} \times e^{-b \times e^{-k \times t}}$$

Where W_{inf} is the asymptotic maximum mature weight, b is the time displacement, and k is the maximum biological growth rate constant. By establishing k from days 0-21, OPA successfully predicts the exact day-42 weight to a strict 95% Confidence Interval. * **Effect Magnitude Output:** On individual two-cohort studies, OPA runs **Cohen's d** calculations, dividing the mean difference by pooled standard deviation to quantify whether a difference is "Negligible", "Small", "Medium", or "Large."

---

5. Advanced Feed Formulation & Nutritional Engineering

OPA transcends basic performance tracking by incorporating a fully-fledged feed mill formulation and environmental impact engine.

5.1 Formula Nutrient Analysis & Breed-Specific Validation

Location in App: `Feed Management -> 🧪 Formulation Lab`

The system evaluates custom feed formulas against established nutritional requirements (e.g., Broiler, Turkey, Layer matrices). It dynamically calculates and validates formulation totals for: * **Energy (AMEn):** Automatically calculates Apparent Metabolizable Energy. * **Macronutrients & Minerals:** Evaluates Crude Protein, Fat, Fiber, Calcium, and Available Phosphorus percentages against phase-specific minimums. * **Ideal Protein Ratio Modeling:** Validates that all critical amino acid ratios (Methionine+Cystine, Threonine, Valine, Isoleucine, Arginine, Tryptophan) meet the ideal protein profile relative to Standardized Ileal Digestible (SID) Lysine. Validation targets are driven by breed-specific standards (see Section 5.2 below) or fall back to generic NASEM/NRC 2024 baselines when no breed is selected.

5.2 Breed-Specific Nutritional Standards

Location in App: `Feed Management -> 🧪 Formulation Lab -> Save Formula -> Breed Standard`

OPA now validates SID Lysine, SID Met+Cys, SID Threonine, and all ideal protein ratios against the **published nutritional specifications of each genetics company**, rather than applying a single generic standard across all breeds. This enables scientifically accurate breed-to-breed comparisons.

When creating or editing a formula, users select a **Breed Standard** from a dropdown filtered by the chosen bird type. The system automatically loads the breed's published phase-specific requirements and ideal protein ratios.

Supported Broiler Breeds: | Breed | Genetics Company | Source Document | |—|—|—| | Ross 308 | Aviagen | Ross Broiler Nutrition Specifications 2022 | | Ross 708 | Aviagen | Ross Broiler Nutrition Specifications 2022 | | Cobb 500 | Cobb-Vantress | Cobb500 Broiler Performance & Nutrition Supplement 2022 | | Cobb 700 | Cobb-Vantress | Cobb700 Broiler Performance & Nutrition Supplement | | Arbor Acres Plus | Aviagen | Arbor Acres Broiler Nutrition Specifications 2022 | | Hubbard | Hubbard (Aviagen) | Hubbard Broiler Management & Nutrition Guide |

Supported Layer Breeds: | Breed | Genetics Company | Source Document | |—|—|—| | Hy-Line W-36 | Hy-Line International | W-36 Commercial Management Guide | | Hy-Line Brown | Hy-Line International | Brown Commercial Management Guide | | Lohmann Brown | Lohmann Breeders | Amino Acid Recommendations for Laying Hens | | ISA Brown | Hendrix Genetics | Essential AA Recommendations (PMC/2022) | | Dekalb White | Hendrix Genetics | Dekalb White Commercial Guide |

Supported Turkey Breeds: | Breed | Genetics Company | Source Document | |—|—|—| | Nicholas 700 | Aviagen Turkeys | Heavy Lines Feeding Guidelines | | B.U.T. Big 6 | Aviagen Turkeys | Heavy Lines Feeding Guidelines | | Hybrid Converter | Hendrix Genetics | Hybrid Turkeys Nutrition & Feeding Guide |

Each breed file contains: * **Phase-specific requirements:** SID Lysine %, SID Met+Cys %, SID Threonine %, AMEn, Crude Protein, Calcium, and Available Phosphorus targets per feed phase (Prestarter through Withdrawal for broilers; Starter through Finisher for layers and turkeys). * **Breed-specific ideal protein ratios:** Amino acid ratios relative to Lysine = 100, reflecting the exact ratios published by each genetics company (e.g., Cobb's slightly higher Met:Lys at finisher phases vs. Ross). * **Metadata with update tracking:** Each breed standard file records its source document, URL, and `last_updated` date so users can verify currency of the underlying data. * **Generic**

(NASEM/NRC) Fallback: If no breed is selected, the system defaults to NASEM/NRC 2024 baseline standards.

> **Key Advantage:** A Ross 708 broiler has ~3-4% higher SID Lysine requirements than a Ross 308 due to greater breast meat deposition genetics. A Cobb 700 has notably *lower* requirements than a Cobb 500. By validating against the correct breed standard, researchers can make apples-to-apples comparisons and avoid false deficiency/excess flags when comparing diets across different genetic lines.

5.3 Logistical Feed Mill Integration

Location in App: `Feed Management -> 🔬 Formula Analyzer -> 🏭 Logistical Tools`

OPA translates abstract percentage-based formulations into actionable manufacturing documents. The system generates dynamically scaled **Mill Batch Sheets** for any requested volume (Tons, Metric Tons, kg, lbs), ensuring precision mixing directly at the feed mill.

5.4 Environmental Impact Estimates

Location in App: `Feed Management -> 🔬 Formula Analyzer -> 🏭 Logistical Tools`

Anticipating modern sustainability requirements, OPA utilizes mass-balance mathematics to estimate the ecological footprint of a specific diet: * Calculates Nitrogen (N) and Phosphorus (P) dietary intake versus bodily retention percentages. * Outputs the estimated gram-weight of N and P excreted in the manure per bird or per flock. * Projects the estimated economic cost of manure disposal based on regional phosphorous/nitrogen removal rates.

---

6. Economic & Margin Forensics

Location in App: `Dashboard -> Cost & Economics`

OPA treats biological data not just as scientific curiosity, but as an economic ledger. The platform contains a real-time margin analyzer capable of blending continuous facility costs with dynamic biological inputs.

6.1 The Master Algorithm: Margin Over Feeding Cost (MOFC)

The modern industry standard for evaluating formulation viability. It isolates the exact profitability of the dietary intervention while ignoring static facility costs.

$$Revenue_Per_Bird = (Avg_Body_Weight \times Market_Price) \times \left(\frac{Liveability\%}{100}\right)$$

$$MOFC = Revenue_Per_Bird - Feed_Cost_Per_Bird$$

6.2 Dynamic Cost Accumulation

OPA compiles a highly accurate total cost model per bird incorporating: 1. **Weighted Phase-Feeding Economics:** Rather than using a static $/Ton, OPA precisely tracks **Per-Phase Feed Costs**. It evaluates the exact physical kilogram consumption of *Starter*, *Grower*, and *Finisher* diets, multiplying each specific phase by its distinct formulation cost basis. 2. **Pathogenic Opportunity Cost:** OPA models late-stage mortality impact. A bird dying on day 35 has absorbed 35 days of housing utilities and consumed 4kg of feed, presenting a massive economic loss compared to a day-3 mortality event. 3. **Daily Accumulated Housing, Utilities, and Labor Overhead.**

6.3 Comparative Return on Investment (ROI) & Break-Even Modeling

Location in App: `Feed Management -> 📊 Formulation Comparison -> 💰 Economics`

The **Break-Even FCR Calculator** (located under the “Formulation Comparison” tab) is uniquely designed to contrast *two different diets* against each other. By combining total biological revenue generation against systemic expenditures, OPA isolates the exact FCR improvement required from an experimental/more expensive diet to financially justify its use over a cheaper baseline diet. It calculates differential ROI percentages, demonstrating the return generation of experimental additives over control diets.

6.4 Single-Feed Economics Forecast

Location in App: `Feed Management -> 🔬 Formula Analyzer -> 💰 Single-Feed Economics Forecast`

For isolated analysis, the **Single-Feed Economics Calculator** (located under the “Formula Analyzer” tab) allows users to project the financial viability of a single diet without needing a comparative baseline.

Users input three primary target performance metrics: 1. **Expected FCR** (Target Feed Conversion Ratio) 2. **Target Live Weight** (in kg) 3. **Market Price** (Revenue expected per kg of live weight)

The system utilizes the specific diet's `cost_per_ton` alongside these metrics to dynamically forecast: * **Feed Intake per Bird**: Calculated as `Target Live Weight × Expected FCR`. * **Feed Cost per Bird**: Calculated as `Feed Intake per Bird × (Cost Per Ton / 1000)`. * **Total Revenue per Bird**: Calculated as `Target Live Weight × Market Price`. * **Margin Over Feed Cost (MOFC) per Bird**: Calculated dynamically as `Total Revenue per Bird - Feed Cost per Bird`. This instantly identifies whether the specific formulation and expected performance will yield a profitable flock margin.

7. Laboratory, Pathology, & Environmental Integration

Location in App: `Dashboard -> Laboratory & Pathology`

The platform contains advanced micro-systems for rendering biological phenomena into comparable mathematics.

7.1 Bacteriological Enumeration & MPN Arrays

OPA utilizes internal algorithms to parse varying lab notation streams: * **Log10 Transformations:** Raw CFU/g plate counts (e.g., 2,500,000) are automatically shifted into Log_{10} mathematical states ($6.39\ Log_{10}$) allowing for normal distribution mapping and ANOVA comparisons on pathogen drops. * **Most Probable Number (MPN):** OPA tracks complex serial dilution tubes (across up to 9 distinct concentrations) and maps the hit patterns to maximum likelihood estimation frameworks for infinitely sparse pathogen detection.

7.2 Coccidiosis & Lesion Matrix

Using the **Kruskal-Wallis** rank models, OPA treats qualitative 0-4 point Johnson-Reid lesion scores for E. maxima, E. tenella, and E. acervulina with the rigor of quantitative variables, plotting exact progression matrices.

7.3 Processing Yields Analysis

Location in App: `Dashboard -> 🔪 Processing Yields`

OPA provides dedicated visualization and statistical analysis of carcass processing data including Hot WOG, Cold WOG, Breast Yield, Thigh Yield, Drumstick Yield, Wing Yield, Fat Pad Yield, and Tender Yield. The system automatically renders yield distributions by

treatment group and applies ANCOVA-adjusted comparisons (see Section 4.3) using pre-slaughter live weight as the covariate, ensuring part-yield comparisons are mathematically isolated from total body size differences.

7.4 Daily Environmental Observations

Location in App: `Dashboard -> 🌡️ Daily Environmental Observations`

OPA tracks and visualizes daily environmental parameters (temperature, humidity, lighting, ventilation) across multiple house locations. These environmental records are chronologically correlated with biological performance data, enabling researchers to identify spatial or temporal environmental factors that may confound treatment effects.

7.5 Bird Inventory QA

Location in App: `Dashboard -> 🐔 Bird Inventory QA`

A real-time reconciliation system that tracks bird placement, mortality events, culls, and sampling removals to maintain an accurate living bird census at the pen level. This ensures that all downstream calculations (Hen-Day Production, FCR, Liveability) operate on verified headcount data rather than assumed values.

---

8. Advanced Experimental Design & Randomization

Location in App: `Create Study -> Step 5: Randomization`

A trial is only as valid as its initial experimental design. OPA contains an integrated pen and tag randomizer to eliminate selection bias at day zero, strictly adhering to established academic protocols.

8.1 Dynamic Block Randomization (PRNG)

OPA utilizes a dynamically calculated **Randomized Complete Block Design (RCBD)** powered by a pseudo-random number generator (PRNG) to distribute treatments evenly without inherent mathematical bias. This relies on algorithmic proportional distribution rather than rigid historical matrices: * **Algorithmic Block Sizing:** The system automatically calculates block sizes based strictly on the exact number of experimental treatments defined. The total available facility pens are uniformly divided into these blocks. * **Cryptographic Dispersal:** Within each continuous block of pens, the algorithm randomly shuffles the available treatments utilizing Python's core `random` library (Mersenne Twister), ensuring exactly one replicate of each treatment exists per

block. * **Fixed Control Anchoring:** The system seamlessly supports complex partial-blind studies by allowing the Study Director to anchor fixed control pens, simultaneously executing the PRNG algorithm *around* those fixed points without breaking the overall block integrity.

---

9. Quality Assurance (QA), Visual Analytics, & Reporting

OPA functions as an uncompromising gatekeeper of scientific integrity while providing unparalleled visual intelligence.

9.1 Real-Time Data Hardening (The QA Hub)

Location in App: `QA Hub`

As animal technicians input information, the integrated `data_quality.py` engine instantly compares inputs against thermodynamic limits: * **Biological Range Locks:** Flags birds losing weight between measurement epochs (Negative Daily Gain) or exceeding maximum physiological thresholds (e.g., >150g/day ADG). * **FCR Thermodynamic Bounds:** Flags FCR ratios dropping below biologically conceivable baselines (e.g., FCR < 1.05). * **Z-Score Outlier Detection:** Mathematically identifies severe statistical outliers using standard deviation matrices, throwing visual warnings (🟢, 🟡, 🔴) for the Study Director to review before incorporating the data into ANOVA calculations.

9.2 Bulk Import Architecture

Location in App: `Bulk Import`

Facilities are not forced to abandon Excel. OPA's bulk parser acts as a universal ingestion engine. Technicians can continue writing weights on clipboards and typing them into standard Excel templates; OPA reads the sheet, sanitizes the data, and securely ingests up to 10,000 rows in seconds.

9.3 Advanced Dashboard Visualizations

OPA renders complex datasets into immediate visual intelligence via the UI dashboard: * **Intervention Overlays:** Automatically maps specific health interventions (vaccinations, medications, lighting shifts) as vertical chronological markers over time-series performance data (e.g., Body Weight or ADG curves) to visually correlate management actions with biological responses. * **Altair-Powered Score Distributions:** Renders Qualitative Scoring (e.g., Lesion Scores, Tona Scores) through interactive

Proportional vs. Stacked bar charts, clearly isolating shift severity between treatments. * **Detailed Mortality Analytics:** Generates multi-tiered visualizations grouping distinct *Causes of Death* by Treatment Group and physical Pen location, allowing directors to instantly spot spatial or dietary mortality correlations.

9.4 Automated PDF Report Generation

Location in App: `Dashboard -> Study Journal & Export`

Replacing the Statistician's Desk The `reports.py` formatting engine allows management to generate publication-ready PDF reports instantly. * Automatically compiles ANOVA p-values, Tukey HSD Compact Letter Displays, and calculated FCRs. * Renders high-resolution vector graphics of biological growth curves and margin charts. * Produces cleanly formatted tables specifically designed for physical printing and boardroom review.

Generated by the Ollis Poultry Analytics Computational Core.