Published August 21, 2014

and 1997-1998. Knowles produced significantly more seed and oil than Mermaid and Floral in both years. The mean seed yield for Knowles was 1014 kg ha^{-1} compared with 783 kg ha^{-1} for Floral and 688 kg ha^{-1} for Mermaid ($LSD_{\alpha=0.05}$ = 83 kg ha^{-1}). The mean seed oil concentration for Knowles was 297 g kg^{-1} compared with 277 g kg^{-1} for Floral and 272 g kg^{-1} for Mermaid ($LSD_{\alpha=0.05}$ = 12 g kg^{-1}). Knowles grows upright, can be machine harvested, and is well adapted as a winter annual in the Willamette Valley of Oregon. The fatty acid profile of Knowles is similar to that of Floral and *L. alba* ssp. *alba* germplasm (4, 5). The mean 20:1Δ5, erucic acid (22:1Δ13), and 22:2Δ5,Δ13 concentrations for Knowles were 647, 126, and 170 g kg^{-1}, respectively, versus 642, 125, and 168 g kg^{-1}, respectively, for Floral. The 1000-seed weight for Knowles was 9.9 g compared with 9.5 g for Floral and 9.2 g for Mermaid. There were no significant differences in days to flowering or days to physiological maturity between Knowles, Mermaid, and Floral.

Knowles is owned by Oregon State University. Foundation, Registered, and Certified seed of Knowles have been produced by the Oregon Meadowfoam Growers, Inc. (Salem, Oregon). Seed multiplication will be restricted to these classes. U.S. plant variety protection of Knowles has been applied for (PVP Certificate no. 9900298). Breeder seed of Knowles is maintained by the Oregon Agricultural Experiment Station. Small quantities of seed for research purposes may be obtained from the corresponding author for at least five years.

J.M. Crane and S.J. Knapp* (6)

References and Notes

1. Calhoun, W., and J.M. Crane. 1984. Registration of Mermaid meadowfoam. Oregon Agric. Exp. Stn., Corvallis.
2. Jolliff, G.D. 1986. Plant variety protection certificate (#8500166) for Mermaid meadowfoam. USDA., Washington, DC.
3. Jolliff, G.D. 1994. Plant variety protection certificate (#9200257) for Floral meadowfoam. USDA., Washington, DC.
4. Knapp, S.J., and J.M. Crane. 1995. Fatty acid diversity of Section Inflexae *Limnanthes* (meadowfoam). Ind. Crops Products 4: 219–227.
5. Knapp, S.J., and J.M. Crane. 1998. A dominant gene decreases erucic acid and increases dienoic acid in meadowfoam. Crop Sci. 38:1541–1544.
6. J.M. Crane and S.J. Knapp, Dep. of Crop and Soil Science, Oregon State Univ., Corvallis, OR 97331. Published as Oregon Agric. Exp. Stn. Technical Paper no. 11,436. Registration by CSSA. Accepted 31 May 1999. *Corresponding author (knapps@css.orst.edu)

Published in Crop Sci. 40:291–292 (2000).

Registration of 'Jeokwangtangkong' Peanut

'Jeokwangtangkong' peanut (*Arachis hypogaea* L. subsp. *fastigiata* var. *vulgaris*) (Reg. no. CV-63, PI 607913) is a spanish-type developed at the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Andhra Pradesh, India. It was introduced into the Republic of Korea in 1989 as ICGV 86326. After five years of evaluation, it was released in 1996 as Jeokwangtangkong for cultivation in Korea.

Jeokwangtangkong was developed following the bulk breeding method from a cross between a spanish genotype ICGV 87124 and a virginia (*A. hypogaea* subsp. *hypogaea* var. *hypogaea*) bunch cultivar G 201 (also known as 'Kaushal'). The former is an advanced breeding line developed at ICRISAT, and the latter is a released cultivar in India (1). The pedigree of Jeokwangtangkong is ICGV 87124 / G 201 F_2-B_1-B_1-B_1-B_1-B_1.

Jeokwangtangkong was evaluated with the local control 'Daekwangtangkong' for five years (1992–1996) in Korea. It produced an average 6.7% greater seed yield than Daekwangtangkong (2.82 t ha^{-1}), and matured 2 wk earlier (149 d). Because of its earlier maturation, the Korean farmers are now able to introduce double cropping (either leafy vegetable crops before peanut or winter barley after peanut) into their farm lands (3).

Jeokwangtangkong has an erect growth habit (2) with large size obovate dark green leaves. The average number of primary and secondary branches is 11. The average height of the main axis is 39 cm, and the average length of primary branches is 43 cm. It has 2-seeded small pods with slight to moderate pod constriction, moderate pod beak, and slight pod reticulation and ridges. It has meat content of 73%. Its seeds are light red in color, with a 100-seed weight of 49 g. The seeds contain 50% oil and 21% protein. The oleic/linoleic fatty acid ratio is 1.4. Jeokwangtangkong is highly resistant to pod rots [caused by *Fusarium solani* (Mart.) Sacc. f. sp. *phaseoli* (Burkholder) W.C. Snyder & H.N. Hans] (3).

Limited quantities of seed of Jeokwangtangkong can be obtained from the National Honam Agricultural Experiment Station, RDA, Iksan 570-080, Korea (fax 0653-840-2111). Seed has also been deposited with the U.S. National Seed Storage Laboratory, 1111 Mason St., Fort Collins, CO 80521-4500. U.S. plant variety protection of Jeokwangtangkong will not be applied for.

Youn-Sup Oh, Young-Keun Cheong, Jong-Chul Ko, Jong-Tae Kim, Myung-Kyu Oh, Jung-Gon Kim, Ki-Hun Park, Young-Sun Jang, Moon-Soo Park, Su-Yeon Cho, S.L. Dwivedi,* and S.N. Nigam (4)

References and Notes

1. Basu, M.S., and P.S. Reddy. 1987. Groundnut varieties of India. Tech. Bull. 2. Nat. Res. Ctr. for Groundnut (Indian Council of Agric. Res.), Junagadh, Gujarat, India.
2. IBPGR and ICRISAT. 1992. Descriptors for groundnut. International Board of Plant Genetic Resources, Rome, and International Crops Research Institute for the Semi-Arid Tropics, Patancheru, AP, India.
3. Oh, Y.S., Y.K. Cheong, J.C. Ko, J.T. Kim, M.K. Oh, J.G. Kim, K.H. Park, Y.S. Jang, M.S. Park, and S.Y. Cho. 1996. A small-seeded, disease-resistant peanut variety "Jeokwangtangkong". Report jointly prepared by Nat. Honam Agric. Exp. Stn., RDA, Iksan 570-080, and Res. Manage. Bureau, RDA, Suwon 441-707, Korea (for limited distribution).
4. Y.S. Oh, Y.K. Cheong, J.C. Ko, J.T. Kim, M.K. Oh, J.G. Kim, Y.S. Jang, M.S. Park, and S.Y. Cho. Nat. Honam Agr. Exp. Stn., RDA, Iksan 570-080, Korea; Ki-Hun Park, Research Management Bureau, RDA, Suwon 441-707, Korea; and S.L. Dwivedi and S.N. Nigam, ICRISAT, Patancheru, A P, India. ICRISAT Journal Article no. 2304. Registration by CSSA. Accepted 30 June 1999. *Corresponding author (s.dwivedi@cgiar.org).

Published in Crop Sci. 40:292 (2000).

Registration of 'Plainsman' Rapeseed

'Plainsman' winter rapeseed [*Brassica napus* L. subsp. *oleifera* (Metzg.) Sinsk. f. *biennis*] (Reg. no. CV-18, PI 604607) was developed by the Kansas Agricultural Experiment Station and released in 1998. Plainsman has edible oil (canola quality) and improved winter survival compared with other rapeseed cultivars tested under Great Plains conditions. Plainsman was selected from the cross WRER12/'Bienvenu' made in 1987 at the Idaho Agricultural Experiment Station, Moscow, ID. Seed of WRER12 is low in erucic acid and in glucosinolates (<20 g kg^{-1} erucic acid in the oil and <30 μmol g^{-1} glucosinolates in the oil-free meal), and was selected in the F_6 generation from the cross 'Indore'/'Sipal'//'Liraglu'. WRER12 is one of