

# Numerical Summaries

# Summarizing data in R

- Have seen `summary` (5-number summary of each column). But what if we want:
  - a summary or two of just one column
  - a count of observations in each category of a categorical variable
  - summaries by group
  - a different summary of all columns (eg. SD)
- To do this, meet pipe operator `%>%`. This takes input data frame, does something to it, and outputs result. (Learn: `Ctrl-Shift-M`.)
- Output from a pipe can be used as input to something else, so can have a sequence of pipes.
- Summaries include: mean, median, min, max, sd, IQR, quantile (for obtaining quartiles or any percentile), n (for counting observations).
- Use our Australian athletes data again.

## Packages for this section

```
library(tidyverse)
```

# Summarizing one column

- Mean height:

```
athletes %>% summarize(m=mean(Ht))
```

m
180.104

or to get mean and SD of BMI:

```
athletes %>% summarize(m=mean(BMI), s=sd(BMI))
```

m	s
22.95589	2.863933

# Quartiles

- quantile calculates percentiles (“fractiles”), so we want the 25th and 75th percentiles:

```
athletes %>% summarize( Q1=quantile(Wt, 0.25),  
                        Q3=quantile(Wt, 0.75))
```

Q1	Q3
66.525	84.125

# Creating new columns

- These weights are in kilograms. Maybe we want to summarize the weights in pounds.
- Convert kg to lb by multiplying by 2.2.
- Create new column and summarize that:

```
athletes %>% mutate(wt_lb=Wt*2.2) %>%  
  summarize(Q1_lb=quantile(wt_lb, 0.25),  
            Q3_lb=quantile(wt_lb, 0.75))
```

Q1_lb	Q3_lb
146.355	185.075

# Counting how many

for example, number of athletes in each sport:

```
athletes %>% count(Sport)
```

Sport	n
BBall	25
Field	19
Gym	4
Netball	23
Row	37
Swim	22
T400m	29
Tennis	11
TSprnt	15
WPolo	17

## Counting how many, variation 2:

Another way (which will make sense in a moment):

```
athletes %>% group_by(Sport) %>%  
  summarize(count=n())
```

```
## `summarise()` ungrouping output (override with `.groups` argument)
```

Sport	count
BBall	25
Field	19
Gym	4
Netball	23
Row	37
Swim	22
T400m	29
Tennis	11
TSprnt	15
WPolo	17



# Summaries by group

- Might want separate summaries for each “group”, eg. mean and SD of height for males and females. Strategy is `group_by` (to define the groups) and then `summarize`:

```
athletes %>% group_by(Sex) %>%  
  summarize(m=mean(Ht), s=sd(Ht))
```

```
## `summarise()` ungrouping output (override with `.groups` argument)
```

Sex	m	s
female	174.5940	8.242203
male	185.5059	7.903487

- This explains second variation on counting within group: “within each sport, how many athletes were there?”

# Summarizing several columns

- Standard deviation of each (numeric) column:

```
athletes %>% summarize_if(is.numeric, sd)
```

RCC	WCC	Hc	Hg	Ferr	BMI	SSF	%Bfat	LBM	Ht
0.457976	4.800549	93.662989	1.362451	147.50124	22.863933	32.565336	18.189826	13.07029	173.449

- Median and IQR of all columns whose name starts with H:

```
athletes %>% summarize_at(vars(starts_with("H")),  
                           list(med=median, iqr=IQR))
```

Hc_med	Hg_med	Ht_med	Hc_iqr	Hg_iqr	Ht_iqr
43.5	14.7	179.7	4.975	2.075	12.175