

Mood's Median Test

Packages

```
library(tidyverse)  
library(smmr)
```

Two-sample test: What to do if normality fails

- If normality fails (for one or both of the groups), what do we do then?
- Again, can compare medians: use the thought process of the sign test, which does not depend on normality and is not damaged by outliers.
- A suitable test called Mood's median test.
- Before we get to that, a diversion.

The chi-squared test for independence

Suppose we want to know whether people are in favour of having daylight savings time all year round. We ask 20 males and 20 females whether they each agree with having DST all year round ("yes") or not ("no"). Some randomly chosen rows of data:

```
my_url <- "http://ritsokiguess.site/datafiles/dst.txt"
dst <- read_delim(my_url, " ")
dst %>% slice_sample(n = 8)
```

```
# A tibble: 8 x 2
```

	gender	agree
	<chr>	<chr>
1	male	yes
2	female	yes
3	female	no
4	female	yes
5	male	yes
6	male	yes
7	male	yes
8	female	yes

... continued

Count up individuals in each category combination, and arrange in contingency table:

```
tab <- with(dst, table(gender, agree))
tab
```

	agree	
gender	no	yes
female	11	9
male	3	17

- Most of the males say “yes”, but the females are about evenly split.
- Looks like males more likely to say “yes”, ie. an association between gender and agreement.
- Test an H_0 of “no association” (“independence”) vs. alternative that there is really some association.
- Done with `chisq.test`.

...And finally

```
chisq.test(tab, correct=FALSE)
```

Pearson's Chi-squared test

data: tab

X-squared = 7.033, df = 1, p-value = 0.008002

- Reject null hypothesis of no association (P-value 0.008)
- therefore there is a difference in rates of agreement between (all) males and females (or that gender and agreement are associated).
- This calculation gives same answers as you would get by hand. (Omitting `correct = FALSE` uses "Yates correction".

Mood's median test

- Earlier: compare medians of two groups.
- Sign test: count number of values above and below something (there, hypothesized median).
- Mood's median test:
 - ▶ Find “grand median” of all the data, regardless of group
 - ▶ Count data values in each group above/below grand median.
 - ▶ Make contingency table of group vs. above/below.
 - ▶ Test for association.
- If group medians equal, each group should have about half its observations above/below grand median. If not, one group will be mostly above grand median and other below.

Mood's median test for reading data

- Find overall median score:

```
kids %>% summarize(med=median(score)) %>% pull(med) -> m  
m
```

```
[1] 47
```

- Make table of above/below vs. group:

```
tab <- with(kids, table(group, score > m))  
tab
```

group	FALSE	TRUE
c	15	8
t	7	14

- Treatment group scores mostly above median, control group scores mostly below, as expected.

The test

- Do chi-squared test:

```
chisq.test(tab, correct=FALSE)
```

Pearson's Chi-squared test

data: tab

X-squared = 4.4638, df = 1, p-value = 0.03462

- This test actually two-sided (tests for any association).
- Here want to test that new reading method *better* (one-sided).
- Most of treatment children above overall median, so do 1-sided test by halving P-value to get 0.017.
- This way too, children do better at learning to read using the new method.

Or by smmr

- median_test does the whole thing:

```
median_test(kids, score, group)
```

```
$grand_median  
[1] 47
```

```
$table  
      above  
group above below  
  c      8     15  
  t     14      7
```

```
$test  
      what      value  
1 statistic 4.46376812  
2         df 1.00000000  
3   P-value 0.03462105
```

- P-value again two-sided.

Comments

- P-value 0.013 for (1-sided) t -test, 0.017 for (1-sided) Mood median test.
- Like the sign test, Mood's median test doesn't use the data very efficiently (only, is each value above or below grand median).
- Thus, if we can justify doing t -test, we should do it. This is the case here.
- The t -test will usually give smaller P-value because it uses the data more efficiently.
- The time to use Mood's median test is if we are definitely unhappy with the normality assumption (and thus the t -test P-value is not to be trusted).


A graph

- Two categorical variables (above/below and group), so grouped bar chart
- group is “explanatory” and above/below is “response”.
- Plot explanatory as x and plot response as fill (y will be frequency):

```
kids %>%  
  mutate(above = (score > m)) %>%  
  ggplot(aes(x = group, fill = above)) + geom_bar(position = 'dodge')
```

The graph

g



- The significant association is because the treatment group children mostly score above the grand median, and the control group children score mostly below.