

# Solving Integrable Boltzmann Equation for Coscattering Dark Matter

(Final Presentation for PHYS-GA-2000)

Xucheng Gan



Catch a glimpse of the equation  
we want to solve ; )

# Catch a glimpse of the equation we want to solve ; )

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2}\Delta^{1/2}y^4\delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$
$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

# Catch a glimpse of the equation we want to solve ; )

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2}\Delta^{1/2}y^4\delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$

$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}(x, R = \frac{m_i}{m_\chi})$$

# Catch a glimpse of the equation we want to solve ; )

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2}\Delta^{1/2}y^4\delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$

$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

$$r \sim 1$$

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}\left(x, R = \frac{m_i}{m_\chi}\right)$$

# Catch a glimpse of the equation we want to solve ; )

$$\begin{aligned}
 \frac{dY_\chi}{dx} = & -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2}\Delta^{1/2}y^4\delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right) \\
 & \times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)
 \end{aligned}$$

↗ ↗ ↗  
\$r \sim 1\$      \$\Delta = \frac{m\_\psi - m\_\chi}{m\_\chi} \sim 1\$

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}\left(x, R = \frac{m_i}{m_\chi}\right)$$

# Catch a glimpse of the equation we want to solve ; )

$$\begin{aligned}
 \frac{dY_\chi}{dx} = & -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2}\Delta^{1/2}y^4\delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right) \\
 & \times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)
 \end{aligned}$$

r ~ 1      Δ =  $\frac{m_\psi - m_\chi}{m_\chi} \sim 1$       y ~ 1

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}\left(x, R = \frac{m_i}{m_\chi}\right)$$

# Catch a glimpse of the equation we want to solve ; )

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2} \Delta^{1/2} y^4 \delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$

$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

$r \sim 1$

$\Delta = \frac{m_\psi - m_\chi}{m_\chi} \sim 1$

$y \sim 1$

$\delta \ll 1$

Small mixing angle

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp(\sqrt{\xi^2 + (Rx)^2}) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}(x, R = \frac{m_i}{m_\chi})$$

# Catch a glimpse of the equation we want to solve ; )

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2} \Delta^{1/2} y^4 \delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$

$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

$r \sim 1$

$\Delta = \frac{m_\psi - m_\chi}{m_\chi} \sim 1$

$y \sim 1$

$\delta \ll 1$

$\frac{M_{pl}}{m_\chi} \sim 10^{21}$

Small mixing angle

$$f(r) = \frac{(r^2 + r + 2)^2}{\sqrt{2}(r - 2)^2 r^{9/2} (r + 1)^{7/2}}$$

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp(\sqrt{\xi^2 + (Rx)^2}) - 1}$$

$$Y_i^{(eq)}(x) = \frac{45}{4\pi^4 \cdot g_{\star S}(m_\chi/x)} I_b^{(eq)}(x, R = \frac{m_i}{m_\chi})$$



# Boltzmann for Coscattering and Coannihilation Process

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \langle \sigma_{ij} v \rangle \left( n_i n_j - n_i^{(eq)} n_j^{(eq)} \right) \right]$$

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \langle \sigma_{ij} v \rangle \left( n_i n_j - n_i^{(eq)} n_j^{(eq)} \right) \right]$$

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \langle \sigma_{ij} v \rangle \left( n_i n_j - n_i^{(eq)} n_j^{(eq)} \right) \right]$$

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -n_\phi^{(eq)} \langle \sigma_{\chi \rightarrow \psi} v \rangle \left( n_\chi - n_\chi^{(eq)} \right)$$

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \cancel{\langle \sigma_{ij} v \rangle (n_i n_j - n_i^{(eq)} n_j^{(eq)})} \right]$$

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -n_\phi^{(eq)} \langle \sigma_{\chi \rightarrow \psi} v \rangle \left( n_\chi - n_\chi^{(eq)} \right)$$

$$\frac{dY_\chi}{dx} = -\frac{S(x)}{H(x)x} \langle \sigma_{\chi \rightarrow \psi} v \rangle Y_\phi^{(eq)} Y_\chi^{(eq)} \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \langle \sigma_{ij} v \rangle \left( n_i n_j - n_i^{(eq)} n_j^{(eq)} \right) \right]$$

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -n_\phi^{(eq)} \langle \sigma_{\chi \rightarrow \psi} v \rangle \left( n_\chi - n_\chi^{(eq)} \right)$$

$$\boxed{\frac{dY_\chi}{dx} = -\frac{S(x)}{H(x)x} \langle \sigma_{\chi \rightarrow \psi} v \rangle Y_\phi^{(eq)} Y_\chi^{(eq)} \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)}$$

# Boltzmann for Coscattering and Coannihilation Process

$$\frac{dn_i}{dt} + 3Hn_i = - \sum_j \left[ n_\phi^{(eq)} \langle \sigma_{i \rightarrow j} v \rangle \left( n_i - n_i^{(eq)} \frac{n_j}{n_j^{(eq)}} \right) + \cancel{\langle \sigma_{ij} v \rangle \left( n_i n_j - n_i^{(eq)} n_j^{(eq)} \right)} \right]$$

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -n_\phi^{(eq)} \langle \sigma_{\chi \rightarrow \psi} v \rangle \left( n_\chi - n_\chi^{(eq)} \right)$$

$$\boxed{\frac{dY_\chi}{dx} = -\frac{S(x)}{H(x)x} \langle \sigma_{\chi \rightarrow \psi} v \rangle Y_\phi^{(eq)} Y_\chi^{(eq)} \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)}$$

$$H(x) = \frac{H(m_\chi)}{x^2} \quad H(m_\chi) = \sqrt{\frac{\pi^2}{90} \frac{g_*(T)}{{M_{pl}}^2} m_\chi^4}$$

$$S(x) = \frac{S(m_\chi)}{x^3} \quad S(m_\chi) = \frac{2\pi^2}{45} g_* S(T) m_\chi^3$$



# The Dark Sector Lagrangian

# The Dark Sector Lagrangian

$$\mathcal{L} \supset -\frac{m_\chi}{2}\chi^2 - \frac{m_\psi}{2}\psi^2 - \boxed{\delta} m \chi \psi - \boxed{\frac{y}{2}} \phi \psi^2 + h.c.$$

# The Dark Sector Lagrangian

$$\mathcal{L} \supset -\frac{m_\chi}{2}\chi^2 - \frac{m_\psi}{2}\psi^2 - \boxed{\delta} m\chi\psi - \boxed{\frac{y}{2}}\phi\psi^2 + h.c.$$

$\delta$  will induce the mixing angle. So we need to diagonalize mass term to get the mass eigenstate. After diagonalizing the mass term,  $\delta$  will also enter the interaction term.

# The Dark Sector Lagrangian

$$\mathcal{L} \supset -\frac{m_\chi}{2}\chi^2 - \frac{m_\psi}{2}\psi^2 - \boxed{\delta} m\chi\psi - \boxed{y} \frac{1}{2}\phi\psi^2 + h.c.$$

$\delta$  will induce the mixing angle. So we need to diagonalize mass term to get the mass eigenstate. After diagonalizing the mass term,  $\delta$  will also enter the interaction term.

Finally using the Lagrangian(Toy model) given, we can calculate the thermal cross section( $x>2$ ).

$$\langle \sigma v_{Mol} \rangle = \frac{\int_{\tilde{s}=2(m_1^2+m_2^2)}^{\tilde{s}=+\infty} d\tilde{s} \quad \sigma(\tilde{s})F(\tilde{s})\sqrt{\tilde{s}-2(m_1^2+m_2^2)}K_1\left(\frac{\sqrt{\tilde{s}}}{T}\right)}{4T\left(m_1^2K_0\left(\frac{m_1}{T}\right)\right)\left(m_2^2K_0\left(\frac{m_2}{T}\right)\right)}$$

# The Dark Sector Lagrangian

$$\mathcal{L} \supset -\frac{m_\chi}{2}\chi^2 - \frac{m_\psi}{2}\psi^2 - \boxed{\delta} m\chi\psi - \boxed{y} \frac{1}{2}\phi\psi^2 + h.c.$$

$\delta$  will induce the mixing angle. So we need to diagonalize mass term to get the mass eigenstate. After diagonalizing the mass term,  $\delta$  will also enter the interaction term.

Finally using the Lagrangian(Toy model) given, we can calculate the thermal cross section( $x > 2$ ).

$$\langle \sigma v_{Mol} \rangle = \frac{\int_{\tilde{s}=2(m_1^2+m_2^2)}^{\tilde{s}=+\infty} d\tilde{s} \quad \sigma(\tilde{s}) F(\tilde{s}) \sqrt{\tilde{s} - 2(m_1^2 + m_2^2)} K_1\left(\frac{\sqrt{\tilde{s}}}{T}\right)}{4T \left(m_1^2 K_0\left(\frac{m_1}{T}\right)\right) \left(m_2^2 K_0\left(\frac{m_2}{T}\right)\right)}$$

$$\langle \sigma_{\chi \rightarrow \psi} v \rangle(x) = \left(\frac{m_\psi}{m_\chi}\right)^{\frac{3}{2}} e^{-\Delta x} \langle \sigma_{\psi \rightarrow \chi} v \rangle \approx \left(\frac{m_\psi}{m_\chi}\right)^{\frac{3}{2}} e^{-\Delta x} f(r) \sqrt{\Delta} \frac{y^4 \delta^2}{2\pi m_\chi^4}$$

# The Dark Sector Lagrangian

$$\mathcal{L} \supset -\frac{m_\chi}{2}\chi^2 - \frac{m_\psi}{2}\psi^2 - \boxed{\delta} m\chi\psi - \boxed{y} \frac{\phi}{2}\psi^2 + h.c.$$

$\delta$  will induce the mixing angle. So we need to diagonalize mass term to get the mass eigenstate. After diagonalizing the mass term,  $\delta$  will also enter the interaction term.

Finally using the Lagrangian(Toy model) given, Exponentially suppressed we can calculate the thermal cross section( $x > 2$ ) in low temperature


$$\langle \sigma v_{Mol} \rangle = \frac{\int_{\tilde{s}=2(m_1^2+m_2^2)}^{\tilde{s}=+\infty} d\tilde{s} \quad \sigma(\tilde{s}) F(\tilde{s}) \sqrt{\tilde{s} - 2(m_1^2 + m_2^2)} K_1(\frac{\sqrt{\tilde{s}}}{T})}{4T \left( m_1^2 K_0(\frac{m_1}{T}) \right) \left( m_2^2 K_0(\frac{m_2}{T}) \right)}$$

$$\langle \sigma_{\chi \rightarrow \psi} v \rangle(x) = \left( \frac{m_\psi}{m_\chi} \right)^{\frac{3}{2}} e^{-\Delta x} \langle \sigma_{\psi \rightarrow \chi} v \rangle \approx \left( \frac{m_\psi}{m_\chi} \right)^{\frac{3}{2}} e^{-\Delta x} f(r) \sqrt{\Delta} \frac{y^4 \delta^2}{2\pi m_\chi^4}$$




# Effective Number of Relativistic Degree of Freedom


# Effective Number of Relativistic Degree of Freedom




# Effective Number of Relativistic Degree of Freedom




# Effective Number of Relativistic Degree of Freedom




# Effective Number of Relativistic Degree of Freedom



# Effective Number of Relativistic Degree of Freedom



# Effective Number of Relativistic Degree of Freedom





# Some Tricky Things About Doing Integration using Python

# Some Tricky Things About Doing Integration using Python

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi - \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

# Some Tricky Things About Doing Integration using Python

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$I_b^{(eq)}(x, R) = \begin{cases} \int_0^{+\Lambda} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1} & 0 \leq x < 23 \\ (Rx)^2 K_2(Rx) & x \geq 23 \end{cases}$$

# Some Tricky Things About Doing Integration using Python

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$I_b^{(eq)}(x, R) = \begin{cases} \int_0^{r+\Lambda} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1} & 0 \leq x < 23 \\ (Rx)^2 K_2(Rx) & x \geq 23 \end{cases}$$

# Some Tricky Things About Doing Integration using Python

$$I_b^{(eq)}(x, R) = \int_0^{+\infty} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1}$$

$$I_b^{(eq)}(x, R) = \begin{cases} \int_0^{r+\Lambda} d\xi \frac{\xi^2}{\exp\left(\sqrt{\xi^2 + (Rx)^2}\right) - 1} & 0 \leq x < 23 \\ (Rx)^2 K_2(Rx) & x \geq 23 \end{cases}$$



**Go Back to the Equation  
We Want to Solve. lol**

# Go Back to the Equation We Want to Solve. lol

$$\frac{dY_\chi}{dx} = -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2} \Delta^{1/2} y^4 \delta^2 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)$$
$$\times I_b^{(eq)}(x, R = r) I_b^{(eq)}(x, R = 1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right)$$

The diagram illustrates the simplifications made to the differential equation. Four parameters are highlighted in boxes and connected by arrows to specific terms in the equation:

- A cyan box labeled  $r \sim 1$  has an arrow pointing to the term  $I_b^{(eq)}(x, R = r)$ .
- A yellow box labeled  $\Delta = \frac{m_\psi - m_\chi}{m_\chi} \sim 1$  has an arrow pointing to the term  $(1 + \Delta)^{3/2}$ .
- A green box labeled  $y \sim 1$  has an arrow pointing to the term  $y^4$ .
- A pink box labeled  $\delta \ll 1$  has an arrow pointing to the term  $\delta^2$ .

On the right side of the equation, there is a red textured box containing the expression  $\frac{M_{pl}}{m_\chi} \sim 10^{21}$ .

# Go Back to the Equation We Want to Solve. lol

$$\begin{aligned}
 \frac{dY_\chi}{dx} = & -\frac{45^{3/2}}{2^{7/2}\pi^8} \frac{1}{g_{\star S}(m_\chi/x)\sqrt{g_\star(m_\chi/x)}} \left( f(r)(1+\Delta)^{3/2} \Delta^{1/2} y^4 \delta^2 \right) \\
 & \times I_b^{(eq)}(x, R=r) I_b^{(eq)}(x, R=1) \left( \frac{Y_\chi}{Y_\chi^{(eq)}} - 1 \right) \left( \frac{M_{pl}}{m_\chi} \right) \left( \frac{e^{-\Delta x}}{x^2} \right)
 \end{aligned}$$


r ~ 1       $\Delta = \frac{m_\psi - m_\chi}{m_\chi} \sim 1$        $y \sim 1$        $\delta \ll 1$        $\frac{M_{pl}}{m_\chi} \sim 10^{21}$

Only when this term =  $10^{-30}$ ,  
my code(RK4) works :-(




# Dark Matter Freeze Out


# Dark Matter Freeze Out



# Dark Matter Freeze Out




# Dark Matter Freeze Out






# The Problem I met

# The Problem I met



# The Problem I met



# The Things I want to do later

- Make the calculation works for arbitrary value of  $\delta$ .
- Reproduce the consequence of WIMP miracle as benchmark.
- Plot the deviation of  $I_b^{(eq)}(x, R)$  using my python code and exact value ( Using Mathematica ).
- Plot the freeze-out number density when parameters  $r$  and  $\Delta$  .
- Solve the non-integrable Boltzmann and compare with integrable one.  
(If TIME permitts)