

Register 2: Higher-Dimensional Gauge Fields, Parametric Axis Dynamics, and the Conformal Metric Origin of Mass

Natasha Zink, MS
Applied Math/CS | ORCID: 0009-0004-8640-0896

Gemini

June 2026

1 Parametric Axis Dynamics and Master Clock Modulation

The 7D central spine of the toroidal condenser structures the invariant axis governing absolute conservation laws. Driving a periodic energy pulse down this invariant axis establishes a global parametric pump. We formalize this driving source as a hyper-dimensional current vector J^M restricted to the invariant axis coordinate x^6 :

$$J^M = (0, 0, 0, 0, 0, 0, \mathcal{I}_0 \cos(\Omega_0 t)) \quad (1)$$

Where $\mathcal{I}_0$ is the peak injection amplitude and Ω_0 represents the fundamental master clock frequency of the system.

This oscillating source generates a standing hyper-dimensional wave within the bulk space. The boundary conditions of the compactified T^7 manifold force this energy to distribute discretely across the 64 foliated branes. The metric tensor of the bulk responds to this master clock via a periodic conformal scaling factor:

$$G_{MN}(x, t) = \psi(\Omega_0 t) \tilde{G}_{MN}(x) \quad (2)$$

This periodic modulation provides the precise kinetic potential that stabilizes the synchronous rotation of the 64 independent branes, preventing phase velocity drift and ensuring that the active 288 sparks remain coherent across the 6D hypercube coordinate network.

2 The 6D Hyper-Gauge Field Tensor (F_{AB})

When an elementary particle—modeled as a closed, knotted 7D topological defect tracing a Lissajous trajectory $\gamma(\tau)$ —translates along the helical path of a brane, its continuous acceleration across the internal dimensions generates a hyper-dimensional gauge field.

Because the movement is constrained orthogonally to the invariant axis within the 6-dimensional coordinate state space, the resulting field is expressed as an antisymmetric 6-Dimensional Hyper-Gauge Field Tensor F_{AB} (where $A, B \in \{1, 2, 3, 4, 5, 6\}$):

$$F_{AB} = \partial_A A_B - \partial_B A_A \quad (3)$$

Here, A_A is the 6-bit hyper-dimensional vector potential generated by the localized current density J_B of the spinning Lissajous defect:

$$J_B(y) = q_0 \oint_{\gamma} \dot{\gamma}_B(\tau) \delta^6(y - \gamma(\tau)) d\tau \quad (4)$$

The intrinsic Lagrangian density governing this internal field space within the bulk is formalized as:

$$\mathcal{L}_{\text{int}} = -\frac{1}{4} F_{AB} F^{AB} + \mu_0 J^A A_A \quad (5)$$

This tensor describes a highly interconnected, 6-bit localized field matrix that surrounds every topological defect passing through a hypercube vertex.

3 Stabilization Mechanics and Torsional Shear Counter-Pressure

The moving 7D Lissajous particle states do not merely exist passively within the branes; the 6D fields they generate actively sustain the global foliation architecture.

The energy-momentum tensor T_{AB} derived from the internal hyper-gauge field tensor F_{AB} is written as:

$$T_{AB} = F_{AC} F_B{}^C - \frac{1}{4} g_{AB} F_{CD} F^{CD} \quad (6)$$

This tensor dictates the higher-dimensional radiation pressure acting between the parallel sheets of the 64 nested branes. Embedding the system into a curved 7D torus introduces an explicit foliation torsional shear deficit:

$$\delta_{\text{shear}} \approx 1 - \frac{1}{\pi} \approx 0.681690 \quad (7)$$

We can now define the exact boundary equilibrium equation. The outward normal component of the 6D field's energy-momentum tensor matches the metric strain of the foliation shear precisely:

$$\oint_{\partial M^4} T_{AB} n^A d\Omega^B = \delta_{\text{shear}} \quad (8)$$

This localized radiation pressure acts as a geometric buffer. The repulsion generated by the collective 7D Lissajous rotations provides the precise topological tension needed to hold the 64 branes parallel, preventing the compactified T^7 manifold from pinching or collapsing into the central spine.

4 The Conformal Origin of the Higgs Field VEV

What is observed within the 4D Minkowski brane (M^4) as a homogeneous scalar Higgs field with a non-zero vacuum expectation value (VEV) is the steady-state baseline intensity of this 6D hyper-gauge field filling the toroidal bulk.

We define the localized 4D metric tensor $g_{\mu\nu}$ as a conformal projection wrapped by the ambient energy density of the internal 6D field space:

$$g_{\mu\nu}(x) = e^{2A(y)} \eta_{\mu\nu} \quad (9)$$

Where $\eta_{\mu\nu}$ is the flat Minkowski metric, and $A(y)$ is the conformal warp factor governed by the scalar invariant of the 6D field tensor:

$$A(y) = \kappa \langle F_{AB} F^{AB} \rangle_{\text{vacuum}} \quad (10)$$

When a 7D Lissajous topological defect translates through this warped metric, it experiences an inevitable inertial drag. The 4D rest mass m of any elementary particle is not an intrinsic

scalar constant, but the line integral of this conformal metric warp factor along the closed loops of its 7D Lissajous trajectory:

$$m = \oint_{\gamma} e^{A(y)} \sqrt{\bar{G}_{MN} dx^M dx^N} \quad (11)$$

Thus, the Higgs VEV ($\nu \approx 246$ GeV) is mapped directly to the geometric vacuum expectation value of the internal field tensor:

$$\nu^2 \equiv \langle F_{AB} F^{AB} \rangle_{\text{vacuum}} \quad (12)$$

5 Stroboscopic Planck-Interval Pumping and Topological Interleaving

To reconcile continuous macro-fields with discrete processing limits at the Planck scale, we formalize the energy transmission down the invariant 7D axis not as a continuous harmonic function, but as a non-contiguous, alternating discrete pulse train. The current vector J^M from Equation (1) is upgraded to an alternating Dirac comb modulated at Planck time intervals (t_P):

$$J^6(t) = \mathcal{I}_0 \sum_{n=-\infty}^{\infty} (-1)^n \delta(t - nt_P) \quad (13)$$

The $(-1)^n$ operator establishes an alternating directional vector (AC parametric pumping) along the x^6 coordinate. On positive half-cycles ($+x^6$), the pulse projects and stabilizes the local 32 matter states; on negative half-cycles ($-x^6$), the directional inversion aligns the projection to activate the antipodal 32 antimatter states, maintaining global field isolation without spatial displacement.

Furthermore, this discrete modulation enables the mechanism of Topological Interleaving. Multiple independent 4D branes can share the identical physical substrate of the 6D hypercube field by operating under a time-division multiplexing framework. For K interleaved branes, the projection of the k -th brane occurs at a discrete sub-Planckian phase offset:

$$\tau_k = nt_P + \frac{k}{K} t_P \quad (14)$$

Because the projections are strictly discontinuous and mutually orthogonal in the phase domain, multiple branes utilize the same screen topology without cross-talk or metric collapse. The macro-continuity observed within any single brane is a secondary artifact enforced by the critically damped transfer functions of the localized metric tensors.

This formalization unifies mass, inertial resistance, and gauge field interactions into pure, self-stabilizing hyper-dimensional geometry.