

量子力学的共鳴を扱うための 統一的な完全WKB解析の枠組み

森川億人

理研iTHEMS


19/9/2025 日本物理学会第80回年次大会 @広島大学

- OM and S. Ogawa, PTEP [arXiv:2503.18741 [hep-th]].
- OM and S. Ogawa, JHEP [arXiv:2505.02301 [hep-th]].
- OM and S. Ogawa, arXiv:2508.09211 [quant-ph].

(Collaboration with nuclear physicist)

Introduction to “Resonance”

- Interesting in **quasi-stationary** potentials with developments of **non-perturbative** approaches
 - ▶ scattering theory, nuclear physics, open system, **non-Hermitian** QM.
- (Usual) Quasi-stationary state in quantum system
 - ▶ There are some local minima (vacua); one decays to an other.
- Unstable state after decay: **Resonant state**




- ▶ Simply say, plane wave in asymptotic region
 - ★ But, discrete and complex: $k = k_R - ik_I \in \mathbb{C}$.

Subtleties of resonance physics

- Resonance appears quite universally!
 - ▶ Traditional viewpoint: pole of S-matrix
- Some puzzling points:
 - ▶ Non-normalizable (divergence of norm)
 - ▶ Completeness?
 - ▶ What is the wave function itself?
- Some regularization schemes in phenomenological senses
 - ▶ Zel'dovich regularization, **complex scaling method**, rigged Hilbert space, etc.
 - ▶ No transparent relation has been found.
- Can resurgence theory (e.g., exact WKB) overcome difficulties as **non-perturbative formulation of QM**?
- If so, *essence in “ $\forall QM$ ” = analyticity in resurgence*

Review 1: Prescription of complex scaling

- Complex scaling method
Radial direction $r \rightarrow re^{i\theta}$



- Extended Hilbert space with resonance and θ
- Normalizable resonant wave function $\psi_R \sim$ “bound”
- $$\sum_B |\psi_B\rangle\langle\psi_B| + \sum_{R_\theta} |\psi_{R_\theta}\rangle\langle\tilde{\psi}_{R_\theta}| + \int_{S_\theta} |\psi_{S_\theta}\rangle\langle\tilde{\psi}_{S_\theta}| = 1$$
- Eigenvalue problem!

[See Myo–Kikuchi–Masui–Katō '14]

- Works well owing to sophisticated mathematical background
[Aguilar—Balslev—Combes (ABC) theorem AC '71, BC '71]
- Physical meaning??? What is observed???

Review 2: WKB ansatz and Borel resummation

- Consider a Schrödinger equation


$$\left(-\frac{d^2}{dx^2} + \hbar^{-2}Q(x)\right)\psi = 0, \quad Q(x) = 2[V(x) - E]$$

- Introduce WKB ansatz as a formal power series

$$\psi(x, \hbar) = e^{\int^x S(x', \hbar) dx'}, \quad S(x, \hbar) = \frac{1}{\hbar}S_{-1}(x) + S_0(x) + \hbar S_1(x) + \dots$$


- WKB ansatz is asymptotic series; where is it Borel summable?

$$\text{Stokes curve : } \left\{ x \left| \operatorname{Im} \int_a^x dx' \sqrt{Q} = 0, \text{ where } Q(x=a) = 0 \right. \right\}$$



Quantization condition from normalizability

- (Usual) QM: normalizable from $x \rightarrow -\infty$ to $x \rightarrow \infty$
 - ▶ E.g., Harmonic oscillator




- ▶ $\psi_{\text{final}} = (\text{analytic continuation, Stokes pheno}) \times \psi_{\text{initial}}$
- ▶ Non-normalizable solution vanishes: $\psi_{\text{initial}}^- \not\rightarrow \psi_{\text{final}}^+$


$$1 + A = 1 + e^{\hbar^{-1} \oint \sqrt{Q}} = 0 \quad \Rightarrow \quad E = \hbar \left(n + \frac{1}{2} \right)$$

How to see resonant state from exact WKB


- Resonant state: quasi-stable in complex region



- ▶ Inverted Rosen–Morse: $V = 1/\cosh^2 x$



Quantization path (leading order) [JPS 9/24]



- Normalizable path:
- Barrier resonant energy from exact WKB analysis (leading)

▶ A-cycle

$$\int_{-\cosh^{-1} \sqrt{\frac{1}{E}}}^{\cosh^{-1} \sqrt{\frac{1}{E}}} \sqrt{2 \left(\frac{1}{\cosh^2 x} - E \right)} = \sqrt{2} \pi (1 + \sqrt{E})$$

quantization condition: $1 - A = 0$ $E = \left(1 - \frac{in}{\sqrt{2}} \right)^2$

Rigorous estimate being identical to exact solution

- $A = 1$:

$$\left[\frac{F\left(-\frac{ik}{\beta} - s, -\frac{ik}{\beta} + s + 1, -\frac{ik}{\beta} + 1, \frac{1-|\xi|}{2}\right)}{F\left(-\frac{ik}{\beta} - s, -\frac{ik}{\beta} + s + 1, -\frac{ik}{\beta} + 1, \frac{1+|\xi|}{2}\right)} \right]^2 = 1,$$

where $|\xi| = \tanh(\beta \cosh^{-1} \sqrt{U_0/E})$.

- Here, we use the formula of the hypergeometric function as


$$\begin{aligned} & F\left(-\frac{ik}{\beta} - s, -\frac{ik}{\beta} + s + 1, -\frac{ik}{\beta} + 1, \frac{1 \pm |\xi|}{2}\right) \\ &= \mathfrak{A} F\left(-\frac{ik}{2\beta} - \frac{s}{2}, -\frac{ik}{2\beta} + \frac{s+1}{2}, \frac{1}{2}, |\xi|^2\right) \mp \mathfrak{B} F\left(-\frac{ik}{2\beta} - \frac{s-1}{2}, -\frac{ik}{2\beta} + \frac{s}{2} + 1, \frac{3}{2}, |\xi|^2\right), \end{aligned}$$

where

$$\mathfrak{A} = \frac{\Gamma\left(-\frac{ik}{\beta} + 1\right) \Gamma\left(\frac{1}{2}\right)}{\Gamma\left(-\frac{ik}{2\beta} - \frac{s-1}{2}\right) \Gamma\left(-\frac{ik}{2\beta} + \frac{s}{2} + 1\right)}, \quad \mathfrak{B} = \frac{\Gamma\left(-\frac{ik}{\beta} + 1\right) \Gamma\left(-\frac{1}{2}\right)}{\Gamma\left(-\frac{ik}{2\beta} - \frac{s}{2}\right) \Gamma\left(-\frac{ik}{2\beta} + \frac{s+1}{2}\right)}.$$

- $\mathfrak{A} = 0, \mathfrak{B} = 0$: odd/even number of nodes

What is problematic in naive path?




Regularization schemes

- Zel'dovich transformation: $S = e^{-\epsilon x^2}$ ['61, Berggren '68]

$$H\psi = E\psi \rightarrow H_S\psi_S = E\psi_S, \quad \psi_S \equiv S\psi, \quad H_S \equiv SHS^{-1}.$$

- ▶ We find $H_S = H - \frac{\hbar^2 \epsilon}{m} \left(\frac{d}{dx} x + x \frac{d}{dx} \right)$
- ▶ Now, $e^{I_\infty} \rightarrow e^{\int (S_{\text{odd}} - 2\epsilon x)}$ becomes finite.

- Complex scaling by θ rotates the Stokes graph as



- ▶ Now, $e^{I_\infty} \rightarrow e^{I'_\infty}$ becomes finite.
- ▶ ABC theorem described in a physical sense.

Backup: Rigged Hilbert space

- We introduce

$$\mathcal{D}_\varepsilon^R = \{x \in \mathbb{C} \mid \varepsilon > 0, \lim_{r \rightarrow \pm\infty} |x - r| < \varepsilon\}$$

- ▶ $\mathcal{D}_\varepsilon^R$ is the most crucial singular region.
- if $x \in \mathbb{C} \setminus \mathcal{D}_\varepsilon^R$, the Hilbert space is well-defined, $\mathcal{H}_\varepsilon$
 - ▶ Def. of norm and inner product is quite different!
 - ▶ Resonance is included as “bound” state.
- (Operator algebra) Set of operators $\{A_i\}$ where A_i is defined on $D(A_i) \subset \mathcal{H}_\varepsilon$; then let us introduce the dense subspace

$$\Phi \equiv \cap_i D(A_i) \subseteq \mathcal{H}_\varepsilon.$$

- The range is defined by the limit as $\varepsilon \rightarrow 0$, say, $\Phi^\times$
- We find the Gelfand multiplet

$$\Phi \subseteq \mathcal{H}_\varepsilon \subset \Phi^\times,$$

and $(\mathcal{H}_\varepsilon, \Phi)$ is the rigged (艀装) Hilbert space.

Summary

- We develop a **unified** framework for analyzing quantum mechanical **resonances** using **exact WKB method**
 - ▶ Serving as a non-perturbative formulation of QM.
 - ▶ Incorporating the Zel'dovich regularization, the complex scaling method, and the rigged Hilbert space.
 - ▶ Demonstrated by examining the inverted Rosen–Morse potential.
- Future works:
 - ▶ Realistic potential (E.g., $1/\cosh^2(x-a) + 1/\cosh^2(x+a)$)
 - ▶ Numerical approach (Quantization cond $\rightarrow$ Estimate cycles)
 - ▶ Higher dim???

Backup: Resonant state and S-matrix


- Let φ_l be regular solution of radial Schrödinger equation
 - φ_l is subject to

$$\varphi_l(k, r) \xrightarrow{r \rightarrow 0} j_l(kr) \quad j_l: \text{Spherical Bessel function}$$

with angular momentum l .

- At $r \rightarrow \infty$, linear combination of Hankel functions h_l :

$$\varphi_l(k, r) \rightarrow \mathcal{F}_l(k) h_l^-(kr) - \mathcal{F}_l^*(k) h_l^+(kr) \quad \mathcal{F}_l: \text{Jost function}$$



- S-matrix is written in terms of Jost function

$$S_l(k) = \frac{\mathcal{F}_l^*(k)}{\mathcal{F}_l(k)} \rightarrow \infty \quad \text{pole singularity}$$

- Some kind of scattering of multiple particles