

Gradient Structures from Classical to Quantum

Oliver Tse

Eindhoven University of Technology

Updated on December 2, 2025

Tutorial notes given at IPAM in Spring 2025

Abstract.

Contents

1	Noncommutative probability	1
1.1	From commutative to noncommutative	1
1.2	Observables	3
2	Stochastic Dilation	5
2.1	Noncommutative random variables	5
2.2	From commutative to noncommutative	7
2.3	Stochastic processes on matrix algebras	13
2.4	Stochastic processes on the unitary group	13
2.4.1	The unitary group and its Lie structure	13
2.4.2	Unitary-valued Brownian motion (rank- r noise)	14
2.4.3	Girsanov transform on $U(n)$ (Karandikar type)	15
2.4.4	Induced Girsanov on pure states	16
2.5	Dilation of semigroups (Sz.Nagy)	17
2.6	Stochastic dilations: How to make a heat bath	17
2.7	Interpretation of LDPs on unitary evolutions	17
2.8	LDP via Gärtner-Ellis revisited	18
2.8.1	Example	19
2.9	Groups to algebras	22
3	Boson Fock spaces and quantum noise	24
3.1	Boson Fock spaces	24
3.1.1	Weyl and field operators	24
3.1.2	Second quantization	26
3.1.3	Free field operators	28

3.1.4	Gaussian states	28
3.1.5	States invariant under free evolutions	28
3.1.6	Boson Fock white noise	30
3.1.7	Gaussian equilibrium states: The KMS condition	31
3.2	Composite systems	31
3.2.1	The reservoir	31
3.2.2	The system Hamiltonian	32
3.2.3	The interaction Hamiltonian	32
3.3	The weak interaction stochastic limit	32
3.4	Damped harmonic oscillator	34

1 Noncommutative probability

1.1 From commutative to noncommutative

Consider the Hilbert space $\mathcal{H} = L^2_{\mathbb{C}}(\Omega, \mu)$ over the complete probability space $(\Omega, \mathcal{F}, \mu)$. Then any function $f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)$ gives rise to a multiplication operator $\mathbf{M}_f \in \mathcal{B}(\mathcal{H})$:

$$\mathbf{M}_f g = fg \in \mathcal{H} \quad \forall g \in \mathcal{H},$$

with $\|\mathbf{M}_f\|_{\infty} = \|f\|_{L^{\infty}(\mu)}$. The collection of all such multiplication operators

$$\mathcal{A} := \{\mathbf{M}_f : f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)\} \subset \mathcal{B}(\mathcal{H})$$

forms a *commutative* subalgebra of $\mathcal{B}(\mathcal{H})$.

In fact, this subalgebra is a possibly noncommutative *von Neumann* algebra:

Definition 1.1 (von Neumann algebra) A (unital) *von Neumann algebra* is a $*$ -subalgebra $\mathcal{A} \subset \mathcal{B}(\mathcal{H})$ that contains $I_{\mathcal{H}}$ and is closed in the *weak operator topology* (WOT), i.e.,

$$\text{WOT-lim } \mathbf{a}_n = \mathbf{a} \iff \langle f, \mathbf{a}_n g \rangle_{\mathcal{H}} \rightarrow \langle f, \mathbf{a} g \rangle_{\mathcal{H}} \quad \forall f, g \in \mathcal{H}.$$

Equivalently, is it a C^* -algebra with a predual \mathcal{A}_* that is a Banach space.

There are sub-families of \mathcal{A} that play a distinguished role, namely,

$$\begin{aligned} \mathcal{A}_+ &:= \{\mathbf{a} \in \mathcal{A} : \mathbf{a} \succeq 0\} && \text{(nonnegative operators)} \\ \mathcal{O} &:= \{\mathbf{a} \in \mathcal{A} : \mathbf{a}^{\dagger} = \mathbf{a}\} && \text{(Hermitian operators)} \\ \mathcal{P} &:= \{\mathbf{a} \in \mathcal{O} : \mathbf{a}^2 = \mathbf{a}\} && \text{(projection operators)} \end{aligned}$$

Here, $\mathbf{a} \succeq 0$ if and only if $\langle f, \mathbf{a} f \rangle_{\mathcal{H}} \geq 0$ for all $f \in \mathcal{H}$. Additionally, for von Neumann algebras, we also have the following useful density result:

Proposition 1.2. *Let \mathcal{A} be a von Neumann algebra. Then*

$$\mathcal{A} = \overline{\text{span}(\mathcal{P})}^{\text{WOT}},$$

i.e., \mathcal{A} is the WOT-closure of the linear span of its projections.

Let us return to our previous example with $\mathcal{A} = \{\mathbf{M}_f : f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)\}$. It is not difficult to see that \mathcal{A} is a W^* -algebra. Notice that since \mathbf{M}_f is self-adjoint for $f \in L^{\infty}_{\mathbb{R}}(\Omega, \mu)$, the family of self-adjoint operators is given by $\mathcal{O} = \{\mathbf{M}_f : f \in L^{\infty}_{\mathbb{R}}(\Omega, \mu)\}$ and the family of projection operators is given by $\mathcal{P} = \{\mathbf{M}_f \in \mathcal{O} : f = \mathbf{1}_A, A \in \mathcal{F}\}$.

On a von Neumann algebra \mathcal{A} , we can talk about special types of continuous linear functionals on \mathcal{A} , called states.

Definition 1.3 A *state* on a von Neumann algebra \mathcal{A} is a linear functional $\psi : \mathcal{A} \rightarrow \mathbb{C}$ that is positive and normalized, i.e., $\psi(\mathbf{a}^{\dagger} \mathbf{a}) \geq 0$ for all $\mathbf{a} \in \mathcal{A}$ and $\psi(I_{\mathcal{B}(\mathcal{H})}) = 1$.

A state ψ is said to be

faithful if $\psi(\mathbf{a}^\dagger \mathbf{a}) = 0 \Leftrightarrow \mathbf{a} = 0$,

tracial if $\psi(\mathbf{ab}) = \psi(\mathbf{ba})$ for all $\mathbf{a}, \mathbf{b} \in \mathcal{A}$, and

normal if $\psi \in \mathcal{A}_*$, i.e., if it is an element of the predual \mathcal{A}_* .

For any probability measure $\nu \ll \mu$, we set

$$\psi_\nu(\mathbf{M}_f) := \int_{\Omega} f \, d\nu, \quad f \in L^\infty_{\mathbb{C}}(\Omega, \mu),$$

we find that ψ is a linear functional that is positive and normalized, i.e., ψ is a state. Moreover, it is tracial. It is normal if $\omega := d\nu/d\mu \in L^1(\Omega, \mu)$ and faithful if $\omega > 0$.

Normal states play an essential role, serving as a counterpart to classical measures, as made explicit by the following proposition.

prop:normal-state

Proposition 1.4. *Let ψ be a state on a von Neumann algebra \mathcal{A} . The following are equivalent:*

(i) ψ is a normal state.

(ii) (σ -additivity) *If $(\mathbf{a}_n) \subset \mathcal{P}$ are mutually orthogonal projections, i.e., $\mathbf{a}_n(\mathcal{H}) \perp \mathbf{a}_m(\mathcal{H})$ for all $n \neq m$, and $\mathbf{a} = \vee_n \mathbf{a}_n$ being the projection on the smallest closed subspace containing $\cup_n \mathbf{a}_n(\mathcal{H})$, then*

$$\psi(\mathbf{a}) = \sum_n \psi(\mathbf{a}_n).$$

(iii) (*Continuity from below*) *For any increasing net $0 \preceq \mathbf{a}_n \uparrow \mathbf{a}$ in \mathcal{A}_+ , one has the increasing limit $\psi(\mathbf{a}_n) \uparrow \psi(\mathbf{a})$.*

(iv) *There exists a family $\{\xi_n\} \subset \mathcal{H}$ with $\sum_n \|\xi_n\|_{\mathcal{H}}^2 = 1$ such that¹*

$$\psi = \sum_n \langle \xi_n, \cdot \xi_n \rangle \quad \text{in the sense of norm convergence.}$$

Consider an arbitrary normal state ψ and set

$$\mu(A) := \psi(\mathbf{M}_{1_A}) \quad \text{for every } A \in \mathcal{F}.$$

Then, clearly, $\mu(\emptyset) = 0$, $\mu(\Omega) = \psi(I_{\mathcal{B}(\mathcal{H})}) = 1$ and μ is σ -additive due to the equivalent characterization of a normal state given by Proposition 1.4(ii). In particular, one obtains a classical measure on (Ω, \mathcal{F}) . In this sense, a state on a noncommutative von Neumann algebra generalizes that of a classical measure.

¹[Theorem 7.1.8, Fundamentals of the Theory of Operator Algebras, V.II, Kadison-Ringrose]

1.2 Observables

In classical probability, we are often interested in computing expressions like $\mathbb{P}(X \in A)$, where X is a random variable and $A \subset \mathbb{R}$ is a Borel set.

In the quantum world, a random variable is modelled by a Hermitian operator $\mathbf{a} \in \mathcal{O}$ and is called an *observable*. Observables have their spectrum in \mathbb{R} and can therefore be *measured*.

Given a ψ on a von Neumann algebra \mathcal{A} , the expectation of an observable \mathbf{a} w.r.t. the state ψ is given by $\psi(\mathbf{a})$. Since every observable $\mathbf{a} \in \mathcal{O}$ has a spectral decomposition

$$\mathbf{a} = \int_{\mathbb{R}} \lambda E_{\mathbf{a}}(d\lambda), \quad E_{\mathbf{a}} \hat{=} \mathcal{P}\text{-valued measure},$$

we can associate a classical probability measure with the observable \mathbf{a} and state ψ :

$$\mathbb{P}_{\psi}(\mathbf{a} \in A) := \psi(E_{\mathbf{a}}(A)) \quad \text{for all Borel set } A \subset \mathbb{R}.$$

For two observables $\mathbf{a}, \mathbf{b} \in \mathcal{O}$ that do not commute, one would like to be able to write $\mathbb{P}_{\psi}(\mathbf{a} \in A, \mathbf{b} \in B)$ for two Borel sets $A, B \subset \mathbb{R}$. The simple argument for this is that if $[\mathbf{a}, \mathbf{b}] \neq 0$, then $E_{\mathbf{a}}(A)$ and $E_{\mathbf{b}}(B)$ may not commute for all Borel sets A, B . In particular,

$$\psi(E_{\mathbf{a}}(A)E_{\mathbf{b}}(B)) \neq \psi(E_{\mathbf{b}}(B)E_{\mathbf{a}}(A)),$$

so there is no consistent way of writing $\mathbb{P}_{\psi}(\mathbf{a} \in A, \mathbf{b} \in B)$. Yet, when they do commute, then $\psi(E_{\mathbf{a}}(A)E_{\mathbf{b}}(B)) = \psi(E_{\mathbf{b}}(B)E_{\mathbf{a}}(A))$ and a joint distribution exists and we are back to the classical scenario.

However, this turns out not to be possible, as the following example portrays.

Example 1.5 (Stern-Gerlach measurements) Take a beam of atoms (each with spin- $\frac{1}{2}$) and perform the following steps:




Figure 1: Stern-Gerlach experiment

SG_z Measure spin in z -direction: Send the beam through a Stern-Gerlach magnet oriented along z . The beam splits into spin-up $|\uparrow\rangle$ and spin-down $|\downarrow\rangle$ paths. Keep only the spin-up branch, which gives a pure state $|\uparrow_z\rangle$.

SG_x Measure spin in x -direction: Now send the filtered beam through a second magnet, oriented along x . The beam splits again into $|\uparrow_x\rangle$ and $|\downarrow_x\rangle$ outcomes with 50%–50% probability. So far, this can *still* be explained with classical probability.

SG_z Measure spin in z -direction again: Send either of the filtered beams $|\uparrow_x\rangle$ or $|\downarrow_x\rangle$ through a z -magnet again. You do *not* get back the original result, i.e., instead of remaining spin-up, you see another 50%–50% split between $|\uparrow_z\rangle$ and $|\downarrow_z\rangle$

Morally, if spin- x and spin- z were classical random variables, measuring x would *erase* knowledge of z . In classical probability, observing one property never randomizes another unless there is *hidden causal disturbance*.

Quantum mechanically, the two observables (or random variables)

$$\sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{spin in } z\text{-direction}, \quad \sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{spin in } x\text{-direction}.$$

do not commute, i.e., $[\sigma_x, \sigma_z] \neq 0$, and measuring σ_x *changes* information about σ_z . In this sense, they cannot be jointly sampled, i.e., there is no classical joint probability $p(\sigma_x, \sigma_z)$ for these observables, which illustrates the need for noncommutative probability.

2 Stochastic Dilation

2.1 Noncommutative random variables

def:random-variable

Definition 2.1 A *noncommutative \mathfrak{X} -valued random variable* on a quantum probability space (\mathfrak{A}, μ) is an identity preserving $*$ -homomorphism

$$\mathfrak{z}: \mathfrak{X} \rightarrow \mathfrak{A}.$$

Correspondingly, a *noncommutative \mathfrak{X} -valued stochastic process* on a quantum probability space (\mathfrak{A}, μ) is a family of random variables

$$\mathfrak{z}_t: \mathfrak{X} \rightarrow \mathfrak{A}, \quad t \in \mathbb{T},$$

with \mathbb{T} being a (possibly uncountable) index set.

Example 2.2 (Classical random variable) Let X be a classical E -valued random variable on (Ω, \mathbb{P}) . Setting

$$\mathfrak{X} := B_b(E), \quad \mathfrak{A} := L^\infty(\Omega, \mathbb{P}), \quad \mu(g) := \int_{\Omega} g \, d\mathbb{P}, \quad g \in \mathfrak{A},$$

we have that

$$\mathfrak{X} \ni f \mapsto \mathfrak{z}(f) := f \circ X \in L^\infty(\Omega, \mathbb{P})$$

is a noncommutative \mathfrak{X} -valued random variable. We see here that the noncommutative notion of a random variable is ‘dual’ to the classical notion of a random variable.

ex:stochastic-classical

Example 2.3 (Classical stochastic process) Consider the space of continuous paths (or trajectories) $\Omega := \mathcal{C}_0(\mathbb{R}_+; E)$ starting at 0 and the Wiener measure \mathbb{R} . Let $X = (X_t)_{t \in \mathbb{T}}$ be the canonical stochastic process $X_t(\omega) = \omega(t)$, $t \in \mathbb{R}_+$. As before, we set

$$\mathfrak{X} := B_b(E), \quad \mathfrak{A} := L^\infty(\Omega, \mathbb{R}), \quad \mu(G) := \int_{\Omega} G \, d\mathbb{R}, \quad G \in \mathfrak{A}.$$

Then,

$$\mathfrak{X} \ni f \mapsto \mathfrak{z}_t(f) := f \circ X_t \in L^\infty(\Omega, \mathbb{R})$$

defines a noncommutative \mathfrak{X} -valued stochastic process on (\mathfrak{A}, μ) .

Definition 2.4 A stochastic process $\mathfrak{z}_t: \mathfrak{X} \rightarrow (\mathfrak{A}, \mu)$, $t \in \mathbb{T}$ admits a *time translation* if there are $*$ -homomorphisms $\alpha_t: \mathfrak{A} \rightarrow \mathfrak{A}$, $t \in \mathbb{T}$ such that

- (1) $\alpha_{t+s} = \alpha_t \circ \alpha_s$ for all $s, t \in \mathbb{T}$, and
- (2) $\mathfrak{z}_t = \alpha_t \circ \mathfrak{z}_0$ for all $t \in \mathbb{T}$.

Example 2.5 Consider the Wiener space (Ω, \mathbb{R}) in Example 2.3. Then, \mathbb{R} is stationary under the time-shift operator $\mathfrak{s}_t \omega = \omega(\cdot - t)$, i.e., $(\mathfrak{s}_t)_\# \mathbb{R} = \mathbb{R}$ for every $t \in \mathbb{T}$. Hence, the $*$ -homomorphism

$$\alpha_t(G) = G \circ \mathfrak{s}_t, \quad G \in \mathfrak{A}, \quad t \in \mathbb{T}$$

is a time translation for the corresponding process $(\mathfrak{z}_t)_{t \in \mathbb{T}}$ and leaves the state μ invariant. Indeed, property (1) holds trivially, while property (2) follows from

$$\alpha_t \circ \mathfrak{z}_0(f) = f \circ X_0 \circ \mathfrak{s}_t = f \circ X_t = \mathfrak{z}_t(f), \quad f \in \mathfrak{X}.$$

Finally, we conclude with

$$\mu(\alpha_t(G)) = \int_{\Omega} \alpha_t(G) dR = \int_{\Omega} G \circ \mathfrak{s}_t dR = \int_{\Omega} G d(\mathfrak{s}_t)_{\#} R = \int_{\Omega} G dR = \mu(G),$$

thus proving the invariance of μ .

For $h \in L^2(\mathbb{T})$ we consider the exponential martingale

$$Z_t = \exp \left(- \int_0^t h_r dB_r - \frac{1}{2} \int_0^t |h_r|^2 dr \right), \quad t \in \mathbb{T}.$$

It is easy to see that $(Z_t)_{t \in \mathbb{T}}$ satisfies

$$dZ_t = -h_t Z_t dB_t, \quad Z_0 = 1.$$

Setting $Q := Z_T P$, we claim that

$$W_t = B_t + \int_0^t h_r dr \quad \text{is a } Q\text{-martingale.}$$

To see this, we note that W is a Q -martingale if and only if $Z_T W$ is a P -martingale. Since

$$\begin{aligned} d(Z_t W_t) &= dZ_t W_t + Z_t dW_t + d[Z, W]_t \\ &= -h_t Z_t dB_t + Z_t dB_t = -(h_t - 1) Z_t dB_t, \end{aligned}$$

we conclude that $Z_T W$ is indeed a P -martingale.

$$dU_t = -(a^* a dt + i\sqrt{2} a dB_t) U_t$$

In the classical case, the notion of adaptedness plays an important role for stochastic processes. In this regard, we choose to simply consider the natural filtration induced by a stochastic process.

Definition 2.6 Let $\mathfrak{z}_t: \mathfrak{X} \rightarrow (\mathfrak{A}, \mu)$, $t \in \mathbb{T}$ be a stochastic process. For $\mathbb{I} \subset \mathbb{T}$, we denote by $\mathfrak{A}_{\mathbb{I}}$ the subalgebra of \mathfrak{A} generated by $\{\mathfrak{z}_t(x) : x \in \mathfrak{X}, t \in \mathbb{I}\}$. The filtration induced by $(\mathfrak{z}_t)_{t \in \mathbb{T}}$ is then given by $\mathfrak{F} = (\mathfrak{A}_{[0,t]})_{t \in \mathbb{T}}$.

2.2 From commutative to noncommutative

Consider the Hilbert space $\mathcal{H} = L^2_{\mathbb{C}}(\Omega, \mu)$ over the complete probability space $(\Omega, \mathcal{F}, \mu)$. Then any function $f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)$ gives rise to a multiplication operator $M_f \in \mathcal{B}(\mathcal{H})$:

$$M_f g = fg \in \mathcal{H} \quad \forall g \in \mathcal{H},$$

with $\|M_f\|_{\infty} = \|f\|_{L^{\infty}(\mu)}$. The collection of all such multiplication operators

$$\mathcal{A} := \{M_f : f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)\} \subset \mathcal{B}(\mathcal{H})$$

forms a commutative subalgebra of $\mathcal{B}(\mathcal{H})$.

In fact, this subalgebra is a *von Neumann algebra*:

Definition 2.7 (von Neumann algebra) A (unital) *von Neumann algebra* (or W^* -algebra) is a $*$ -subalgebra $\mathcal{A} \subset \mathcal{B}(\mathcal{H})$ that contains $I_{\mathcal{H}}$ and is closed in the *weak operator topology* (WOT), i.e.,

$$\text{WOT-lim } a_n = a \iff \langle f, a_n g \rangle_{\mathcal{H}} \rightarrow \langle f, a g \rangle_{\mathcal{H}} \quad \forall f, g \in \mathcal{H}.$$

Equivalently, is it a C^* -algebra with a predual \mathcal{A}_* that is a Banach space.

There are sub-families of \mathcal{A} that play a distinguished role, namely,

$$\begin{aligned} \mathcal{A}_+ &:= \{a \in \mathcal{A} : a \succeq 0\} && \text{(nonnegative operators)} \\ \mathcal{O} &:= \{a \in \mathcal{A} : a^* = a\} && \text{(self-adjoint operators)} \\ \mathcal{P} &:= \{a \in \mathcal{O} : a^2 = a\} && \text{(projection operators)} \end{aligned}$$

Here, $a \succeq 0$ if and only if $\langle f, a f \rangle_{\mathcal{H}} \geq 0$ for all $f \in \mathcal{H}$.

On a von Neumann algebra \mathcal{A} , we can talk about special types of continuous linear functionals on \mathcal{A} , called states.

Definition 2.8 A *state* on a von Neumann algebra \mathcal{A} is a linear functional $\psi : \mathcal{A} \rightarrow \mathbb{C}$ that is positive and normalized, i.e., $\psi(a^*a) \geq 0$ for all $a \in \mathcal{A}$ and $\psi(I_{\mathcal{B}(\mathcal{H})}) = 1$.

A state ψ is said to be

faithful if $\psi(a^*a) = 0 \iff a = 0$,

tracial if $\psi(ab) = \psi(ba)$ for all $a, b \in \mathcal{A}$, and

normal if $\psi \in \mathcal{A}_*$, i.e., if it is an element of the predual \mathcal{A}_* .

Let us return to our previous example with $\mathcal{A} = \{M_f : f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu)\}$. It is not difficult to see that \mathcal{A} is a W^* -algebra. Notice that since M_f is self-adjoint for $f \in L^{\infty}_{\mathbb{R}}(\Omega, \mu)$, the family of self-adjoint operators is given by $\mathcal{O} = \{M_f : f \in L^{\infty}_{\mathbb{R}}(\Omega, \mu)\}$ and the family of projection operators is given by $\mathcal{P} = \{M_f : f = \mathbf{1}_A, A \in \mathcal{F}\}$. Additionally,

For any probability measure $\nu \ll \mu$, we set

$$\psi_{\nu}(M_f) := \int_{\Omega} f d\nu, \quad f \in L^{\infty}_{\mathbb{C}}(\Omega, \mu),$$

we find that ψ is a linear functional that is positive and normalized, i.e., ψ is a state. Moreover, it is tracial. It is normal if $\omega := d\nu/d\mu \in L^1(\Omega, \mu)$ and faithful if $\omega > 0$.

Normal states play an essential role, serving as a counterpart to classical measures, as made explicit by the following proposition.

Proposition 2.9. *Let $\psi : \mathcal{A} \rightarrow \mathbb{C}$ be a state on a von Neumann algebra \mathcal{A} . The following are equivalent:*

- (i) ψ is a normal state.
- (ii) (σ -additivity) If $(a_n) \subset \mathcal{P}$ are mutually orthogonal projections, i.e., $a_n(\mathcal{H}) \perp a_m(\mathcal{H})$ for all $n \neq m$, and $a = \vee_n a_n$ being the projection on the smallest closed subspace containing $\cup_n a_n(\mathcal{H})$, then

$$\psi(a) = \sum_n \psi(a_n).$$

- (iii) (Continuity from above) For any increasing net $0 \preceq a_n \uparrow a$ in $\mathcal{A}_+ \Rightarrow \psi(a_n) \uparrow \psi(a)$.

Consider an arbitrary normal state ψ and set

$$\mu(A) := \psi(\mathbf{M}_{1_A}) \quad \text{for every } A \in \mathcal{F}.$$

In particular, we recover the probability \mathbb{P} by

$$\mathbb{P}(A) = \psi(\mathbf{M}_{1_A})$$

$$\psi(\mathbf{M}_{1_A} \mathbf{M}_{1_B}) = \psi(\mathbf{M}_{1_{A \cap B}}) = \mathbb{P}(A \cap B)$$

$$\mathbb{P}(\cup_i A_i) = \psi(\mathbf{M}_{1_{\cup_i A_i}}) = \psi(\sum_i \mathbf{M}_{1_{A_i}}) = \sum_i \psi(\mathbf{M}_{1_{A_i}}) = \sum_i \mathbb{P}(A_i)$$

Let $(X_t)_{t \in \mathbb{T}}$ be a stochastic process on the Wiener space (Ω, \mathbb{R})

Consider a path measure \mathbf{P} on $\Omega := \mathcal{D}(\mathbb{R}_+; E)$ and the canonical process $(X_t)_{t \geq 0}$ given by $X_t(\omega) = \omega(t)$, $\omega \in \Omega$, $t \in \mathbb{R}_+$. Suppose that for every $f \in B_b(E)$,

$$f(X_t) - f(X_0) - \int_0^t Lf(X_{r-}) dr \quad \text{is an } (\mathfrak{F}, \mathbf{P})\text{-martingale,} \quad (2.1)$$

where $\mathfrak{F} = (\mathcal{F}_t)_{t \geq 0}$ is the canonical filtration $\mathcal{F}_t = \sigma(X_s : s \leq t)$, and $L : B_b(E) \rightarrow B_b(E)$ is a bounded Markov generator such that $L^*\pi = 0$ for some stationary measure $\pi \in \mathcal{P}(E)$.

We now write (2.1) in the form above. Set $\mathfrak{X} := L^\infty(E, \pi)$ and $\mathfrak{A} := L^\infty(\Omega, \mathbf{P}_\pi)$, which are both commutative von Neumann algebras and where

$$\mathbf{P}_\pi(A) = \int_E \mathbf{P}(x + A) \pi(dx).$$

Now, define the stochastic process

$$\mathfrak{z}_t : \mathfrak{X} \rightarrow (\mathfrak{A}, \mu), \quad \mathfrak{z}_t(f) = f \circ X_t,$$

with the state

$$\mu(G) := \int_\Omega G(\omega) \mathbf{P}_\pi(d\omega), \quad G \in \mathfrak{A},$$

which makes (\mathfrak{A}, μ) a quantum probability space.

$$\mathbf{E}_t[\mathfrak{z}]$$

and the $*$ -homomorphism

$$\begin{aligned} \mathfrak{z}_t(f) - \mathfrak{z}_0(f) - \int_0^t \mathfrak{z}_{r-}(Lf) dr \\ \alpha_t(F) = F \circ \mathfrak{s}_t \\ \alpha_t \circ \mathfrak{z}_0(f) = \alpha_t(f \circ X_0 \otimes \mathbf{1}_\Omega) = f \circ X_0 \circ \mathfrak{s}_t(\omega) \\ \alpha_t : \mathcal{A} \times \Omega \rightarrow \mathcal{A}; \quad \alpha_t(M_f, \omega) = M_{f \circ X_t(\omega)}. \end{aligned}$$

The martingale identity (2.1) above can then be expressed as

$$\mathfrak{m}_t[M_f] := \alpha_t(M_f) - \alpha_0(M_f) - \int_0^t \alpha_{r-}(M_{Lf}) dr.$$

Setting $\mathcal{L}M_f := M_{Lf}$, the previous identity allows one to write (2.1) as

$$\mathfrak{m}_t[A] = \alpha_t(A) - \alpha_0(A) - \int_0^t \alpha_{r-}(\mathcal{L}A) dr, \quad A \in \mathcal{A},$$

thereby generalizing the martingale problem to a non-commutative setting.

$$\mathfrak{m}_t[A] - \mathfrak{m}_s[A] = \alpha_t(A) - \alpha_s(A) - \int_s^t \alpha_{r-}(\mathcal{L}A) dr$$

$$\mathcal{L}_a = a^*[\bullet, a] + [a^*, \bullet]a$$

Example 2.10 (Diffusion) Consider the formal example of the (possibly degenerate) diffusion process with the generator

$$Lf = \operatorname{div}(A\nabla f) - A\nabla V \cdot \nabla f, \quad f \in \mathcal{C}_c^\infty(\mathbb{R}^d),$$

i.e., it is the generator of the Itô diffusion

$$dX_t = -\sigma(X_t)\sigma^\top(X_t)\nabla V(X_t)dt + \sigma(X_t) \circ dB_t,$$

with $A = \sigma\sigma^\top \in \mathbb{R}^{d \times d}$, $\sigma \in \mathbb{R}^{d \times m}$

$$a_i = \sum_{j=1}^d \sigma_{ji} \partial_j, \quad a_i^* = - \sum_{j=1}^d \partial_j(\sigma_{ji} \bullet), \quad i = 1, \dots, m.$$

It is not difficult to see that $[a_i^*, M_f] = -[a_i, M_f]$. Moreover, for any $g \in C_c^\infty(\mathbb{R}^d)$,

$$\begin{aligned} a_i^*[M_f, a_i]g &= \sum_{j=1}^d \partial_j(\sigma_{ji}g) \sum_{k=1}^d \sigma_{ki} \partial_k f \\ &= \sum_{j=1}^d g \partial_j(\sigma_{ji}) \sum_{k=1}^d \sigma_{ki} \partial_k f + \sum_{j,k=1}^d \sigma_{ji} \sigma_{ki} \partial_j g \partial_k f \\ &= \sum_{j=1}^d g \partial_j(\sigma_{ji}) \sum_{k=1}^d \sigma_{ki} \partial_k f - [M_f, a_i]a_i g. \end{aligned}$$

Consequently, we find that

$$\sum_{i=1}^m \mathcal{L}_{a_i}(M_f) = \sum_{i=1}^m (a_i^*[M_f, a_i] + [a_i^*, M_f]a_i) = M_{\operatorname{div}(A\nabla f)}$$

Moreover, setting

$$H_i = \sum_{j,k=1}^d \sigma_{ji} \sigma_{ki} \partial_j V \partial_k, \quad i = 1, \dots, m,$$

we have that

$$\begin{aligned} \sum_{i=1}^m [H_i, M_f] &= M_{\sigma\sigma^\top \nabla V \cdot \nabla f} \\ &\quad [a_i, M_V][a_i, \bullet] \end{aligned}$$

Therefore, we obtain

$$\mathcal{L}M_f = \sum_{i=1}^m ([H_i, M_f] + \mathcal{L}_{a_i}M_f) = M_{L_f} \quad \forall f \in \mathcal{C}_c^\infty(\mathbb{R}^d).$$

However, we know that the diffusion process constitutes a gradient flow with driving energy $\mathcal{F} = \operatorname{Ent}(\bullet|\pi)$, where $\pi = e^{-V} \operatorname{Leb}$ is the invariant measure. Formally, one defines a state on $\mathcal{C}_c^\infty(\mathbb{R}^d)$ by

$$\tau(M_f) := \int_{\mathbb{R}^d} f d\pi, \quad f \in \mathcal{C}_c^\infty(\mathbb{R}^d).$$

$$\begin{aligned}\tau_\pi(M_f M_g) &= \tau(M_f M_g M_\pi) = \tau(M_{fg\pi}) = \tau_\pi(M_g M_f) \\ \Delta_\pi A &= M_\pi A M_\pi^{-1}\end{aligned}$$

$$\begin{aligned}(\Delta_\pi V_j)(g) &= M_\pi V_j M_\pi^{-1} g = \pi \sum_{j=1}^d \sigma_{ji} \partial_j (\pi^{-1} g) - \sigma_{ji} \partial_j V (\pi^{-1} g) \\ &= \sum_{j=1}^d \sigma_{ji} [g \partial_j V + \partial_j g] - \sigma_{ji} \partial_j V g =\end{aligned}$$

$$V_j = \sum_{j=1}^d \sigma_{ji} \partial_j - M_{\sigma_{ji} \partial_j V}$$

$$\tau_\sigma(M_g \mathcal{L} M_f) = \hat{\sigma}(M_{gLf}) = \int_{\mathbb{R}^d} g Lf \, \mathrm{d}\sigma =$$

$$\begin{aligned}\mathcal{L} M_{e^{-V}} g &= (\operatorname{div}(A \nabla e^{-V}) + A \nabla V \cdot \nabla e^{-V}) g \\ &= (-\operatorname{div}(A \nabla V e^{-V}) - A \nabla V \cdot \nabla V e^{-V}) g\end{aligned}$$

$$[a_i, M_f]g = \sum_{j=1}^d \sigma_{ji} \partial_j (fg) - \sum_{j=1}^d f \sigma_{ji} \partial_j g = \sum_{j=1}^d \sigma_{ji} g \partial_j f = M_{a_i(f)}g$$

$$[a_i^*, M_f]g = - \sum_{j=1}^d \partial_j (\sigma_{ji} fg) + \sum_{j=1}^d f \partial_j (\sigma_{ji} g) = - \sum_{j=1}^d \sigma_{ji} \partial_j fg = -M_{a_i(f)}g = -[a_i^*, M_f]g$$

Consequently, we find that

$$\mathcal{L}_{a_i} M_f =$$

$$[a_i, M_f]a_i g = \sum_{j,k=1}^d \sigma_{ji} \sigma_{ki} \partial_j f \partial_k g$$

$$[a_i, M_V] = \sum_{j=1}^d \sigma_{ji} \partial_j V$$

$$[a_i, [a_i, M_f]] = [a_i, M_{a_i(f)}] = M_{a_i(a_i(f))},$$

where we used the fact that $[a_i, M_f] = M_{a_i(f)}$ for every $i = 1, \dots, d$.

$$\begin{aligned} \sum_{i=1}^m a_i \circ a_i(f) &= \sum_{i=1}^m \sum_{j=1}^d \sigma_{ji} \partial_j \left(\sum_{k=1}^d \sigma_{ki} \partial_k f \right) = \sum_{i=1}^m \sum_{j,k=1}^d \sigma_{ji} \partial_j (\sigma_{ki} \partial_k f) \\ &= \sum_{i=1}^m \sum_{j,k=1}^d \sigma_{ji} \partial_j (\sigma_{ki} \partial_k f) \end{aligned}$$

$$\begin{aligned} \operatorname{div}(\sigma \sigma^\top \nabla f) &= \sum_{j=1}^d \sum_{i=1}^m \sum_{k=1}^d \partial_j (\sigma_{ji} \sigma_{ki} \partial_k f) \\ &= \sum_{j=1}^d \sum_{i=1}^m \sum_{k=1}^d \partial_j (\sigma_{ji} \sigma_{ki} \partial_k f) \end{aligned}$$

$$\begin{aligned} \langle \psi | M_{L_f} | \psi \rangle &= \int (L_f)(x) |\psi(x)|^2 \pi(dx) \\ &= - \int \sigma^2 \partial_x f \partial_x (\psi^2 \pi) dx - \int \sigma^2 \partial_x f \partial_x V \psi^2 d\pi \\ &= -2 \int \sigma^2 \partial_x f \psi \partial_x \psi d\pi \\ &= -2 \int \\ &\quad a\psi = \sigma \partial_x \psi \end{aligned}$$

$$\begin{aligned}
\langle \varphi | a \psi \rangle &= \int \varphi \sigma \partial_x \psi \, d\pi \\
&= - \int \partial_x (\sigma \varphi \pi) \psi(x) \, dx \\
&= - \int \partial_x (\sigma \varphi) \psi \, d\pi + \int \varphi \psi \sigma \partial_x V \, d\pi \\
&= \int [-\partial_x (\sigma \varphi) + \varphi \sigma \partial_x V] \psi \, d\pi = \langle a^* \varphi | \psi \rangle
\end{aligned}$$

$$\begin{aligned}
(a^* M_f a \psi)(x) &= -\partial_x (f \sigma^2 \partial_x \psi) + f \sigma^2 \partial_x \psi \partial_x V \\
(a^* a M_f \psi)(x) &= -\partial_x (\sigma^2 \partial_x (f \psi)) + \sigma^2 \partial_x (f \psi) \partial_x V \\
(M_f a^* a \psi)(x) &= -f \partial (\sigma^2 \partial_x \psi) + \sigma^2 f \partial_x \psi \partial_x V
\end{aligned}$$

$$\int \psi(x) (a^* M_f a \psi)(x) \, d\pi = \int f |\sigma \partial_x \psi|^2 \, d\pi$$

2.3 Stochastic processes on matrix algebras

In this section, we consider $\mathfrak{X} = \text{Mat}(\mathbb{C}, n)$. Let $a \in \mathcal{O}(\mathfrak{X})$ be an observable on \mathfrak{X} and set

$$\mathcal{L}x = a^*[x, a] + [a^*, x]a, \quad x \in \mathfrak{X}.$$

We then consider the solution of the

$$d\mathcal{U}_t = (-a^* a \, dt + i\sqrt{2}a \, dB_t) \mathcal{U}_t,$$

which, due to $a = a^*$, can be explicitly expressed as

$$\mathcal{U}_t(x, e(f)) := \exp(i\sqrt{2}a B_t(\omega))x, \quad t \in \mathbb{T},$$

which is a $*$ -automorphism on \mathfrak{X} , which extends to an isometry

$$\mathcal{U}_t: \mathfrak{X} \otimes \mathcal{A}$$

$$B \in \mathcal{A}$$

$$\mathfrak{z}_t(x) - \mathfrak{z}_0(x) - \int_0^t \mathfrak{z}_r(\mathcal{L}x) \, dr$$

2.4 Stochastic processes on the unitary group

2.4.1 The unitary group and its Lie structure

The unitary group is defined as

$$U(n) = \{U \in \mathbb{C}^{n \times n} : U^* U = I\}.$$

It is a compact Lie group with Lie algebra (tangent space at identity)

$$\mathcal{U}(n) = \mathbb{T}_I U(n) = \{\mathfrak{a} \in \mathbb{C}^{n \times n} : \mathfrak{a}^* = -\mathfrak{a}\},$$

i.e., the space of skew-Hermitian matrices. A vector field $\mathfrak{a} : U(n) \rightarrow \mathcal{U}(n)$ is *left-invariant* if $\mathfrak{a}(U) = U\mathfrak{a}(I) \in \mathcal{U}(n)$ for all $U \in U(n)$.

The Lie algebra $\mathcal{U}(n)$ can be equipped with a real inner product given by the Hilbert-Schmidt inner product

$$\langle \mathfrak{a}, \mathfrak{b} \rangle_{\mathcal{U}(n)} = -\operatorname{tr}(\mathfrak{a}^* \mathfrak{b}) = \Re \operatorname{tr}(\mathfrak{a} \mathfrak{b}^*),$$

which is positive-definite on $\mathcal{U}(n)$. The associated norm is then

$$|\mathfrak{a}|_{\mathcal{U}(n)}^2 = \langle \mathfrak{a}, \mathfrak{a} \rangle_{\mathcal{U}(n)} = \operatorname{Tr}(\mathfrak{a} \mathfrak{a}^*).$$

2.4.2 Unitary-valued Brownian motion (rank- r noise)

Let $\mathfrak{a}_1, \dots, \mathfrak{a}_r \in \mathcal{U}(n)$ be orthonormal under the inner product $\langle \cdot, \cdot \rangle_{\mathcal{U}(n)}$, where $1 \leq r \leq n^2$. Further, let $\beta_t^1, \dots, \beta_t^r$ be independent standard real Brownian motions.

Define the $\mathcal{U}(n)$ -valued (possibly degenerate) Brownian driver

$$\mathfrak{w}_t = \sum_j \mathfrak{a}_j \beta_t^j.$$

If $r = n^2$, the driver is elliptic (non-degenerate). If $r < n^2$, the covariance has rank r and the process is *degenerate* (hypoelliptic), exploring only the connected subgroup

$$H = \exp(\mathcal{H}) \subset U(n), \quad \mathcal{H} = \operatorname{Lie}\{\mathfrak{a}_1, \dots, \mathfrak{a}_r\} \subset \mathcal{U}(n).$$

The intrinsic $U(n)$ -valued diffusion process solves the Stratonovich SDE

$$dU_t = U_t \circ d\mathfrak{w}_t, \quad U_0 = I,$$

or in components

$$dU_t = \sum_j U_t \mathfrak{a}_j \circ d\beta_t^j.$$

The matrix-valued quadratic variation of \mathfrak{w} is given by

$$d[\mathfrak{w}] = \sum_j \mathfrak{a}_j \otimes \mathfrak{a}_j dt.$$

In Itô form, the equivalent SDE reads

$$dU_t = U_t (d\mathfrak{w}_t + \mathfrak{l} dt), \quad \mathfrak{l} := \frac{1}{2} \sum_j \mathfrak{a}_j^2,$$

where \mathfrak{l} is the Laplace-Beltrami generator associated with the left-invariant connection.

In the following, we denote its law on path space $C([0, T]; U(n))$ by \mathbf{P} .

2.4.3 Girsanov transform on $U(n)$ (Karandikar type)

Let $\mathbf{u} \in L^2((0, T); \mathcal{U}(n))$ be adapted and square-integrable.

Define the drifted group motion by

$$dU_t^{\mathbf{u}} = U_t^{\mathbf{u}} \circ (d\mathbf{w}_t + \mathbf{u}_t dt), \quad U_0^{\mathbf{u}} = I.$$

Assuming a Novikov condition in the active noise directions,

$$\mathbb{E} \left[\exp \left(\frac{1}{2} \int_0^T |\mathbf{u}_s|^2 ds \right) \right] < +\infty,$$

the process defined below is a true martingale.

Decompose the drift against the active algebra frame:

$$\mathbf{u}_t = \sum_j \mathbf{a}_j v_t^j, \quad v = (v^1, \dots, v^r) \in L^2((0, T); \mathbb{R}^r).$$

Define the exponential Cameron-Martin density process

$$\begin{aligned} M_T &= \exp \left(\int_0^T \langle \mathbf{u}_t, d\mathbf{w}_t \rangle - \frac{1}{2} \sum_j \int_0^T \langle \mathbf{u}_t, \mathbf{a}_j \otimes \mathbf{a}_j \mathbf{u}_t \rangle dt \right) \\ &= \exp \left(\sum_j \int_0^T v_t^j d\beta_t^j - \frac{1}{2} \sum_j \int_0^T |v_t^j|^2 dt \right). \end{aligned}$$

This expression depends only on the directions with nonzero quadratic variation. It is the standard Cameron-Martin shift density on \mathbb{R}^r lifted to the group via left multiplication.

We then define a new path measure on the same probability space by

$$\mathbf{Q} = M_T \mathbf{P}.$$

Under \mathbf{Q} :

(i) The shifted drivers

$$\beta_t^{j, \mathbf{Q}} := \beta_t^j - \int_0^t v_s^j ds$$

are Brownian motions again on \mathbb{R} .

(ii) The algebra noise

$$d\mathbf{w}_t^{\mathbf{Q}} := d\mathbf{w}_t - \mathbf{u}_t dt = \sum_j \mathbf{a}_j d\beta_t^{j, \mathbf{Q}}$$

is a $\mathcal{U}(n)$ -valued Brownian motion again with same covariance

$$d[\mathbf{w}^{\mathbf{Q}}] = d[\mathbf{w}] = \sum_j \mathbf{a}_j \otimes \mathbf{a}_j dt.$$

(iii) The group paths satisfy the same *driftless* intrinsic SDE

$$dU_t^{\mathbf{u}} = U_t^{\mathbf{u}} \circ dW_t^{\mathbf{Q}} = \sum_j U_t^{\mathbf{u}} \mathbf{a}_j \circ d\beta_t^{j, \mathbf{Q}},$$

so $U_t^{\mathbf{u}}$ is an intrinsic Brownian motion on the same symmetry group H .

Hence, M_T is the Radon-Nikodym derivative between the laws of the drifted and driftless group diffusions on path space.

2.4.4 Induced Girsanov on pure states

Rank-1 projections evolve by the conjugation map

$$P_t = U_t P_0 U_t^*, \quad P_0 = \psi_0 \psi_0^*.$$

This gives a valid reference probability on paths of $C([0, T]; \mathcal{M})$, where

$$\mathcal{M} = \{V P_0 V^* : V \in H\} \subset \mathcal{P}_1(n) \simeq \mathbb{CP}^{n-1}$$

is the homogeneous orbit that the degenerate noise can reach.

Linearising conjugation and applying the Stratonovich chain rule yields the SDE

$$dP_t^u = [\circ dW_t^Q, P_t^u] = \sum_j [\mathfrak{a}_j, P_t^u] \circ d\beta_t^{j,Q},$$

so the pure-state path law remains Brownian on the same orbit manifold \mathcal{M} .

If a drift \mathbf{u} has components outside $\text{span}\{\mathfrak{a}_j\}$, only the projection of the drift onto that span can be removed. Directions with zero quadratic variation cannot absorb drifts.

$$\int F(U_t P_0 U_t^*) \mathbf{P}(dU) =$$

2.5 Dilation of semigroups (Sz.Nagy)

2.6 Stochastic dilations: How to make a heat bath

2.7 Interpretation of LDPs on unitary evolutions

The task of this subsection is to reinterpret the large deviations of empirical measures in a functional analytic framework.

Consider a family of iid E -valued random variables $(X_i)_{i \in \mathbb{N}}$ with $\sigma = \text{Law} X_1$. Setting $\Omega = E^{\mathbb{N}}$ and $\mathbf{R} = \sigma^{\otimes \mathbb{N}}$, we see that $\text{Law}(X_i)_{i \in \mathbb{N}} = \mathbf{R}$. Moreover, we can disregard the initial probability space for X_i and consider instead the canonical random variable on Ω .

For every $f \in \mathfrak{X} := \mathcal{C}_b(E)$ and $n \in \mathbb{N}$, we define the noncommutative random variable

$$\mathfrak{X} \ni f \mapsto \mathfrak{z}^n(f) = \frac{1}{n} \sum_{i=1}^n f(X_i) \in \mathfrak{A} := L^\infty(\Omega, \mathbf{R}),$$

where \mathfrak{A} is equipped with the state $\mu(G) = \mathbf{E}_{\mathbf{R}}[G]$, $G \in \mathfrak{A}$.

$$\mu(\mathfrak{z}^n(f)) = \frac{1}{n} \sum_{i=1}^n \mathbf{E}_{\mathbf{R}}[f(X_i)] = \frac{1}{n} \sum_{i=1}^n \int_E f \, d\sigma = \int_E f \, d\sigma = \mu(\mathfrak{z}_1(f)).$$

$$\begin{aligned} \mathbf{R}(\sup_f \{\mu^n(f) - \nu(f)\} > \varepsilon) &= \mathbf{R}(n\mu^n(f) > n(\nu(f) + \varepsilon)) \\ &= \mathbf{R}(e^{n\mu^n(f)} > e^{n(\nu(f) + \varepsilon)}) \leq e^{-n(\nu(f) + \varepsilon)} \int_{\Omega} e^{n\mu^n(f)} \, d\mathbf{R} \end{aligned}$$

$$\mathfrak{z}: \mathfrak{X} \rightarrow (\mathfrak{X} \otimes \mathfrak{C}, \text{tr} \otimes \mu)$$

$$\mathfrak{A} = \mathfrak{X} \otimes \mathfrak{C}^{\otimes \mathbb{Z}}, \quad \varphi = \text{tr} \otimes \mu^{\otimes \mathbb{Z}}$$

Let X_i

$$\mathfrak{z}_i(x) = x \otimes c_i, \quad c_i = \cdots 1 \otimes c$$

$$\mathfrak{z}^n = \frac{1}{n} \sum_{i=1}^n \mathfrak{z}_i, \quad \mathfrak{z}_i = \cdots \otimes \mathbb{1} \otimes \mathfrak{z} \otimes \mathbb{1} \otimes \cdots$$

$$\varphi(\mathfrak{z}^n(x)) = \frac{1}{n} \sum_{i=1}^n \varphi(\mathfrak{z}_i(x)) = \varphi(\mathfrak{z}_1(x)) = \mu(\mathfrak{z}(x))$$

$$\mathfrak{z}_i(x) = \mathbb{1} \otimes \cdots \otimes \mathbb{1} \otimes \mathfrak{z}(x) \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1}$$

$$\mathfrak{z}^n(x) \in \mathfrak{A} \cong \mathfrak{X} \otimes \mathfrak{Y}$$

$$\mu = \text{tr} \otimes \mu_{\mathfrak{C}}^{\otimes \mathbb{N}}$$

$$\mu(\mathbf{1}_A(\mathfrak{z}^n(x) - \mathfrak{y}_1(x))) = \mu_{\mathfrak{C}}^{\otimes \mathbb{N}}(\text{tr}[\mathbf{1}_A(\mathfrak{z}^n(x) - \mathfrak{y}(x) \otimes \mathbb{1}_{\mathfrak{C}^{\otimes \mathbb{N}}})])$$

$$\sup_f \{\mu^n(f) - \nu(f)\} > \varepsilon \quad \Rightarrow \quad \mu^n(f^\delta) - \nu(f^\delta) > \varepsilon - \delta$$

$$\begin{aligned} \mathbf{R}(\mu^n(f) - \nu(f) > \varepsilon) &= \mathbf{R}(e^{n\mu^n(f^\delta)} > e^{n(\nu(f^\delta) + \varepsilon - \delta)}) \\ &\leq e^{-n(\nu(f^\delta) + \varepsilon - \delta)} \mathbf{E}_{\mathbf{R}}[e^{n\mu^n(f^\delta)}] \end{aligned}$$

$$\begin{aligned} -\frac{1}{n} \log \mathbf{R}(\mu^n(f) - \mu(f) > \varepsilon) &\geq \mu(f) + \varepsilon - \frac{1}{n} \log \int_{\Omega} e^{n\mu^n(f)} d\mathbf{R} \\ &= \mu(f) + \varepsilon - \log \int_E e^f d\mu \\ -\frac{1}{n} \log \mathbf{R}(\mu^n(f) - \mu(f) > \varepsilon) &\geq \end{aligned}$$

Taking the sup

$$\begin{aligned} \int_{\Omega} e^{n\mu^n(f)} d\mathbf{R} &= \int_{\Omega} \prod_{i=1}^n e^{f(X_i)} d\mathbf{R} \\ &= \int_{E^N} \prod_{i=1}^n e^{f(x_i)} \mu(dx_1) \cdots \mu(dx_n) = \left(\int_E e^{f(x)} \mu(dx) \right)^n \end{aligned}$$

2.8 LDP via Gärtner-Ellis revisited

Let $\Omega = \mathcal{C}(I; \mathfrak{h})$ be the space of \mathfrak{h} -valued continuous path and consider the canonical process $(X_t)_{t \in I}$ defined by $X_t(\omega) := \omega(t)$ for all $t \in I$. Further, let $\mathcal{F} = \cup_{t \geq 0} \mathcal{F}_t$, where the filtration $\mathfrak{F} := \{\mathcal{F}_t := \sigma(X_s : 0 \leq s \leq t)\}_{t \in I}$ is generated by $(X_t)_{t \in I}$. Throughout, we consider a reference path measure $\mathbf{R} \in \mathcal{P}(\Omega)$ such that X admits the \mathbf{R} -semimartingale decomposition

$$X = X_0 + B^{\mathbf{R}} + M^{\mathbf{R}} \quad \mathbf{R}\text{-almost surely,}$$

where $B^{\mathbf{R}}$ is an adapted process with absolutely continuous sample paths \mathbf{R} -almost surely, and $M^{\mathbf{R}}$ is an \mathbf{R} -martingale. For simplicity of presentation, we assume that the quadratic variation of $M^{\mathbf{R}}$ is an $\mathcal{S}_+(\mathfrak{X})$ -valued random measure on I with Lebesgue density, i.e.,

$$\frac{d[M^{\mathbf{R}}, M^{\mathbf{R}}]}{d\lambda}(t) = \mathbf{a}_t(X_t) \in \mathcal{S}_+(\mathfrak{h}) \quad \mathbf{R}\text{-almost surely,}$$

and also that $t \mapsto \mathbf{a}_t(X_t)$ is an \mathfrak{F} -adapted process.

and has the quadratic variation $\langle M^{\mathbf{R}}[\varphi] \rangle$ for every smooth function φ .

Let X^1, X^2, \dots be independent copies of (X^1, \dots, X^n)

$$\begin{aligned} \frac{dQ^n}{dP^{\otimes n}} &= \exp \left(\sum_{i=1}^n \int_I \langle \beta_t^i(X_t), dM_t^{\mathbf{R}} \rangle - \frac{1}{2} \sum_{i=1}^n \int_I \langle \beta_t^i(X_t), \mathbf{a}_t(X_t) \beta_t^i(X_t) \rangle dt \right) =: Z^{(n)} \\ Y_t^{(n)} &:= \sum_{i=1}^n \int_0^t \langle \beta_s^i(X_s), dM_s^{\mathbf{R}} \rangle - \frac{1}{2} \sum_{i=1}^n \int_0^t \langle \beta_s^i(X_s), \mathbf{a}_s(X_s) \beta_s^i(X_s) \rangle ds \end{aligned}$$

$$dZ_t^{(n)} = dY_t^{(n)} = Z_t^{(n)} \left(dY_t^{(n)} + \frac{1}{2} d[Y^{(n)}, Y^{(n)}]_t \right) = \sum_{i=1}^n Z_t^{(n)} \langle \beta_t^i(X_t), dM_t^{\mathbf{R}} \rangle$$

Define $\rho^n := \frac{1}{n} \sum_{i=1}^n X^i$. Then,

$$d\rho^n = \frac{1}{n} \sum_{i=1}^n dX_t^i = \frac{1}{n} \sum_{i=1}^n b_t^{\mathbf{R}}(X_t^i) dt + \frac{1}{n} \sum_{i=1}^n dM_t^{\mathbf{R},i}$$

$$\begin{aligned} \mathbf{P}^{\otimes n} \left(\frac{1}{n} \sum_{i=1}^n X^i \in A \right) &= \int_A e^{-\int_I \langle \beta_t^i(X_t), dM_t^{\mathbf{R}} \rangle + \frac{1}{2} \int_I \langle \beta_t^i(X_t), \mathbf{a}_t(X_t) \beta_t^i(X_t) \rangle dt} d\mathbf{Q}^{(n)} \\ &= \int_A e^{-\int_I \langle \beta_t^i(X_t), dM_t^{\mathbf{R}} \rangle + \frac{1}{2} \int_I \langle \beta_t^i(X_t), \mathbf{a}_t(X_t) \beta_t^i(X_t) \rangle dt} d\mathbf{Q}^{(n)} \end{aligned}$$

$$A := \{\rho : \partial_t \rho + \overline{\text{div}} j = L\rho\}$$

$$D_{\beta^i} \exp \left(\int_I \langle \beta_t^i(X_t), dM_t^{\mathbf{R}} \rangle + \frac{1}{2} \int_I \langle \beta_t^i(X_t), \mathbf{a}_t(X_t) \beta_t^i(X_t) \rangle dt \right) [h^i]$$

$$\int_I \langle \beta_t, j_t \rangle dt - \log \int_A Z^{(n)} d\mathbf{P}^{\otimes n}$$

2.8.1 Example

We consider the Hilbert space $\mathfrak{h} = \mathbb{M}$ equipped with the scalar product $\langle A, B \rangle = \text{tr}[A^* B]$. Consider the stochastic process

$$P_t = P_0 + \int_0^t LP_s ds + \int_0^t \overline{\nabla} P_s dW_s,$$

where $\overline{\nabla} P = [i\sigma_X, P]$ and $LP = (1/2)[i\sigma_X, [i\sigma_X, P]]$. In other words,

$$B_t^{\mathbf{R}} = \int_0^t LP_s ds, \quad M_t^{\mathbf{R}} = \int_0^t \overline{\nabla} P_s dW_s,$$

where $M^{\mathbf{R}}$ has the quadratic variation

$$\mathbf{a}(P) = \langle (\overline{\nabla} P)^*, \bullet \rangle \overline{\nabla} P \quad \text{for every } P \in \mathfrak{h}.$$

Let P^1, P^2, \dots be independent copies and define $\rho^n := \frac{1}{n} \sum_{i=1}^n P^i$. Then,

$$d\rho_t^n = L\rho_t^n dt + \frac{1}{n} \sum_{i=1}^n \overline{\nabla} P_t^i dW_t^i = L\rho_t^n dt + \frac{1}{n} dM_t^{\mathbf{R},n}.$$

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n \bar{\nabla} P_t^i dW_t^i &= \frac{1}{n} \sum_{i=1}^n \left(\bar{\nabla} P_t^i dW_t^i - \mathbf{a}(P_t^i) H_t dt \right) + \frac{1}{n} \sum_{i=1}^n \mathbf{a}(P_t^i) H_t dt \\ d\rho^n &= L\rho^n dt + \frac{1}{n} \sum_{i=1}^n \mathbf{a}(P_t^i) H_t dt + \frac{1}{n} dM_t^{\mathbf{Q},n}\end{aligned}$$

For each $n \in \mathbb{N}$, we set

$$\frac{d\mathbf{Q}^n}{d\mathbf{R}^{\otimes n}} = \exp \left(\sum_{i=1}^n \int_I \langle \bar{\nabla} P_t^i, H_t \rangle dW_t^i - \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) =: Z^{(n)}.$$

Then, for curves η satisfying $\partial_t \eta + \bar{\text{div}} J^\eta = L\eta$ for some J^η , we find that

$$J^\eta = \sum \langle \sigma_j, J^\eta \rangle \sigma_j$$

Suppose

$$-\bar{\text{div}} = \sum_{j \in J} \alpha_j [i\sigma_j, \bullet],$$

then

$$-\bar{\text{div}} J^\eta = \sum_{j,k} \alpha_j \langle \sigma_k, J^\eta \rangle [i\sigma_j, \sigma_k] = \sum_{j,k \neq j} \alpha_j \langle \sigma_k, J^\eta \rangle [i\sigma_j, \sigma_k] \in \text{span}\{[i\sigma_j, \sigma_k] : j \in J, k \neq j\}$$

$$\begin{aligned}\mathbf{R}^{\otimes n}(\rho^n \in A) &= \int_A \exp \left(- \sum_{i=1}^n \int_I \langle \bar{\nabla} P_t^i, H_t \rangle dW_t^i + \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) d\mathbf{Q}^{(n)} \\ &= \int_A \exp \left(- \int_I \langle dM_t^{\mathbf{R},n}, H_t \rangle + \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) d\mathbf{Q}^{(n)} \\ &= \int_A \exp \left(- \int_I \langle dM_t^{\mathbf{Q},n}, H_t \rangle - \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) d\mathbf{Q}^{(n)}\end{aligned}$$

$$H^\eta \in \underset{H}{\text{argmin}} \left\{ \mathbb{E}_{\mathbf{Q}} \left[\frac{1}{2} \int_I \langle H_t, \mathbf{a}(P_t) H_t \rangle dt \right] - \int_I \langle H_t, \bar{\text{div}} J_t^\eta \rangle dt \right\}$$

$$\mathbb{E}_{\mathbf{Q}} \left[\int_I (\text{tr}[P_t \bar{\nabla} H_t])^2 dt \right] = \int_I \mathbb{E}_{\mathbf{Q}} \left[(\text{tr}[P_t \bar{\nabla} H_t])^2 \right] dt$$

$$\text{tr}[\rho_t^{\mathbf{Q}} \bar{\nabla} H_t] = \sum \lambda_i(t) \langle v_i(t), \bar{\nabla} H_t v_i(t) \rangle \geq \gamma \sum \lambda_i(t)$$

$$\mathbb{E}_{\mathbf{R}} \left[\int_I \langle G_t, \mathbf{a}(P_t) H_t \rangle dt \right] = \int_I \langle G_t, \bar{\text{div}} J_t \rangle dt$$

$$\int_I \langle H_t, \bar{\text{div}} J_t \rangle dt - \log \mathbb{E}_{\mathbf{R}} \left[\exp \left(\int_I \langle H_t, dM_t^{\mathbf{R},n} \rangle - \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) \right]$$

$$\int_I \langle G_t, \overline{\text{div}} J_t \rangle dt = -\mathbb{E}_{\mathbb{R}^{\otimes n}} \left[\left(\int_I \langle G_t, dM_t^{\mathbb{R},n} \rangle - \sum_{i=1}^n \int_I \langle G_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) \exp \left(\int_I \langle H_t, dM_t^{\mathbb{R},n} \rangle - \frac{1}{2} \sum_{i=1}^n \int_I \langle H_t, \mathbf{a}(P_t^i) H_t \rangle dt \right) \right]$$

Let $Y_t^{(n)} = \sum_{i=1}^n \int_0^t \langle \overline{\nabla} H_s, \overline{\nabla} P_s^i \rangle dW_s^i$ and $Z_t^{(n)} := \exp(Y_t^{(n)})$. Then,

$$dZ_t^{(n)} = Z_t^{(n)} \left(\sum_{i=1}^n \langle \overline{\nabla} H_t, \overline{\nabla} P_t^i \rangle dW_t^i + \frac{1}{2} \sum_{i=1}^n \langle \overline{\nabla} H_t, \mathbf{a}(P_t^i) \overline{\nabla} H_t \rangle dt \right)$$

$$Z_t^{(n)} = 1 + \sum_{i=1}^n \int_0^t Z_s^{(n)} \langle \overline{\nabla} H_s, \overline{\nabla} P_s^i \rangle dW_s^i + \frac{1}{2} \sum_{i=1}^n \int_0^t Z_s^{(n)} \langle \overline{\nabla} H_s, \mathbf{a}(P_s^i) \overline{\nabla} H_s \rangle ds$$

Let $\varrho \in \mathcal{S}(\mathcal{H})$ be state and $(\varrho^n)_{n \in \mathbb{N}} \subset \mathcal{S}(\mathcal{H})$ be a sequence of states. We say that $\varrho^n \rightarrow \varrho$ setwise if

$$\varrho^n(E) \rightarrow \varrho(E) \quad \text{for all projections } E \in \mathcal{P}(\mathcal{H}).$$

Further, let $(\mathfrak{z}_t)_{t \in \mathbb{T}}$ be a stochastic process in the sense of Definition 2.1 with

$$\mathfrak{z}_t: \mathfrak{X} \rightarrow (\mathfrak{X} \otimes \mathfrak{C}, \varphi), \quad \varphi = \varrho \otimes \mu.$$

$$\mathfrak{z}_t(x) = \alpha_t^* x \alpha_t$$

and $\mathfrak{z}_0(x) = x \otimes 1$, $x \in \mathfrak{X}$. By construction, we have that

$$\varphi(\mathfrak{z}_0(x)) = \varphi(x \otimes 1) = \varrho(x)\mu(1) = \varrho(x) \quad \forall x \in \mathfrak{X}.$$

We suppose that

$$\begin{aligned} \mathcal{S}(\mathfrak{X} \otimes \mathfrak{C}^{\otimes n}) &\rightarrow \mathcal{S}(\mathfrak{X} \otimes \mathfrak{C}) \\ \mu_\varphi &= \frac{1}{n} \sum_{i=1}^n \varphi_i = \frac{1}{n} \sum_{i=1}^n \rho \otimes \varphi \\ \mathbf{x}_i: \mathfrak{C}^{\otimes n} &\rightarrow \mathfrak{C}; \quad \mathbf{x}_i(\omega_1, \dots, \omega_n) = \omega_i \\ \delta_{\mathbf{x}_i}(E) & \end{aligned}$$

2.9 Groups to algebras

Let \mathbb{T} be the 1-dimensional torus. Further, let \mathcal{H} be a Hilbert space and $\mathfrak{h} \in \mathcal{O}(\mathcal{H})$ be an observable on \mathcal{H} . Consider the unitary $\mathbf{u}: \mathbb{T} \rightarrow \mathcal{U}(\mathcal{H})$ such that $\mathbf{u}(\exp(tX)) = e^{i\mathfrak{h}t}$.

$$\left. \frac{d}{dt} f(\exp(tX)) \right|_{t=0} = X(f)$$

$$dX_t =$$

$$\mathcal{B}(\mathcal{H}) \rightarrow L^\infty(\mathbb{T}; \mathcal{B}(\mathcal{H})); \quad a \mapsto \mathbf{u}^*(z) a \mathbf{u}(z)$$

Let $\rho_0 \in \mathcal{S}(\mathcal{H})$ be a given fixed state

$$\begin{aligned} \varrho_\mu(x) &= \int_{\mathbb{T}} \text{tr}[\mathbf{u}^*(z) x \mathbf{u}(z) \varrho_0] \mu(dz) = \int_{\mathbb{T}} f_{\rho_0, x}(z) \mu(dz) \\ &= \text{tr} \left[x \int_{\mathbb{T}} \mathbf{u}(z) \varrho_0 \mathbf{u}^*(z) \mu(dz) \right] = \text{tr}[x \varrho_\mu] \end{aligned}$$

Consider iid random variables $(Z^i)_{i \in \mathbb{N}}$, $Z^i \sim \mu \in \mathcal{P}(\mathbb{T})$ and the corresponding sequence of empirical measures

$$\mu^n = \frac{1}{n} \sum_{i=1}^n \delta_{Z^i}, \quad \varrho_{\mu^n}(x) = \frac{1}{n} \sum_{i=1}^n \mathbf{u}(Z^i) \varrho_0 \mathbf{u}^*(Z^i)$$

$$\begin{aligned}\frac{1}{n} \log \mathbb{E}[e^{n\langle f, \mu^n \rangle}] &= \frac{1}{n} \log \mathbb{E} \left[\exp \left(\sum_{i=1}^n f(Z^i) \right) \right] \\ &= \log \left(\int_{\mathbb{T}} e^{f(z)} \mu(dz) \right)\end{aligned}$$

$$\sup_{f=f_{\varrho_0, x}} \left\{ \langle f, \nu \rangle - \log \int_{\mathbb{T}} e^{f(z)} \mu(dz) \right\} = \sup_{x \in \mathcal{O}(\mathcal{H})} \left\{ \text{tr}[x \varrho_\nu] - \log \int_{\mathbb{T}} e^{\text{tr}[\mathbf{u}^*(z) x \mathbf{u}(z) \varrho_0]} \mu(dz) \right\}$$

$$\begin{aligned}& \sum_{j,k} x_{jk} \text{tr}[E_{jk} \varrho_\nu] - \log \int_{\mathbb{T}} e^{\sum_{j,k} x_{jk} \text{tr}[\mathbf{u}^*(z) E_{jk} \mathbf{u}(z) \varrho_0]} \mu(dz) \\ &= \sum_{j,k} x_{jk} \text{tr}[E_{jk} \varrho_\nu] - \log \int_{\mathbb{T}} \prod_{j,k} e^{x_{jk} \text{tr}[\mathbf{u}^*(z) E_{jk} \mathbf{u}(z) \varrho_0]} \mu(dz)\end{aligned}$$

$$\text{tr}[E_{jk} \varrho_\nu] = \frac{1}{c} \int_{\mathbb{T}} \text{tr}[\mathbf{u}^*(z) E_{jk} \mathbf{u}(z) \varrho_0] e^{\sum_{j,k} x_{jk} \text{tr}[\mathbf{u}^*(z) E_{jk} \mathbf{u}(z) \varrho_0]} \mu(dz)$$

$$b = \frac{1}{c} \int_{\mathbb{T}} v(z) e^{\langle \mathbf{x}, v(z) \rangle} \mu(dz)$$

Consider the map

$$\mathbf{x} \mapsto \Lambda(\mathbf{x}) = \int_{\mathbb{T}} v(z) e^{\langle \mathbf{x}, v(z) \rangle} \mu(dz)$$

$$\langle \Lambda(\mathbf{x}) - \Lambda(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle = \int_0^1 \langle D\Lambda((1-\tau)\mathbf{x} + \tau\mathbf{y})[\mathbf{x} - \mathbf{y}], \mathbf{x} - \mathbf{y} \rangle d\tau \geq 0$$

$$\frac{d}{d\lambda} \int_{\mathbb{T}} v(z) e^{\langle \lambda, v(z) \rangle} \mu(dz) = \int_{\mathbb{T}} v(z) \otimes v(z) e^{\langle \lambda, v(z) \rangle} \mu(dz)$$

$$\text{tr}[E_{jk} \mathbf{u}(z) \varrho_0 \mathbf{u}^*(z)] = \langle \psi_k | \mathbf{u}(z) \varrho_0 \mathbf{u}^*(z) | \psi_j \rangle = \sum_i \lambda_i \langle \psi_k | \mathbf{u}(z) | \psi_i \rangle \langle \psi_i | \mathbf{u}^*(z) | \psi_j \rangle$$

If $\varrho_0 = \mathbb{1}$, then

$$\text{tr}[E_{jk} \mathbf{u}(z) \varrho_0 \mathbf{u}^*(z)] = \delta_{kj}.$$

Hence,

$$\text{tr}[E_{jk} \varrho_\nu] = \frac{1}{c} \delta_{kj} e^{\sum_j x_j}$$

3 Boson Fock spaces and quantum noise

3.1 Boson Fock spaces

Consider a (complex) Hilbert space \mathcal{H} with scalar product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. The symmetric Fock space associated with \mathcal{H} is

$$\mathfrak{F} = \mathfrak{F}_{\text{sym}}(\mathcal{H}) := \bigoplus_{n \in \mathbb{N}_0} \mathcal{H}^{\odot n}, \quad \mathcal{H}^{\odot 0} = \mathbb{C},$$

where \odot denotes the symmetric tensor product such that

$$\mathcal{H}^{\odot n} = \left\{ f \in \mathcal{H}^{\otimes n} : f(x_{\sigma_1}, \dots, x_{\sigma_n}) = f(x_1, \dots, x_n) \text{ for every permutation } \sigma \right\}.$$

The Fock space \mathfrak{F} inherits the scalar product from \mathcal{H} defined by

$$\langle \oplus f^{(n)}, \oplus g^{(n)} \rangle_{\mathfrak{F}} = \sum_{n \in \mathbb{N}_0} \langle f^{(n)}, g^{(n)} \rangle_{\mathcal{H}^{\otimes n}}.$$

We define the *vacuum vector* $\Omega = 1 \oplus 0 \oplus 0^{\otimes 2} \oplus \dots \in \mathfrak{F}$, and the *exponential vectors*

$$\mathbf{e}(f) = \bigoplus_{n \in \mathbb{N}_0} \frac{1}{\sqrt{n!}} f^{\otimes n}, \quad f \in \mathcal{H}.$$

It turns out that the family of exponential vectors \mathfrak{E} is *total* in \mathfrak{F} , i.e., the linear span of \mathfrak{E} is dense in \mathfrak{F} . This fact will be helpful for us in the future. Since,

$$\langle \mathbf{e}(f), \mathbf{e}(g) \rangle_{\mathfrak{F}} = \sum_{n \in \mathbb{N}_0} \frac{1}{n!} \langle f, g \rangle_{\mathcal{H}}^n = e^{\langle f, g \rangle_{\mathcal{H}}}, \quad f, g \in \mathcal{H},$$

the exponential vectors are normalizable. These normalized exponential vectors

$$\psi(f) = e^{-\frac{1}{2}\|f\|_{\mathcal{H}}^2} \mathbf{e}(f), \quad f \in \mathcal{H},$$

are commonly known as *coherent vectors*.

3.1.1 Weyl and field operators

For any $f \in \mathcal{H}$, we define the *Weyl operator* on exponential vectors by

$$\mathbf{W}(f)\mathbf{e}(g) := \exp\left(-\langle f, g \rangle_{\mathcal{H}} - \frac{1}{2}\|f\|_{\mathcal{H}}^2\right) \mathbf{e}(f+g), \quad g \in \mathcal{H}.$$

Weyl operators play an essential role in the setup of Fock spaces. For one, they generate coherent states by acting on the vacuum state, i.e.,

$$\mathbf{W}(f)\Omega = e^{-\frac{1}{2}\|f\|_{\mathcal{H}}^2} \mathbf{e}(f) = \psi(f), \quad f \in \mathcal{H}.$$

Moreover, they give the means to map any element $f \in \mathcal{H}$ to unitary operators on \mathfrak{F} that satisfy the *canonical commutation relation* (CCR):

Proposition 3.1. *The Weyl operator $W(f)$ is a unitary operator and satisfies*

$$(i) \quad W^\dagger(f) = W(-f).$$

$$(ii) \quad W(f)W(g) = e^{-i\text{Im}(\langle f, g \rangle_{\mathcal{H}})} W(f+g) = e^{-2i\text{Im}(\langle f, g \rangle_{\mathcal{H}})} W(g)W(f).$$

Property (ii) is the Weyl form of the canonical commutation relation (CCR).

Proof. For any $g \in \mathcal{H}$,

$$\begin{aligned} \langle W(f)e(g), W(f)e(g) \rangle_{\mathfrak{F}} &= \exp(-2\langle f, g \rangle_{\mathcal{H}} - \|f\|_{\mathcal{H}}^2) \langle e(f+g), e(f+g) \rangle_{\mathfrak{F}} \\ &= \exp(-2\langle f, g \rangle_{\mathcal{H}} - \|f\|_{\mathcal{H}}^2 + \|f+g\|_{\mathcal{H}}^2) \\ &= e^{\|g\|_{\mathcal{H}}^2} = \langle e(g), e(g) \rangle_{\mathfrak{F}}. \end{aligned}$$

Hence, $W(f)$ preserves inner products on \mathfrak{E} . Since \mathfrak{E} is dense in \mathfrak{F} , $W(f)$ extends uniquely to an isometry on \mathfrak{F} .

In a similar fashion, we compute

$$W(-f)W(f)e(g) = e(g) = W(f)W(-f)e(g) \quad \forall g \in \mathcal{H},$$

i.e., $W(-f)W(f)$ is the identity on the dense set \mathfrak{E} . In particular, $W(f)$ is surjective and an isometry, i.e., $W(f)$ is unitary with $W^\dagger(f) = W(-f)$.

As for the last property, we observe that

$$\begin{aligned} W(f)W(g)W(-(f+g))e(h) &= e^{\langle f+g, h \rangle_{\mathcal{H}} - \frac{1}{2}\|f+g\|_{\mathcal{H}}^2} W(f)W(g)e(-(f+g)+h) \\ &= e^{\langle f+g, h \rangle_{\mathcal{H}} - \frac{1}{2}\|f\|_{\mathcal{H}}^2 - \langle g, -f+h \rangle_{\mathcal{H}} - \text{Re}(\langle g, f \rangle_{\mathcal{H}})} W(f)e(-f+h) \\ &= e^{-i\text{Im}(\langle f, g \rangle_{\mathcal{H}})} e(h), \end{aligned}$$

and hence, $W(f)W(g)W(-(f+g)) = e^{-i\text{Im}(\langle f, g \rangle_{\mathcal{H}})} I_{\mathfrak{F}}$. We then conclude by using property (i) of Weyl operators. \square

Since $W(f)$ is unitary for every $f \in \mathcal{H}$, the family $\{W(tf)\}_{t \in \mathbb{R}}$ forms a one-parameter (strongly continuous) group of unitaries. In particular, due to Stone's theorem, it has a corresponding Hermitian operator $P(f)$ such that

$$W(tf) = \exp(itP(f)).$$

We further define the following operators

$$Q(f) := -P(if), \quad A^-(f) := \frac{Q(f) + iP(f)}{2}, \quad A^+(f) := \frac{Q(f) - iP(f)}{2}.$$

The operators A^\pm are called the *field operators* and will play an essential role as they form the creation/annihilation operators on Fock spaces.

Proposition 3.2. *The following are true: For any $f, g \in \mathcal{H}$,*

$$(i) \quad \mathfrak{E} \text{ is a core for } P(f) \text{ and } [P(f), P(g)] = 2i\text{Im}(\langle f, g \rangle_{\mathcal{H}})I_{\mathfrak{F}}.$$

$$(ii) \quad A^-(f)e(g) = \langle f, g \rangle_{\mathcal{H}} e(g), \quad A^+(f)e(g) = \frac{d}{dt} e(g + tf)|_{t=0}.$$

$$(iii) \quad W^\dagger(f)A^-(g)W(f) = A^-(g) + \langle g, f \rangle_{\mathcal{H}} \mathbf{l}_{\mathfrak{F}}, \quad W^\dagger(f)A^+(g)W(f) = A^+(g) + \langle f, g \rangle_{\mathcal{H}} \mathbf{l}_{\mathfrak{F}}.$$

$$(iv) \quad [A^-(f), A^-(g)] = [A^+(f), A^+(g)] = 0, \quad [A^-(f), A^+(g)] = \langle f, g \rangle_{\mathcal{H}} \mathbf{l}_{\mathfrak{F}},$$

i.e., the field operators A^\pm satisfy the canonical commutation relation.

Remark 3.3 On the finite particle vectors, the field operators act as

$$A^-(f)g^{\otimes n} = \sqrt{n} \langle f, g \rangle_{\mathcal{H}} g^{\otimes(n-1)}, \quad A^+(f)g^{\otimes(n-1)} = \frac{1}{\sqrt{n}} \sum_{k=0}^{n-1} g^{\otimes k} \otimes f \otimes g^{\otimes(n-1-k)}.$$

Example 3.4 Let $\mathcal{H} = L_{\mathbb{C}}^2([-\pi, \pi])$ be the space of square-integrable functions. Then the countable set $\{\psi_\ell(x) = e^{i\ell x} : \ell \in \mathbb{Z}\}$ forms an orthonormal basis for \mathcal{H} .

$$\psi(x) = \sum_{\ell} e^{i\ell x} A_{\ell}^-$$

3.1.2 Second quantization

The term *second quantization* is associated with the action of lifting operators on an k -particle space $\mathcal{H}^{\otimes k}$ to an associated operator on the Fock space.

We begin our discussion with the 1-particle case. For any bounded operator $A \in \mathcal{B}(\mathcal{H})$ one defines the map

$$\Gamma(A) := I \oplus \bigoplus_{n \in \mathbb{N}} A^{\otimes n},$$

which acts on $\mathcal{H}^{\otimes n}$ by

$$\Gamma(A)g_1 \otimes \cdots \otimes g_n = Ag_1 \otimes \cdots \otimes Ag_n.$$

Clearly, if A is unitary, then so is $\Gamma(A)$. Indeed, in this case, one has

$$\langle \Gamma(A)e(g), \Gamma(A)e(g) \rangle_{\mathfrak{F}(\mathcal{H})} = \sum_{n \in \mathbb{N}_0} \frac{1}{n!} \langle Ag, Ag \rangle_{\mathcal{H}}^n = \sum_{n \in \mathbb{N}_0} \frac{1}{n!} \|g\|_{\mathcal{H}}^n = \langle e(g), e(g) \rangle_{\mathfrak{F}(\mathcal{H})}.$$

Now let H be a self-adjoint operator on \mathcal{H} and $U_t := \exp(itH)$ be its unitary evolution. Then, $\Gamma(U_t)$ is a one-parameter group of unitary operators on $\mathfrak{F}(\mathcal{H})$. Stone's theorem then provides the existence of a densely defined Hermitian operator $d\Gamma(H)$ such that

$$\Gamma(U_t) = \exp(it d\Gamma(H)).$$

The generator $d\Gamma(H)$ is called the *second quantization of H* , and takes the explicit form

$$d\Gamma(H)g^{(n)} = \sum_{j=1}^n g_1 \otimes \cdots \otimes g_{j-1} \otimes Hg_j \otimes g_{j+1} \otimes \cdots \otimes g_n = \sum_{j=1}^n H_j g^{(n)},$$

for any $g^{(n)} = g_1 \otimes \cdots \otimes g_n \in \mathcal{H}^{\otimes n}$ with $g_j \in D(H)$ and

$$H_j = I_{\mathcal{H}} \otimes \cdots \otimes I_{\mathcal{H}} \otimes H \otimes I_{\mathcal{H}} \otimes \cdots \otimes I_{\mathcal{H}},$$

where H acts on the j -th tensor product. The special case $H = I_{\mathcal{H}}$ yields

$$\mathbf{N}g^{(n)} := d\Gamma(I_{\mathcal{H}})g^{(n)} = ng^{(n)}, \quad n \in \mathbb{N},$$

and is called the *number operator* due to its diagonal nature, with eigenvalues representing the number of particles in each configuration. Its domain is given by

$$D(\mathbf{N}) = \left\{ \{f^{(n)}\}_{n \in \mathbb{N}_0} : \sum_{n \in \mathbb{N}_0} n^2 \|f^{(n)}\|_{\mathcal{H}^{\otimes n}}^2 < +\infty \right\}.$$

$$\mathbf{A}_j^- \mathbf{A}_j^+ \psi_j^{\otimes n} = \sqrt{n} \mathbf{A}_j^- \psi_j^{\otimes (n-1)} = \sum_{k=0}^{n-1} \psi_j^{\otimes k} \otimes \psi_j \otimes \psi_j^{\otimes (n-1-k)} = n \psi_j^{\otimes n}$$

$$\mathbf{A}_j^+ \mathbf{A}_j^- \mathbf{e}(\psi_j) = \delta_{jk} \mathbf{N}_j \mathbf{e}(\psi_j)$$

This construction can be performed similarly for the general k -particle case.

$$d\Gamma(H^{(k)})f^{(n)} = \sum_{j_1 \neq \dots \neq j_k} H_{j_1 \dots j_k} f^{(n)},$$

where H_{j_1, \dots, j_k} denotes the operator where $H^{(k)}$ acts on the (j_1, \dots, j_k) -th tensor product.

Example 3.5 Consider the Hilbert space $\mathcal{H} = L_{\mathbb{C}}^2(\mathbb{T})$ and the Hamiltonians,

$$H^{(1)} = -\Delta \in \mathcal{O}(\mathcal{H}), \quad H^{(2)} = \mathbf{M}_W \in \mathcal{O}(\mathcal{H}^{\otimes 2}),$$

where $W : \mathbb{T} \times \mathbb{T} \rightarrow \mathbb{R}$ is an interaction potential, and \mathbf{M}_f denotes the multiplication operator corresponding to f .

Then, their second quantization is given by

$$d\Gamma(H^{(1)}) = \sum_{j=1}^n H_j^{(1)}, \quad d\Gamma(H^{(2)}) = \sum_{j \neq \ell} H_{j\ell}^{(2)} \quad \text{on } \mathcal{H}^{\otimes n}.$$

$$d\Gamma(H^{(1)})\psi_j^{\otimes n} = nj^2\psi_j^{\otimes n} = j^2\mathbf{A}_j^+\mathbf{A}_j^-\psi_j^{\otimes n} = j^2\mathbf{A}_j^+\mathbf{A}_j^-\psi_j^{\otimes n}$$

$$\psi^{(n)} = \psi_{k_1} \otimes \dots \otimes \psi_{k_n}$$

$$d\Gamma(H^{(1)})\psi^{(n)} = \sum_{j=1}^n k_j^2 \psi^{(n)}$$

3.1.3 Free field operators

For each $f \in \mathcal{H}$, we consider the pair $\{a^-(f), a^+(f)\}$ of operators on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ defined by

$$a^-(f)\Psi =$$

satisfying the commutation relations: For any $f, g \in \mathcal{H}$,

$$[a^\pm(f), a^\pm(g)] = 0, \quad [a^-(f), a^+(g)] = \langle f, g \rangle,$$

where $f \mapsto a^-(f)$ is conjugate linear and $f \mapsto a^+(f)$ is linear. Moreover, if $\Omega \in \mathfrak{F}_{\text{sym}}(\mathcal{H})$ is the *vacuum vector*, then $a^-(f)\Omega = 0$ for every $f \in \mathcal{H}$. The *field operators* a^- and a^+ are called *creation* and *annihilation* operators, respectively. On appropriate domains, the field operators are adjoints of one another, i.e., $(a^-(f))^\dagger = a^+(f)$ for every $f \in \mathcal{H}$.

It is common in the physics literature to consider operator-valued distributions $\{a_x^-, a_x^+\}$ instead, where if $\mathcal{H} = L_{\mathbb{C}}^2(\mathbb{R}^d)$, then $x \in \mathbb{R}^d$ and

$$a^-(f) = \int_{\mathbb{R}^d} \overline{f(x)} a_x \, dx, \quad a^+(f) = \int_{\mathbb{R}^d} f(x) a_x \, dx.$$

The commutation relations then simply read

$$[a_x^\pm, a_y^\pm] = 0, \quad [a_x^-, a_y^+] = \delta(x - y).$$

The *number operator* is formally defined by $N_x = a_x^+ a_x$, $x \in \mathbb{R}^d$.

On the rigorous not, if \mathcal{H} is separable with orthonormal basis $\{\psi_i\}$, then one obtains a family of field operators $\{a_i^-, a_i^+\}$ with $a_i^\pm : a^\pm(\psi_i)$.

3.1.4 Gaussian states

Definition 3.6 (Gaussian states) A state Ψ on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ is said to be a mean-zero *Gaussian* (or *quasi-free*) state if it can be uniquely determined from the field operators $\{a, a^\dagger\}$ by its covariance

$$\Sigma_\Psi(f, g) := \begin{pmatrix} \Psi(a^+(f) a^-(g)) & \Psi(a^-(f) a^-(g)) \\ \Psi(a^+(f) a^+(g)) & \Psi(a^-(f) a^+(g)) \end{pmatrix}, \quad f, g \in \mathcal{H}.$$

If the off-diagonal elements of the covariance are zero, the state Ψ is called *gauge-invariant* since it is invariant under the so-called gauge transformations of the first kind, i.e.,

$$a^\pm(f) = e^{\pm i\alpha} a^\pm(f) \quad \text{for any } \alpha \in \mathbb{R}.$$

If the off-diagonal elements of the covariance are nonzero, the state Ψ is called *squeezed*.

3.1.5 States invariant under free evolutions

Consider a Hamiltonian H on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ with $\mathcal{H} = L_{\mathbb{C}}^2(\mathbb{R}^d)$ and its associated 1-parameter automorphism group

$$\mathbf{u}_t(a) = e^{itH} a e^{-itH}, \quad t \in \mathbb{R}.$$

Definition 3.7 The Hamiltonian H on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ is called *free* if there exists a real-valued function $\omega: \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$\mathbf{u}_t(a_x^-) = e^{-it\omega(x)} a_x^-, \quad x \in \mathbb{R}^d.$$

In this case, the function ω is called the *free 1-particle Hamiltonian*, and H is said to be the *second quantization* of ω . Accordingly, \mathbf{u}_t is called a *free evolution*.

Example 3.8 Consider a Hamiltonian on $\mathcal{H} = L_{\mathbb{C}}^2(\mathbb{R})$ and its eigensystem $\{(\lambda_i, \psi_i)\}_i$ such that $\{\psi_i\}_i$ forms an orthonormal basis for \mathcal{H} . Setting $a_i^{\pm} = a^{\pm}(\psi_i)$, we see that

$$[a_i^{\pm}, a_j^{\pm}] = 0, \quad [a_i^-, a_j^+] = \delta_{ij},$$

i.e., a_i^{\pm} define field operators on the symmetric Fock space $\mathfrak{F}_{\text{sym}}(\mathcal{K})$ with $\mathcal{K} = \ell_{\mathbb{C}}^2$. Defining the field operators

$$a^- = \frac{1}{\sqrt{2}}(Q + iP), \quad a^+ = \frac{1}{\sqrt{2}}(Q - iP), \quad N := a^+ a^-$$

such that $[a^-, a^+] = I$, we find that

$$H = a^+ a^- + \frac{1}{2} = N + \frac{1}{2}.$$

Clearly, the eigenvectors of N and H coincide

Definition 3.9 A Gaussian state Ψ on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ with $\mathcal{H} = L^2(\mathbb{R}^d)$ is said to be invariant under a free evolution \mathbf{u}_t if

$$\Psi(\mathbf{u}_t(a_x^-) \mathbf{u}_s(a_y^+)) = \Psi(\mathbf{u}_{t-s}(a_x^-) a_y^+) = \Psi(a_x^- \mathbf{u}_{s-t}(a_y^+)) \quad \text{for all } t \in \mathbb{R}, x \in \mathbb{R}^d.$$

A *Gaussian free state* is a Gaussian state that is invariant under *all* free evolutions.

Theorem 3.10. A Gaussian state Ψ on $\mathfrak{F}_{\text{sym}}(\mathcal{H})$ is a Gaussian free state if and only if it is gauge-invariant and its diagonal correlations are supported on the diagonal, i.e.,

$$\Psi(a_x^- a_y^+) = m(x) \delta(x - y), \quad \Psi(a_x^+ a_y^-) = n(x) \delta(x - y).$$

Theorem 3.11. The field operators $\{a_x^{\pm}, a_x^{\mp}\}$ satisfying the commutation relations

$$[a_x^{\pm}, a_y^{\pm}] = 0, \quad [a_x^-, a_y^+] = m(x) \delta(x - y).$$

are mean-zero Gaussian random variables w.r.t. the Fock vacuum state $\Psi_{\Omega} = \langle \Omega, \cdot \Omega \rangle$, where $\Omega \in \mathfrak{F}_{\text{sym}}(\mathcal{H})$ is the Fock vacuum vector, with covariance

$$\Sigma_{\Psi}(x, y) = \begin{pmatrix} 0 & 0 \\ 0 & m(x) \end{pmatrix} \delta(x - y), \quad x, y \in \mathbb{R}^d.$$

Conversely, if $\{a_x^{\pm}, a_x^{\mp}\}$ are random variables with these properties, then they satisfy the commutation relations above.

3.1.6 Boson Fock white noise

def:boson-white-noise

Definition 3.12 A boson Fock white noise on \mathbb{R}^d is a boson Fock field $\{b_{t,x}^+, b_{t,x}^-\}$ on \mathbb{R}^{d+1} with vacuum vector Ω satisfying the commutation relations

$$[b_{t,x}^\pm, b_{s,y}^\pm] = 0, \quad [b_{t,x}^-, b_{s,y}^+] = \delta(t-s)m(x)\delta(x-y), \quad b_{t,x}^-\Omega = 0.$$

thm:stochastic-limit

Theorem 3.13. Let $\{a_x^+, a_x^-\}$ be Gaussian free fields w.r.t. the Fock vacuum state Ψ_Ω with

$$\mathbf{u}_t(a_x^-) = e^{-it\omega(x)}a_x^-.$$

Then the rescaled field operators

$$b_{t,x}^{\lambda,\pm} := \frac{1}{\lambda} \mathbf{u}_{t/\lambda^2}(a_x^\pm)$$

converges in the sense of correlator distributions to a boson Fock white noise, i.e.,

$$\lim_{\lambda \rightarrow 0} \Psi(b_{t,x}^{\lambda,\varepsilon_1} b_{t,y}^{\lambda,\varepsilon_2}) = \Psi(b_{t,x}^{\varepsilon_1} b_{t,y}^{\varepsilon_2}) \quad \varepsilon_1, \varepsilon_2 \in \{+, -\},$$

where $b_{t,x}^\pm$ as defined in Definition 3.12 with $m = 2\pi\delta(\omega)$.

Proof. Using the invariance of Ψ_Ω under free evolutions, we find

$$\begin{aligned} \Psi_\Omega(b_{t,x}^{\lambda,-} b_{s,y}^{\lambda,+}) &= \frac{1}{\lambda^2} \Psi_\Omega(\mathbf{u}_{(t-s)/\lambda^2}(a_x^-) a_y^+) \\ &= \frac{1}{\lambda^2} e^{-i\omega(x)(t-s)/\lambda^2} \Psi_\Omega(a_x^- a_y^+) = \frac{1}{\lambda^2} e^{-i\omega(x)(t-s)/\lambda^2} \delta(x-y). \end{aligned}$$

Passing to the limit $\lambda \rightarrow 0$ recovers the desired limit. All the other terms vanish. \square

Remark 3.14 Let $\mathcal{K} \subset L^2(\mathbb{R}^d)$ be a set of functions for which

$$\int_{\mathbb{R}} |\langle f, e^{it\omega} g \rangle| dt < +\infty \quad \forall f, g \in \mathcal{K}.$$

Since $t \mapsto \langle f, e^{it\omega} f \rangle$ is positive definite for each $f \in \mathcal{K}$, Bochner's theorem implies that the sesquilinear form

$$\langle f, 2\pi\delta(\omega)g \rangle := \int_{\mathbb{R}} \langle f, e^{it\omega} g \rangle dt,$$

is a pre-scalar product. With $(\cdot|\cdot)$, the set \mathcal{K} becomes a pre-Hilbert space, which can be completed to obtain a Hilbert space, still denoted by \mathcal{K} . The function $m = 2\pi\delta(\omega)$ has to be understood in this sense, and only makes sense for functions in \mathcal{K} .

The operators

$$B_t^-(f) := \int_0^t \int_{\mathbb{R}^d} \overline{f(x)} b_{s,x}^- dx ds, \quad B_t^+(f) := \int_0^t \int_{\mathbb{R}^d} f(x) b_{s,x}^+ dx ds,$$

define *quantum Brownian motions*. The self-adjoint (*momentum*) operators

$$P_t(f) := \frac{1}{i} [B_t^-(f) - B_t^+(f)],$$

form a commuting family of classical random variables whose statistics in the Fock vacuum state $\Psi_\Omega = \langle \Omega, \cdot \Omega \rangle$ is completely determined by the relation

$$\Psi(e^{iP_t(f)}) = \exp\left(-\frac{t}{2} \|f\|_{L^2(\mathbb{R}^d)}^2\right). \quad \text{check!}$$

3.1.7 Gaussian equilibrium states: The KMS condition

For any states $a, b \in \mathcal{A}$, the map $t \mapsto \Psi(a \mathbf{u}_t(b))$ can be analytically continued and satisfies the so-called *KMS condition* at inverse temperature $\beta > 0$:

$$\Psi(a \mathbf{u}_{t+i\beta}(b)) = \Psi(\mathbf{u}_t(a) b) \quad \forall a, b \in \mathcal{A}. \quad \text{check!}$$

3.2 Composite systems

Definition 3.15 A composite system of two given quantum dynamical systems $S = \{\mathcal{H}_S, H_S\}$, $R = \{\mathcal{H}_R, H_R\}$ is a quantum dynamical system of the form

$$\{\mathcal{H}_S \otimes \mathcal{H}_R, H_{SR}\}, \quad H_{SR} = H_S \otimes 1_R + 1_S \otimes H_R + H_I,$$

where H_I is called the *interaction Hamiltonian* and contains all the new physics of the composite system, while $H_0 := H_S \otimes 1_R + 1_S \otimes H_R$ is called the free Hamiltonian.

We will consider *scaled* total Hamiltonians

$$H^\lambda := H_0 + \lambda H_I,$$

and the following unitary evolutions:

$$\begin{aligned} \text{free evolution} \quad V_t^0 &= e^{-itH_0}, & \text{total evolution} \quad V_t^\lambda &= e^{-itH^\lambda}, \\ \text{interacting representation evolution} \quad U_t^\lambda &= (V_t^0)^\dagger V_t^\lambda, \end{aligned}$$

where U_t^λ satisfies the Schrödinger equation in the interaction picture:

$$\partial_t U_t^\lambda = -i\lambda H_I(t) U_t^\lambda, \quad U_0^\lambda = I,$$

with the time dependent Hamiltonian $H_I(t) = (V_t^0)^\dagger H_I V_t^0$.

For simplicity, we will make the following assumptions on system $S = \{\mathcal{H}_S, H_S\}$, the reservoir $R = \{\mathcal{H}_R, H_R\}$, and the interaction Hamiltonian H_I .

3.2.1 The reservoir

The reservoir $R = \{\mathcal{H}_R, H_R\}$ is given by the Hilbert space $\mathfrak{H}_{\text{sym}}(\mathcal{H})$ with $\mathcal{H} = L_{\mathbb{C}}^2(\mathbb{R}^d)$, a free Hamiltonian H_R with continuous spectrum \mathbb{R} , and a mean-zero Gaussian free state Ψ_R such that

$$\int_{\mathbb{R}} |\Psi_R(a_x^{\varepsilon_1} \mathbf{u}_t(a_y^{\varepsilon_2}))| dt < +\infty, \quad \varepsilon_1, \varepsilon_2 \in \{+, -\}.$$

In particular, Ψ_R is characterized by the covariances

$$\Psi_R(a_x^- a_y^+) = m(x) \delta(x - y), \quad \Psi_R(a_x^+ a_y^-) = n(x) \delta(x - y).$$

Since H_R is free, there exists a function ω , for which

$$\mathbf{u}_t(a_x^-) = e^{itH_R} a_x^- e^{-itH_R} = e^{-it\omega(x)} a_x^-,$$

where ω describes the 1-particle evolution.

An example of a free reservoir Hamiltonian is given by

$$H_R = \int_{\mathbb{R}^d} \omega(x) a_x^+ a_x^- dx,$$

where ω is a smooth cutoff function.

3.2.2 The system Hamiltonian

For simplicity, we shall assume that the system Hamiltonian H_S has a discrete spectrum such that

$$H_S = \sum_j \lambda_j P_j,$$

where λ_j are the eigenvalues and P_j are their corresponding spectral projections.

3.2.3 The interaction Hamiltonian

We consider dipole-type interaction Hamiltonians of the form

$$H_I = \int_{\mathbb{R}^d} [D(x) \otimes a_x^+ + D^\dagger(x) \otimes a_x^-] dx,$$

where $\{D(x) : x \in \mathbb{R}^d\}$ is a family of system operators called the *response terms*.

With the spectral projections of H_S , we may express H_I as

$$H_I = \sum_{j,k} \int_{\mathbb{R}^s} [P_j D(x) P_k \otimes a_x^+ + P_k D^\dagger(x) P_j \otimes a_x^-] dx,$$

and hence, the time-dependent Hamiltonian reads

$$\begin{aligned} H_I(t) &= \sum_{j,k} \int_{\mathbb{R}^d} [P_j D(x) P_k \otimes e^{it(\omega(x)+\lambda_j-\lambda_k)} a_x^+ + P_k D^\dagger(x) P_j \otimes e^{-it(\omega(x)+\lambda_j-\lambda_k)} a_x^-] dx \\ &= \sum_q \sum_{\lambda_k-\lambda_j=\eta_q} \int_{\mathbb{R}^d} [P_j D(x) P_k \otimes e^{it(\omega(x)-\eta_q)} a_x^+ + P_k D^\dagger(x) P_j \otimes e^{-it(\omega(x)-\eta_q)} a_x^-] dx \\ &= \sum_q \int_{\mathbb{R}^d} [D_q(x) \otimes e^{it(\omega(x)-\eta_q)} a_x^+ + D_q^\dagger(x) \otimes e^{-it(\omega(x)-\eta_q)} a_x^-] dx, \end{aligned}$$

where the system operators

$$D_q(x) := \sum_{\lambda_k-\lambda_j=\eta_q} P_j D(x) P_k \quad \text{satisfy} \quad e^{itH_S} D_q(x) e^{-itH_S} = e^{-it\eta_q} D_q(x).$$

To simplify things drastically, we assume that $q = 1$ and $D_1(x) = \chi(x)D$ for some smooth cutoff function χ and a fixed system operator D . In this case, we obtain

$$H_I(t) = \int_{\mathbb{R}^d} [D \otimes \chi(x) e^{it(\omega(x)-\eta)} a_x^+ + D^\dagger \otimes \overline{\chi(x)} e^{-it(\omega(x)-\eta)} a_x^-] dx.$$

3.3 The weak interaction stochastic limit

Altogether, we arrive at the rescaled Schrödinger equation in the interaction picture

$$U_{t/\lambda^2}^\lambda = I - i \int_0^t H_I^\lambda(s) U_{s/\lambda^2}^\lambda ds, \quad \text{eq:rescaled-schrodinger-interaction} \quad (3.1)$$

with $H_I^\lambda(t) = D \otimes b_t^{\lambda,+} + D^\dagger \otimes b_t^{\lambda,-}$, where, due to Theorem 3.13,

$$b_t^{\lambda,\pm} = \frac{1}{\lambda} a^\pm(\chi e^{i(t/\lambda^2)(\omega-\eta)}) \longrightarrow b_t^\pm \quad \text{in the sense of correlators,}$$

and therefore,

$$H_I^\lambda(t) \longrightarrow H_t = D \otimes b_t^+ + D^\dagger \otimes b_t^-, \quad U_{t/\lambda^2}^\lambda \longrightarrow U_t,$$

where U_t satisfies the formal stochastic differential equation

$$dU_t = -i(D \otimes dB_t^+ + D^\dagger \otimes dB_t^-)U_t, \quad B_t^\pm = \int_0^t b_s^\pm ds. \quad (\text{eq:sde} \quad (3.2))$$

Under certain assumptions on ω , this SDE has a unique solution.

The SDE (3.2) in natural-time order (or Itô form) is the SDE given by

$$dU_t = -i(D dB_t^+ U_t + D^\dagger U_t dB_t^-) - \gamma_- D^\dagger D U_t dt, \quad (\text{eq:sde-normal} \quad (3.3))$$

obtained by commuting $dB_t^- U_t$ and using the fact that the solution U_t to (3.2) satisfies

$$\begin{aligned} [b_t^-, U_t] &= -i\gamma_- D U_t, \\ [U_t^\dagger, b_t^+] &= i\bar{\gamma}_- U_t^\dagger D^\dagger, \quad \text{where } \gamma_- := \int_{-\infty}^0 \langle \chi, e^{-it(\omega-\eta)} \chi \rangle dt. \\ [b_t^-, U_t^\dagger] &= i\gamma_- U_t^\dagger D, \end{aligned}$$

The additional term is known as the Itô correction term in quantum stochastic calculus. Morally, the natural-time order is the order induced by the filtration for which one expects U_t to be adapted to. **Expand on the concept of normal order!**

With the unitary evolution U_t one can derive an evolution for any system observable $X_t = U_t^\dagger (X \otimes 1_{\mathcal{H}_R}) U_t$ given by

$$dX_t = -i dB_t^+ [D, X_t] + i[X_t, D^\dagger] dB_t^- + LX_t dt$$

where

$$LX = 2\Re(\gamma_-) D^\dagger X D - \gamma_- D^\dagger D X - \bar{\gamma}_- X D^\dagger D,$$

is the corresponding Lindblad operator.

Remark 3.16 The rescaled equation (3.1) may be formally expressed as

$$U_{t/\lambda^2}^\lambda = I + \sum_{n=1}^{\infty} (-i)^n \lambda^n \int_0^{t/\lambda^2} \cdots \int_0^{t_{n-1}} H_I(t_1) \cdots H_I(t_n) dt_1 \cdots dt_n.$$

In the case when $H_I(t)$ commutes for every $t \geq 0$, the integrand would be a symmetric function of t_1, \dots, t_n , and gives

$$U_{t/\lambda^2}^\lambda = I + \sum_{n=1}^{\infty} (-i)^n \frac{\lambda^n}{n!} \left(\int_0^{t/\lambda^2} H_I(s) ds \right)^n = \exp \left(-i\lambda \int_0^{t/\lambda^2} H_I(s) ds \right),$$

i.e., the expectation of U_{t/λ^2}^λ is the characteristic function of the process

$$W_t^\lambda := \lambda \int_0^{t/\lambda^2} H_I(s) ds.$$

3.4 Damped harmonic oscillator

We consider a simple setup in which a single atom interacts with an electromagnetic field. The total system is given by the composite of the atom and the reservoir systems

$$S = \{\mathfrak{F}_{\text{sym}}(\mathbb{C}^2), H_S\}, \quad R = \{\mathfrak{F}_{\text{sym}}(L^2(\mathbb{R}^d)), H_R\},$$

with the free Hamiltonians

$$H_S = \omega_0 c^+ c^-, \quad H_R = \int_{\mathbb{R}^d} \omega(x) a_x^+ a_x^- dx,$$

where $\{c^+, c^-\}$ is the field operator for $\mathcal{H}_S = \mathfrak{F}_{\text{sym}}(\mathbb{C}^2)$, $\{a_x^+, a_x^-\}$ are the field operators for $\mathcal{H}_R = \mathfrak{F}_{\text{sym}}(L^2(\mathbb{R}^d))$, and ω is a suitable cutoff function.

For the interaction Hamiltonian H_I , we consider a dipole approximation of the form

$$H_I = \int_{\mathbb{R}^d} \chi(x) [c^- \otimes a_x^+ + c^+ \otimes a_x^-] dx = c^- \otimes A^+ + c^+ \otimes A^-,$$

where χ are suitable cutoff function and

$$A^\pm = \int_{\mathbb{R}^d} \chi(x) a_x^\pm dx.$$

such that the rescaled total Hamiltonian is given by

$$H^\lambda = H_S \otimes 1_{\mathcal{H}_R} + 1_{\mathcal{H}_S} \otimes H_R + \lambda H_I.$$

Define the evolution

$$\mathbf{u}_t^\lambda(a) = e^{itH^\lambda} a e^{-itH^\lambda}.$$

Then, the Heisenberg equation for $c(t) = \mathbf{u}_t^\lambda(c \otimes 1_{\mathcal{H}_R})$ and $a_x^-(t) = \mathbf{u}_t^\lambda(1_{\mathcal{H}_S} \otimes a_x^-)$ reads

$$\begin{aligned} \frac{d}{dt} c^-(t) &= -i\omega_0 c^-(t) - i\lambda \int_{\mathbb{R}^d} \chi(x) a_x^-(t) dx \\ \frac{d}{dt} a_x^-(t) &= -i\omega(x) a_x^-(t) - \lambda \chi(x) c(t). \end{aligned}$$

Solving for $a_x^-(t)$, one obtains

$$a_x^-(t) = a_x^- e^{-it\omega(x)} - i\lambda \chi(x) \int_0^t e^{-i(t-s)\omega(x)} c^-(s) ds.$$

Inserting $a_x^-(t)$ into the equation for $c(t)$, we find

$$\frac{d}{dt} c^-(t) = -i\omega_0 c^-(t) - \int_0^t \gamma(t-s) c^-(s) ds - i\xi_t^-.$$

where we defined the quantities

$$\gamma(r) := \lambda^2 \int_{\mathbb{R}^d} e^{-ir\omega(x)} \chi^2(x) dx, \quad \xi_t^- := \lambda \int_{\mathbb{R}^d} e^{-it\omega(x)} \chi(x) a_x^- dx.$$

Observe that ξ_t^- depends only on the reservoir field operator a_x^- , and therefore, acts as an *external force* to the atomic system.

Now let us consider the statistics of ξ_t^- for the Fock vacuum state free state $\Psi_\Omega = \langle \Omega, \cdot \Omega \rangle$ on \mathcal{H}_R . Since Ψ_Ω is a Gaussian free state, the statistics of ξ_t^- are uniquely determined by the 2-points correlation functions. Clearly, Ψ_Ω has zero mean, and therefore,

$$\Psi_\Omega(\xi_t^-) = \lambda \int_{\mathbb{R}^d} e^{-it\omega(x)} \chi(x) \Psi_\Omega(a_x^-) dx = 0, \quad \Psi_\Omega((\xi_t^-)^\dagger) = 0.$$

Moreover, we find

$$\Psi_\Omega((\xi_t^-)^\dagger \xi_s^-) = \Psi_\Omega(\xi_t^- \xi_s^-) = \Psi_\Omega((\xi_t^-)^\dagger (\xi_s^-)^\dagger) = 0,$$

and

$$\begin{aligned} \Psi_\Omega(\xi_t^- (\xi_s^-)^\dagger) &= \lambda^2 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} e^{-it(\omega(x)-\omega(y))} \chi(x) \chi(y) \Psi_\Omega(a_x^- a_y^+) dx dy \\ &= \lambda^2 \int_{\mathbb{R}^d} e^{-i(t-s)\omega(x)} \chi^2(x) dx = \gamma(t-s). \end{aligned}$$

Hence, ξ_t^- is a mean-zero γ -correlated *random process* under Ψ .

To obtain the stochastic limit, we start by rescaling time $t \mapsto t/\lambda^2$ and consider the rescaled quantity $c^{\lambda,\pm}(t) := c^\pm(t/\lambda^2)$, which yields

$$\dot{c}^{\lambda,-}(t) = -i\omega_0 c^{\lambda,-}(t) - \int_0^t \gamma^\lambda(t-s) c^{\lambda,-}(s) ds - i \xi_t^{\lambda,-}, \quad \text{eq:quantum-pre-langevin} \quad (3.4)$$

with

$$\gamma^\lambda(r) := \frac{1}{\lambda^2} \langle \chi, e^{-i(r/\lambda^2)\omega} \chi \rangle, \quad \xi_t^{\lambda,-} := \lambda^{-2} \xi_{t/\lambda^2}^-.$$

From Theorem 3.13, we then establish that

$$\gamma^\lambda(r) \longrightarrow \gamma \delta(r), \quad \xi_t^{\lambda,-} \longrightarrow b_t^- \quad \text{in the sense of correlators,}$$

with $\gamma = \langle \chi, \delta(\omega) \chi \rangle$. Consequently, we obtain

$$dc_t^- = -(i\omega_0 + \gamma) c_t^- dt - i dB_t^-, \quad B_t^- = \int_0^t b_s^- ds.$$

$$dU_t = -i(c^- dB_t^+ U_t + c^- U_t dB_t^-) - \gamma_- c^+ c^- U_t dt$$

$$\begin{aligned} d(U_t^\dagger c^- U_t) &= i(c^- dB_t^+ U_t + c^+ U_t dB_t^-)^\dagger c^- U_t - \bar{\gamma}_- U_t^\dagger c^- c^+ c^- U_t dt \\ &\quad - iU_t^\dagger c^- (c^- dB_t^+ U_t + c^+ U_t dB_t^-) - \gamma_- U_t^\dagger c^- c^+ c^- U_t dt \\ &= iU_t^\dagger dB_t^- c^+ c^- U_t + i dB_t^+ U_t^\dagger c^- c^- U_t - \bar{\gamma}_- U_t^\dagger c^- c^+ c^- U_t dt \end{aligned}$$