

References

1. Ascher, U.M. and Petzold, L.R.: Computer Methods for Ordinary Differential Equations and Differential-Algebraic Equations, SIAM (1998)
2. Azamov, A.A. and Bekimov, M.A.: An approximation algorithm for quadratic dynamic systems based on N. Chomsky's grammar for Taylor's formula, Proc. Steklov Inst. Math. **293**, S1–S5 (2016)
3. Bashforth, F. and Adams, J.C.: An Attempt to Test the Theories of Capillary Action by Comparing the Theoretical and Measured Forms of Drops of Fluid, with an Explanation of the Method of Integration Employed in Constructing the Tables which Give the Theoretical Forms of Such Drops. Cambridge University Press, Cambridge (1883)
4. Brouder, C.: Runge–Kutta methods and renormalization, Eur. Phys. J. C. **12**, 521–534 (2000)
5. Burrage, K.: A special family of RungeKutta methods for solving stiff differential equations, BIT **18** 22–41 (1978)
6. Burrage K. and Butcher, J.C.: Non-linear stability of a general class of differential equation methods. BIT **20**, 185–203 (1980)
7. Butcher, J.C.: Coefficients for the study of Runge–Kutta integration processes. J. Austral. Math. Soc. **3**, 185–201 (1963)
8. Butcher, J.C.: On the integration processes of A. Hut'a, J. Austral. Math. Soc. **3**, 202–206 (1963)
9. Butcher, J.C.: Implicit RungeKutta processes, Math. Comp. **18**, 50–64 (1964)
10. Butcher, J.C.: A modified multistep method for the numerical integration of ordinary differential equations. J. Assoc. Comput. Mach. **12**, 124–135 (1965)
11. Butcher, J.C.: On the attainable order of Runge–Kutta methods, Math. Comp. **19**, 408–417 (1965)
12. Butcher, J.C.: On the convergence of numerical solutions to ordinary differential equations, Math. Comp. **20**, 1–10 (1966)
13. Butcher, J.C.: The effective order of Runge–Kutta methods, Lecture Notes in Math. **109**, 133–139 (1969)
14. Butcher, J.C.: An algebraic theory of integration methods. Math. Comp. **26**, 79–106 (1972)
15. Butcher, J.C.: A stability property of implicit Runge–Kutta methods. BIT **15**, 358–381 (1975)
16. Butcher, J.C.: The nonexistence of ten-stage eighth order explicit Runge–Kutta methods, BIT **25**, 521–540 (1985)
17. Butcher, J.C.: The Numerical Analysis of Ordinary Differential Equations, Runge–Kutta and General Linear Methods, John Wiley & Sons Ltd, Chichester (1987)
18. Butcher, J.C.: An introduction to “Almost Runge–Kutta” methods. Appl. Numer. Math. **24**, 331–342 (1997)
19. Butcher, J.C.: General linear methods, Acta Numerica **15**, 157–256 (2006)
20. Butcher, J.C.: Numerical Methods for Ordinary Differential Equations (Third Edition), John Wiley & Sons, Chichester (2016)

21. Butcher, J.C., Habib, Y., Hill, A.T. and Norton, T.J.T.: The control of parasitism in G-symplectic methods, *SIAM J. Numer. Anal.* **52**, 2440–2465 (2014)
22. Butcher, J.C. and Hill, A.T.: Linear multistep methods as irreducible general linear methods. *BIT Numerical Mathematics* **46**, 5–19 (2006)
23. Butcher, J.C., Hill, A.T. and Norton T.J.T.: Symmetric general linear methods. *BIT Numerical Mathematics* **56**, 1189–1212 (2016)
24. Butcher, J.C. and Imran, G.: Order conditions for G-symplectic methods. *BIT* **55**, 927–948 (2015)
25. Butcher, J.C., Imran, G. and Podhaisky, H.: A G-symplectic method with order 6. *BIT Numerical Mathematics* **57**, 313–328 (2017)
26. Butcher, J.C. and Sanz-Serna, J.M.: The number of conditions for a Runge–Kutta method to have effective order p . *Appl. Numer. Math.* **22**, 103–111 (1996)
27. Byrne, G.D. and Lambert, R.J.: Pseudo-Runge–Kutta methods involving two points. *J. Assoc. Comput. Mach.* **13**, 114–123 (1966)
28. Cayley, A.: On the theory of the analytical forms called trees. *Phil. Mag.* **13**, 172–176 (1857)
29. Celledoni, E., McLachlan, R.I., McLaren, D.I., Owren, B., Quispel, G.R.W. and Wright, W.M.: Energy-preserving Runge–Kutta methods, *Math. Model. Numer. Anal.* **43**, 645–649 (2009)
30. Coddington, E.A. and Levinson, N.: *Theory of Ordinary Differential Equations*, McGraw–Hill, New York (1955)
31. Cohen, D. and Hairer E.: Linear energy-preserving integrators for Poisson systems, *BIT* **51**, 91–101 (2011)
32. Cooper, G.J.: Stability of Runge–Kutta methods for trajectory problems, *J. Numer. Anal.* **7**, 1–13 (1987)
33. Cooper, G.J. and Verner, J.H.: Some explicit Runge–Kutta methods of high order, *SIAM J. Numer. Anal.* **9**, 389–405 (1972)
34. Creutz, M. and Gocksch, A.: Higher order hybrid Monte-Carlo algorithms, *Phys. Rev. Lett.* **63**, 912 (1989)
35. Curtiss, C.F. and Hirschfelder, J.O.: Integration of stiff equations. *Proc. Nat. Acad. Sci. U.S.A.* **38**, 235–243 (1952)
36. Dahlquist, G.: Convergence and stability in the numerical integration of ordinary differential equations. *Math. Scand.* **3**, 33–53 (1963)
37. Dahlquist, G.: A special stability property of linear multistep methods. *BIT* **4**, 27–43 (1956)
38. Dahlquist, G.: Error analysis of a class of methods for stiff nonlinear initial value problems, *Numerical Analysis, Dundee, Lecture Notes in Mathematics* **506**, 60–74 (1976)
39. Dahlquist, G. and Jeltsch, R.: Reducibility and contractivity of Runge–Kutta methods revisited, *BIT* **46**, 567–587 (2006)
40. D’Ambrosio, R. and Hairer, E.: Long-term stability of multi-value methods for ordinary differential equations, *J. Sci. Comput.* **60**, 627–640 (2014)
41. Donelson, J. and Hansen, E.: Cyclic composite multistep predictor–corrector methods. *SIAM J. Numer. Anal.* **8**, 137–157 (1971)
42. Euler, L.: De integratione aequationum differentialium per approximationem, In: *Opera Omnia*, 1st series, Vol. 11, *Institutiones Calculi Integralis*, pp 424–434. Teubner, Leipzig and Berlin (1913)
43. Gear, C.W.: Hybrid methods for initial value problems in ordinary differential equations. *SIAM J. Numer. Anal.* **2**, 69–86 (1965)
44. Gear, C.W.: The numerical integration of ordinary differential equations, *Math. Comp.* **21**, 146–156 (1967)
45. Gill, S.: A process for the step-by-step integration of differential equations in an automatic computing machine. *Proc. Cambridge Philos. Soc.* **47**, 96–108 (1951)
46. Gragg, W.B. and Stetter, H.J.: Generalized multistep predictor–corrector methods. *J. Assoc. Comput. Mach.* **11**, 188–209 (1964)
47. Hairer, E.: A Runge–Kutta method of order 10, *J. Inst. Math. Appl.* **21**, 47–59 (1978)
48. Hairer, E.: Energy-preserving variant of collocation methods, *J. Numer. Anal. Ind. Appl. Math.* **5**, 73–84 (2010)

49. Hairer, E., Lubich, C. and Wanner, G.: Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations, second edition, Springer-Verlag, Berlin (2006)
50. Hairer, E., Nørsett, S.P. and Wanner, G.: Solving Ordinary Differential Equations I: Nonstiff Problems, Springer, Berlin (1993)
51. Hairer, E. and Wanner, G.: Multistep-multistage-multiderivative methods for ordinary differential equations. *Computing* **11**, 287–303 (1973)
52. Hairer, E. and Wanner, G.: On the Butcher group and general multi-value methods. *Computing* **13**, 1–15 (1974)
53. Hairer, E. and Wanner, G.: Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems, Springer, Berlin (1996)
54. Hammer, P.C. and Hollingsworth, J.W.: Trapezoidal methods of approximating solutions of differential equations. *Math. Tables Aids Comput.* **9**, 92–96 (1955)
55. Henrici, P.: Discrete Variable Methods in Ordinary Differential Equations. John Wiley & Sons Inc, New York (1962)
56. Heun, K.: Neue Methode zur approximativen Integration der Differentialgleichungen einer unabhängigen Veränderlichen. *Z. Math. Phys.* **45**, 23–38 (1900)
57. Hull, T.E., Enright, W.H., Fellen, B.M. and Sedgwick, A.E.: Comparing numerical methods for ordinary differential equations. *SIAM J Numer. Anal.* **9**, 603–637 (1972)
58. Hundsdorfer W.H. and M. N. Spijker M.N.: A note on B-stability of Runge–Kutta methods. *Numer. Math.* **36**, 319–333 (1981)
59. Huťa, A.: Une amélioration de la méthode de Runge–Kutta–Nyström pour la résolution numérique des équations différentielles du premier ordre. *Acta Fac. Nat. Univ. Comenian. Math.* **1**, 201–224 (1956)
60. Huťa, A.: Contribution à la formule de sixième ordre dans la méthode de Runge–Kutta–Nyström. *Acta Fac. Nat. Univ. Comenian. Math.* **2**, 21–24 (1957)
61. Iserles, A.: A First Course in the Numerical Analysis of Differential Equations (Cambridge Texts in Applied Mathematics), second edition (2008)
62. Iserles, A., Munthe-Kaas, H.Z. Nørsett, S.P. and Zanna, A.: Lie-group methods, in *Acta Numerica* **9**, 215–365 (2001)
63. Jackiewicz, Z.: General Linear Methods for Ordinary Differential Equations, John Wiley & Sons, Hoboken (2009)
64. Kirchgraber, U.: Multistep methods are essentially one-step methods. *Numer. Math.* **48**, 85–90 (1986)
65. Kuntzmann, J.: Neuere Entwicklungen der Methode von Runge–Kutta. *Z. Angew. Math. Mech.* **41**, T28–T31 (1961)
66. Kutta, W.: Beitrag zur näherungsweisen Integration totaler Differentialgleichungen. *Z. Math. Phys.* **46**, 435–453 (1901)
67. Lambert, J.D.: Numerical Methods for Ordinary Differential Systems: The Initial Value Problem, Wiley, Chichester (1991)
68. Lasagni, F.M.: Canonical Runge–Kutta methods, *Z. Angew. Math. Phys.* **39**, 952–953 (1988)
69. López-Marcos, M.A., Skeel, R.D and Sanz-Serna, J.M.: Cheap enhancement of symplectic integrators, in *Numerical Analysis*, D.F. Griffiths and G.A. Watson (eds.), Pitman Res. Notes Math. Ser. **344**, Longman, Harlow, 107–122 (1996)
70. Łukasiewicz, J. and Tarski, J.: Investigations into the sentential calculus. *Comp. rend. Soc. Sci. Lett. Warsaw* **23**, Class III, 31–32 (1930)
71. McLachlan, R. and Quispel, R.: Six lectures on the geometric integration of ODEs. *Foundations of Computational Mathematics* (Oxford, 1999) London Math. Soc. Lecture Note Ser. 284: Cambridge Univ. Press, Cambridge 155–210 (2001)
72. Merson, R.H.: An operational method for the study of integration processes. *Proc. Symp. Data Processing*, Weapons Research Establishment, Salisbury, S. Australia (1957)
73. Miyatake Y: An energy-preserving exponentially-fitted continuous stage Runge–Kutta method for Hamiltonian systems, *BIT* **54**, 777–799 (2014)
74. Moulton, F.R.: New Methods in Exterior Ballistics. University of Chicago Press (1926)
75. Naur, P.: Report on the algorithmic language ALGOL 60, *Comm. ACM*, **6** (1), 1–17 (1963)

76. Nordsieck, A.: On numerical integration of ordinary differential equations. *Math. Comp.* **16**, 22–49 (1962)
77. Nyström, E.J.: Über die numerische Integration von Differentialgleichungen. *Acta Soc. Sci. Fennicae* **50** (13), 1–55 (1925)
78. Prince, P.J. and Dormand, J.R.: High order embedded Runge–Kutta formulae, *J. Comput. Appl. Math.* **7**, 67–75 (1981)
79. Prothero, A. and Robinson, A.: On the stability and accuracy of one-step methods for solving stiff systems of ordinary differential equations. *Math. Comp.* **28**, 145–162 (1974)
80. Quispel, G.R.W. and McLaren, D.I.: A new class of energy-preserving numerical integration methods, *J. Phys. A* **41**, 045206 (2008)
81. Rudin, W.: *Principles of Mathematical Analysis*. McGraw-Hill, 3rd Edition (1976)
82. Runge, C.: Über die numerische Auflösung von Differentialgleichungen. *Math. Ann.* **46**, 167–178 (1895)
83. Sanz-Serna, J.M.: Runge–Kutta schemes for Hamiltonian systems, *BIT* **28**, 877–883 (1988)
84. Sanz-Serna, J.M. and Abia, L.: Order conditions for canonical RK schemes. *SIAM J Numer. Anal.* **28**, 1081–1096 (1991)
85. Sanz-Serna, J.M. and Calvo, M.P.: *Numerical Hamiltonian Problems*, Chapman & Hall, London (1994)
86. Söderlind, G., Jay L. and Calvo M.: Stiffness 1952–2012: Sixty years in search of a definition, *BIT Numer Math* **55**, 531–558 (2015)
87. Stoffer, D.: General linear methods: connection to one step methods and invariant curves. *Numer. Math.* **64**, 395–408 (1993)
88. Suzuki, M.: Fractal decomposition of exponential operators with applications to many-body theories and Monte Carlo simulations, *Phys. Lett. A* **146**, 319–323 (1990)
89. Underwood, R.G.: *An introduction to Hopf algebras*, Springer-Verlag, Berlin, (2011)
90. van der Houwen, P.J. and Sommeijer, B.P.: On the internal stability of explicit m -stage Runge–Kutta methods for large m -values. *Z. Angew. Math. Mech.* **60**, 479–485 (1980)
91. Verner, J. H.: Some Runge–Kutta formula pairs: *SIAM J. Numer. Anal.* **28**, 496–511 (1991)
92. Yoshida, H.: Construction of higher order symplectic integrators, *Phys. Lett. A* **150**, 262–268 (1990)

Index

- \mathcal{A} (generating function), 54
- algebra
 - co-, 82
 - Hopf, 44, 88
 - linear, 95, 97
- α (combinatorial), 60, 61
- antipode, 43, 88, 91, 145
 - B^+ recursion, 93
 - involution property, 88, 91
- arborescence, 2
- atom, 77
- autonomous, 4, 24
- \mathcal{B} (generating function), 54
- B, B^*, B^0 , 113
- B-group = Butcher group, 165
- B^+ , 44–47, 50, 59, 86
- B-series, 3, 33, 41, 65, 100, 110, 113, 125
 - central, 134
 - composition, 100, 133
 - fractional powers, 147
 - inverse, 145
- balanced parentheses, 45
- $\underline{\beta}$ (combinatorial), 60, 61
- $\bar{\beta}$ (combinatorial), 60, 61
- beta-product, 45, 47, 84, 85
 - iterated, 46
 - recursion, 92
- BNF (formal languages), 45
- \mathcal{C} (generating function), 54
- combinatorics, 39
- composition, 43, 59, 72
- conformability, 259
 - weak, 259
- conjugacy test, 241
- conjugate order, 219
- conservation of angular momentum, 13, 247
- conservation of energy, 13, 247
- consistency, 29, 32
- convergence, 29, 32
- covariance, 101
- $D \in B^0$, 116
- Dahlquist barrier, 216
 - second, 217
- Δ (Sweedler notation), 82
- DETEST, 12
- differential equation, 1, 4
 - linear, 8
- distributive, 54
- $E \in B$, 115
- edge, 39
- effective order, 219
- elementary differential, 35, 41, 100, 103, 110, 112–114, 127, 131
 - attainable value, 129
 - perturbed, 137
- elementary weight, 100, 124, 126
 - attainable value, 129
- energy-preserving, 25
- equivalence class, 48
- evolution, 65
- evolve, 70
- exponential growth and decay, 8
- exponential integrator, 79
- flow, 1
- forest, 39, 41, 50, 75
 - antipode, 90
 - labelled, 75
 - space, 53
 - span $\mathbf{I} \cdot \mathbf{I}$, 52

- universal, 76
- formal languages, 45
- formal Taylor series, 34
- Fréchet derivative, 3, 34, 100, 106
- generating function, 54
- ∇ (gradient), 3, 4
- graph, 2, 39
 - connected, 39, 40
 - loop in, 39
 - order, 40
 - path in, 39
- graph theory, 39
- group, 58
 - abelian, 54
- Hamiltonian function, 250
- Heaviside function, 24
- index set, 24
- initial value, 5
- invariant subspace, 240
- involution, 91
- isomer, 77
- J (operator on B^0), 175
- Jacobian matrix, 3
- Kronecker product, 100, 127
- λ , 141, 142
- Λ , 95, 145
- Lipschitz
 - condition, 6
 - local, 7
 - constant, 6
- loop, 39
- mapping
 - central, 102, 113, 134
 - composition, 133
 - differential, 102
 - Euler*, 105, 115
 - flow*, 105, 111, 115, 117, 122, 123
 - flow-slope*, 105, 111
 - id*, 105, 115
 - implicit*, 105, 110, 111, 117, 121, 123, 124
 - mid-point*, 105
 - Runge-I*, 105, 111
 - Runge-II*, 105, 111
 - slope*, 105, 115, 116
- Merson, 74
- method
 - Adams–Bashforth, 29, 30, 211, 224
 - Adams–Moulton, 29, 211
 - adjoint, 23
 - Almost Runge–Kutta, 227
 - average vector field, 25, 153, 248, 285
 - backward difference, 211
 - backward difference representation, 230
 - continuous Runge–Kutta, 152
 - cyclic composite, 211, 218, 226
 - energy preserving, 281
 - Euler, 3, 14, 30, 33
 - explicit, 7
 - implicit, 7, 22
 - G-symplectic, 248, 262, 263, 266, 270
 - G4123, 266, 277, 278
 - G6245, 272, 276, 277, 279
 - Gauss, 23
 - general linear, 31, 212
 - algebraically stable, 263
 - conformability, 264
 - consistency, 222
 - convergence, 223
 - finishing, 232
 - order algorithm, 239
 - pre-consistency, 222
 - reducibility, 152
 - stability, 222
 - starting, 232
 - time-reversal symmetric, 271
 - weak conformability, 265
 - general linear method
 - order, 231
 - implicit mid-point, 22
 - integration, 151, 152
 - A-equivalent stages, 159
 - composition, 163
 - elementary weight, 154, 163
 - ϕ -equivalent stages, 159
 - reduced, 158
 - inverse, 23
 - inverting, 161
 - linear multistep, 28, 213, 217, 224
 - A-stability, 215
 - consistency, 213
 - convergence, 214
 - formulation, 220
 - order, 214
 - stability, 214, 215
 - multivalued, 28
 - Nordsieck representation, 230
 - numerical, 1
 - accuracy of, 2
 - off-step, 217, 226
 - off-step predictor, 211
 - one-leg, 216

- Picard, 153
- predictor-corrector, 211
- pseudo Runge–Kutta, 212
- re-use, 212, 219, 225
- Rosenbrock, 79
- Runge–Kutta, 1, 19, 75, 132, 148, 154, 170, 172, 224, 235
 - Hammer and Hollingsworth, 202
 - ambiguous order, 183
 - block diagonally implicit, 258
 - canonical, 247, 252, 259
 - Chebyshev, 190
 - composition, 152, 160
 - continuous stage, 25
 - cyclic composite, 220
 - diagonally implicit, 257
 - effective order, 205, 219
 - elementary weight, 170
 - equivalence, 25, 155
 - equivalent, 25, 26
 - explicit, 189
 - Gauss, 202, 203, 256
 - Gill, 195
 - Hairer, 199
 - high order, 197
 - Hu’a, 186
 - implementable, 203
 - implicit, 22, 201
 - Kutta’s classification, 195
 - low order, 190
 - order, 21, 78, 99, 124, 177
 - order barrier, 199
 - order conditions (symplectic), 254
 - P-equivalence, 156
 - φ -equivalence, 156
 - pre-reduced, 157
 - processing, 247
 - Radau IIA, 27, 202, 203
 - reduced, 158
 - reduced tableau, 193
 - reducibility, 151
 - reducible, 155
 - scalar, 178
 - simplifying assumptions, 197
 - singly-implicit, 202
 - stability, 177, 187
 - stability function, 187
 - stage-order, 172
 - symplectic, 41, 247, 252
 - tableau, 21
 - transformed, 204
- starting, 213, 235, 238
- Taylor series, 14, 18, 33
- theta, 16, 22, 34, 35
 - transformation, 228
 - underlying one-step, 213, 240
- moments of inertia, 250
- monoid, 41
 - commutative, 54
- multi-dimensional, 107, 108
- multinomial coefficient, 69
- multiplicative function, 94
- Newton iteration, 23, 145
- $1 + O_{p+1}$ = order-defining subgroup, 152, 166
- operator
 - bilinear, 103, 106, 108
 - linear, 106, 107
- order with processing, 258
- parasitism, 264
- partition
 - evolution, 70
 - of number, 65
 - of set, 65
- Picard–Lindelöf theorem, 153
- Polish notation, 46, 82, 106, 108
- polynomial
 - Laguerre, 205
 - Legendre, 203
- preconsistency, 29, 32
- problem, 100, 101
 - diamond, 12, 17
 - Euler rigid body, 250
 - Hénon–Heiles, 277
 - Hamiltonian, 248, 250
 - harmonic oscillator, 9
 - initial value, 1, 5, 99
 - autonomous, 110
 - Kepler, 13, 16, 26
 - Poisson, 8, 250
 - Prothero–Robinson, 12
 - Runge, 11
 - scalar, 5
 - simple pendulum, 9, 277
 - stiff, 10
 - symplectic, 252
 - variational, 252
 - well-posed, 6
- pruning, 82, 84, 94
- p-weight, 68, 69
- quadratic identity, 253
- quadratic invariant, 248
- question, 100, 101
 - easy set, 105

- recursion, 44, 84
- ρ (polynomial), 28, 213
- ring, 54
 - commutative, 54
- root condition, 29
- semi-group, 41
- σ (polynomial), 28, 213
- σ (symmetry), 58
- stability, 29, 32
- stability interval, 188
- stability region, 188
- stage, 19, 22, 24, 25, 28, 31
- stepsize, 19
- Stone–Weierstrass theorem, 129
- stump, 76
 - atomic, 77
 - product, 77
 - sub-atomic, 80
 - uni-valent, 79
- subtree, 84, 94
- supertree, 84
- Sweedler notation, 82, 95, 139
- symbolic matrix, 95
 - antipode, 96
- symbolic vector, 95
- symplectic flow, 252
- Taylor series, 2, 21, 100, 103, 107–112
- trapezoidal rule, 20
- tree, 2, 35, 39, 44, 50
 - antipode, 88
 - automorphism, 58
 - Cayley, 42
 - density, 59
 - empty, 40, 75
 - enumeration, 54
 - evolution, 72
 - $t!$ (factorial of t), 59
 - free, 40, 44
 - height, 52
 - isomeric, 77, 183
 - kinship, 42
 - labelled, 81
 - leaf, 52
 - N -, 48
 - offcut, 80, 81
 - $|t|$ (order of t), 2, 44
 - partition, 50
 - order, 50
 - pruning, 43, 80
 - root, 40, 52
 - rooted, 2, 40, 41, 44
 - S -, 48
 - space, 53
 - standard numbering, 61
 - sub, 43, 80, 87
 - subtree, 43
 - super, 43, 80, 87
 - superfluous, 48
 - supertree, 43
 - symmetry, 2, 58, 113
 - τ , 44
 - unrooted, 40, 41, 44, 48
 - width, 52
- truncation error, 21
- tuple, 75
- variable stepsize, 15
- vertex, 39
 - ancestor, 52
 - child, 52
 - dependant, 52
 - descendant, 52
 - parent, 52
 - valency, 76