

Solution Manual to *Linear Algebra Done Right*, 4th Edition by Sheldon Axler

Chenyu Tang



CONTENTS

Contents	3
1 Vector Spaces	5
1 \mathbb{R}^n and \mathbb{C}^n	5
2 Definition of Vector Space	5
3 Subspaces	7

CHAPTER 1

VECTOR SPACES

1 \mathbb{R}^n and \mathbb{C}^n

We skip this section.

2 Definition of Vector Space

Exercise 2.1: (1B-1)

Prove that $-(-v) = v$ for every $v \in V$.

Solution:

For $v \in V$, we have

$$-(-v) = -(-v) + (-v) + v = v. \quad (1.1)$$

Thus we know the additive inverse of the additive inverse of v is itself. ■

Exercise 2.2: (1B-2)

Suppose $a \in \mathbf{F}$, $v \in V$, and $av = 0$. Prove that $a = 0$ or $v = 0$.

Solution:

If $a = 0$, then we are done.

If $a \neq 0$, then

$$v = \left(\frac{1}{a} \cdot a\right)v = \frac{1}{a}(av) = 0. \quad (1.2)$$

■

Exercise 2.3: (1B-3)

Suppose $v, w \in V$. Explain why there exists a unique $x \in V$ such that $v + 3x = w$.

Solution:

Let $x = \frac{w-v}{3}$, then

$$v + 3x = w. \quad (1.3)$$

This show existence. Now we show the uniqueness. Suppose there is an x' that satisfies $v + 3x' = w$, then

$$3(x - x') = 3x - 3x' = (w - v) - (w - v) = 0. \quad (1.4)$$

By Exercise 2.2, we must have $x - x' = 0$, thus $x = x'$. ■

Exercise 2.4: (1B-4)

The empty set is not a vector space. The empty set fails to satisfy only one of the requirements listed in the definition of a vector space (1.20). Which one?

Solution:

Additive identity. In an empty set \emptyset , there does not exist an element 0 that $v + 0 = v$ for all $v \in \emptyset$. ■

Exercise 2.4 shows that the additive identity condition can be replaced with the condition that the set is not empty (because then taking $u \in U$ and multiplying it by 0 would imply that $0 \in U$).

Exercise 2.5: (1B-6)

Let ∞ and $-\infty$ denote two distinct objects, neither of which is in \mathbf{R} . Define an addition and scalar multiplication on $\mathbf{R} \cup \{\infty, -\infty\}$ as you could guess from the notation. Specifically, the sum and product of two real numbers is as usual, and for $t \in \mathbf{R}$ define

$$t\infty = \begin{cases} -\infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ \infty & \text{if } t > 0, \end{cases} \quad t(-\infty) = \begin{cases} \infty & \text{if } t < 0, \\ 0 & \text{if } t = 0, \\ -\infty & \text{if } t > 0, \end{cases}$$

and

$$\begin{aligned} t + \infty &= \infty + t = \infty + \infty = \infty, \\ t + (-\infty) &= (-\infty) + t = (-\infty) + (-\infty) = -\infty, \\ \infty + (-\infty) &= (-\infty) + \infty = 0. \end{aligned}$$

With these operations of addition and scalar multiplication, is $\mathbf{R} \cup \{\infty, -\infty\}$ a vector space over \mathbf{R} ? Explain.

Solution:

We can notice that

$$\infty = (2 + (-1))\infty = 2\infty + (-1)\infty = \infty + (-\infty) = 0. \quad (1.5)$$

For $\infty \neq 0$, the set doesn't follow the distributive property. Thus $\mathbf{R} \cup \{\infty, -\infty\}$ is not a vector space. ■

Exercise 2.6: (1B-8)

Suppose V is a real vector space.

- The *complexification* of V , denoted by $V_{\mathbf{C}}$, equals $V \times V$. An element of $V_{\mathbf{C}}$ is an ordered pair (u, v) , where $u, v \in V$, but we write this as $u + iv$.
- Addition on $V_{\mathbf{C}}$ is defined by

$$(u_1 + iv_1) + (u_2 + iv_2) = (u_1 + u_2) + i(v_1 + v_2)$$

for all $u_1, v_1, u_2, v_2 \in V$.

- Complex scalar multiplication on $V_{\mathbf{C}}$ is defined by

$$(a + bi)(u + iv) = (au - bv) + i(av + bu)$$

for all $a, b \in \mathbf{R}$ and all $u, v \in V$.

Prove that with the definitions of addition and scalar multiplication as above, $V_{\mathbf{C}}$ is a complex vector space.

Solution:

Just verify the six properties of vector spaces. For example:

commutativity

$$\begin{aligned} (u_1 + iv_1) + (u_2 + iv_2) &= (u_1 + u_2) + i(v_1 + v_2) \\ &= (u_2 + u_1) + i(v_2 + v_1) \\ &= (u_2 + iv_2) + (u_1 + iv_1) \end{aligned} \quad (1.6)$$

for all $u_1, u_2, v_1, v_2 \in V$. The remaining five properties are the same. Thus we have the complex vector space $V_{\mathbf{C}}$. ■

3 Subspaces

Exercise 3.1: (1C-5)

Is \mathbf{R}^2 a subspace of the complex vector space \mathbf{C}^2 ?

Solution:

Notice that subspaces of \mathbf{C}^2 are closed under scalar multiplication in \mathbf{C} , then

$$i(1, 1) = (i, i) \notin \mathbf{R}^2.$$

Thus \mathbf{R}^2 is not a subspace of \mathbf{C}^2 . ■

Exercise 3.2: (1C-6)

(a) Is $\{(a, b, c) \in \mathbf{R}^3 : a^3 = b^3\}$ a subspace of \mathbf{R}^3 ?

(b) Is $\{(a, b, c) \in \mathbf{C}^3 : a^3 = b^3\}$ a subspace of \mathbf{C}^3 ?

Solution:

(a)

The equation $a^3 = b^3$ has the only solution $a = b$ in \mathbf{R} , hence

$$\{(a, b, c) \in \mathbf{R}^3 : a^3 = b^3\} = \{(a, b, c) \in \mathbf{R}^3 : a = b\} \quad (1.7)$$

is obviously a subspace of \mathbf{R}^3 .

(b)

In \mathbf{C}^3 , we have

$$\left(1, \frac{-1 + \sqrt{3}i}{2}, 0\right) \in \{(a, b, c) \in \mathbf{C}^3 : a^3 = b^3\} \quad (1.8)$$

and

$$\left(1, \frac{-1 - \sqrt{3}i}{2}, 0\right) \in \{(a, b, c) \in \mathbf{C}^3 : a^3 = b^3\}. \quad (1.9)$$

However,

$$\left(1, \frac{-1 + \sqrt{3}i}{2}, 0\right) + \left(1, \frac{-1 - \sqrt{3}i}{2}, 0\right) = (2, -1, 0) \notin \{(a, b, c) \in \mathbf{C}^3 : a^3 = b^3\}.$$

Hence $\{(a, b, c) \in \mathbf{C}^3 : a^3 = b^3\}$ is not a vector space. ■

Exercise 3.3: (1C-11)

Prove that the intersection of every collection of subspaces of V is a subspace of V .

Solution:

Assume U_i is a subspace of V for $i \in I$, then prove $\cap_{i \in I} U_i$ is a subspace of V .

Additive identity

$$0 \in U_i \quad \forall i \in I \Rightarrow 0 \in \cap_{i \in I} U_i. \quad (1.10)$$

Closed under addition For all $u, v \in \cap_{i \in I} U_i$, $u, v \in U_i$, $\forall i \in I$, thus

$$u + v \in U_i, \forall i \in I \Rightarrow u + v \in \cap_{i \in I} U_i \quad (1.11)$$

which shows $\cap_{i \in I} U_i$ is closed under addition.

Closed under scalar multiplication For all $w \in \cap_{i \in I} U_i$, $w \in U_i$, $\forall i \in I$, thus

$$\lambda w \in U_i, \forall i \in I \Rightarrow \lambda w \in \cap_{i \in I} U_i \quad (1.12)$$

which shows $\cap_{i \in I} U_i$ is closed under scalar multiplication. ■

Exercise 3.4: (1C-12)

Prove that the union of two subspaces of V is a subspace of V if and only if one of the subspaces is contained in the other.

Solution:

Let U_1, U_2 be subspaces of V . Suppose $u \in U_1$ and $u \notin U_2$ while $v \in U_2$ and $v \notin U_1$, then we have $u, v \in U_1 \cup U_2$ but $u + v \notin U_1$ or U_2 , thus not in $U_1 \cup U_2$. Hence $U_1 \cup U_2$ is not closed under addition, and we get a contradiction. So either $\forall u \in U_1, u \in U_2$ or $\forall v \in U_2, v \in U_1$, or $U_1 = U_2$. ■

Exercise 3.5: (1C-13)

Prove that the union of three subspaces of V is a subspace of V if and only if one of the subspaces contains the other two.

Solution:

Let U_1, U_2, U_3 be subspaces of V , consider $u \in U_1$ and $v \in U_2$. If $U \cup V \neq U$ or V , then we can assume there exist $u \notin U_2$ and $v \notin U_1$. From the assumption we know $u + v \in \cup_{i=1}^3 U_i$ and $u + v$ not in U_1 or U_2 , thus $u + v \in U_3$. For the same reason we know $u + 2v$ and $2u + v$ are in U_3 , and because U_3 is closed under addition and additive inverse, we have $u, v \in U_3$. Hence $\mathbb{C}_{U_1} U_1 \cap U_2$ and $\mathbb{C}_{U_2} U_1 \cap U_2$ are contained by U_3 .

For $w \in U_1 \cap U_2$, taking $u \in \mathbb{C}_{U_1} U_1 \cap U_2$, then $u + w$ in U_1 but not in U_2 , and we come back to the above condition. Thus we can prove $U_1 \cap U_2 \subset U_3$, which means U_3 contains U_1 and U_2 .

Now consider the condition that $U_1 \subset U_2$. If $U_2 \subset U_3$, then we are done. If $U_1 \subset U_3$, then consider $u \in \mathbb{C}_{U_2} U_1$ and $v \in \mathbb{C}_{U_3} U_1$, because the union of U_1, U_2, U_3 is closed under addition, $u + v \in \cup_{i=1}^3 U_i$. Evidently $u + v \notin U_1$, if $u + v \in U_2$, then we can prove $U_2 = U_3$; the other side is the same.

From the conditions above, we prove the proposition in the question. ■

Remark.

Problem 15-19 in Exercises 1C prove some properties of the addition in the set of subspaces of V , including additive identity, commutativity, associativity. From these propositions we know the group of subspaces of V is not an abel group.

Exercise 3.6: (1C-21)

Suppose

$$U = \{(x, y, x + y, x - y, 2x) \in \mathbf{F}^5 : x, y \in \mathbf{F}\}.$$

Find a subspace W of \mathbf{F}^5 such that $\mathbf{F}^5 = U \oplus W$.

Solution:

Let $W = \{(0, 0, u, v, w) \in \mathbf{F}^5 : u, v, w \in \mathbf{F}\}$. Consider $(x, y, x + y, x - y, 2x) \in U \cap W$, then we have $x = y = 0$, which means $U \cap W = \{0\}$. Hence $\mathbf{F}^5 = U \oplus W$ by 1.45. ■

Problem 20-22 in Exercises 1C are the same type. To solve these problems, we first write a subspace W that the union of W and the given subspaces is the vector space asked for. Then prove the intersection of the subspaces is $\{0\}$.

Exercise 3.7: (1C-23)

Prove or give a counterexample: If V_1, V_2, U are subspaces of V such that

$$V = V_1 \oplus U \quad \text{and} \quad V = V_2 \oplus U,$$

then $V_1 = V_2$.

Solution:

Consider the example in Exercise 3.6. We change W into $W' = \{(0, z, 0, v, w) \in \mathbf{F}^5 : u, v, w \in \mathbf{F}\}$, and it is easy to verify $\mathbf{F}^5 = U \oplus W'$. Hence we give a counterexample. ■

Exercise 3.8: (1C-24)

A function $f : \mathbf{R} \rightarrow \mathbf{R}$ is called *even* if

$$f(-x) = f(x)$$

for all $x \in \mathbf{R}$. A function $f : \mathbf{R} \rightarrow \mathbf{R}$ is called *odd* if

$$f(-x) = -f(x)$$

for all $x \in \mathbf{R}$. Let V_e denote the set of real-valued even functions on \mathbf{R} and let V_o denote the set of real-valued odd functions on \mathbf{R} . Show that $\mathbf{R}^{\mathbf{R}} = V_e \oplus V_o$.

Solution:

For $h : \mathbf{R} \rightarrow \mathbf{R} \in \mathbf{R}^{\mathbf{R}}$, suppose $h(x) = f(x) + g(x)$, where $f \in V_e$ and $g \in V_o$. Then we must verify $h(-x) = f(-x) + g(-x) = f(x) - g(x)$. From the above discription, let

$$f(x) = \frac{h(x) + h(-x)}{2} \quad (1.13)$$

and

$$g(x) = \frac{h(x) - h(-x)}{2}. \quad (1.14)$$

Then we have $\mathbf{R}^{\mathbf{R}} = V_e + V_o$.

Consider $f_0 \in V_e \cap V_o$,

$$f_0(x) = f_0(-x) = -f_0(x), \quad \forall x \in \mathbf{R}. \quad (1.15)$$

Thus $f_0(x) \equiv 0$, $V_e \cap V_o = \{0\}$. By 1.45, $\mathbf{R}^{\mathbf{R}} = V_e \oplus V_o$. ■

From Exercise 3.8 we know every real-valued function can be divided into the sum of an even function and an odd function.