

The actual and probably very long title of the thesis

Author

Advisors: Duh, Dih, Dah

The date


- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets



And an appearing figure.

- Some
  - Appearing
  - And disappearing
  - Bullets
    - With sub-bullets
  - That appear and disappear with their parent

- Some
- **Appearing**
- And disappearing
- Bullets
  - With sub-bullets
- That appear and disappear with their parent



- Some
- Appearing
- **And disappearing**
- Bullets
  - With sub-bullets
- That appear and disappear with their parent

- Some
- Appearing
- And disappearing
- **Bullets**
  - With sub-bullets
- That appear and disappear with their parent

- Some
- Appearing
- And disappearing
- Bullets
  - With sub-bullets
- That appear and disappear with their parent

- ❶ Some
- ❷ Numbers
  - With sub-bullets
- ❸ That appear and the same time
- ❹ Nicely spaced on the slide

- Do not put all references at the end, it is impossible to remember
- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*
- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*
- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)
- Can also be put in footline <sup>1</sup>

---

<sup>1</sup> O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Do not put all references at the end, it is impossible to remember
- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)
- Can also be put in footline <sup>1</sup>

---

<sup>1</sup> O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*

- Do not put all references at the end, it is impossible to remember
- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)
- Can also be put in footnote <sup>1</sup>

---

<sup>1</sup> O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*

- Do not put all references at the end, it is impossible to remember
- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)
- Can also be put in footnote <sup>1</sup>

---

<sup>1</sup> O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*



- Do not put all references at the end, it is impossible to remember
- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*
- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)
- Can also be put in footline <sup>1</sup>

---

<sup>1</sup> O. S. Pythagoras (Feb. -580). “Theorem”. In: *Some old journal*

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

- $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

- $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references
- Can also be included in the text / bullets
  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references
- Can also be included in the text / bullets

- $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references
- Can also be included in the text / bullets
  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with `\begin{equation}`

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \quad (1)$$

- Do not have a number if used with `\begin{equation*}`

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply `\begin{center}`

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

- Probably more suited to slides as we use less equation references
- Can also be included in the text / bullets
  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$