The actual and probably very long title of the thesis

Author

Advisors: Duh, Dih, Dah

The date

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
    - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

- Some
- Appearing
- Bullets
  - With sub-bullets

And an appearing figure.

- Some
- Appearing
- And disappearing
- Bullets
  - With sub-bullets
- That appear and disappear with their parent

- Some

- **Appearing**
- And disappearing
- Bullets
    - With sub-bullets

- That appear and disappear with their parent

- Some
- Appearing
- **And disappearing**
- Bullets
  - With sub-bullets
- That appear and disappear with their parent

- Some
- Appearing
- And disappearing
- Bullets
  - With sub-bullets
- That appear and disappear with their parent

- Some
- Appearing
- And disappearing
- Bullets
    - With sub-bullets
- That appear and disappear with their parent

1. Some

2. Numbers
   - With sub-bullets

3. That appear and the same time

4. Nicely spaced on the slide

- Do not put all references at the end, it is impossible to remember

- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)

- Can also be put in footline [1]

---

[1] O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Do not put all references at the end, it is impossible to remember

- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)

- Can also be put in footline [1]

---

[1] O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Do not put all references at the end, it is impossible to remember

- Should be done with \fullcite
    - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- You may also use \smallcite
    - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Check all imported references for:
    - Name / Journal name / year / editor (journals)
    - Carefully check conference name (IEEExplore)

- Can also be put in footline [1]

---

[1] O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Do not put all references at the end, it is impossible to remember

- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)

- Can also be put in footline [1]

---
[1] O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Do not put all references at the end, it is impossible to remember

- Should be done with \fullcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- You may also use \smallcite
  - O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Check all imported references for:
  - Name / Journal name / year / editor (journals)
  - Carefully check conference name (IEEEexplore)

- Can also be put in footline [1]

---

[1] O. S. Pythagoras (Feb. -580). "Theorem". In: *Some old journal*

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets

  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$

- Have a number if used with \begin{equation}

$$\forall \phi : \quad \cos^2 \phi + \sin^2 \phi = 1 \tag{1}$$

- Do not have a number if used with \begin{equation*}

$$\forall a, b : \quad (a + b)^2 = a^2 + 2ab + b^2$$

- Another useful environment is simply \begin{center}

$$\forall a, b : \quad (a - b)^2 = a^2 - 2ab + b^2$$

  - Probably more suited to slides as we use less equation references

- Can also be included in the text / bullets
  - $\forall \phi : \quad (\cos \phi + \sin \phi)^2 = 2 \cos \phi \sin \phi + 1$