

Assignment 3: Data Exploration

Cal Oakley, Section #2

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

Directions

1. Change “Student Name, Section #” on line 3 (above) with your name and section number.
2. Work through the steps, **creating code and output** that fulfill each instruction.
3. Be sure to **answer the questions** in this assignment document.
4. When you have completed the assignment, **Knit** the text and code into a single PDF file.
5. After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai. Add your last name into the file name (e.g., “FirstLast_A03_DataExploration.Rmd”) prior to submission.

The completed exercise is due on <>.

Set up your R session

1. Check your working directory, load necessary packages (tidyverse), and upload two datasets: the ECOTOX neonicotinoid dataset (ECOTOX_Neonicotinoids_Insects_raw.csv) and the Niwot Ridge NEON dataset for litter and woody debris (NEON_NIWO_Litter_massdata_2018-08_raw.csv). Name these datasets “Neonics” and “Litter”, respectively. **Be sure to add the stringsAsFactors = TRUE parameter to the function when reading in the CSV files.**

```
setwd("K:/GradSchool/Spring2022/EnvironmentalDataAnalytics/Environmental_Data_Analytics_2022/Data/Raw")  
#setting WD to raw data folder for this chunk, so I can load csv's easily  
library(tidyverse) #loads tidyverse package  
Neonics <- read.csv("ECOTOX_Neonicotinoids_Insects_raw.csv",  
                   stringsAsFactors = TRUE)  
#reads "ECOTOX_Neonicotinoids_Insects_raw.csv" into working environment  
Litter <- read.csv("NEON_NIWO_Litter_massdata_2018-08_raw.csv",  
                  stringsAsFactors = TRUE)  
#reads "NEON_NIWO_Litter_massdata_2018-08_raw.csv" into working environment
```

Learn about your system

2. The neonicotinoid dataset was collected from the Environmental Protection Agency’s ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: Neonicotinoids are manufactured chemicals that are similar to the natural compound nicotine. They are generally considered to be more harmful to insects than mammals. Since they are often used to control insects that attack crops, they tend to be broadcast into the environment in a manner that could expose non-target insect species which has implications for

insect populations beyond the agro-ecosystem, especially if the chemicals are persist for extended periods.

3. The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: Sampling forest litter and woody debris is important for a variety of reasons. It can be used to study nutrient cycling within forest ecosystems, assess wildfire hazard potential, and estimate carbon storage in soil.

4. How is litter and woody debris sampled as part of the NEON network? Read the NEON_Litterfall_UserGuide.pdf document to learn more. List three pieces of salient information about the sampling methods here:

Answer: It is sampled using elevated and ground traps to intercept litter and woody debris on its way from the canopy to the forest floor. Sampling are taken at terrestrial NEON sites with woody vegetation greater than 2 meters in height.

Different plant organs (leaves, needles, twigs/branches, cones, flowers, seeds, fruit, bark, etc.) are weighed separately to determine the mass of each group. Elevated and ground traps are different sizes and are designed to capture different types of plant organs. *All samples are oven dried before being weighed by technicians.

Obtain basic summaries of your data (Neonics)

5. What are the dimensions of the dataset?

```
dim(Neonics) #returns the dimensions of 'Neonics'
```

```
## [1] 4623 30
```

6. Using the summary function on the “Effect” column, determine the most common effects that are studied. Why might these effects specifically be of interest?

```
summary(Neonics$Effect) #returns the different unique values in 'Effect' column
```

##	Accumulation	Avoidance	Behavior	Biochemistry
##	12	102	360	11
##	Cell(s)	Development	Enzyme(s)	Feeding behavior
##	9	136	62	255
##	Genetics	Growth	Histology	Hormone(s)
##	82	38	5	1
##	Immunological	Intoxication	Morphology	Mortality
##	16	12	22	1493
##	Physiology	Population	Reproduction	
##	7	1803	197	

```
#and counts of each
```

Answer: From looking at these data, I think the most common effects are ‘Population’, ‘Mortality’, ‘Behavior’, ‘Feeding behavior’, ‘Reproduction’, and ‘Development’. These effects could be of interest because, as is evident from the names alone, they all relate to population dynamics. Some very directly, like ‘Population’, ‘Mortality’, & ‘Reproduction’; and other more indirectly, like ‘Behavior’, ‘Feeding behavior’, & ‘Development’. These can all be used to observe the scale and degree to which pesticide spillover from agriculture effects non-target insect species populations (the ‘direct’ effects) and the casual pathways involved (the ‘indirect’ effects).

7. Using the `summary` function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

```
summary(Neonics$Species.Common.Name) #does the same as previous chunk, but for
```

```
##              Honey Bee              Parasitic Wasp
##              667              285
##      Buff Tailed Bumblebee      Carniolan Honey Bee
##              183              152
##              Bumble Bee              Italian Honeybee
##              140              113
##      Japanese Beetle              Asian Lady Beetle
##              94              76
##      Euonymus Scale              Wireworm
##              75              69
##      European Dark Bee              Minute Pirate Bug
##              66              62
##      Asian Citrus Psyllid              Parastic Wasp
##              60              58
##      Colorado Potato Beetle              Parasitoid Wasp
##              57              51
##      Erythrina Gall Wasp              Beetle Order
##              49              47
##      Snout Beetle Family, Weevil              Sevenspotted Lady Beetle
##              47              46
##      True Bug Order              Buff-tailed Bumblebee
##              45              39
##      Aphid Family              Cabbage Looper
##              38              38
##      Sweetpotato Whitefly              Braconid Wasp
##              37              33
##      Cotton Aphid              Predatory Mite
##              33              33
##      Ladybird Beetle Family              Parasitoid
##              30              30
##      Scarab Beetle              Spring Tiphia
##              29              29
##      Thrip Order              Ground Beetle Family
##              29              27
##      Rove Beetle Family              Tobacco Aphid
##              27              27
##      Chalcid Wasp              Convergent Lady Beetle
##              25              25
##      Stingless Bee              Spider/Mite Class
##              25              24
##      Tobacco Flea Beetle              Citrus Leafminer
##              24              23
##      Ladybird Beetle              Mason Bee
##              23              22
##      Mosquito              Argentine Ant
##              22              21
##      Beetle              Flatheaded Appletree Borer
##              21              20
##      Horned Oak Gall Wasp              Leaf Beetle Family
```

##	20	20
##	Potato Leafhopper	Tooth-necked Fungus Beetle
##	20	20
##	Codling Moth	Black-spotted Lady Beetle
##	19	18
##	Calico Scale	Fairyfly Parasitoid
##	18	18
##	Lady Beetle	Minute Parasitic Wasps
##	18	18
##	Mirid Bug	Mulberry Pyralid
##	18	18
##	Silkworm	Vedalia Beetle
##	18	18
##	Araneoid Spider Order	Bee Order
##	17	17
##	Egg Parasitoid	Insect Class
##	17	17
##	Moth And Butterfly Order	Oystershell Scale Parasitoid
##	17	17
##	Hemlock Woolly Adelgid Lady Beetle	Hemlock Woolly Adelgid
##	16	16
##	Mite	Onion Thrip
##	16	16
##	Western Flower Thrips	Corn Earworm
##	15	14
##	Green Peach Aphid	House Fly
##	14	14
##	Ox Beetle	Red Scale Parasite
##	14	14
##	Spined Soldier Bug	Armoured Scale Family
##	14	13
##	Diamondback Moth	Eulophid Wasp
##	13	13
##	Monarch Butterfly	Predatory Bug
##	13	13
##	Yellow Fever Mosquito	Braconid Parasitoid
##	13	12
##	Common Thrip	Eastern Subterranean Termite
##	12	12
##	Jassid	Mite Order
##	12	12
##	Pea Aphid	Pond Wolf Spider
##	12	12
##	Spotless Ladybird Beetle	Glasshouse Potato Wasp
##	11	10
##	Lacewing	Southern House Mosquito
##	10	10
##	Two Spotted Lady Beetle	Ant Family
##	10	9
##	Apple Maggot	(Other)
##	9	670

#the 'Species.Common.Name' column

Answer: The top six most commonly studied species are all bees (with the exception of 'Parasitic

Wasp’, which to me looks like a polyphyletic group; but still, close enough). These species are of particular interest relative to other insects because of their value in crop production (pollination -> plant reproduction -> fruit+veg). In agro-ecosystems there can be just as many desirable insects as there are undesirable ones. So, if a farmer is going to broadcast insecticides over their farm to control an undesirable species they need to be assured that it won’t have a negative impact on the species they need to generate enough of a harvest.

8. Concentrations are always a numeric value. What is the class of Conc.1..Author. in the dataset, and why is it not numeric?

```
class(Neonics$Conc.1..Author.) #returns class of 'Neonics' column 'Conc.1..Author.'
```


```
## [1] "factor"
```

Answer: ‘Conc.1..Author.’ is classed as “factor” in this dataset. I think what has happened here is that the values, although numeric, were classed as “string” in the raw CSV, so when I imported the dataset with the ‘stringsAsFactors = TRUE’ they were converted to “factor”.

Explore your data graphically (Neonics)

9. Using `geom_freqpoly`, generate a plot of the number of studies conducted by publication year.

```
library(ggplot2) #loads ggplot2 package for plotting
ggplot(Neonics) +
  geom_freqpoly(aes(x = Publication.Year))
```




```
#greats frequency plot for 'Publication.Year'
```

10. Reproduce the same graph but now add a color aesthetic so that different Test.Location are displayed

as different colors.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Publication.Year, color = Test.Location))
```




```
#greats frequency plot for 'Publication.Year', symbolized by color
```

Interpret this graph. What are the most common test locations, and do they differ over time?

Answer: The most common test locations are 'Lab' & 'Field natural'. The number of studies classified as 'Field natural' grew steadily between 1990 and the early 2000's and saw a peak around 2009-2010, but have since fallen. The number of studies classified as 'Lab' also began growing after 1990 albeit at a faster rate than 'Field natural'. 'Lab' studies peaked around 2015, but have also dropped off since then.

11. Create a bar graph of Endpoint counts. What are the two most common end points, and how are they defined? Consult the ECOTOX_CodeAppendix for more information.

```
ggplot(Neonics) +  
  geom_bar(aes(x = Endpoint, fill = Endpoint)) #creates bar chart displaying
```



#frequency of different ecotoxicology studies endpoints

Answer: The two most common endpoints are ‘NOEL’ and ‘LOEL’. According to ‘ECO-TOX_CodeAppendix’, ‘NOEL’ represents “no-observable-effect-level” which means the study reported the highest dose at which the insecticide in question produced no effects that differed significantly from control results. In other words, these studies reported a safe dose limit for the insecticide they were studying. ‘LOEL’ represents “Lowest Observed Effects Residue” which means the study reported the smallest dose at which observers found a response that differed significantly from the control.

Explore your data (Litter)

- Determine the class of collectDate. Is it a date? If not, change to a date and confirm the new class of the variable. Using the `unique` function, determine which dates litter was sampled in August 2018.

```
# class(Litter$collectDate) #checking class of 'collectDate' column
library(lubridate) #load package lubridate to force class of 'collectDate'
```

```
## Warning: package 'lubridate' was built under R version 4.1.2
```

```
##
```

```
## Attaching package: 'lubridate'
```

```
## The following objects are masked from 'package:base':
```

```
##
```

```
## date, intersect, setdiff, union
```

```
#from factor to date
```

```
Litter$collectDate <- ymd(Litter$collectDate) #force class of 'collectDate'
```

```
#from factor to date
unique(Litter$collectDate) #check which dates litter were sampled
```

```
## [1] "2018-08-02" "2018-08-30"
```

13. Using the `unique` function, determine how many plots were sampled at Niwot Ridge. How is the information obtained from `unique` different from that obtained from `summary`?

```
unique(Litter$plotID) #returns each unique value in 'plotID' column, last line
```


```
## [1] NIWO_061 NIWO_064 NIWO_067 NIWO_040 NIWO_041 NIWO_063 NIWO_047 NIWO_051
## [9] NIWO_058 NIWO_046 NIWO_062 NIWO_057
## 12 Levels: NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 ... NIWO_067
```

```
#of return reports the number of different values, 12 in this case
```

Answer: ‘summary’ differs from ‘unique’ in that ‘summary’ looks at a field and returns the count of each unique type of answer, while ‘unique’ just returns a count of those unique answer types.

14. Create a bar graph of `functionalGroup` counts. This shows you what type of litter is collected at the Niwot Ridge sites. Notice that litter types are fairly equally distributed across the Niwot Ridge sites.


```
ggplot(Litter) +
  geom_bar(aes(x = functionalGroup, fill = functionalGroup)) #creates a bar chart
```



```
#of how many samples there are of each litter type ('functionalGroup'),
#symbolized by color
```

15. Using `geom_boxplot` and `geom_violin`, create a boxplot and a violin plot of `dryMass` by `functionalGroup`.


```
ggplot(Litter) +
  geom_boxplot(aes(x = functionalGroup, y = dryMass, fill = functionalGroup))
```




```
#creates boxplot of the mass ('dryMass') of all the samples in each litter
#type
```

```
ggplot(Litter) +
  geom_violin(aes(x = functionalGroup, y = dryMass, fill = functionalGroup),
    draw_quantiles = c(0.25, 0.5, 0.75))#creates violin plot of
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```

```
## Warning in regularize.values(x, y, ties, missing(ties), na.rm = na.rm):
## collapsing to unique 'x' values
```



the mass ('dryMass') of all the samples in each litter type

Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer: In this case, a boxplot is a more effective visualization option, compared to the violin plot, because the distribution of 'dryMass' within each 'functionalGroup' differs too much between each 'functionalGroups'. For example, 'Needles' 'dryMass' ranges between 0.0 and 7.5, with samples appearing to occur in similar, low frequencies throughout said range. Meanwhile, 'Seeds' 'dryMass' ranges between 0.0 and something just above that, with all samples occurring there. By asking ggplot() to put even just those two 'functionalGroups' on the same figure obfuscates all patterns in the data.

What type(s) of litter tend to have the highest biomass at these sites?

Answer: At this site, 'Needles' and 'Mixed' litters tend to have the greatest biomass.