

Structure-Activity Study on Antiviral 5-Vinylpyrimidine Nucleoside Analogs Using Wiener's Topological Index

Seema Mendiratta and A. K. Madan*

College of Pharmacy, University of Delhi, Pushp Vihar, New Delhi 110 017, India

Received November 21, 1993*

The relationship between Wiener's topological index and the antiviral activity of a series of 5-vinylpyrimidine nucleoside analogs has been investigated. Values for more than 100 compounds were computed, and an active range was identified. The predicted activity of each compound was compared with reported antiviral activity against herpes simplex virus type I. Due to significant correlation between antiviral activity and Wiener's topological index, it was possible to predict antiviral activity with an accuracy of ~83%.

INTRODUCTION

The interest in the influence of molecular topology on molecular properties has grown remarkably during the past few years. The objective of all such studies is to explore the role of connectedness of atoms in the expression of biological activities of molecules.¹ Thus, molecular structures are translated into characteristic numerical descriptors known as topological indices, which may then be used in the development of structure-activity relationship (SAR) studies.² The use of topological indices in SAR seems to play an important role in situations where the biological activity is determined predominantly by topological architecture of molecular structure, i.e. where simple connectivity among neighboring atoms, without considering the chemical nature of atoms or the nature of chemical bonding, may be the major determinant of biological activity of a molecule.¹ Such an approach³⁻⁶ was initiated by H. Wiener in 1947 with the use of indices based on topological distances. The path number developed by Wiener, now known as Wiener's topological index, is equal to one-half the sum of elements of the distance matrix $D(G)$.

$$W(G) = \frac{1}{2} \sum_{ij} D(G)$$

where $D(G)$ represents off-diagonal elements of $D(G)$. The smaller the Wiener's number, the greater the compactness of molecules.²

Many indices have been devised.^{2,7-9} From them, Randic's molecular connectivity index,^{10,11} Hosoya's index,^{12,13} and Balaban's index J^{14-16} appear to be the most useful. Bonchev and Trinajstić devised information theoretical indices¹⁷⁻¹⁹ and proposed a superindex²⁰ consisting of a sum of 10 different topological indices. Some other indices based on adjacency and distance matrix include the Zagreb group index,^{21,22} the comparability index,²³ Platt's number,^{24,25} Gordon-Scantlebury's index,²⁶ Altenburg's polynomial,²⁷ Smolenskii's additivity index,²⁸ and the centric index.⁸ Various topological indices have different correlating abilities with physical, chemical, or biological properties of molecules.²⁹

In the present studies, an attempt has been made to correlate Wiener's index with the antiviral activity of 5-vinylpyrimidine nucleoside analogs. Antiviral chemotherapy is at the same stage today as antibacterial chemotherapy was half a century ago.³⁰ Although there are many compounds that have potent antiviral activity in cell cultures, only 10 synthetic compounds




Figure 1. 5-Vinyl-2-pyrimidones.

Table 1. List of R⁵ Substituents

base number	name
I	β-2'-deoxyribosyl
II	α-2'-deoxyribosyl
III	2'-fluoroarabinosyl
IV	arabinosyl
V	(2-hydroxyethoxy)methyl
VI	5'-chloro-2',5'-dideoxyribosyl
VII	5'-bromo-2',5'-dideoxyribosyl
VIII	5'-iodo-2',5'-dideoxyribosyl
IX	5'-azido-2',5'-dideoxyribosyl
X	5'-amino-2',5'-dideoxyribosyl
XI	2'-deoxyxylosyl
XII	3'-chloro-2',3'-dideoxyribosyl
XIII	3'-azido-2',3'-dideoxyribosyl
XIV	3'-amino-2',3'-dideoxyribosyl
XV	3'-azido-2',3'-dideoxyribosyl
XVI	3'-amino-2',3'-dideoxyxylosyl
XVII	4-hydroxybutyl

have been approved by FDA for clinical use.³¹ None of these drugs, however, are without toxicity, and hence there is a strong need for improved drugs not only to improve efficiency but also to circumvent these problems of toxicity. There is also a need to find an effective therapy for viral infections for which we do not at present have clinically useful drugs, an important infection being acquired immune deficiency syndrome (AIDS).³²⁻³⁴ The search for new antiviral drugs has been boosted by the advent of AIDS and identification of HIV as the etiological agent.³⁵

Consequently, the need for treatment of viral infection is enormous, as currently available drugs barely scratch the surface, but the development of antiviral compounds has been hampered by a number of problems,³³ the major ones being the following:

(a) Selectivity, or inhibition of virus infection and reproduction without causing adverse effects to host cells; that is, the drug must be designed rationally on exploitable biochemical differences that exist between virus-specific processes and cellular biosynthetic processes.

* Abstract published in *Advance ACS Abstracts*, April 1, 1994.

Table 2. Relationship between Wiener's Topological Index and Antiviral Activity of 5-Vinylpyrimidine Nucleoside Analogs and Congeners (Boxes Indicate the Substituents Replacing the 5-Vinyl Group)

compd no.	R ¹	R ²	R ³	R ⁴	R ⁵	Wiener's index	antiviral activity		references
							predicted	reported	
1	5-acetyl			OH	H	150	— ^a	—	45, 47
2	5-ethynyl			OH	H	116	—	—	45, 47
3	5-ethynyl			NH ₂	H	116	—	—	46, 47
4	H	H	H	OH	H	116	—	—	47
5	Cl	H	H	OH	H	150	—	—	47
6	H	Br	H	OH	H	158	—	—	47
7	5-acetyl			OH	I	683	+ ^b	+	47
8	5-ethynyl			OH	I	593	+	+	38, 48, 49
9	5-ethynyl			OH	IV	668	+	+	48
10	5-ethynyl			NH ₂	I	593	+	+	50
11	H	H	H	OH	I	593	+	+	38, 49, 51
12	H	H	H	NH ₂	I	593	+	+	51
13	H	H	H	OH	III	668	+	+	52
14	H	H	H	OH	IV	668	+	+	53–55
15	H	Br	H	OH	I	699	+	+	38, 49, 56
16	H	H	Br	OH	I	699	+	+	57
17	H	I	H	OH	I	699	+	+	38, 49, 56
18	H	Cl	H	OH	I	699	+	+	56
19	H	Br	H	NH ₂	I	699	+	+	50
20	H	I	H	NH ₂	I	699	+	+	50
21	H	H	F	OH	I	699	+	—	58
22	H	F	H	OH	I	699	+	+	58
23	H	Br	H	OH	IV	782	+	+	53, 55, 59, 60, 81
24	H	Br	H	OH	IV	782	+	+	52
25	H	I	H	OH	III	782	+	+	52
26	H	Cl	H	OH	III	782	+	+	52
27	H	Cl	H	OH	IV	782	+	+	53, 55, 81
28	H	Br	H	OH	V	478	—	—	47, 61
29	Cl	H	H	OH	I	683	+	—	38
30	Cl	Br	H	OH	I	802	+	+	62
31	H	COOCH ₃	H	OH	I	1096	—	+	63
32	H	Br	H	OH	VI	699	+	+	64
33	H	Br	H	OH	VII	699	+	+	64
34	H	Br	H	OH	VIII	699	+	+	64, 56
35	H	Br	H	OH	IX	963	± ^c	—	64
36	H	Br	H	OH	X	782	+	+	64
37	H	Br	H	OH	XI	782	+	—	64
38	H	Br	H	OH	XII	782	+	—	64
39	H	Br	H	OH	XIII	935	±	+	64, 56
40	H	Br	H	OH	XIV	782	+	+	64, 65
41	H	Br	H	OH	XV	935	±	+	64
42	H	Br	H	OH	XVI	782	+	+	64
43	H	CN	H	OH	I	824	+	+	63, 66
44	H	CN	H	OH	IV	916	±	—	38
45	H	CH ₃	H	OH	I	782	+	+	67–69
46	H	C ₂ H ₅	H	OH	I	824	+	+	69
47	H	CH=CH ₂	H	OH	I	824	+	+	70
48	H	SCH ₃	H	OH	I	824	+	+	63
49	H	H	CH ₃	OH	I	782	+	+	63
50	H	CH ₃	H	SH	I	782	+	+	63
51	H	CH ₃	H	NH ₂	I	782	+	+	63
52	H	H	CH ₃	NH ₂	I	782	+	+	63
53	H	C ₃ H ₇	H	OH	I	967	±	—	63
54	H	C ₄ H ₉	H	OH	I	1133	—	—	63
55	H	C(CH ₃) ₃	H	OH	I	1078	—	+	63
56	5-allyl			OH	I	782	+	+	69, 71
57	5-(2-chloroallyl)			OH	I	807	+	+	63
58	5-(2-butenyl)			OH	I	824	+	+	69
59	5-(but-3-en-1-ynyl)			OH	I	824	+	+	47
60	5-allylmercapto			OH	I	824	+	+	72
61	5-allyloxy			OH	I	824	+	+	73
62	H	CF ₃	H	OH	I	1078	—	+	74
63	H	C ₂ H ₄ CH ₂ Cl	H	OH	I	1133	—	+	75
64	H	C ₂ H ₄ CH ₂ CN	H	OH	I	1322	—	+	75
65	H	C ₈ H ₁₇	H	OH	I	2028	—	+	75
66	H	CH=CHCOOCH ₃	H	OH	I	1486	—	—	63
67	H	C ₂ H ₄ C=CH	H	OH	I	1133	—	+	75
68	H	<i>p</i> -methoxyphenyl	H	OH	I	1828	—	—	75
69	H	OC ₂ H ₅	H	OH	I	967	±	+	58, 75
70	H	C ₂ H ₄ COOH	H	OH	I	1298	—	—	75
71	H	C ₅ H ₁₁	H	OH	I	1531	—	—	75
72	H	C ₈ H ₁₆ COOCH ₃	H	OH	I	3278	—	—	75
73	H	C(CH ₃)=CH ₂	H	OH	I	864	+	—	75

Table 2 (Continued)

compd no.	R ¹	R ²	R ³	R ⁴	R ⁵	Wiener's index	antiviral activity		references
							predicted	reported	
74	H	1-chloro-2,2,3,3-tetra-methylcyclopropyl	H	OH	I	1732	—	—	75
75	H	1-hydroxycyclopentenyl	H	OH	I	1398	—	+	75
76	H	H	C ₄ H ₉	OH	I	1132	—	—	47
77	H	CH ₃	H	OH	IV	782	+	+	76
78	H	C ₂ H ₅	H	OH	IV	915	±	+	76
79	H	CH ₃	CH ₃	OH	I	807	+	+	63
80	CH ₃	CH ₃	H	OH	I	802	+	+	63
81	CH ₃	H	CH ₃	OH	I	802	+	—	63
82	CH ₃	COOCH ₃	H	OH	I	1203	—	—	63
83	H	COOCH ₃	CH ₃	OH	I	1219	—	—	63
84	H	Br	CH ₃	OH	I	807	+	+	63
85	H	CH ₃	Br	OH	I	807	+	+	63
86	CH ₃	Br	H	OH	I	802	+	+	63
87	H	Br	Br	OH	I	807	+	+	63
88	H	COOH	F	OH	I	1065	—	—	58
89	CH ₃	Cl	H	OH	I	802	+	—	63
90	C ₂ H ₅	C ₂ H ₅	H	OH	I	1039	—	—	75
91	F	Cl	Cl	OH	I	913	±	+	77
92	F	Cl	Cl	OH	II	913	±	—	77
93	F	F	Cl	OH	I	913	±	+	77
94	F	Cl	F	OH	I	913	±	—	77
95	F	F	CF ₃	OH	I	1307	—	—	77
96	F	CF ₃	F	OH	I	1307	—	—	77
97	H	phenyl	H	OH	I	1415	—	+	78
98	H	<i>m</i> -nitrophenyl	H	OH	I	1813	—	—	78
99	H	<i>m</i> -azidophenyl	H	OH	I	2012	—	—	78
100	H	H	H	OH	XVII	393	—	—	80
101	H	Br	H	OH	XVII	478	—	—	80
102	H	Br	Br	OH	XVII	563	±	+	80
103	H	H	H	OH	V	478	—	—	80
104	H	Br	Br	OH	V	563	±	—	80
105	H	I	C ₄ H ₃	OH	V	782	+	+	80
106	5-phenyl 5-(2-phenylethyl) 5-phenylethynyl 5-(3-nitrophenyl) 5-anilino CH(OH)CH ₂ Br CH(OH)CH ₂ Cl CH(OCH ₃)CH ₂ Br CH(OCH ₃)CH ₂ Cl CH(OH)CH(Br)COOCH ₃ CH(OH)CH ₂ I CH(Br)CH(OH)I CH(OH)CH(OH)I			OH	I	1040	—	—	47
107				OH	I	1415	—	+	78
108				OH	I	1415	—	—	65
109				OH	I	1440	—	—	47
110				OH	I	1040	—	+	79
111				OH	I	802	+	+	82
112				OH	I	802	+	+	82
113				OH	I	915	±	+	82
114				OH	I	915	±	+	82
115				OH	I	1607	—	+	82
116				OH	I	802	+	+	82
117				OH	I	913	±	+	82
118				OH	I	913	±	+	82

^a Negative antiviral activity. ^b Positive antiviral activity. ^c Transitional range where activity could not be specifically assigned.

(b) Development of drug resistance resulting from selection and overgrowth of viral populations, e.g. in treatment of HSV infection with acyclovir.³⁶

(c) Latency, perhaps the most difficult problem of all; presently no solution seems to be in sight.

(d) Drug delivery and drug metabolism, which are being approached with the use of liposomes and prodrugs, respectively.

Despite the above-mentioned limitations, antiviral chemotherapy has witnessed rapid progress, especially in the late 1980s and early 1990s. Recent trends in antiviral drug design have been focused on the development of derivatives of existing drugs as well as on the synthesis of entirely new antiviral agents. The first approach is aimed at either increasing potency, decreasing the toxicity, widening the antiviral spectrum, or improving pharmacokinetic profiles of currently available drugs.³⁷ This is the approach where the concept of the structure–function relationship can be successfully exploited. To assist in the identification and development of more effective antiviral agents, the SAR of 5-vinylpyrimidine nucleoside analogs has been studied in the present investiga-

tions. These compounds have been shown to have selective and potent antiviral properties.³⁸ The compounds included in the study have been reportedly evaluated against herpes simplex virus (HSV) type I in cell culture.

METHODOLOGY

Antiviral agents have been extensively reviewed.^{31–35,37,39–43} A set of active and inactive analogs based on 5-vinylpyrimidine as depicted in Figure 1 was selected. Various sugar substituents of R⁵ are listed in Table 1. As for R⁴ substitutions, compounds with R⁴ = OH will exhibit tautomerism with NHCO instead of the N=C(OH) group, as indicated by the formula. The Wiener's index value for each compound was computed from the hydrogen-suppressed molecular structure of the compounds. The index values so obtained were classified into active, transitional, and inactive ranges. These ranges were then employed to predict activity of molecules on the basis of computed Wiener's index values. For 5-vinylpyrimidine nucleoside analogs, Wiener's index values of 576–900 were selected as an active range. Compounds with values less than 500 and greater than 975 have been predicted to be

Table 3. Relative Distribution of Test Compounds in Various Ranges of Wiener's Topological Index Values

S no.	ranges of Wiener's index value	nature of range	total number of compounds falling within the range	compounds predicted correctly	accuracy of prediction
1	500 or less	inactive	10	10	100%
2	501–575	lower transitional range	2	not applicable	not applicable
3	576–900	active	60	52	86.66%
4	901–975	upper transitional range	15	not applicable	not applicable
5	976 or more	inactive	31	19	61.29%

inactive against HSV-I infection. The intermediate ranges, i.e. 501–575 and 901–975, were selected as transitional or borderline ranges where antiviral activity could not be specifically assigned on the basis of their Wiener's index values.

RESULTS AND DISCUSSION

A retrofit study of Table 2 shows that a total of 81 out of 101 compounds were classified correctly within active and inactive ranges. The relative distribution of test compounds falling under various ranges of Wiener's topological index along with predicted and reported activities has been compiled in Table 3.

Wiener's index was found to be a promising topological descriptor as the results obtained in the above work showed 82.65% accuracy in prediction. However, when the upper transitional range of Wiener's index values was modified to 900–1100, the correlation between the Wiener's index value and antiviral activity resulted in an enhancement of prediction accuracy to 85.22%.

Some of the mispredictions in the above work may be due to high degeneracy for Wiener's index values, probably due to the following: (i) Wiener's index gives the same value for both α and β anomers, while as a general rule, all α anomers are inactive or have relatively weak activity, i.e. compounds 91 and 92.

(ii) *E* and *Z* isomers give the same value for Wiener's index, while C-5 compounds with only *E* stereochemistry are found to be more active, i.e. compounds 21, 22, 51, 52, 80, and 81.

However despite these shortcomings, Wiener's index was found to have a highly significant correlating ability with biological properties of 5-vinylpyrimidines. The initial results obtained from this study show considerable promise regarding the usefulness of topological indices in the SAR of antiviral agents. Further studies on the SAR of antiviral agents with the main emphasis on anti-AIDS activity are under progress and will be communicated shortly.

REFERENCES AND NOTES

- Klopman, G.; Raychaudhury, C. A novel approach to use of graph theory in structure activity relationship studies. Application to the qualitative evaluation of mutagenicity in a series of nonfused ring aromatic compounds. *J. Comput. Chem.* **1988**, *9*, 232–243.
- Sabljic, A.; Trinajstić, N. Quantitative structure activity relationship: The role of topological indices. *Acta Pharm. Jugosl.* **1981**, *31*, 189–214.
- Wiener, H. Correlation of heat of isomerization and difference in heat of vaporisation of isomers, among paraffin hydrocarbons. *J. Am. Chem. Soc.* **1947**, *69* (17), 2636.
- Wiener, H. Influence of intratomic forces on paraffin properties. *J. Chem. Phys.* **1947**, *15*, 766.
- Wiener, H. Vapour pressure–temperature relations among the branched paraffin hydrocarbons. *J. Chem. Phys.* **1948**, *52*, 425–430.
- Wiener, H. Relationship of physical properties of isomeric alkanes to molecular structure surface tension, specific dispersion and critical solution temperature in aniline. *J. Phys. Colloid Chem.* **1948**, *52*, 1082–1089.
- Balaban, A. T.; Chiriac, A.; Motoc, I.; Simon, Z. Steric Fit in QSAR. *Lect. Notes Chem.* **1980**, *15*, 22.
- Balaban, A. T. Chemical graphs XXXIV. 5 new topological indices for the branching of tree like graphs. *Theor. Chim. Acta* **1979**, *53*, 355.
- Kier, L. B.; Hall, L. H.; Murray, W. J.; Randić, M. Molecular Connectivity I: Relationship to nonspecific local anaesthetics. *J. Pharm. Sci.* **1975**, *64* (12), 1971–1974.
- Kier, L. B.; Hall, L. H. *Molecular Connectivity In Chemistry And Drug Research*; Academic Press: New York, 1976.
- Randić, M. On characterization of molecular branching. *J. Am. Chem. Soc.* **1965**, *97*, 6609–6615.
- Hosoya, H. Topological index: Newly proposed quantity characterizing the topological nature of structure of isomers of saturated hydrocarbons. *Bull. Chem. Soc. Jpn.* **1971**, *44*, 2332.
- Hosoya, H. Topological index as a sorting device for coding chemical structures. *J. Chem. Doc.* **1972**, *12*, (3), 181–183.
- Balaban, A. T. *Chem. Phys. Lett.* **1982**, *89*, 399.
- Balaban, A. T. Computer programme for topological index J (average distance sum connectivity). *Math. Chem.* **1984**, *16*, 163–190.
- Balaban, A. T.; Quintar, L. B. The smallest graphs, trees and 4-trees with degenerate topological index *J. Math. Chem.* **1983**, *14*, 213–233.
- Bonchev, D.; Trinajstić, N. Information theory, distance matrix and molecular branching. *J. Chem. Phys.* **1977**, *67*, 4517–4533.
- Bonchev, D. *Theoretic Information, Indices for Characterization of Chemical Structures*; Wiley-Research Studies: Chichester, England, 1983.
- Bonchev, D. On topological characterization of molecular branching. *Int. J. Quantum Chem. Symp.* **1978**, *12*, 293.
- Motoc, I.; Balaban, A. T. Topological index: Intercorrelation, physical meaning and correlational ability. *Rev. Roum. Chim.* **1981**, *26*, 539.
- Gutman, I.; Trinajstić, N. Graph theory and molecular orbitals, total electron energy of alternate hydrocarbons. *Chem. Phys. Lett.* **1972**, *17*, 535.
- Gutman, I.; Ruscic, B.; Trinajstić, N.; Wilcox, C. F. Graph theory and molecular orbitals XII. Acyclic polyenes. *J. Chem. Phys.* **1975**, *62*, 3399.
- Gutman, I.; Randić, M. Algebraic characterization of skeletal branching. *Chem. Phys. Lett.* **1977**, *47*, 15–19.
- Platt, J. R. *J. Chem. Phys.* **1947**, *15*, 419.
- Platt, J. R. *J. Phys. Chem.* **1952**, *56*, 328.
- Gordon, M.; Scantlebury, C. R. *Trans. Faraday Soc.* **1964**, *60*, 604.
- Attenburg, K. *Kolloidn. Zh.* **1961**, *178*, 112.
- Smolenskii, E. A. *Zh. Fiz. Khim.* **1964**, *38*, 1288.
- Balaban, A. T. Application of graph theory in chemistry. *J. Chem. Inf. Comput. Sci.* **1985**, *25*, 334–343.
- Nagarajan, R.; Naik, S. R. Recent developments in antiviral agents. *Hind. Antibiot. Bull.* **1985**, *31*, 90–113.
- Bean, B. Antiviral therapy: Current concepts and practices. *Clin. Microbiol. Rev.* **1992**, *5*, 146–182.
- De Clercq, E. New acquisitions in the development of anti-HIV agents. *Antiviral Res.* **1989**, *12*, 1–20.
- Montgomery, J. A. Approaches to antiviral chemotherapy. *Antiviral Res.* **1989**, *12*, 113–132.
- Broder, S. Antiretroviral therapy in AIDS. *Ann. Int. Med.* **1990**, *113*, 604–618.
- De Clercq, E. Recent advances in search for selective antiviral agents. *Adv. Drug Res.* **1988**, *17*, 1–59.
- Field, H. J. The development of antiviral drug resistance in antiviral agents. In *The development and assessment of antiviral chemotherapy*. Field, H. J., Ed.; CRC Press: Boca Raton, FL, 1988; Vol. 1, pp 127–149.
- De Clercq, E. Recent trends and development in antiviral chemotherapy. *Antiviral Res.* **1985**, Suppl. 1, 11–19.
- De Clercq, E.; Descamps, J.; De Somer, P.; Barr, P. J.; Jones, A. S.; Walker, R. T. (E)-5-(2-Bromovinyl)-2'-deoxyuridine: a potent and selective anti-herpes agent. *Proc. Natl. Acad. Sci. U.S.A.* **1979**, *76*, 2947–2951.
- DeClercq, E.; Walker, R. T. Synthesis and antiviral properties of 5-vinylpyrimidine nucleoside analogs. *Pharmacol. Ther.* **1984**, *26*, 1–57.
- Prusoff, W. H.; Lin, T. S.; August, E. M.; Wood, T. G.; Marongiu, M. E. Approaches to antiviral drug development. *Yale J. Biol. Med.* **1989**, *62*, 215–225.
- Diana, G. D.; Pevear, D.; Young, D. C.; Antiviral agents. *Annu. Rep. Med. Chem.* **1989**, *24*, 129–135.
- Yarchoan, R.; Misuya, H.; Myers, C. E.; Broder, S. Clinical pharmacology of 3'-azido-2',3'-dideoxythymidine and related dideoxynucleosides. *The New Eng. J. Med.* **1989**, *321*, 11, 726–738.
- Nasr, M.; Litterst, C.; McGowan, J. Computer-assisted SAR of dideoxynucleoside analogs as potential anti-HIV drugs. *Antiviral Res.* **1990**, *14*, 125–148.

- (44) Bobek, M.; Bloch, A. The chemistry and biology of some new nucleoside analogs active against tumor cells. In *Chemistry and Biology of Nucleosides and Nucleotides*; Harmon, R. E., Robins, R. K., Townsend, L. B., Eds.; Academic Press: New York, 1978; pp 135-148.
- (45) Barr, P. J.; Jones, A. S.; Walker, R. T. Incorporation of 5-substituted uracil derivatives into nucleic acids. Part IV. The synthesis of 5-ethynyluracil. *Nucleic Acids Res.* **1976**, *3*, 2845-2849.
- (46) Barr, P. J.; Jones, A. S.; Serafinowski, P.; Walker, R. T. The synthesis of nucleosides derived from 5-ethynyluracil and 5-ethynylcytosine. *J. Chem. Soc., Perkin Trans. 1* **1978**, 1263-1267.
- (47) De Clercq, E. Synthesis and antiviral properties of 5-vinyl pyrimidine nucleoside analogs. *Pharmacol. Ther.* **1984**, *26*, 1-58.
- (48) De Clercq, E.; Descamps, J.; Balzarini, J.; Giziewicz, J.; Barr, P. J.; Robins, M. J. Nucleic acid related compounds. 40. Synthesis and biological activities of 5-alkynyluracil nucleosides. *J. Med. Chem.* **1983**, *26*, 661-666.
- (49) De Clercq, E.; Descamps, J.; Verhelst, G.; Jones, A. S.; Walker, R. T.; Torrence, P. F.; Shugar, D. Comparative efficacy of antiherpes drugs against different strains of herpes simplex virus. *J. Infect. Dis.* **1980**, *141*, 563-574.
- (50) De Clercq, E.; Balzarini, J.; Torrence, P. F.; Mertes, M. P.; Schmidt, C. L.; Shugar, D.; Barr, P. J.; Jones, A. S.; Verhelst, G.; Walker, R. T. Antiviral, antimetabolic and cytotoxic activities of 5-substituted 2'-deoxy cytidines. *Mol. Pharmacol.* **1982**, *21*, 217-223.
- (51) Rahim, S. G.; Duggan, M. J. H.; Walker, R. T.; Jones, A. S.; Dyer, R. L.; Balzarini, J.; De Clercq, E. Synthesis and biological properties of 2'-deoxy-5-vinyl uridine and 2'-deoxy-5-vinyl cytidine. *Nucleic Acids Res.* **1982**, *10*, 5285-5295.
- (52) Watanabe, K. A.; Su, T.-L.; Reichman, U.; Greenberg, N.; Lopez, C.; Fox, J. J. Nucleosides. 129. Synthesis of antiviral nucleosides: 5-alkenyl-1-(2-deoxy-2-fluoro- β -D-arabinofuranosyl)uracils. *J. Med. Chem.* **1984**, *27*, 91-94.
- (53) Sakata, S.; Shibuya, S.; Machida, H.; Yoshino, H.; Hirota, K.; Senda, S.; Ikeda, K.; Mizuno, Y. Synthesis and antiherpesviral activity of 5-C-substituted uracil nucleosides. *Nucleic Acids Res. Symp. Ser.* **1980**, *8*, s39-s42.
- (54) Machida, H.; Kuninaka, A.; Yoshino, H.; Ikeda, K.; Mizuno, Y. Antiherpesviral activity and inhibitory action on cell growth of 5-alkenyl derivatives of 1- β -D-arabinouranosyluracil. *Antimicrob. Agents Chemother.* **1980**, *17*, 1030-1031.
- (55) Machida, H.; Sakata, S.; Kuninaka, A.; Yoshino, H. Antiherpesviral and anticellular effects of 1- β -D-arabinofuranosyl-(E)-5-(2-halogenovinyl)-uracils. *Antimicrob. Agents Chemother.* **1981**, *20*, 47-52.
- (56) De Clercq, E.; Descamps, J.; Verhelst, G.; Walker, R. T.; Jones, A. S. Antiviral activity of 5-(2-halogenovinyl)-2'-deoxyuridines. In *Current Chemotherapy and Infectious Diseases*; Nelson, J. D., Grassi, G. C., Eds.; American Society of Microbiology: Washington, DC, 1980; pp 1372-1374.
- (57) Jones, A. S.; Rahim, S. G.; Walker, R. T.; De Clercq, E. Synthesis and antiviral properties of (Z)-5-(2-bromovinyl)-2'-deoxyuridine. *J. Med. Chem.* **1981**, *24*, 759-760.
- (58) Reefschrager, J.; Barwolff, D.; Langen, P. Differential antiherpes activity of (E) and (Z) isomers of 5-(2-fluorovinyl)-2'-deoxyuridine. *Acta Virol.* **1984**, *28*, 282-286.
- (59) De Clercq, E.; Busson, R.; Colla, L.; Descamps, J.; Balzarini, J.; Vanderhaeghe, H. Antiviral activity of sugar modified derivatives of (E)-5-(2-bromovinyl)-2'-deoxyuridine. In *Current Chemotherapy and Immunotherapy*; Periti, P., Grassi, G. C., Eds.; American Society for Microbiology: Washington, DC, 1982; pp 1065-1066.
- (60) Reefschrager, J.; Herrmann, G.; Barwolff, D.; Schwarz, B.; Cech, D.; Langen, P. Antiherpes activity of (E)-5-(2-bromovinyl) and 5-vinyl-1- β -D-arabinofuranosyl uracil and some other 5-substituted uracil arabinosyl nucleosides in two different cell lines. *Antiviral Res.* **1983**, *3*, 175-187.
- (61) Colla, L.; Busson, R.; De Clercq, E.; Vanderhaeghe, H. Synthesis of aliphatic nucleoside analogs with potential antiviral activity. *Eur. J. Med. Chem.* **1982**, *17*, 569-576.
- (62) Verhelst, G.; De Clercq, E.; Balzarini, J.; Descamps, J.; Jones, A. S.; Walker, R. T. Synthesis, antiviral and antitumor activity of 5-(1-chloro-2-bromovinyl)-2'-deoxyuridine. Presented at VIIth International Symposium On Medicinal Chemistry, Torremolinos, Malaga, Spain, Sept 2-5, 1980; Abstract 18.
- (63) Goodchild, J.; Porter, R. A.; Raper, R. H.; Sim, I. S.; Upton, R. M.; Viney, J.; Wadsworth, H. J. Structural requirements of olefinic 5-substituted deoxyuridines for antiherpes activity. *J. Med. Chem.* **1983**, *26*, 1252-1257.
- (64) Busson, R.; Colla, L.; Vanderhaeghe, H.; De Clercq, E. Synthesis and antiviral activity of some sugar modified derivatives of (E)-5-(2-bromovinyl)-deoxyuridine. *Nucleic Acids Res. Symp. Ser.* **1981**, *9*, 49-52.
- (65) De Clercq, E.; Descamps, J.; Balzarini, J.; Fukui, T.; Allaudeen, H. S. Antiviral activity of 3'-amino derivative of (E)-5-(2-bromovinyl)-2'-deoxyuridine. *Biochem. J.* **1983**, *211*, 439-445.
- (66) De Clercq, E.; Verhelst, G.; Descamps, J.; Bergstrom, D. E. Differential inhibition of herpes simplex virus type-1 and type-2 by (E)-5-(2-X-vinyl)-2'-deoxyuridines. *Acta Microbiol. Acad. Sci. Hung.* **1981**, *28*, 307-312.
- (67) Bergstrom, D. E.; Ruth, J. L.; Reddy, P. A.; De Clercq, E. Synthesis of (E)-5-(3,3,3-trifluoro-1-propenyl)-2'-deoxyuridine and related analogs: Potent and unusually selective antiviral activity of (E)-5-(3,3,3-trifluoro-1-propenyl)-2'-deoxyuridine against HSV-1. *J. Med. Chem.* **1984**, *27*, 279-284.
- (68) Stening, G.; Gotthammar, B.; Larsson, A.; Alenius, S.; Johansson, N. G.; Oberg, B. Antiherpes activity of (E)-5-(1-propenyl)-2'-deoxyuridine and 5-(1-propenyl)-10- β -D-arabinofuranosyluracil. *Antiviral Res.* **1981**, *1*, 213-223.
- (69) Sim, I. S.; Goodchild, J.; Meredith, D. M.; Porter, R. A.; Raper, R. H.; Viney, J.; Wadsworth, H. J. Possible molecular basis for antiviral activity of certain 5-substituted deoxyuridines. *Antimicrob. Agents Chemother.* **1983**, *23*, 416-421.
- (70) Sim, I. S.; Raper, R. H. 5-Substituted deoxyuridines. Structural requirements for antiviral activity against HSV-1 and 2 and possible biochemical basis for relative potency. *Antiviral Res.* **1984**, *4*, 159-168.
- (71) Cheng, Y. C.; Domin, B. A.; Sharma, R. A.; Bobek, M. Antiviral action and cellular toxicity of four thymidine analogs: 5-ethyl, 5-vinyl, 5-propyl and 5-allyl-2'-deoxyuridine. *Antimicrob. Agents Chemother.* **1976**, *10*, 119-122.
- (72) Dinan, F. J.; Bardos, T. J. Synthesis of some new S-alkylated derivatives of 5-mercapto-2'-deoxyuridine as potential antiviral agents. *J. Med. Chem.* **1980**, *23*, 569-572.
- (73) Torrence, P. F.; Spencer, J. W.; Bobst, A. M.; Descamps, J.; De Clercq, E. 5-o-Alkylated derivatives of 5-hydroxy-2'-deoxyuridine as potential antiviral agents. Antiherpes activity of 5-propenyloxy-2'-deoxyuridine. *J. Med. Chem.* **1978**, *21*, 228-231.
- (74) Wataya, Y.; Matsuda, A.; Santi, D. V.; Bergstrom, D. E.; Ruth, J. L. Trans-5-(3,3,3-trifluoro-1-propenyl)-2'-deoxyuridylate: a mechanism based inhibitor of thymidylate synthetase. *J. Med. Chem.* **1979**, *22*, 339-340.
- (75) Vincent, P.; Beaucourt, J. P.; De Clercq, E.; Balzarini, J.; Pichat, L. Syntheses et activites biologiques de nouvelles (E)-alcenyl-5-désoxy-2'-uridines. *Nucleosides Nucleotides* **1985**, *4* (4), 429-45.
- (76) Machida, H.; Kuninaka, A.; Yoshino, H.; Ikeda, K.; Mizuno, Y. Anti herpes viral activity and inhibitory action on cell growth of 5-alkenyl derivatives of 1- β -D-arabinofuranosyluracil. *Antimicrob. Agents Chemother.* **1980**, *17*, 1030-1031.
- (77) Coe, P. L.; Harnden, M. R.; Jones, A. S.; Noble, S. A.; Walker, R. T. Synthesis and antiviral properties of some 2'-deoxy-5-(fluoroalkenyl) uridines. *J. Med. Chem.* **1982**, *25*, 1329-1334.
- (78) De Clercq, E.; Balzarini, J.; Descamps, J.; Bigge, C. F.; Chang, C. T.-C.; Kalaritis, P.; Mertes, M. P. Antiviral, antitumor and thymidylate synthetase inhibition studies of 5-substituted styryl derivatives of 2'-deoxyuridine and their 5'-phosphates. *Biochem. Pharmacol.* **1981**, *30*, 495-502.
- (79) Bobst, A. M.; Ozinskar, A. J.; De Clercq, E. 46. Synthesis and biological activities of C(5)-N-spinlabelled uridines and related derivatives. *Helv. Chim. Acta* **1983**, *66*, 534-541.
- (80) Reefschrager, J.; Herrmann, G.; Jennrich, H.; Frost, N. Antiherpes simplex virus and cytostatic activity of some new 5-substituted 1-(4-hydroxybutyl)- and 1-(2-hydroxyethoxymethyl) uracil nucleoside analogs. *Acta Virol.* **1985**, *29*, 185-193.
- (81) Machida, H.; Nishitani, M.; Suzutani, J.; Kozaburo, H. Different antiviral potencies of BV-araU and related nucleoside analogs against HSV-1 in human cell lines and Vero cells. *Microbiol. Immunol.* **1991**, *35*, 963-973.
- (82) Kumar, R.; Wiebe, L. I.; Hall, T. W.; Knaus, E. E.; Towell, D. R.; Tyrell, D. L.; Allen, T. M. Synthesis of 5-(1-hydroxy(or methoxy)-2-bromo-(or chloro) ethyl 2'-deoxyuridines and related halohydrin analogs with antiviral and cytotoxic activity. *J. Med. Chem.* **1989**, *32*, 941-944.