

- 2:30 (1264) Electrochemical Hydride Generation Sample Introduction into ICPMS. **I. Marawi**, University of Cincinnati, T. J. Ntozakhe, K. Khalema, T. H. Ridgeway, J. A. Caruso
- 2:50 (1265) Recent Advances in the Design of Thermospray-Based Sample Introduction for AES. **T. S. Conner**, Southern Illinois University, W. Howe, J. A. Koropchak
- 3:25 (1266) Elimination of Acid Matrix and Organic Solvent Interferences in Ultrasonic Nebulization ICP-AES. **J. J. Zhu**, CETAC Technologies
- 3:45 (1267) Use of an Ultrasonic Nebulizer in the Determination of Boron by ICP Spectrometry. **D. R. Wieder**, CETAC Technologies, F. G. Smith
- 4:05 (1268) Determination of Trace Elements by IC(Argon)P Spectrometry with a Modified Ultrasonic Nebulizer. **C. M. Harris**, U. S. Geological Survey, C. J. Litteral, D. L. Milne
- 4:25 (1269) ICP with Ultrasonic Nebulization at 5, 10, and 20 $\mu\text{L}/\text{min}$ Liquid Flow Rates. **M. A. Tarr**, U. S. Environmental Protection Agency, G. Zhu, R. F. Browner
- 4:45 (1270) Utilizing Fluorescence for Hydride Elements with a Commercial Detector. **J. Poling**, Questron, A. C. Grillo, P. Stockwell

Separation Sciences II

Room 269

I. P. Ho, USDA, *Presiding*

- 1:30 (1271) New Sample Preparation Method for the Determination of Sucralose in Ketchup Using Dialysis, Trace Enrichment, and HPLC. **N. M. Rafel**, McNeil Specialty Products, G. P. Cosgrove, C. M. Merkel
- 1:50 (1272) Interfacial Component in the Complexation of Metal Ions by Dithizone in Microemulsions. **Y. X. Xia**, University of Arizona, S. Muralidharan, H. Freiser
- 2:10 (1273) Kinetics and Equilibrium of the Extraction and Back Extraction of Ni(II) and Cu(II) by Aromatic Oximes. **E. Ma**, University of Arizona, S. Muralidharan, R. P. Sperline, H. Freiser
- 2:30 (1274) Fundamental Aspects of the Separation of Platinum Group Metals by CPC. **H. Ma**, University of Arizona, S. Muralidharan, H. Freiser
- 2:50 (1275) Automated SPE/HPLC for the Extraction and Analysis of Drugs from Biological Fluids. **K. Fogelman**, Hewlett Packard, W. Miles, P. Castelli
- 3:25 (1276) HPLC/Diode Array Detection Analysis of Unstable Analytes Using Automated Sample Preparation. **R. Gard**, Hitachi Instruments
- 3:45 (1277) Systematic Chiral Separation Development on Bonded R versus S versus RS Naphthylethyl Carbamate Derivatives of β - and γ -Cyclodextrin. **T. E. Beesley**, Advanced Separation Technologies, T. Zakszewski
- 4:05 (1278) Analysis of Ligand/Protein Complexation Using CE Focusing. **A. C. Cater**, Furman University, C. V. Thomas, D. B. Meetze, J. F. Wheeler

- 4:25 (1279) Determination of Ammonia and Inorganic Nitrogen Using the Diffusion-Conductivity Method. **I. K. Henderson**, Alltech Associates, R. Saari-Nordhaus, J. M. Anderson, Jr.
- 4:45 (1280) Rate of Extraction and Separation Efficiency in CPC for Some Tervalent Lanthanide Metals. **K. Inaba**, University of Arizona, S. Muralidharan, H. Freiser


Surface Phenomena with Analytical Potentialities


Room 363

A. J. Sharkins, ALCOA, *Presiding*

- 1:30 (1281) Metalloporphyrin-Based Anion-Selective Optical Sensors. **E. Wang**, University of Michigan, M. E. Meyerhoff
- 1:50 (1282) Voltammetry in CO_2 -Based Fluids with Platinum Microdisk Electrodes Coated with Polymer Electrolytes. **E. F. Sullenberger**, University of Pittsburgh, M. G. Garguilo, A. C. Michael
- 2:10 (1283) Response of Graphite Surfaces with Surfactants as Molecular Spacers. **A. Marino**, University of Florida, A. Brajter-Toth
- 2:30 (1284) Characterization of Plasma-Modified Fluoropolymer Surfaces. **M. Li**, State University of New York, F. V. Bright
- 2:50 (1285) Characterization of Doped Sol-Gel Materials for Use as Chemical Sensors and Tunable Glass Lasers. **U. Narang**, State University of New York, R. A. Dunbar, P. N. Prasad, F. V. Bright

Dr. Smith
found
two peaks...





Using PeakFit™,
Dr. Jones found
three peaks...

Guess who got the grant?

Because PeakFit™ uses sophisticated non-linear curve fitting techniques, it is far more effective than other data analysis tools in reducing noise and separating unresolved peaks! In fact, PeakFit is one of the most powerful software packages available today for chromatography and spectroscopy data analysis. For chromatographers, PeakFit extracts important system suitability information such as peak areas, asymmetries, and resolutions – plus column efficiency (number of theoretical plates) from raw data. For spectroscopists, PeakFit calculates peak amplitudes, areas, centers and widths – with complete uncertainty analysis, too!

Use PeakFit's 30 built-in peak functions (including Gaussian, Exponentially-Modified Gaussian, Voigt and many others specifically suited for chromatography and spectroscopy) or input and use your own! Output both graphs and data to over 150 printers, including PostScript®. Of course we can't promise that PeakFit will win you a grant (your competition may already be using PeakFit!), but we can promise that PeakFit will enhance your current analysis methods. Order PeakFit today and see just how many peaks there really are in your data!

Manufacturers: Call for OEM info!

CIRCLE 88 ON READER SERVICE CARD

New Version 3.0!

PeakFit™
Peak Analysis Software
for the IBM PC

For free brochure:

800-874-1888 (U.S.)
415-453-6700 FAX 415-453-7769

From the makers of SigmaPlot®!

Jandel
SCIENTIFIC

2591 Kerner Blvd
San Rafael, CA 94901
Europe: Schimmelbuschstr. 25
4006 Erkrath, Germany

