

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231443340>

# Intramolecular $S_N1$ cleavage of allylic ethers by enolate anions

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MARCH 1989

Impact Factor: 12.11 · DOI: 10.1021/ja00188a073

---

CITATIONS

35

---

READS

15

4 AUTHORS, INCLUDING:



Christopher A. Teleha

Johnson & Johnson

36 PUBLICATIONS 822 CITATIONS

SEE PROFILE

These and other questions regarding this unusual pathway<sup>18</sup> are currently under investigation.

**Acknowledgment.** Dr. David Martin and Shirley Gerpheide of The Upjohn Co. are thanked for providing samples of acivicin and 4-hydroxyacivicin, slants of *Streptomyces svicens*, and the design of the baffled flask used in fermentations. P. C. Prabhakaran and Rodger Kohnert are thanked for obtaining the <sup>2</sup>H NMR spectra. The work was supported by a grant from the Public Health Service GM 32110 to S.J.G. NMR spectra were obtained on a Bruker AM 400 spectrometer purchased in part by grants from the National Science Foundation (CHE-8216190) and from the M. J. Murdock Charitable Trust to Oregon State University.


### Intramolecular S<sub>N</sub> Cleavage of Allylic Ethers by Enolate Anions

Leo A. Paquette,<sup>\*,1a</sup> Jeff Reagan,<sup>1b</sup> Stuart L. Schreiber,<sup>\*,1b</sup> and Christopher A. Teleha<sup>1a</sup>

Evans Chemical Laboratories  
The Ohio State University, Columbus, Ohio 43210  
Yale University, Sterling Chemistry Laboratory  
New Haven, Connecticut 06511

Received November 21, 1988


A fundamental property of allylic systems relates to their ability to engage nucleophiles in S<sub>N</sub> reaction. Extensive theoretical<sup>2</sup> and experimental investigations<sup>3</sup> over many years attest to the significance attached to the phenomenon. Despite the wide-ranging nature of these studies, however, no attention has been purposefully directed to bimolecular nucleophilic substitution of allyl alkyl ethers, perhaps because of a predetermination that these systems would prove as inert toward displacement as dialkyl ethers. One interesting example known to us is due to Farnum and Monego who showed that dimetalation of **1** proceeds with subsequent proton abstraction from solvent to give **2**, which then experiences intramolecular S<sub>N</sub> displacement of methoxide.<sup>4</sup>



As part of ongoing investigations of anionic oxy-Cope rearrangements,<sup>5,6</sup> our research groups have independently examined


the ability of medium-ring enolates to undergo transannular cyclization concomitant with S<sub>N</sub> displacement of a methoxyl leaving group. The result is rapid and efficient construction of structurally intricate polycyclic systems.

For example, when alcohol **4**<sup>7</sup> was stirred in anhydrous tetrahydrofuran solution with 1.1 equiv of potassium hexamethyldisilazide at 20 °C for 4 days, smooth conversion to ketone **6** was



observed. Chromatographically purified material, isolated in 51% yield, crystallized as colorless, rectangular plates well suited to X-ray analysis.<sup>8</sup> The suggested pathway to the product diquinane involves initial [3,3]sigmatropic electron reorganization via a chair-like transition state to generate **5**. This process establishes three stereocenters and the double bond geometry. The β-configuration of the methoxyl-substituted carbon results in proper alignment of the C-OCH<sub>3</sub> bond with the flanking π orbital, thereby allowing for the onset of the intramolecular S<sub>N</sub> ring closure.

Support for this mechanistic analysis was gained by subjecting **7** to comparable ring expansion. In this instance, **8** was produced efficiently (88%) after only 5.5 h at room temperature. Since crystals of **8** of suitable quality could not be grown, saturation



of its double bond was undertaken. The structure of **9** was subsequently established by crystallographic methods to be as shown.<sup>8</sup> These data indicate the **7** → **8** conversion to be mediated by an


- (1) (a) The Ohio State University. (b) Yale University.  
(2) (a) Houk, K. N.; Paddon-Row, M. N.; Rondan, N. G. *THEOCHEM* 1983, 12, 197; *Chem. Abstr.* 1983, 99, 211818. (b) Chen, N.; Zhou, B. *Youji Huaxue* 1983, 257; *Chem. Abstr.* 1983, 99, 211825. (c) Dewar, M. J. S. *J. Am. Chem. Soc.* 1984, 106, 209. (d) Carrion, F.; Dewar, M. J. S. *Ibid.* 1984, 106, 3531. (e) Stohrer, W.-D. *Angew. Chem., Int. Ed. Engl.* 1983, 22, 613. (f) Houk, K. N. *Pure Appl. Chem.* 1983, 55, 277. (g) Murphy, W. S.; O'Mahony, B. *Tetrahedron Lett.* 1981, 22, 585.  
(3) (a) Magid, R. M. *Tetrahedron* 1980, 36, 1901. (b) Overton, K. H. *Chem. Soc. Rev.* 1979, 8, 447. (c) Cane, D. E. *Tetrahedron* 1980, 36, 1109.  
(4) Farnum, D. G.; Monego, T. *Tetrahedron Lett.* 1983, 24, 1361.  
(5) (a) Schreiber, S. L.; Hawley, R. C. *Tetrahedron Lett.* 1985, 26, 5971. (b) Schreiber, S. L.; Santini, C. *J. Am. Chem. Soc.* 1984, 106, 4038. (c) Schreiber, S. L.; Santini, C. *Tetrahedron Lett.* 1981, 22, 4651.

- (6) (a) Paquette, L. A.; Poupart, M.-A. *Tetrahedron Lett.* 1988, 29, 273. (b) Paquette, L. A.; DeRussy, D. T.; Rogers, R. D. *Tetrahedron* 1988, 44, 3139. (c) Paquette, L. A.; DeRussy, D. T.; Cottrell, C. E. *J. Am. Chem. Soc.* 1988, 110, 890. (d) Paquette, L. A.; Learn, K. S.; Romine, J. L.; Lin, H.-S. *Ibid.* 1988, 110, 879. (e) Oplinger, J. A.; Paquette, L. A. *Tetrahedron Lett.* 1987, 28, 5441. (f) Paquette, L. A.; Pierre, F.; Cottrell, C. E. *J. Am. Chem. Soc.* 1987, 109, 5731. (g) Paquette, L. A.; Learn, K. S.; Romine, J. L. *Tetrahedron* 1987, 43, 4889. (h) Paquette, L. A.; Romine, J. L.; Lin, H.-S. *Tetrahedron Lett.* 1987, 28, 31.

- (7) Details relating to the synthesis of this compound will appear elsewhere.  
(8) We thank Dr. J. C. Gallucci of the Ohio State University Crystallographic Facility for these analyses.

exo-boat transition state.<sup>9</sup> Since the methoxyl substituent in the initially generated enolate anion is consequently  $\alpha$ -oriented, i.e., positioned nearly orthogonal to the  $\pi$  electrons, further transannular C-C bond formation is now stereoelectronically disfavored and does not operate.


In yet another fascinating reaction, heating of **10** with potassium hydride and 18-crown-6 in dry tetrahydrofuran afforded **14** (40%).



Because detailed spectroscopic analysis of this tricyclic ketone did not allow for definitive determination of the position of its cyclobutene double bond, suitable degradation was undertaken. Thus, ozonolysis of **14** and subsequent treatment of stable ozonide **15** with  $\text{FeSO}_4/\text{Cu}(\text{OAc})_2$ <sup>10</sup> in methanol gave the cyclopentene derivative **17**. The formation of **17** unequivocally establishes the site of unsaturation to be as in **14** rather than **13**.

This transformation is believed to originate from enolate anion **11** as the first-formed intermediate. Due to the large amount of strain associated with the bicyclo[3.2.0]hept-1(6)-ene part structure of **13**, intramolecular displacement of methoxide in **11** is energetically disfavored. Instead, enolate isomerization of **11** to **12** operates, with tetrahydrofuran or 18-crown-6 perhaps acting as the proton source, followed by  $\text{S}_{\text{N}}'$  transannular "scaffolding" to deliver the observed tricyclic product.

In order to scrutinize the kinetic consequences of methoxyl stereochemistry, alcohol **18** was also heated as its potassium salt. Under the same conditions, **19** was isolated in ca. 60% yield.



Charge annihilation within the intermediate enolate anion by  $\text{S}_{\text{N}}'$  ejection of methoxide must consequently also be facile. This is because modest conformational flexing within cyclobutenes conveniently permits attainment of the necessary stereoalignment from either  $\pi$  face.

(9) The various mechanistic pathways open to these alcohols will be discussed in detail at a later date.

(10) (a) Schreiber, S. L. *J. Am. Chem. Soc.* **1980**, *102*, 6163. (b) Schreiber, S. L.; Liew, W.-F. *Tetrahedron Lett.* **1983**, *24*, 2363. (c) Schreiber, S. L.; Liew, W.-F. *J. Am. Chem. Soc.* **1985**, *107*, 2980. (d) Schreiber, S. L.; Sammakia, T.; Hulin, B.; Schulte, G. *Ibid.* **1986**, *108*, 2106. (e) Schreiber, S. L.; Hulin, B.; Liew, W.-F. *Tetrahedron* **1986**, *42*, 2945.

Among the many interesting facets of these reactions is their mixed stereochemical course. Whereas the **5**  $\rightarrow$  **6** and **18**  $\rightarrow$  **19** processes involve a trans-antiplanar arrangement of the entering and leaving groups, the conversion of **12** to **14** occurs in syn fashion (as does **2**  $\rightarrow$  **3**). Further, the preceding observations show the  $\text{S}_{\text{N}}'$  displacement of methoxide by enolate ions to proceed with reasonable efficiency and to constitute an exceptionally powerful synthetic method.

**Acknowledgment.** This investigation was supported by NIH Grants CA-12115 (to L.A.P.) and GM-32527 (to S.L.S.).


## An Efficient Silylformylation of Alkynes Catalyzed by $\text{Rh}_4(\text{CO})_{12}$

Isamu Matsuda,\* Akira Ogiso, Susumu Sato, and Yusuke Izumi

Department of Synthetic Chemistry, Faculty of Engineering, Nagoya University  
Chikusa, Nagoya 464-01, Japan

Received December 6, 1988

A combination of transition-metal complexes and hydrosilanes is a useful tool for highly selective transformations of alkenes, alkynes, and carbonyl groups.<sup>1</sup> In particular, cobalt-catalyzed incorporation of carbon monoxide is interesting as a unique method for the homologation of alkenes, aldehydes, and cyclic ethers, in which  $\text{R}_3\text{Si-Co}(\text{CO})_4$  plays an important role.<sup>2</sup> On the other hand, while hydroformylation is one of the most useful processes for the practical synthesis of aldehydes,<sup>3</sup> no analogous operation toward acetylenic compounds has been developed because of the concomitant formation of undesired products.<sup>4</sup> An excellent precedent is the selective acylation of 1-alkynes at the terminal carbon with the assistance of  $\text{Rh}_4(\text{CO})_{12}$ ,<sup>5</sup> or with a stoichiometric amount of  $\text{R-Mn}(\text{CO})_5$ ,<sup>6</sup> to give  $\alpha,\beta$ -unsaturated enones. A formal silylformylation of 1-alkynes to give 3-silyl-2-alkenals **2'** can represent a useful synthetic approach to these valuable building blocks, due to the ready accessibility and elaboration of alkynes. We have designed a simple formylation reaction of alkynes in the presence of carbon monoxide, employing dimethylphenylsilane ( $\text{Me}_2\text{PhSiH}$ ) instead of  $\text{H}_2$ , and  $\text{Rh}_4(\text{CO})_{12}$  as catalyst and have found a novel and selective formylation pattern of alkynes to give **2**. We report here the scope and limitations of this new catalytic silylformylation of alkynes.



When an alkyne **1** was allowed to react with 1 equiv each of  $\text{Me}_2\text{PhSiH}$  and  $\text{Et}_3\text{N}$ , using a catalytic amount (1 mol %) of  $\text{Rh}_4(\text{CO})_{12}$  in benzene solution under CO (30 kg/cm<sup>2</sup>) at 100 °C

(1) Colvin, E. W. *Silicon in Organic Synthesis*; Butterworths: London, 1981; pp 325-336.

(2) Murai, S.; Sonoda, N. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 837-846.

(3) (a) Pino, P.; Braca, G. *Organic Synthesis via Metal Carbonyls*; Wender, I.; Pino, P., Eds.; Wiley-Interscience: New York, 1977; Vol. 2, pp 41-231. (b) Prueit, R. L. *Advanced Organometallic Chemistry*; Stone, F. G. A., West, R., Eds.; Academic Press: New York, 1979; Vol. 17, pp 1-60. (c) Tkatchenko, I.; *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G. A., Abel, E. W., Eds.; Pergamon Press: Oxford, 1982; Vol. 8, pp 101-223.

(4) Pino, P.; Braca, G. *Organic Synthesis via Metal Carbonyls*; Wender, I.; Pino, P., Eds.; Wiley-Interscience: New York, 1977; Vol. 2, pp 419-516.

(5) Hong, P.; Mise, T.; Yamazaki, H. *J. Organomet. Chem.* **1987**, *334*, 129-140.

(6) Deshong, P.; Sidder, D. R.; Rybczynski, P. J.; Slough, G. A.; Rheingold, A. L. *J. Am. Chem. Soc.* **1988**, *110*, 2575-2585.

(7) Otera, J.; Mandai, T.; Shiba, M.; Saito, T.; Shimohata, K.; Takemori, K.; Kawasaki, Y. *Organometallics* **1983**, *2*, 332-336.