

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11436710>

Ruthenium–Mediated Cycloaromatization of Acyclic Enediynes and Dienynes at Ambient Temperature

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MAY 2002

Impact Factor: 12.11 · DOI: 10.1021/ja017873t · Source: PubMed

CITATIONS

46

READS

27

3 AUTHORS, INCLUDING:



Joseph M O'Connor

University of California, San Diego

101 PUBLICATIONS 2,071 CITATIONS

SEE PROFILE

Ruthenium-Mediated Cycloaromatization of Acyclic Enediynes and Dienynes at Ambient Temperature


Joseph M. O'Connor,* Seth J. Friese, and Mark Tichenor

Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, California 92093

Received December 23, 2001


Potential applications of the Bergman¹ cycloaromatization in synthetic² and medicinal chemistry³ have stimulated research into methods for promoting enediyne cycloaromatization under mild conditions.^{4–7} This is particularly desirable in the case of non-strained acyclic 3-ene-1,5-diyne substrates which often require elevated temperatures for onset of thermal cycloaromatization. Of particular note in this regard is Finn's report on the conversion of diyne **1** to vinylidene **2**, which undergoes cycloaromatization at 100 °C in the presence of 1,4-cyclohexadiene to give **3** (Scheme 1).^{6a} When 1,4-cyclohexadiene-*d*₄ was used as a D-atom donor, 30% deuterium incorporation was observed at both the C4- and C5-hydrogen positions of the benz[*e*]indene product, leading the authors to propose a Myers–Saito-type mechanism⁷ involving the diradical intermediate **I**.

Scheme 1



We previously reported that $[(\eta^5\text{-C}_5\text{Me}_5)\text{Ru}(\text{CH}_3\text{CN})_3]\text{OTf}$ (**4**)⁸ mediates the room-temperature cycloaromatization of *strained-ring* benzoenediynes, such as the conversion of enediyne **5** to **6** (Scheme 2).⁹ We were disappointed to find that the strain-free acyclic enediyne **7** failed to cyclize, but instead gave only the uncyclized η^6 -arene complex **8**.


Scheme 2



We now report that complex **4** does indeed mediate the cycloaromatization reaction of *acyclic*¹⁰ enediynes, as well as the cycloaromatization of conjugated dienynes.


When a THF-*d*₈ solution of **9-TMS** and 1,4-cyclohexadiene was heated at 150 °C for 14 d, there was no evidence for the formation of a dihydroindene derivative by ¹H NMR spectroscopic analysis of the sample. However, reaction of **9-TMS** (103 mg, 0.4 mmol) and the ruthenium complex **4** (0.4 mmol) in THF (10 mL) at 100 °C led to isolation of the η^6 -(2,3-dihydro-1H-inden-5-yl)-trimethylsilane] complex **10** in 69% yield (Scheme 3). Under similar conditions the enediynes with bulky alkyne substituents, **9-Bu'**, **9-TIPS**, and **9-Me**, failed to undergo a detectable (by NMR spectroscopy) cycloaromatization reaction.

Scheme 3



The work of Finn suggested that loss of a TMS substituent in the conversion of **9-TMS** to **10** may have generated a terminal alkyne capable of cycloaromatization via a vinylidene mechanism. We therefore examined the reaction of 1-ethynyl-2-(1-propynyl)-cyclopentene (**11**; 0.048 mmol, 4.8 mM) with **4** (0.047 mmol) in THF solvent and observed the *room-temperature* formation of **12** in 92% isolated yield (Scheme 4). When the reaction was carried

Scheme 4




out in THF-*d*₈ and monitored by ¹H NMR spectroscopy, the deuterium-enriched arene **12-d**₂ was formed within 10 min at room temperature. Integration of the ¹H NMR signals for **12-d**₂ indicated ca. 90% deuterium enrichment at both the C4- and C7-hydrogen positions. Furthermore, reaction of the deuterium-labeled analogue

* To whom correspondence should be addressed. E-mail: jmoconno@ucsd.edu.

11-*d*₁ (83% deuterium enrichment at the ethynyl hydrogen) and **4** led to the formation of **12-*d*₁** with 63% deuterium incorporation at the C6-hydrogen position and no isotopic enrichment at either the C4- or C7-hydrogen sites.


These isotopic labeling results are consistent with the formation of a *p*-benzynes intermediate, possibly arene complex **II**, in the conversion of **11** to **12**. The absence of deuterium incorporation at the C7-hydrogen position of **12-*d*₁** rules out a vinylidene-based mechanism proceeding via diradical **III** (Chart 1).

Chart 1




Encouraged by the results with enediyne **11**, the reactions of internal enediynes **13-Me**, **13-*Pr*^{*i*}**, **13-Bu^{*i*}**, and **14** with **4** were examined (Scheme 5). In all cases, a rapid reaction with **4** occurred within minutes at room temperature to give good yields of the η^6 -dihydroindene complexes **15** and **16**.

Scheme 5




The substantial driving force exhibited by the [Cp**Ru*] cation for enediyne cycloaromatization suggested that conjugated dienes may be susceptible to a ruthenium-mediated Hopf cyclization.¹¹ As shown in Scheme 6, the Hopf cyclization involves the high temperature (200–250 °C) conversion of hexadienynes, **17**, to benzene derivatives **18**. As is the case for the thermal Bergman cycloaromatization, the Hopf cyclization proceeds via a cyclic intermediate of diradical character (**IV**).

Scheme 6



In a preliminary experiment, treatment of dienyne **19** (0.029 mmol) with **4** (0.029 mmol) in THF-*d*₈ solvent (0.23 mL) at room temperature led to the formation of the η^6 -dihydroindene complex **20** within 10 min (52% NMR yield; Scheme 7). In contrast to the

Scheme 7



reactions of **4** with enediynes in THF-*d*₈, there is no deuterium enrichment (<5% by ¹H NMR analysis) at any dihydroindene hydrogen position in **20**. The location of the *n*-propyl substituent at C5 excludes a vinylidene intermediate in the formation of **20**.^{12,13}

The lack of significant D-atom abstraction from THF-*d*₈ and the rapid rate of reaction suggested that CDCl₃ may also serve as a solvent.¹⁴ Indeed, reaction of **19** (0.023 mmol) and **4** (0.023 mmol) in CDCl₃ (0.44 mL) at room temperature (50 min) resulted in the formation of **20** in 96% NMR yield, with no significant deuterium enrichment. By analogy with intermediates **II** and **IV**, the Ru(III) cyclohexadienyl cation **V** must be considered as a *potential* intermediate in the conversion of **19** to **20**. However, intermediate **V** requires a H-atom transfer, possibly intramolecular, which is rapid relative to the rate of D-atom abstraction from solvent.

Finally, we note that the lack of cycloaromatization of enediyne **7** may be the result of a more rapid Ru-arene formation, which is not possible with the cyclopentene substrates reported herein. Studies are currently underway to determine the detailed mechanism and scope of these new metal-mediated cycloaromatization reactions.

Acknowledgment. We gratefully acknowledge financial support from the National Science Foundation (CHE-9970480) and the University of California Cancer Coordinating Committee. S.J.F. acknowledges the award of a 2000 Urey Fellowship.

Supporting Information Available: Characterization data for compounds **13–16**, **19**, **20** and tables of crystallographic data for **10** and **15-*Pr*^{*i*}** (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Bergman, R. G. *Acc. Chem. Res.* **1973**, *6*, 25. (b) Lockhart, T. P.; Bergman, R. G. *J. Am. Chem. Soc.* **1981**, *103*, 4091.
- For leading references: (a) Chen, X.; Tolbert, L. M.; Hess, D. W.; Henderson, C. *Macromolecules* **2001**, *34*, 4104. (b) Bowles, D. M.; Palmer, G. J.; Landis, C. A.; Scott, J. L.; Anthony, J. E. *Tetrahedron* **2001**, *57*, 3753. (c) Shah, H. V.; Brittain, S. T.; Huang, Q.; Hwu, S. J.; Whitesides, G. M.; Smith, D. W., Jr. *Chem. Mater.* **1999**, *11*, 2623. (d) Grissom, J. W.; Gunawardena, G. U.; Klingberg, D.; Huang, D. H. *Tetrahedron* **1996**, *52*, 6453.
- For leading references: (a) Xi, Z.; Goldberg, I. DNA damaging enediyne compounds. In *Comprehensive Natural Products Chemistry*; Barton, D. H. R.; Nakanishi, K., Eds.; Pergamon: Oxford, 1999; Vol 7, p 553. (b) Liu, W.; Shen, B. *Antimicrob. Agents Chemother.* **2000**, *44*, 382. (c) Smith, A. L.; Nicolaou, K. C. *J. Med. Chem.* **1996**, *39*, 2103.
- (a) Jones, G. B.; Warner, P. M. *J. Am. Chem. Soc.* **2001**, *123*, 2134. (b) König, B.; Pitsch, W.; Klein, M.; Vasold, R.; Prall, M.; Schreiner, P. R. *J. Org. Chem.* **2001**, *66*, 1742 and references therein.
- (a) Rawat, D. S.; Zaleski, J. M. *J. Am. Chem. Soc.* **2001**, *123*, 9675. (b) Schmitt, E. W.; Huffman, J. C.; Zaleski, J. M. *Chem. Commun.* **2001**, 167. (c) Chandra, T.; Pink, M.; Zaleski, J. M. *Inorg. Chem.* **2001**, *40*, 5878. (d) Benites, P. J.; Rawat, D. S.; Zaleski, J. M. *J. Am. Chem. Soc.* **2000**, *122*, 7208. (e) Coalter, N. L.; Concolino, T. E.; Streib, W. E.; Hughes, C. G.; Rheingold, A. L.; Zaleski, J. M. *J. Am. Chem. Soc.* **2000**, *122*, 3112. (f) Rawat, D. S.; Zaleski, J. M. *Chem. Commun.* **2000**, 2493. (g) König, B. *Eur. J. Org. Chem.* **2000**, 381. (h) Basak, A.; Shain, J. C.; Khamrai, U. K.; Rudra, K. R.; Basak, A. *J. Chem. Soc., Perkin Trans. 1* **2000**, 1955. (i) Warner, B. P.; Millar, S. P.; Broene, R. D.; Buchwald, S. L. *Science* **1995**, *269*, 814. (j) König, B.; Hollnagel, H.; Ahrens, B.; Jones, P. G. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 2538.
- (a) Wang, Y. S.; Finn, M. G. *J. Am. Chem. Soc.* **1995**, *117*, 8045. (b) Manabe, T.; Yanagi, S. I.; Ohe, K.; Uemura, S. *Organometallics* **1998**, *17*, 2942.
- (a) Myers, A. G.; Kuo, E. Y.; Finney, N. S. *J. Am. Chem. Soc.* **1989**, *111*, 8057. (b) Saito, K.; Watanabe, T.; Takahashi, K. *Chem. Lett.* **1989**, 2099.
- The [Cp**Ru*] cation is a potent areneophile and aromatization reagent for polyenes: (a) Older, C. M.; Stryker, J. M. *J. Am. Chem. Soc.* **2000**, *122*, 2784. (b) Chaudret, B. *Bull. Soc. Chim. Fr.* **1995**, *132*, 268. (c) Fagan, P. J.; Ward, M. D.; Calabrese, J. C. *J. Am. Chem. Soc.* **1989**, *111*, 1698.
- O'Connor, J. M.; Lee, L. I.; Gantzel, P.; Rheingold, A. L.; Lam, K. C. *J. Am. Chem. Soc.* **2000**, *122*, 12057.
- Here *acyclic* refers to the structural feature in which the two alkynes are not located within a ring.
- (a) Hopf, H. *Angew. Chem., Int. Ed. Engl.* **1969**, *8*, 680. (b) Zimmermann, G. *Eur. J. Org. Chem.* **2001**, 457.
- Merlic previously observed a ruthenium-catalyzed cycloaromatization of dienyne which proceeds via a metal-vinylidene intermediate: (a) Merlic, C. A.; Pauly, M. E. *J. Am. Chem. Soc.* **1996**, *118*, 11319. (b) Maeyama, K.; Iwasawa, N. *J. Org. Chem.* **1999**, *64*, 1344.
- For a metal-catalyzed conversion of electron-rich aromatic enynes to substituted naphthalenes: Dankwardt, J. W. *Tetrahedron Lett.* **2001**, *42*, 5809.
- Complex **4** decomposes over the course of hours in CHCl₃.

JA017873T