

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231439743>

Evidence for Fe(IV):O in the Molecular Mechanism of Action of the Trioxane Antimalarial Artemisinin

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MAY 1995

Impact Factor: 12.11 · DOI: 10.1021/ja00126a042

CITATIONS

116

READS

43

4 AUTHORS, INCLUDING:



Poonsakdi Ploypradith

Chulabhorn Research Institute

61 PUBLICATIONS 1,364 CITATIONS

SEE PROFILE

Evidence for Fe(IV)=O in the Molecular Mechanism of Action of the Trioxane Antimalarial Artemisinin

Gary H. Posner,* Jared N. Cumming,
Poonsakdi Ploypradith, and Chang Ho Oh†

Department of Chemistry, The Johns Hopkins University
Baltimore, Maryland 21218

Received March 10, 1995

Among the 300–500 million people worldwide who are currently infected with malaria, about 2 million deaths, many of them children, occur each year.¹ As malaria parasites develop ever-increasing multidrug resistance to traditional alkaloidal antimalarial drugs,² artemisinin (qinghaosu, **1**), a non-alkaloidal endoperoxide natural product discovered in China, and related 1,2,4-trioxanes are increasingly being used for effective chemotherapy of malaria.^{3,4} These organic peroxides, causing oxidative stress⁵ to malaria parasites, apparently are reduced by the iron-rich parasites to form cytotoxic radical intermediates.⁶ Using an oxygen-18-labeled trioxane⁷ and some mechanism-based synthetic analogs,⁸ we have shown that a carbon-centered radical, formed from an oxy radical via an intramolecular 1,5-hydrogen atom shift, is important for antimalarial activity.⁹ We now report several kinds of evidence supporting the intermediacy of a high-valent, non-heme, iron–oxo species resembling that characteristic of monooxygenase metalloenzymes and known to cause oxidative damage to biological macromolecules.^{10,11} It is proposed that such a high-valent iron–oxo species is formed via homolytic oxygen–carbon bond scission¹² from a β -ferryloxyethyl radical and that a highly electrophilic epoxide (e.g., **5**, Scheme 1), a potent alkylating agent,¹³ also is formed. A molecular mechanism representing these transformations and the types of evidence we have accumulated for a high-valent iron–oxo intermediate are summarized in Scheme 1. This molecular mechanism represents the first report of generation of Fe(IV)=O during ferrous ion activation of a 1,2,4-trioxane rather than, as usual, by heme protein (i.e., cytochrome P₄₅₀) or metalloporphyrin model compounds activating dioxygen or hydrogen peroxide.^{10,11}

† On leave from the Department of Chemistry, Inje University, Kimhae 621-749, Korea.

(1) TDR News (News from the WHO Division of Control of Tropical Diseases), 1994, 46, 5.

(2) (a) Peters, W. *Chemotherapy and Drug Resistance in Malaria*, 2nd ed.; Academic Press: London, 1987. (b) Hien, T. T.; White, N. J. *Lancet* 1993, 341, 603.

(3) For reviews, see: (a) Klayman, D. L. *Science* 1985, 228, 1049. (b) White, N. J. *Trans. R. Soc. Trop. Med. Hyg.* 1994, 88 (Suppl. 1), 1. (c) Zhou, W.-S.; Xu, X.-X. *Acc. Chem. Res.* 1994, 27, 211. (d) Jung, M. *Curr. Med. Chem.* 1994, 1, 45.

(4) For a recent pharmacokinetic study of artemisinin used to treat malaria in humans, see: Duc, D. D.; DeVries, P. J.; Khanh, N. X.; Binh, L. N.; Kager, D. A.; Van Bostel, C. J. *Am. J. Trop. Med. Hyg.* 1994, 51, 785.

(5) (a) Greenspan, H. C.; Aruoma, O. I. *Immunol. Today* 1994, 15, 209. (b) Harel, R.; Marva, E.; Chevion, M.; Golenser, J. *Free Rad. Res. Commun.* 1993, 18, 279. (c) Gutteridge, J. M. C. In *Oxidative Damage and Repair*; Davies, K. J. A., Ed.; Pergamon Press: New York, 1991; pp 355–363. (d) Aruoma, O. I., Ed. *Free Radicals in Tropical Diseases*; Harwood Academic Publishers: Chur, Switzerland, 1993.


(6) For the first reports on the importance of iron in triggering the cytotoxic effects of trioxanes, see: (a) Meshnick, S. R.; Thomas, A.; Ranz, C. X.; Pan, H. Z. *Mol. Biol. Parasit.* 1991, 49, 181. (b) Meshnick, S. R.; Yang, Y.-Z.; Lima, V.; Kuypers, F.; Kamchonwongpaisan, S.; Yathavong, Y. *Antimicrob. Agents Chemother.* 1993, 37, 1108. (c) Hong, Y.-L.; Yang, Y.-Z.; Meshnick, S. R. *Mol. Biochem. Parasitol.* 1994, 63, 121.

(7) Posner, G. H.; Oh, C. H. *J. Am. Chem. Soc.* 1992, 114, 8328.

(8) Posner, G. H.; Oh, C. H.; Wang, D.; Gerena, L.; Milhous, W. K.; Meshnick, S. R.; Asawamahasadka, W. *J. Med. Chem.* 1994, 37, 1256. See, also: Avery, M. A.; Fan, P.; Karle, J. M.; Miller, R.; Goins, K. *Tetrahedron Lett.* In press.

(9) For recent reports on the cytotoxic effects of free radicals, see: (a) Gutteridge, J. M. C. *Free Rad. Res. Commun.* 1993, 19, 141. (b) Hiramoto, K.; Johkoh, H.; Sako, K.-I.; Kikugawa, K. *Free Rad. Res. Commun.* 1993, 19, 323.

Scheme 1



Evidence for a high-valent iron–oxo intermediate is provided by reactions characteristic of such a species, as follows: (1) hexamethyl Dewar benzene was rearranged into hexamethylbenzene (in 40% yield),^{14,15} while the amount of C₄-hydroxylated artemisinin product **6** was diminished from 15–20% to 5–7% (Table 1); (2) methyl phenyl sulfide was oxygenated to the corresponding sulfoxide; and (3) tetralin (1,2,3,4-tetrahydronaphthalene) was oxidized into hydroxytetralin.¹⁶ Three

(10) For excellent overviews with leading references, see: (a) Valentine, J. S.; Nam, W.; Ho, R. Y. N. In *The Activation of Dioxygen and Homogeneous Catalytic Oxidation*; Barton, D. H. R., Martell, A. E., Sawyer, D. T., Eds.; Plenum Press: New York, 1993; p 183. (b) Ortiz de Montellano, P. R. In *The Activation of Dioxygen and Homogeneous Catalytic Oxidation*; Barton, D. H. R., Martell, A. E., Sawyer, D. T., Eds.; Plenum Press: New York, 1993; p 257. (c) Leising, R. A.; Kojima, T.; Que, L., Jr. In *The Activation of Dioxygen and Homogeneous Catalytic Oxidation*; Barton, D. H. R., Martell, A. E., Sawyer, D. T., Eds.; Plenum Press: New York, 1993; p 321. (d) Arasasingham, R. D.; Bruce, T. C. In *The Activation of Dioxygen and Homogeneous Catalytic Oxidation*; Barton, D. H. R., Martell, A. E., Sawyer, D. T., Eds.; Plenum Press: New York, 1993; p 147. (e) Mansuy, D. In *The Activation of Dioxygen and Homogeneous Catalytic Oxidation*; Barton, D. H. R., Martell, A. E., Sawyer, D. T., Eds.; Plenum Press: New York, 1993; p 347. (f) Feig, A. L.; Lippard, S. J. *Chem. Rev.* 1994, 94, 759.

(11) For more recent studies, see: (a) Dexter, A. F.; Hager, L. P. *J. Am. Chem. Soc.* 1995, 117, 817. (b) Minisci, F.; Fontana, F.; Araneo, S.; Recupero, F.; Banfi, S.; Quici, S. *J. Am. Chem. Soc.* 1995, 117, 226. (c) Tian, Z.-Q.; Richards, J. L.; Traylor, T. G. *J. Am. Chem. Soc.* 1995, 117, 21. (d) Harris, D. L.; Loew, G. H. *J. Am. Chem. Soc.* 1994, 116, 11671. (e) Akhtar, M.; Wright, J. N.; Shyadehi, A.-Z.; Rochicaud, P. *Pure Appl. Chem.* 1994, 66, 2387.

(12) (a) Kochi, J. K.; Singleton, D. M. *J. Am. Chem. Soc.* 1968, 90, 1582. (b) For leading references to related heteroatom β -scissions, see: Santoyo-Gonzalez, F.; Calvo-Flores, F. G.; Herando-Mateo, F.; Garcia-Mendoza, P.; Isac-Garcia, J.; Perez-Alvarez, M. D. *Synlett* 1994, 454. (c) For the absence of β -scission in β -acyloxyalkyl radicals generated using tin hydride, see: Foubelo, F.; Lhoret, F.; Yus, M. *Tetrahedron* 1994, 50, 5131.

(13) (a) For coupling reactions of structurally analogous 1,2-anhydro-pyranose systems as reactive glycosyl donors, see: Gordon, D. M.; Danishefsky, S. J. *J. Org. Chem.* 1994, 59, 3713 and references therein. (b) For synthesis of analogous 1,2-anhydrofuranose systems, see: Du, Y.; Kong, F. *Tetrahedron Lett.* 1995, 36, 427 and references therein.

(14) Traylor, T. G.; Mikszals, A. R. *J. Am. Chem. Soc.* 1987, 109, 2770.