

# Bond Ordering Relation. Incompatibility between Bonds, Essential Single/Double Bonds, and the Nonexistence of Kekulé Structures in Molecular Subgraphs

Tetsuo Morikawa

Chemistry Department, Joetsu University of Education, Yamayashiki, Joetsu, Niigata Prefecture 943, Japan

Received November 21, 1995<sup>®</sup>

An ordering relation between single and/or double bonds, such that one bond gives an order to another, on a set of Kekulé structures (perfect matchings) in hydrogen-suppressed molecular graphs is defined. By use of this binary relation, lemmas relating to both incompatibility between bonds and essential single/double bonds can be described. It is thus possible to estimate the nonexistence of Kekulé structures in a given molecular subgraph.

## 1. INTRODUCTION

We follow graphic terminology of ref 1. Let  $v_i$  and  $e_i$  be a vertex and an edge of a hydrogen-suppressed molecular graph  $G$ ; in a path  $v_i e_i v_{i+1}$ ,  $e_i$  joins  $v_i$  and  $v_{i+1}$  together. When a double bond connects  $v_i$  and  $v_{i+1}$  in a Kekulé structure (a perfect matching) of  $G$ , the bond is denoted by  $d_i$ ; and when a single bond, by  $s_i$ . We use  $b_i$  to indicate  $s_i$  or  $d_i$ ;  $\bar{b}_i$  represents  $d_i$  if  $b_i = s_i$ ;  $\bar{b}_i$  represents  $s_i$  if  $b_i = d_i$ ;  $\bar{\bar{b}}_i = b_i$ ; also refer to Table 1.

Let  $K\{G\}$  be the number of Kekulé structures in  $G$ , and let  $K\{b_i, r\}$  be the number of Kekulé structures containing  $b_i$  in  $G$ . Here  $\{r\}$  is an abbreviation for the rest of  $G$ . Two identity equations in Kekulé structure counting<sup>1</sup> are thus expressed in the present notation as

$$K\{e_i, r\} = K\{b_i, r\} + K\{\bar{b}_i, r\} \quad (1)$$

$$K\{e_i, e_j, r\} = K\{b_i, b_j, r\} + K\{b_i, \bar{b}_j, r\} + K\{\bar{b}_i, b_j, r\} + K\{\bar{b}_i, \bar{b}_j, r\} (e_i \neq e_j) \quad (2)$$

This note is concerned with local properties relating to  $e_i$ ,  $b_i$ , and  $\bar{b}_i$  in Kekulé structures.

It is a well-known fact in Kekulé structure counting that the selection of a bond  $b_i$  in paths and cycles uniquely determines the local conjugated structures containing this  $b_i$ . In a hexagon (benzene), for example, a bond  $b_i$  fixes all the other bonds in a Kekulé structure. This propagation of bonds can be interpreted as an ordering relation in mathematics. An edge  $e_i$  between  $v_i$  and  $v_{i+1}$  in polyhexes is called *forcing*<sup>2,3</sup> if it decides Kekulé structures. We first introduce the bond ordering as a mathematical binary relation  $\leq$ , and secondly define both incompatibility between bonds and essential single/double bonds by using this binary relation. We shall prove lemmas relating these concepts, and, as a result, establish two theorems on the nonexistence of Kekulé structures in hydrogen-suppressed molecular subgraphs.

## 2. BOND ORDERING RELATION

We begin by defining a binary relation  $b_i \leq b_j$ , and read it as “ $b_i$  precedes or equals  $b_j$ ”. Notice that a Kekulé structure contains only one of  $b_i$  and  $\bar{b}_i$ .

**Table 1.** Glossary of Symbols

$b_i$	$s_i$ or $d_i$
$\bar{b}_i$	$s_i$ if $b_i = d_i$ , or $d_i$ if $b_i = s_i$
$d_i$	double bond, connecting $v_i$ and $v_{i+1}$ in Kekulé structures
$e_i$	edge, connecting $v_i$ and $v_{i+1}$ in $G$
$G$	hydrogen-suppressed molecular graph
$H, H_i$	subgraph of $G$
$K\{b_i, b_j, r\}$	number of Kekulé structures with $b_i, b_j$ in $G$
$K\{G\}$	number of Kekulé structures in $G$
$q$	abbreviation for graphic part such that $K\{H, q\} > 0$
$r$	abbreviation for the rest of $G$
$s_i$	single bond, connecting $v_i$ and $v_{i+1}$ in Kekulé structures
$t$	abbreviation for the rest of $H$
$v_i$	vertex in $G$
$\rightarrow$	imply (implies)
$\leftarrow$	inverse of $\rightarrow$
$\leftrightarrow$	$\rightarrow$ and $\leftarrow$
$\leq$	precede or equal (precedes or equals)


**Definition 1 (Bond Ordering).** A relation  $\leq$  between  $b_i$  and  $b_j$  is written as  $b_i \leq b_j$ , when (1), and when either (2-1) or (2-2). (1) There is a Kekulé structure with a bond  $b_i$ . (2-1) ( $e_i = e_j$ ) Every Kekulé structure containing  $b_i$  becomes a Kekulé structure containing  $\bar{b}_i$  by the replacement of  $b_i$  with  $\bar{b}_i$ . (2-2) ( $e_i \neq e_j$ ) Every Kekulé structure containing  $b_i$  has a bond  $b_j$ .

The restriction  $K\{G\} > 0$  is an implicit assumption in Definition 1. It is easy to extend this assumption to  $K\{G\} \geq 0$ . Let  $H = \{e_i, e_j, t\}$  be a subgraph of  $G = \{H, r\}$ . In a manner similar to Definition 1, it is possible to define  $b_i \leq b_j$  in  $H$  if there is graphic part  $\{q\}$  such that  $K\{H, q\} > 0$ , because  $\{H, q\}$  can be chosen as a hydrogen-suppressed molecular graph. This extension makes it possible for us to estimate whether or not a given  $G$  is non-Kekuléan by means of  $\leq$ . A bond ordering relation  $\leq$  hereafter means the one for such  $H$ .

Clearly the relation  $\leq$  satisfies  $b_i \leq b_i$  (reflexivity). If  $b_i \leq b_j$ , and if  $b_j \leq b_k$ , then  $b_i \leq b_k$  (transitivity). In the chemical meaning, the ordering relation  $b_i \leq b_j$  is not antisymmetric; i.e.,  $b_i$  is not equivalent to  $b_j$ , even if  $b_i \leq b_j$  and  $b_j \leq b_i$ .

Let us define a binary relation  $\equiv$  by  $b_i \equiv b_j$  if and only if  $b_i \leq b_j$  and  $b_j \leq b_i$ . This relation  $\equiv$  has, needless to say, no valid meaning of chemistry but is a proper definition from the point of view of Kekulé structure counting. We then have  $b_i \leq b_i \rightarrow b_i \equiv b_i$  (reflexivity);  $b_i \equiv b_j$  and  $b_j \equiv b_k \rightarrow b_i \leq b_j$ ,  $b_j \leq b_i$ ,  $b_j \leq b_k$  and  $b_k \leq b_j \rightarrow b_i \leq b_k$  and  $b_k \leq b_i \rightarrow b_i \equiv b_k$  (transitivity);  $b_i \equiv b_j \rightarrow b_i \leq b_j$  and  $b_j \leq b_i \rightarrow b_j \equiv b_i$ .

<sup>®</sup> Abstract published in *Advance ACS Abstracts*, March 1, 1996.



**Figure 1.** All 10 hexagonal subgraphs of  $G$ , each of which has at least one  $b_i$  that precedes or equals all the bonds in the hexagon.  $r$  indicates the rest of  $G$ . Mirror images of the subgraph about the vertical line and/or the horizontal line are all omitted.

$\leq b_i$  and  $b_i \leq b_j \rightarrow b_j \equiv b_i$  (symmetry); namely,  $\equiv$  is an equivalence relation. On a set of  $\{b_i\}$  divided by  $\equiv$ , i.e., on the quotient set  $\{b_i\}/\equiv$ , a binary relation  $\leq^*$  is defined by  $C(b_i) \leq^* C(b_j) \leftrightarrow b_i \leq b_j$ , where  $C(b_i)$  is a class containing  $b_i$  in the quotient set. Clearly  $\leq^*$  satisfies reflexivity and transitivity, because  $\leq$  satisfies reflexivity and transitivity. Suppose that  $C(b_i) \leq^* C(b_j)$  and  $C(b_j) \leq^* C(b_i)$ , then,  $b_i \leq b_j$  and  $b_j \leq b_i \rightarrow b_i \equiv b_j$ ; therefore,  $C(b_i) = C(b_j)$ ; namely, the antisymmetry for  $\leq^*$  holds; i.e.,  $\leq^*$  is a partial ordering relation.<sup>5</sup> The failure of antisymmetry is illustrated by two subpolyhexes (called *alternate cycles* below) in the last column of Figure 1. We use only  $\leq$  hereafter.

Definition 1 leads to Lemmas 1 and 2.

**Lemma 1.** ( $e_i = e_j$ )  $b_i \leq b_j \leftrightarrow K\{b_i, r\} \leq K\{b_j, r\}$ .

**Lemma 2.** ( $e_i \neq e_j$ )  $b_i \leq b_j \rightarrow K\{b_i, e_j, r\} = K\{b_i, b_j, r\} \leq K\{e_i, b_j, r\} \leftrightarrow K\{e_i, b_j, r\} = K\{b_i, \bar{b}_j, r\} \leq K\{b_i, e_j, r\}$ .

Note the direction of arrows in Lemmas 1 and 2,<sup>4</sup> and  $e_i \neq e_j$  means  $b_j \neq b_i, \bar{b}_i$ . It is plain that  $b_j \leq \bar{b}_k$  implies  $K\{e_j, b_k, r\} \leq K\{b_j, e_k, r\}$ . Hence we get Lemma 3.

**Lemma 3.**  $b_i \leq b_k$  and  $b_j \leq \bar{b}_k$  ( $e_i \neq e_j$ )  $\rightarrow K\{b_i, e_j, r\} \leq K\{e_i, \bar{b}_j, r\}$ .

The application of Lemma 2 to a set of bond ordering relations,  $b_i \leq b_j$ ,  $b_i \leq b_k$ , ..., and  $b_i \leq b_m$ , gives

$$K\{b_i, e_j, e_k, \dots, e_m, r\} = K\{b_i, b_j, b_k, \dots, b_m, r\}$$

and

$$K\{\bar{b}_i, \bar{b}_j, \bar{b}_k, \dots, \bar{b}_m, r\} \leq K\{\bar{b}_i, e_j, e_k, \dots, e_m, r\}$$

Using Lemma 1, we thus have Lemma 4.

**Lemma 4.** A set of bond ordering relations,  $b_i \leq b_j$ ,  $b_i \leq b_k$ , ..., and  $b_i \leq b_m$  ( $e_i \neq e_j, e_k, \dots, e_m$ ), such that


$$K\{b_i, b_j, b_k, \dots, b_m, r\} = K\{\bar{b}_i, \bar{b}_j, \bar{b}_k, \dots, \bar{b}_m, r\}$$

is given. Then  $b_i \leq \bar{b}_i$ .

Figure 1 shows all 10 hexagonal subgraphs, each of which fulfills all the sufficient conditions of Lemma 4; i.e.,  $b_i \leq \bar{b}_i$  for each hexagonal subgraph.

Lemma 1 and eq 1 yield the upper and lower bonds for  $K\{e_i, r\}$ .

**Lemma 5.** If  $b_i \leq \bar{b}_i$ , then  $2K\{b_i, r\} \leq K\{e_i, r\} \leq 2K\{\bar{b}_i, r\}$ . Repeating three times the combination of Lemmas 4 and 5 and the elimination of single bonds, we can derive the subgraph (right) from the polyhex subgraph (left) in Figure 2, where



**Figure 2.** A subgraph (polyhex, left) is reducible to another subgraph (right) by means of Lemmas 4 and 5. Three edges, indicated by  $i$ ,  $j$ , and  $k$ , of the subgraphs (left and right) are all essential single bonds for any  $\{r\}$ .

$K\{\text{given subgraph}(\text{left}), r\} \leq$

$$2^3 K\{\text{reduced subgraph}(\text{right}), r\} \quad (3)$$

The right-hand side of this inequality will be estimated in the last section.

### 3. INCOMPATIBILITY BETWEEN BONDS

Let us consider, as an example, a path  $v_1 e_1 v_2 e_2 v_3 e_3 \dots v_{2k} e_{2k} v_{2k+1}$  ( $k \geq 1$ ) in  $G$  such that

$$s_1 \leq d_2 \leq s_3 \leq \dots \leq s_{2k-1} \leq d_{2k}$$

Such a path may be called *alternate*,<sup>6</sup> because single and double bonds appear one after the other. A path beginning and ending at vertices with degree 3 in the subgraph (right) of Figure 2 is an example of alternate paths. For an alternate path, we get  $s_1 \leq d_{2k}$  and  $d_1 \geq s_{2k}$ , then Lemma 2 gives

$$K\{s_1, v_2 e_2 v_3 e_3 \dots v_{2k}, s_{2k}, r\} = 0$$

In other words, we can say that  $s_1$  is incompatible with  $s_{2k}$ . A pair of bonds,  $d_k$  and  $d_{k+1}$ , where the bonds are both incident with a common vertex, is another example; clearly  $K\{d_k, d_{k+1}, r\} = 0$ . In general, we define the incompatibility between bonds by the following.

**Definition 2 (Incompatibility).** Two distinct bonds are said to be incompatible (with each other) when they both are in no Kekulé structure.

A mathematical expression of incompatibility for  $G$  such that  $K\{G\} \geq 0$  is given by the following lemma.

**Lemma 6.** A subgraph  $H = \{e_i, e_j, t\}$  ( $e_i \neq e_j$ ) of  $G = \{H, r\}$ , such that  $K\{H, q\} > 0$  is given. If  $K\{b_i, b_j, t, q\} = 0$ , then  $b_i$  and  $b_j$  are incompatible in  $G$ ; if  $b_i$  and  $b_j$  are incompatible in  $G$ , then  $K\{b_i, b_j, t, q\} = 0$ .

Since a Kekulé structure contains only one of  $b_i$  and  $\bar{b}_i$ , we may say as a special case ( $e_i = e_j$ ) that  $b_i$  and  $\bar{b}_i$  are incompatible. Lemma 2 suggests  $K\{b_i, \bar{b}_j, r\} = 0$  for  $e_i \neq e_j$ . Hence using Lemma 6, we have the following.

**Lemma 7.** If  $b_i \leq b_j$ , then  $b_i$  is incompatible with  $\bar{b}_j$ .

Lemma 7 and eqs 1 and 2 suggest the following.

**Lemma 8.** If  $b_i \leq b_j$  and  $\bar{b}_i \leq b_j$  ( $e_i \neq e_j$ ), then  $K\{e_i, e_j, r\} = K\{e_i, b_j, r\}$ .

Two conditions,  $b_i \leq b_k$  and  $b_j \leq b_l$  ( $e_i \neq e_j$ ), for Lemmas 1 and 2, give

$$K\{b_i, b_j, r\} \leq K\{b_k, b_j, r\} \leq K\{b_k, b_l, r\}$$

Hence we obtain the following.

**Lemma 9.** If  $b_i \leq b_k$  and  $b_j \leq b_l$  ( $e_i \neq e_j$ ), and if  $b_k$  is incompatible with  $b_l$ , then  $b_i$  is incompatible with  $b_j$ . This lemma is useful for determining whether  $b_i$  and  $b_j$  are incompatible in a subgraph  $H$  of  $G$ . It is clear that if the  $K$

value in the right-hand side of  $\leq$  is equal to zero, then that in the left-hand side also equals zero. We can thus state the following.

**Lemma 10.** A subgraph  $H$  of  $G$  is given:  $H$  is decomposed into  $n$  subgraphs,  $H_1, H_2, \dots$ , and  $H_n$ , such that

$$0 < K\{H, r\} \leq cK\{H_1, r\}K\{H_2, r\} \dots K\{H_n, r\}$$

$H$  and one  $H_k$  of the subgraphs both have two bonds  $b_i$  and  $b_j$  ( $c$  being a positive number). Then, if  $b_i$  and  $b_j$  are incompatible in a subgraph  $H_k$ , then they are also incompatible in  $H$ .

**Lemma 11.** A subgraph  $H$  of  $G$  is given:  $H$  is decomposed into  $n$  subgraphs,  $H_1, H_2, \dots$ , and  $H_n$ , such that

$$0 < K\{H, r\} \leq c_1K\{H_1, r\} + c_2K\{H_2, r\} + \dots + c_nK\{H_n, r\}$$

$H$  and all the  $n$  subgraphs have two bonds  $b_i$  and  $b_j$  ( $c_k$  with  $k = 1, 2, \dots, n$ , being a positive number). Then, if  $b_i$  and  $b_j$  are incompatible in  $H_k$  for  $k = 1, 2, \dots, n$ , then they are also incompatible in  $H$ .

#### 4. ESSENTIAL SINGLE/DOUBLE BONDS

Certain bonds for a given conjugated molecule are single or double in all the possible Kekulé structures that one can write; such a bond is called an essential single or essential double bond.<sup>7</sup> We define an essential bond by the following definition.

**Definition 3 (Essential).** A bond is essential when it is in every Kekulé structure. Definition 3 for  $G$  such that  $K\{G\} \geq 0$  is rewritten by use of eq 1 as Lemma 12.

**Lemma 12.** A subgraph  $H = \{e_i, t\}$  of  $G = \{H, r\}$ , such that  $K\{H, q\} > 0$ , is given. If  $K\{\{b_i, t\}, q\} = 0$ , then  $\bar{b}_i$  is essential in  $G$ ; if  $b_i$  is essential in  $G$ , then  $K\{\{b_i, t\}, q\} = 0$ .

As an example, let us consider a cycle  $[v_1e_1v_2e_2v_3e_3 \dots v_{2k}e_{2k}]$ , in which the last edge  $e_{2k}$  connects  $v_{2k}$  and  $v_1$ , such that

$$d_1 \leq s_2 \leq d_3 \leq s_4 \leq \dots \leq d_{2k-1} \leq s_{2k} \leq d_1$$

Such a cycle may be called *alternate*,<sup>6</sup> because alternating single and double bonds occurs. We can observe three alternate cycles in the subgraph (right) of Figure 2. For an alternate cycle, we have

$$\begin{aligned} K\{[d_1, v_2e_2v_3e_3 \dots v_{2k}e_{2k}], r\} \\ &= K\{[d_1, s_2, d_3, s_4, \dots, d_{2k-1}, s_{2k}], r\} \\ &= K\{[s_1, d_2, s_3, d_4, \dots, s_{2k-1}, d_{2k}], r\} \\ &= K\{[s_1, v_2e_2v_3e_3 \dots v_{2k}e_{2k}], r\} \end{aligned}$$

This equality suggests that the  $K$  value of an alternate cycle vanishes if an external edge attached to the cycle is double; i.e., every external edge of alternate cycles is single; hence, such a single bond becomes essential if the  $K$  value is positive.

If  $b_i$  is essential, then  $K\{\bar{b}_i, r\} = 0 \leq K\{b_i, r\}$ ; therefore, Lemma 1 suggests the following.

**Lemma 13.** If  $b_i$  is essential, then  $\bar{b}_i \leq b_i$ .

Remembering that  $b_i \leq b_j$  implies  $K\{e_i, \bar{b}_j, r\} \leq K\{\bar{b}_i, e_j, r\}$ , and using Lemmas 1, 2, and 12, we get the following two lemmas.

**Lemma 14.** If  $b_i$ , such that  $b_i \leq b_j$ , is essential, then  $b_j$  is also essential.

**Lemma 15.** If  $\bar{b}_j$ , such that  $b_i \leq b_j$ , is essential, then  $\bar{b}_i$  is also essential.

Easily Lemma 12 and eq 2 give Lemma 16.

**Lemma 16.** If both of  $b_i$  and  $b_j$  ( $e_i \neq e_j$ ) are essential, then

$$K\{e_i, e_j, r\} = K\{b_i, b_j, r\}$$

Two ordering relations,  $b_i \leq b_k$  and  $b_i \leq b_l$ , imply

$$k\{b_i, e_k, e_l, r\} = K\{b_i, b_k, e_l, r\} \leq K\{e_i, b_k, b_l, r\}$$

which suggests Lemma 17 below. Lemma 17 is useful when we determine whether  $b_i$  is essential in  $H$ . This lemma is also applicable to the case when either  $b_i \leq b_j$  and  $b_i \leq \bar{b}_j$  ( $e_i \neq e_j$ ) or  $b_i \leq d_k$  and  $b_i \leq d_{k+1}$  ( $e_i \neq e_k, e_{k+1}$ ), where  $d_k$  and  $d_{k+1}$  are adjacent. Lemma 18 follows at once from repeated use of Lemma 17. A single polygon with an odd number of vertices has the relations  $b_i \leq \bar{b}_i$  and  $\bar{b}_i \leq b_i$ ; for such a single polygon,  $K = 0$ .

**Lemma 17.** If  $b_i \leq b_k$  and  $b_i \leq b_l$  ( $e_i \neq e_k, e_l$ ), and if  $b_k$  is incompatible with  $b_l$ , then  $\bar{b}_i$  is essential.

**Lemma 18.** If  $b_i \leq \bar{b}_j$ ,  $b_i \leq \bar{b}_j$ ,  $\bar{b}_i \leq b_j$  and  $\bar{b}_i \leq \bar{b}_j$  ( $e_i \neq e_j$ ), then  $K\{b_i, r\} = K\{\bar{b}_i, r\} = K\{b_j, r\} = K\{\bar{b}_j, r\} = 0$ .

It is similar to the derivation of Lemmas 10 and 11 to get the following two lemmas.

**Lemma 19.** A subgraph  $H$  of  $G$  is given:  $H$  is decomposed into  $n$  subgraphs,  $H_1, H_2, \dots$ , and  $H_n$ , such that

$$0 < K\{H, r\} \leq cK\{H_1, r\}K\{H_2, r\} \dots K\{H_n, r\}$$

both of  $H$  and one  $H_k$  of the subgraphs have  $b_i$  ( $c$  being a positive number). Then, if  $b_i$  is essential in a subgraph  $H_k$ , then  $b_i$  is also essential in  $H$ .

**Lemma 20.** A subgraph  $H$  of  $G$  is given:  $H$  is decomposed into  $n$  subgraphs,  $H_1, H_2, \dots$ , and  $H_n$ , such that

$$0 < K\{H, r\} \leq c_1K\{H_1, r\} + c_2K\{H_2, r\} + \dots + c_nK\{H_n, r\}$$

$H$  and the subgraphs all have  $b_i$  ( $c_k$  with  $k = 1, 2, \dots, n$ , being a positive number). Then, if  $b_i$  is essential in  $H_k$  for  $k = 1, 2, \dots, n$ , then  $b_i$  is also essential in  $H$ .

#### 5. ALGORITHMS FOR DETERMINING BOND ORDERING RELATIONS

A vertex  $v_k$  of a conjugated path has three different kinds of external bonds that are attached to  $v_k$ ; namely, double, single, and unfixed;  $(=)_k$ ,  $(-)_k$ , and  $( )_k$ , respectively, denote such bonds in this section. An algorithm below tries to construct a sequence, composed of  $b_k$  and  $\bar{b}_k$ , from a given path  $(v_ie_iv_{i+1} \dots v_ke_kv_{k+1} \dots v_je_jv_{j+1})$ , and also tries to fix  $( )_k$  for  $v_k$ .

- (1) Make  $b_i$  ( $d_i$  or  $s_i$ ).
- (2) Set  $i + 1$  to be  $k$ .
- (3) Loop:
  - (a) If  $b_{k-1} = d_{k-1}$ , and
    - i. if  $(=)_k$ , then Failure,
    - ii. if  $(-)_k$ , then make  $s_k$  and set all  $( )_k$  to be  $(-)_k$ ,

- iii. if  $( )_k$ , then make  $s_k$  and set all  $( )_k$  to be  $(-)_k$ ,
- (b) If  $b_{k-1} = s_{k-1}$ , and
  - i. if  $(=)_k$ , then make  $s_k$ ,
  - ii. if  $(-)_k$ , then Failure,
  - iii. if  $( )_k$ , then Failure.
- (c) Increase  $k$  by 1.
- (4) Repeat the loop until either  $k = j$  or Failure.

If no sequence beginning at  $b_i$  and ending at  $b_j$  for a given path is complete, then search for another path. We can conclude that  $b_i \leq b_j$  if there is a sequence from  $b_i$  to  $b_j$ .

When the algorithm above-mentioned is not applicable to a subgraph  $H$  of  $G$ , we have to reduce  $H$  by use of Lemmas 4, 5, 10, 11, 19, and 20. In  $H$ , if one cycle shares only  $s_i$  with another, then the elimination of  $s_i$  from  $H$  means the fusion of the two cycles into one cycle; if  $\{H, r\}$  is a bipartite graph, then the resultant subgraph by the fusion is also a bipartite graph; see Figure 2.

## 6. NONEXISTENCE OF KEKULE STRUCTURES IN MOLECULAR SUBGRAPHS

The argument of the foregoing sections is summarized as two theorems.

**Theorem 1.** A subgraph  $H = \{e_i, t\}$  of  $G = \{H, r\}$  is given. Then  $(1) \leftrightarrow (2) \rightarrow (3)$ :

- (1) Both  $b_i$  and  $\bar{b}_i$  are essential.
- (2)  $b_i$ , such that  $b_i \leq \bar{b}_i$ , is essential.
- (3)  $K\{H, r\} = 0$ .

In Theorem 1, Lemma 13 derives item (2) from item (1); inversely, by use of Lemma 14, item (2) gives item (1). Equation (1) and Lemma 12 derives item (3) from (1).

**Theorem 2.** A subgraph  $H = \{e_i, e_j, t\}$  of  $G = \{H, r\}$  is given ( $e_i \neq e_j$ ). Then one of the following statements implies  $K\{H, r\} = 0$ .

- (1) Both  $b_i$  and  $\bar{b}_j$ , such that  $b_i \leq b_j$ , are essential.
- (2)  $b_i$  is essential, and  $b_i$  and  $b_j$  ( $b_i \leq b_j$ ) are incompatible.
- (3)  $b_i, \bar{b}_i, b_j$  and  $\bar{b}_j$  are all essential.
- (4)  $b_i \leq b_j, b_i \leq \bar{b}_j, \bar{b}_i \leq b_j$  and  $\bar{b}_i \leq \bar{b}_j$ .
- (5)  $\bar{b}_j$ , such that  $\bar{b}_i \leq b_j$  and  $b_i \leq b_j$ , is essential.

In Theorem 2, by means of Lemmas 14 and 15, item (1) leads to item (3). Lemma 7 and eq 2 give the sufficiency of item (2). Item (4) implies item (3) (Lemma 18); eq 2 suggests the sufficiency of item (3). The sufficiency of item (5) is proved by Lemma 8. We can observe that item (1) is satisfied in the subgraph (right) of Figure 2; i.e., the subgraph (right) is non-Kekuléan. Equation 3 therefore suggests that the subgraph (left) of Figure 2 is non-Kekuléan for any  $\{r\}$ .

## REFERENCES AND NOTES

- (1) Trinajstić, N. *Chemical Graph Theory*, 2nd ed.; CRC Press: Boca Raton, 1992.
- (2) Harary, F.; Klein, D. J.; Živković, T. P. Graphical Properties of Polyhexes: Perfect Matching Vector and Forcing. *J. Math. Chem.* **1991**, 6, 295–306.
- (3) Randić, M.; Klein, D. J. *Kekulé Valence Structures Revisited. Innate Degrees of Freedom of Pi-Electron Couplings In Mathematical and Computational Concepts in Chemistry*; Trinajstić, N., Ed.; Ellis Horwood: New York, 1986; Chapter 23.
- (4) Morikawa, T. *Comm. Math. Chem.* **1995**, 32, 147–157. In this note, the statement, “ $d_i \leq d_j$  implies  $s_i \geq s_j$ ”, should be read as “ $d_i \leq d_j$  implies  $K\{s_i, r\} \geq K\{s_j, r\}$ ”.
- (5) Liu, C. L. *Elements of Discrete Mathematics*, 2nd ed.; McGraw-Hill: New York, 1985; Chapter 4.
- (6) Morikawa, T. Upper and Lower Bounds for the Number of Conjugated Patterns in Carbocyclic and Heterocyclic Compounds. *Z. Naturforsch.* **1994**, 49a, 973–976.
- (7) Dewar, M. J. S.; Dougherty, R. C. *The PMO Theory of Organic Chemistry*; Plenum: New York, 1975; p 99.

CI950174D