

On the Relation between the P'/P Index and the Wiener Number[†]

Dejan Plavšić,^{*,‡} Milan Šoškić,[§] Irena Landeka,[⊥] and Nenad Trinajstić[‡]

The Rugjer Bošković Institute, P.O.B. 1016, HR-10001 Zagreb, The Republic of Croatia, and Department of Chemistry, Faculty of Agriculture, and Department of Biochemistry, Faculty of Science, University of Zagreb, HR-10000 Zagreb, The Republic of Croatia

Received May 6, 1996[Ⓢ]

It is shown analytically that the recently introduced P'/P index and the Wiener number are closely related graph-theoretical invariants for connected undirected acyclic structures and cycles.

INTRODUCTION

Molecular structure is the central theme of chemistry.^{1–3} According to the principle of molecular structure, properties and behaviour of molecules follow from their structures. This statement has also been a subject of some criticisms.^{4,5} If one considers nonmetric properties of a molecule, then the molecule can be represented by a (molecular) graph, which is essentially a nonnumerical mathematical object. Measurable properties of a molecule are usually expressed by means of numbers. Hence, to correlate property or activity of a molecule with its topology, one must first convert by an algorithm the information contained in the graph to a numerical characteristic. A scalar numerical descriptor uniquely determined by a molecular graph is named a topological (graph-theoretical) index.^{6,7}

In the past the selection of graph matrices used for deriving of molecular indices was limited to the adjacency matrix A and the distance matrix D .^{8,9} The situation has been changed in the last few years, and quite a few novel graph-theoretical matrices have been proposed.^{10–15} Randić has recently put forward a novel structure - explicit graph matrix P as well as the novel molecular index P'/P derived from it.^{16,17} He also tested the new index by examining the octane numbers of octanes and empirically found a linear relationship between the P'/P index and the Wiener number^{18,19} in octanes.

The representation of a molecule by a single number (topological index) entails a considerable loss of information concerning the molecular structure. In search of new invariants which would improve the graph-theoretical characterization of molecular structure, a great number of indices have been proposed so far.^{20,21} To make the evaluation of the existing and the future indices easier, Randić put forward a list of desirable attributes for topological indices.¹⁶ A particularly important requirement is that an index is not trivially related to, or highly intercorrelated with, other indices. If an index does not fulfill this condition then its informational content is either entirely or in major part comprised in other indices, as for instance in case of the Schultz index and the Wiener number^{22,23} or the Hosoya Z index and the 1Z index.²⁴

In this article we will discuss the relationship between the P'/P index and the Wiener number for connected undirected acyclic graphs and cycles.

DEFINITIONS

P Matrix. The P matrix of a labeled connected undirected graph G with N vertices, $P = P(G)$ is the square symmetric matrix of order N whose entry in the i th row and j th column is defined as

$$(P)_{ij} = \begin{cases} p'_{ij}/p & \text{if the vertices } v_i \text{ and } v_j \text{ are adjacent in } G \\ 0 & \text{otherwise} \end{cases} \quad (1)$$

where p'_{ij} is the total number of paths in the subgraph G' obtained by the removal of the edge e_{ij} from G , and p is the total number of paths in G . "Otherwise" means that either the vertices v_i and v_j are not adjacent or $i = j$. If G' is disjoint then the contributions of each component could be for instance, added¹⁶ or multiplied.²⁵ Here we follow the Randić's route, i.e., the addition of contributions. The P matrix can be the source of quite a few novel graph invariants-molecular indices. The process of finding P matrix is illustrated for 2,3-dimethylpentane in Figure 1.

P'/P Index. The P'/P index, $P'/P = P'/P(G)$, of a graph G is defined by means of the P matrix entries as

$$P'/P = \sum_{i=1}^{N-1} \sum_{j>i}^N (P)_{ij} \quad (2)$$

The quantity p'_{ij}/p could be understood as a graphical bond order,²⁶ $\pi_{e_{ij}}$ of the edge (bond) e_{ij} of G . It is a measure of relative "importance" of an edge in a graph. Using the sum over all edges in G of these local quantities

$$P'/P = \sum_{e_{ij}} \pi_{e_{ij}} \quad (3)$$

one obtains the graph invariant—molecular index P'/P .

Wiener Number. The Wiener number, $W = W(G)$, of a graph G was introduced as the path number.¹⁸ H. Wiener defined the path number as the number of bonds between all pairs of atoms in an acyclic molecule. Later W was defined in the framework of graph theory¹⁹ by means of the

[†] Reported in part at the 1995 International Chemical Congress of Pacific Basin Societies (Pacifichem '95) - Frontiers in Mathematical Chemistry, Honolulu, Dec 17–22, 1995.

[‡] The Rugjer Bošković Institute.

[§] Department of Chemistry, Faculty of Agriculture.

[⊥] Department of Biochemistry, Faculty of Science.

[Ⓢ] Abstract published in *Advance ACS Abstracts*, November 1, 1996.




Figure 1. Calculation of the elements of the P matrix for 2,3-dimethylpentane. The broken lines represent erased bonds.

elements of the distance matrix $D = D(G)$ of G as

$$W = \sum_{i=1}^{N-1} \sum_{j>i}^N (D)_{ij} \quad (4)$$

RELATIONSHIP BETWEEN P'/P INDEX AND W NUMBER

Let T be a connected undirected acyclic graph (tree) with N vertices. Consider a given edge e_{ij} of T . By deletion of e_{ij} , T breaks into two components, whose numbers of vertices are $X_{e_{ij}}$ and $N - X_{e_{ij}}$, respectively. It is well-known that the Wiener number of T is given by

$$W = \sum_{e_{ij}} W_{e_{ij}} \quad (5)$$

where

$$W_{e_{ij}} = X_{e_{ij}}(N - X_{e_{ij}}); 1 \leq X_{e_{ij}} \leq N - 1 \quad (6)$$

The total number of paths in T is

$$p = (N^2 - N)/2 \quad (7)$$

The graphical bond order of an edge e_{ij} of T , $\pi_{e_{ij}}$ is given by

$$\pi_{e_{ij}} = [(N - X_{e_{ij}})^2 - (N - X_{e_{ij}}) + X_{e_{ij}}^2 - X_{e_{ij}}]/(N^2 - N) \quad (8)$$

Combining eq 8 with eq 6 one finds the link between $\pi_{e_{ij}}$ and $W_{e_{ij}}$,

$$\pi_{e_{ij}} = 1 - [2/(N^2 - N)]W_{e_{ij}} \quad (9)$$

and the relationship between the P'/P index and the Wiener number of T immediately follows:

$$P'/P = (N - 1) - [2/(N^2 - N)]W \quad (10)$$

To calculate the P'/P index of a tree it is necessary and sufficient to have knowledge of the number of vertices and

the value of W . The Wiener number of T can be computed in many ways.^{18,23,27-36} If one selects the route²⁷

$$W = \sum_{n=1}^{n \leq N-1} n {}^n p \quad (11)$$

where ${}^n p$ is the number of paths of length n in T , then it is possible to find out the explicit formula for computing the Wiener number of T . We will show in the further text that one can derive the analytical expressions for ${}^n p$'s of T . The problem has been also considered by Kier, Hall et al. but from the aspect of graph reconstruction and only for paths up to length three.³⁷⁻³⁹

We have derived the explicit formulas for computing the number of paths of a given length in a tree with N vertices:

$${}^1 p = N - 1 \quad (12)$$

$${}^2 p = (1/2) \sum_i v_i^2 - (N - 1) \quad (13)$$

$${}^3 p = \sum_{\substack{d(i,j)=1 \\ i < j}} v_i v_j - \sum_i v_i^2 + (N - 1) \quad (14)$$

$${}^4 p = \sum_{\substack{d(i,j)=2 \\ i < j}} v_i v_j - 2 \sum_{\substack{d(i,j)=1 \\ i < j}} v_i v_j + (3/2) \sum_i v_i^2 - (N - 1) \quad (15)$$

$${}^5 p = \sum_{\substack{d(i,j)=3 \\ i < j}} v_i v_j - 2 \sum_{\substack{d(i,j)=2 \\ i < j}} v_i v_j + 3 \sum_{\substack{d(i,j)=1 \\ i < j}} v_i v_j - 2 \sum_i v_i^2 + (N - 1) \quad (16)$$

etc. where v_i is the valency of the vertex i , and $d(i,j) = 1, 2, 3, \dots$ denotes that the distances between the vertices i and j are 1, 2, 3, ..., respectively. By noting that

$$\sum_{d(i,j)=0} v_i v_j = \sum_i v_i^2 \quad (17)$$

the general analytical expression for the number of paths of length n in T reads as follows:

$$^n p = \sum_{m=0}^{n-2+k} (-1)^{m+1} (n-m-1) \sum_{\substack{d(i,j)=m \\ i < j}} v_i v_j + (-1)^{n+1} \{ [(n-1)/2] \sum_{d(i,j)=0} v_i v_j + (N-1) \} \quad (18)$$

The parameters k and l take the following values:

$$k = \begin{cases} 1 & \text{for } n = 1 \\ 0 & \text{for } n \geq 2 \end{cases} \quad (19)$$

and

$$l = \begin{cases} 1 & \text{for } n = \text{odd} \\ 0 & \text{for } n = \text{even} \end{cases} \quad (20)$$

A closer examination of eqs 12–16 reveals that for the number of paths of a given length, $n \geq 3$, the following recursion relation holds:

$$^n p = \sum_{\substack{d(i,j)=n-2 \\ i < j}} v_i v_j - 2(^{n-1} p) - ^{n-2} p \quad (21)$$

where the initial conditions are given by eqs 12 and 13.

Combining eqs 10 and 11, with eq 18 one obtains the general formula for computing the P'/P index of T

$$P'/P = (N-1) - [2/(N^2-N)] \sum_{n=1}^{n \leq N-1} n \{ \sum_{m=0}^{n-2+k} (-1)^{m+1} (n-m-1) \sum_{\substack{d(i,j)=m \\ i < j}} v_i v_j + (-1)^{n+1} \{ [(n-1)/2] \sum_{d(i,j)=0} v_i v_j + (N-1) \} \} \quad (22)$$

where the parameters k and l take the values from eqs 19 and 20.

Chains. A special case of connected undirected acyclic graphs is the chain graph, L_N .⁷ The relationship between P'/P and W for chains is rather simple and can be given only in terms of N . Since the Wiener number of L_N is given by²⁷

$$W = (N^3 - N)/6 \quad (23)$$

the P'/P index for a chain is

$$P'/P = (2/3)(N-2); \quad N \geq 2 \quad (24)$$

Cycles. For a general connected undirected graph, what means a graph containing ring(s) as well, it is not possible to find out the explicit formula for the P'/P index. However, it is possible to derive the analytical expressions for P'/P in some special cases if ring(s) are present in G . For cycles (regular graphs of degree 2)⁷ the formula is simple, and is given by

$$P'/P = N/2 \quad (25)$$

The relationship between P'/P and W for cycles depends on the parity of N . The expression for W reads as follows:^{23,36}

$$W = \begin{cases} (N/2)^3 & \text{for } N = \text{even} \\ (N/8)(N^2 - 1) & \text{for } N = \text{odd} \end{cases} \quad (26)$$

Hence, the relationship between P'/P and W for cycles is given by

$$P'/P = \begin{cases} (4/N^2)W & \text{for } N = \text{even} \\ [4/(N^2 - 1)]W & \text{for } N = \text{odd} \end{cases} \quad (27)$$

CONCLUDING REMARKS

In this report we have shown that the novel P'/P index is functionally related to the Wiener number in simple classes of chemical graphs such as trees and cycles. It is also demonstrated that P'/P for these classes can be expressed simply in terms of the number of vertices and their valences.

ACKNOWLEDGMENT

This work was supported by the Ministry of Science and Technology of the Republic of Croatia. We are grateful to Professor Ante Graovac (Zagreb) for useful comments.

REFERENCES AND NOTES

- (1) Turro, N. J. Geometric and Topological Thinking in Organic Chemistry. *Angew. Chem., Int. Ed. Engl.* **1986**, 25, 882–901.
- (2) Seybold, P. G.; May, M.; Bagal, U. A. Molecular Structure–Property Relationships. *J. Chem. Educ.* **1987**, 64, 575–581.
- (3) Randić, M. Chemical Structure - What is “She”? *J. Chem. Educ.* **1992**, 69, 713–718.
- (4) Woolley, R. G. On the Description of High-Resolution Experiments in Molecular Physics. *Chem. Phys. Lett.* **1976**, 44, 73–75.
- (5) Woolley, R. G. Must a Molecule Have a Shape? *J. Am. Chem. Soc.* **1978**, 100, 1073–1078.
- (6) Gutman, I.; Polansky, O. *Mathematical Concepts in Organic Chemistry*; Springer-Verlag: Berlin, 1986.
- (7) Trinajstić, N. *Chemical Graph Theory*, 2nd ed.; CRC Press: Boca Raton, FL, 1992.
- (8) Mihalić, Z.; Veljan, D.; Amić, D.; Nikolić, S.; Plavšić, D.; Trinajstić, N. The Distance Matrix in Chemistry. *J. Math. Chem.* **1992**, 11, 223–258.
- (9) Cvetković, D. M.; Doob, M.; Sachs, H. *Spectra of Graphs. Theory and Applications*, 3rd ed.; J. A. Barth Verlag: Heidelberg-Leipzig, 1995.
- (10) Hall, L. H. Computational Aspects of Molecular Connectivity and its Role in Structure-Property Modeling. In *Computational Chemical Graph Theory*; Rouvray, D. H., Ed.; Nova Press: New York, 1990; pp 202–233.
- (11) Tratch, S. S.; Stankevitch, M. I.; Zefirov, N. S. Combinatorial Models and Algorithms in Chemistry. The Expanded Wiener Number - A Novel Topological Index. *J. Comput. Chem.* **1990**, 11, 899–908.
- (12) Klein, D. J.; Randić, M. Resistance distance. *J. Math. Chem.* **1993**, 12, 81–95.
- (13) Randić, M.; Guo, X.; Oxley, T.; Krishnapriyan, H. Wiener Matrix: Source of Novel Graph Invariants. *J. Chem. Inf. Comput. Sci.* **1993**, 33, 709–716.
- (14) Trinajstić, N.; Babić, D.; Nikolić, S.; Plavšić, D.; Amić, D.; Mihalić, Z. The Laplacian Matrix in Chemistry. *J. Chem. Inf. Comput. Sci.* **1994**, 34, 368–376.
- (15) Randić, M. Hosoya Matrix - A Source of New Molecular Descriptors. *Croat. Chem. Acta* **1994**, 67, 415–429.
- (16) Randić, M. Generalized Molecular Descriptors. *J. Math. Chem.* **1991**, 7, 155–168.
- (17) Randić, M.; Trinajstić, N. In Search for Graph Invariants of Chemical Interest. *J. Mol. Struct.* **1993**, 300, 551–571.
- (18) Wiener, H. Structural Determination of Paraffin Boiling Points. *J. Am. Chem. Soc.* **1947**, 69, 17–20.
- (19) Hosoya, H. Topological Index. A Newly Proposed Quantity Characterizing the Topological Nature of Structural Isomers of Saturated Hydrocarbons. *Bull. Chem. Soc. Jpn.* **1971**, 44, 2332–2339.
- (20) Rouvray, D. H. The Modeling of Chemical Phenomena Using Topological Indices. *J. Comput. Chem.* **1987**, 8, 470–480.
- (21) Mihalić, Z.; Trinajstić, N. A Graph-Theoretical Approach to Structure–Property Relationships. *J. Chem. Educ.* **1992**, 69, 701–712.

- (22) Klein, D. J.; Mihalić, Z.; Plavšić, D.; Trinajstić, N. Molecular Topological Index: A Relation with the Wiener Index. *J. Chem. Inf. Comput. Sci.* **1992**, 32, 304–305.
- (23) Plavšić, D.; Nikolić, S.; Trinajstić, N.; Klein, D. J. Relation between the Wiener Index and the Schultz Index for Several Classes of Chemical Graphs. *Croat. Chem. Acta* **1993**, 66, 345–353.
- (24) Plavšić, D.; Šoškić, M.; Landeka, I.; Gutman, I.; Graovac, A. On the Relation between the Path Numbers 1Z , 2Z and the Hosoya Z Index. *J. Chem. Inf. Comput. Sci.*, in press.
- (25) Mekenyan, O.; Bonchev, D.; Balaban, A. Topological Indices for Molecular Fragments and New Graph Invariants. *J. Math. Chem.* **1988**, 2, 347–375.
- (26) Randić, M.; Mihalić, Z.; Nikolić, S.; Trinajstić, N. Graphical Bond Orders: Novel Structural Descriptors. *J. Chem. Inf. Comput. Sci.* **1994**, 34, 403–409.
- (27) Bonchev, D.; Trinajstić, N. Information Theory, Distance Matrix and Molecular Branching. *J. Chem. Phys.* **1977**, 67, 4517–4533.
- (28) Bersohn, M. A Fast Algorithm for Calculation of the Distance Matrix. *J. Comput. Chem.* **1983**, 4, 110–113.
- (29) Canfield, E. R.; Robinson, R. W.; Rouvray, D. H. Determination of the Wiener Molecular Branching Index for the General Tree. *J. Comput. Chem.* **1985**, 6, 598–609.
- (30) Barysz, M.; Plavšić, D.; Trinajstić, N. A Note on Topological Indices. *Math. Chem. (Mulheim/Ruhr)* **1986**, 19, 89–116.
- (31) Muller, W. R.; Szymanski, K.; Knop, J. V.; Trinajstić, N. An Algorithm for Construction of the Molecular Distance Matrix. *J. Comput. Chem.* **1987**, 8, 170–173.
- (32) Mohar, B.; Pisanski, T. How to Compute the Wiener Index of a Graph. *J. Math. Chem.* **1988**, 2, 267–277.
- (33) Senn, P. The Computation of the Distance Matrix and the Wiener Index for Graphs of Arbitrary Complexity with Weighted Vertices and Edges. *Comput. Chem.* **1988**, 12, 219–227.
- (34) Gutman, I.; Yeh, Y. N.; Lee, S. L.; Luo, Y. L. Some Recent Results in the Theory of the Wiener Number. *Indian J. Chem.* **1993**, 32A, 651–661.
- (35) Gutman, I. A New Method for the Calculation of the Wiener Number of Acyclic Molecules. *J. Mol. Struct. (Theochem)* **1993**, 285, 137–142.
- (36) Nikolić, S.; Trinajstić, N.; Mihalić, Z. The Wiener Index: Development and Applications. *Croat. Chem. Acta* **1995**, 68, 105–129.
- (37) Kier, L. B.; Hall, L. H.; Frazer, J. W. Design of Molecules from Quantitative Structure-Activity Relationship Models. 1. Information Transfer between Path and Vertex Degree Counts. *J. Chem. Inf. Comput. Sci.* **1993**, 33, 143–147.
- (38) Hall, L. H.; Kier, L. B.; Frazer, J. W. Design of Molecules from Quantitative Structure-Activity Relationship Models. 2. Derivation and Proof of Information Transfer Relating Equations. *J. Chem. Inf. Comput. Sci.* **1993**, 33, 148–152.
- (39) Hall, L. H.; Dailey, R. S.; Kier, L. B. Design of Molecules from Quantitative Structure-Activity Relationship Models. 3. Role of Higher Order Path Counts: Path 3. *J. Chem. Inf. Comput. Sci.* **1993**, 33, 598–603.

CI9603691