

Catalytic Enantioselective Cyclization/Cross-Coupling with Alkyl Electrophiles

Huan Cong^{†,‡} and Gregory C. Fu^{*,†,‡}

[†]Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States


[‡]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

S Supporting Information

ABSTRACT: As part of our ongoing effort to expand the scope of cross-coupling reactions of alkyl electrophiles, we have pursued a strategy wherein the nucleophilic coupling partner includes a pendant olefin; after transmetalation by such a substrate, if β -migratory insertion proceeds faster than direct cross-coupling, an additional carbon–carbon bond and stereocenter can be formed. With the aid of a nickel/diamine catalyst (both components are commercially available), we have established the viability of this approach for the catalytic asymmetric synthesis of 2,3-dihydrobenzofurans and indanes. Furthermore, we have applied this new method to the construction of the dihydrobenzofuran core of fasiglifam, as well as to a cross-coupling with a racemic alkyl electrophile; in the latter process, the chiral catalyst controls two stereocenters, one that is newly generated in a β -migratory insertion and one that begins as a mixture of enantiomers.

In recent years, significant progress has been reported on the development of methods for the transition-metal-catalyzed cross-coupling of alkyl electrophiles to generate carbon–carbon bonds, including enantioselective processes.¹ To date, most investigations of asymmetric catalysis have focused on stereoconvergent reactions of racemic secondary electrophiles,² although an advance has also been described with a racemic secondary nucleophile (top of Figure 1).³

Previous work:



This study:




Figure 1. Asymmetric cross-couplings of alkyl electrophiles.

As part of our ongoing effort to expand the scope of enantioselective cross-couplings of alkyl electrophiles, we are pursuing an approach wherein an organometallic reagent that bears a pendant olefin is employed as the nucleophilic coupling partner (bottom of Figure 1).^{4–6} In the presence of a chiral catalyst, transmetalation and then β -migratory insertion (left side of Figure 2), followed by alkyl–alkyl coupling, could lead




Figure 2. Complementary approaches to generating a precursor (A) for catalytic enantioselective cyclizations.


to the formation of two carbon–carbon bonds and a new stereocenter (bottom of Figure 1). This strategy complements asymmetric coupling processes wherein an intermediate of type A is generated through oxidative addition of an electrophile (right side of Figure 2).⁷

In this report, we establish that a transmetalation–insertion sequence can indeed be used to generate two, rather than one, carbon–carbon bonds in a cross-coupling with an alkyl electrophile and that this process can be achieved with good enantioselectivity. Specifically, we describe couplings of arylboron reagents that bear a pendant olefin with unactivated alkyl halides, thereby furnishing 2,3-dihydrobenzofurans^{8,9} and indanes^{10,11} in high ee (eq 1).


In order to enhance the likelihood of cyclization (β -migratory insertion) prior to coupling with the electrophile, we chose to focus on an organometallic coupling partner that could form a five-membered ring upon insertion, since such cyclizations are often facile. At the outset, it was unclear what catalyst would enable the desired sequence of bond-forming processes, much less achieve high enantioselectivity.

Received: January 22, 2014


Published: February 27, 2014



Scheme 1. Catalytic Asymmetric Synthesis of the 2,3-Dihydrobenzofuran Core of Fasiglifam



In view of the similarity of the optimized conditions for this new asymmetric cyclization/cross-coupling process to those for our stereoconvergent cross-coupling of racemic γ -haloamides,^{12d} we investigated the possibility that a single chiral catalyst could accomplish two distinct enantioselective transformations: create a new stereocenter through the cyclization of an achiral nucleophile, as well as control the absolute stereochemistry of a second stereocenter through an enantioconvergent coupling of a racemic electrophile. As illustrated in eq 4, this objective can indeed be achieved (minor diastereomer: 86% ee).



In summary, we have expanded the scope of cross-coupling reactions of alkyl electrophiles by incorporating an olefin in the nucleophilic partner, which leads to the formation of an additional carbon–carbon bond and stereocenter, when compared with a simple cross-coupling. With the aid of a nickel/diamine catalyst (both components are commercially

available), we have established that this strategy enables the synthesis of highly enantioenriched 2,3-dihydrobenzofurans and indanes through couplings with a range of alkyl halides. We have applied this new method to the generation of the dihydrobenzofuran core of fasiglifam, as well as to a transformation wherein the chiral catalyst controls the stereochemistry of two rather different processes: a β -migratory insertion and an enantioconvergent coupling of a racemic alkyl halide. Ongoing studies are directed at further enlarging the scope of cross-coupling reactions of alkyl electrophiles, as well as elucidating the mechanisms of these transformations.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

gcfu@caltech.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Support has been provided by the National Institutes of Health (National Institute of General Medical Sciences: R01-GM62871). We thank Yufan Liang, Dr. Allen G. Oliver (University of Notre Dame), Dr. Nathan D. Schley, and Dr. Scott C. Virgil (Caltech Center for Catalysis and Chemical Synthesis, supported by the Gordon and Betty Moore Foundation) for assistance.

■ REFERENCES

- (1) For reviews and leading references, see: (a) Glasspoole, B. W.; Crudden, C. M. *Nat. Chem.* **2011**, 3, 912–913. (b) Rudolph, A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, 48, 2656–2670. (c) Glorius, F. *Angew. Chem., Int. Ed.* **2008**, 47, 8347–8349. (d) Swift, E. C.; Jarvo, E. R. *Tetrahedron* **2013**, 69, 5799–5817.
- (2) For a few examples and leading references, see: (a) Do, H.-Q.; Chandrashekar, E. R. R.; Fu, G. C. *J. Am. Chem. Soc.* **2013**, 135, 16288–16291. (b) Schmidt, T.; Kirschning, A. *Angew. Chem., Int. Ed.* **2012**, 51, 1063–1066. (c) Tsuji, T.; Yorimitsu, H.; Oshima, K. *Angew. Chem., Int. Ed.* **2002**, 41, 4137–4139. (d) Shields, J. D.; Ahneman, D. T.; Graham, T. J. A.; Doyle, A. G. *Org. Lett.* **2014**, 16, 142–145. (e) Caeiro, J.; Sestelo, J. P.; Sarandeses, L. A. *Chem.—Eur. J.* **2008**, 14, 741–746.
- (3) Cordier, C. J.; Lundgren, R. J.; Fu, G. C. *J. Am. Chem. Soc.* **2013**, 135, 10946–10949.
- (4) For reviews on asymmetric carbometallation, see: (a) Ojima, I.; Kaloko, J. J.; Chaterpaul, S. J.; Teng, Y.-H. G.; Lin, C.-F. In *Catalytic Asymmetric Synthesis*; Ojima, I., Ed.; Wiley: Hoboken, NJ, 2010; pp 643–681. (b) Negishi, E.-i.; Tan, Z. *Top. Organomet. Chem.* **2005**, 8, 139–176. (c) Hoveyda, A. H.; Heron, N. M. In *Comprehensive Asymmetric Catalysis*; Jacobsen, E. N., Pfaltz, A., Yamamoto, H., Eds.; Springer-Verlag: Berlin, 1999; pp 431–454.
- (5) McDonald, R. I.; Liu, G.; Stahl, S. S. *Chem. Rev.* **2011**, 111, 2981–3019.
- (6) For recent examples of related catalytic asymmetric couplings wherein the nucleophilic site is a nitrogen, rather than a carbon, and the electrophile is not an alkyl halide, see: (a) Mai, D. N.; Wolfe, J. P. *J. Am. Chem. Soc.* **2010**, 132, 12157–12159. (b) Ingalls, E. L.; Sibbald, P. A.; Kaminsky, W.; Michael, F. E. *J. Am. Chem. Soc.* **2013**, 135, 8854–8856.

(7) For example, asymmetric Heck reactions: (a) McCartney, D.; Guiry, P. J. *Chem. Soc. Rev.* **2011**, *40*, 5122–5150. (b) Dounay, A. B.; Overman, L. E. In *Mizoroki–Heck Reaction*; Oestreich, M., Ed.; Wiley: Chichester, U.K., 2009; pp 533–568. (c) Shibasaki, M.; Vogl, E. M.; Ohshima, T. *Adv. Synth. Catal.* **2004**, *346*, 1533–1552.

(8) For reviews of methods for the synthesis of 2,3-dihydrobenzofurans, as well as descriptions of their significance, see: (a) Sheppard, T. D. *J. Chem. Res.* **2011**, *35*, 377–385. (b) Bertolini, F.; Pineschi, M. *Org. Prep. Proced. Int.* **2009**, *41*, 385–418. (c) Lachia, M.; Moody, C. J. *Nat. Prod. Rep.* **2008**, *25*, 227–253.

(9) Codeine is an example of a bioactive compound that includes a 2,3-dihydrobenzofuran.

(10) For a review of methods for the synthesis of indanes, see: Hong, B.-c.; Sarshar, S. *Org. Prep. Proced. Int.* **1999**, *31*, 1–86.

(11) Rasagiline, which is used for the treatment of Parkinson's disease, is an example of a simple bioactive compound that includes an indane: Hoy, S. M.; Keating, G. M. *Drugs* **2012**, *72*, 643–669.

(12) (a) Saito, B.; Fu, G. C. *J. Am. Chem. Soc.* **2008**, *130*, 6694–6695. (b) Owston, N. A.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 11908–11909. (c) Lu, Z.; Wilsily, A.; Fu, G. C. *J. Am. Chem. Soc.* **2011**, *133*, 8154–8157. (d) Zultanski, S. L.; Fu, G. C. *J. Am. Chem. Soc.* **2011**, *133*, 15362–15364. (e) Wilsily, A.; Tramutola, F.; Owston, N. A.; Fu, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 5794–5797.

(13) The failure to observe a significant amount of the cross-coupling product in the absence of *i*-BuOH (entry 4 of Table 1) could be due to less effective transmetalation in the absence of a less bulky alkoxide.

(14) Some of the nucleophile is consumed in the reduction of the Ni(II) precatalyst to the active catalyst. A small amount also undergoes protodeborylation under the reaction conditions.

(15) For example, see: (a) Pybox ligand: Fischer, C.; Fu, G. C. *J. Am. Chem. Soc.* **2005**, *127*, 4594–4595. (b) Bis(oxazoline) ligand: Lou, S.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 1264–1266.

(16) The alkyl chloride is largely intact at the end of the reaction (>95%).

(17) Notes: Under our standard conditions (Table 1): (a) The coupling illustrated in Table 2, entry 1 proceeded in 96% ee and 67% yield on a gram scale (1.07 g of product). (b) An initial attempt to form a six-membered ring through cyclization/cross-coupling of a homologated arylboron reagent was not successful. (c) In general, the primary undesired side reactions are reduction (hydrodehalogenation) and electrophile homocoupling. (d) PhBr is not a suitable electrophile. (e) An indoline can be generated with promising enantioselectivity and yield (54% ee, 40% yield).

(18) For a discussion and leading references, see: (a) Takeda web page. http://www.takeda.com/news/2013/20131227_6117.html (accessed January 18, 2014). (b) Negoro, N.; Sasaki, S.; Mikami, S.; Ito, M.; Suzuki, M.; Tsujihata, Y.; Ito, R.; Harada, A.; Takeuchi, K.; Suzuki, N.; Miyazaki, J.; Santou, T.; Odani, T.; Kanzaki, N.; Funami, M.; Tanaka, T.; Kogame, A.; Matsunaga, S.; Yasuma, T.; Momose, Y. *ACS Med. Chem. Lett.* **2010**, *1*, 290–294. (c) Kaku, K. *Expert Opinion on Pharmacotherapy* **2013**, *14*, 2591–2600. (d) de Lartigue, J. *Drugs of the Future* **2011**, *36*, 813–818.