

Probing Mode and Site of Substrate Water Binding to the Oxygen-Evolving Complex in the S_2 State of Photosystem II by ^{17}O -HYSCORE Spectroscopy [Journal of the American Chemical Society 2008, 130, 786–787 DOI: 10.1021/ja076620i]. Ji-Hu Su Wolfgang Lubitz* and Johannes Messinger*

Additional experiments (see Figure 1) have revealed that the HYSCORE signal reported for the oxygen-evolving complex (OEC) poised in the S_2 state of spinach BBY samples in the presence of ^{17}O -water was incorrectly assigned to the coupling of ^{17}O to the Mn cluster. This signal should instead be assigned to the hyperfine coupling of the axial ^{14}N ligand of the low-spin Fe^{III} center of oxidized cytochrome b559.¹ We have concluded that

cytb559 was partially oxidized during 200 K illumination in samples incubated with ^{17}O -water, but to a lesser extent in the ^{16}O control samples.

Figure 1A shows the X-band HYSCORE spectrum of the OEC of *Thermosynechococcus elongatus* poised in the S_2 state, measured at the center field of the S_2 EPR spectrum. The two sharp peaks previously assigned to the ^{17}O nucleus of a water-derived ligand to the OEC were observed (left side of panel A). However, they were also seen in the S_1 state and in the absence of ^{17}O -water (data not shown). This signal must therefore arise from the above-mentioned ^{14}N couplings in oxidized cytochrome b559 and/or c550.

To exclude the possibility that the ^{17}O signal seen in the original study (using spinach samples naturally lacking c550) and the cytochrome signal identified above appear in exactly the same spectral position, universal ^{15}N -labeling was employed. Figure 1B,C shows that the two sharp cytochrome peaks shift as compared to panel A, as expected for $^{14}\text{N}/^{15}\text{N}$ exchange. No additional signal was observed in the S_2 state spectrum (panel C, red boxes), demonstrating that our original assignment was incorrect.

ACKNOWLEDGMENT

The authors are indebted to Alain Boussac (CEA Saclay), Nicholas Cox (MPI Muelheim), Julia Sander, Marc Nowaczyk, and Matthias Rögnér (all Ruhr University Bochum) for their help in correcting this assignment.

REFERENCES

- (1) Garcia-Rubio, I.; Martinez, J. I.; Picorel, R.; Yruela, I.; Alonso, P. J. *J. Am. Chem. Soc.* 2003, 125, 15846–15854.

DOI: 10.1021/ja205377n

Published on Web 07/15/2011




Figure 1. X-band HYSCORE spectra of *T. elongatus* photosystem II core preparations suspended in either buffered H_2^{16}O or H_2^{17}O medium: (A) S_2 state of ^{14}N -PSII in H_2^{17}O medium, (B) S_1 state of ^{15}N -PSII in H_2^{16}O medium, and (C) S_2 state of ^{15}N -PSII in H_2^{17}O medium. The red boxes show the region where the ^{17}O signal reported in the full article was observed. The S_2 state was generated by 200 K white light illumination for 5 s. All spectra were obtained at the center field of the S_2 multiline EPR spectrum ($B_0 = 335$ mT). Experimental parameters: (A) $\pi/2 = 24$ ns; $\tau = 196$ ns; t_1, t_2 were varied from 60 to 6720 ns (24 ns steps); shots per point = 50; shot repetition rate = 5 ms. (B,C) $\pi/2 = 6$ ns; $\tau = 196$ ns; t_1, t_2 were varied from 100 to 3172 ns (24 ns steps); shots per point = 100 (B) and 400 (C); shot repetition rate = 1 ms; temperature = 4.8 K.