

Correction to (1-Nosyl-5-nitroindol-3-yl)methyl Ester: A Novel Protective Group for Carboxylic Acids

Takuya Nishimura, Kouhei Yamada, Tohru Takebe, Satoshi Yokoshima, and Tohru Fukuyama* *Org. Lett.* **2008**, *10*(12), 2601–2604. DOI: 10.1021/ol8008655

Supporting Information

Yields in the manuscript were reported incorrectly for compound 13 (Scheme 4) and compound 18 (Table 3, entry 6). Correct yields are 81% and 88%.

Solvent peaks had been removed from the ¹H NMR spectra reported for compounds 13, 9a, 9b, 9c, 9d, 14, 9f, 9g, 9i, 9j, 9k, 6a, 6b, 6h, 6j, 6k, 6n, 17, 18, and 19. Original FIDs were located, and the spectra were reprocessed and have been replaced for the above compounds in the revised Supporting Information submitted with this correction. The spectra editing did not affect any of the conclusions of the published paper. The purities calculated on the basis of the revised spectra are as follows: 13 (98%), 9a (97%), 9b (99%), 9c (99%), 9d (99%), 14 (98%), 9f (94%), 9g (88%), 9i (99%), 9j (99%), 9k (99%), 6a (98%), 6b (97%), 6h (91%), 6j (99%), 6k (97%), 6n (97%), 17 (97%), 18 (98%), and 19 (95%).

ASSOCIATED CONTENT

S Supporting Information

Revised Supporting Information including reprocessed spectra. This material is available free of charge via the Internet at http://pubs.acs.org.

Received: December 25, 2013 Published: February 3, 2014

