

# OFF-ON-OFF Fluorescence Switch with T-Latch Function


Vânia F. Pais,<sup>†</sup> Patricia Remón,<sup>†</sup> Daniel Collado,<sup>‡</sup> Joakim Andréasson,<sup>§</sup> Ezequiel Pérez-Inestrosa,<sup>\*,‡</sup> and Uwe Pischel<sup>\*,†</sup>

Centro de Investigación en Química Sostenible (CIQSO) and Departamento de Ingeniería Química, Química Física y Química Orgánica, Universidad de Huelva, Campus de El Carmen, E-21071 Huelva, Spain, Departamento de Química Orgánica, Universidad de Málaga, Campus Teatinos, E-29071 Málaga, Spain, and Department of Chemical and Biological Engineering, Physical Chemistry, Chalmers University of Technology, SE-41296 Göteborg, Sweden

uwe.pischel@diq.uhu.es; inestrosa@uma.es

Received August 25, 2011

## ABSTRACT



A novel molecular system with characteristics of an OFF-ON-OFF fluorescence switch was designed to integrate the function of a T-latch. In detail, a receptor<sub>1</sub>-fluorophore-receptor<sub>2</sub> architecture was adopted to achieve fluorescence switching upon addition of protons.

The idea to use molecular systems for information processing has attracted a great deal of interest during the recent years.<sup>1–5</sup> This has been manifested by the availability of molecular mimics for all essential logic gates (AND, OR, NOR, NAND, INH, XOR, etc.)<sup>1,3</sup> and for rather complex logic devices such as adders/subtractors, encoders/decoders, and multiplexers/demultiplexers.<sup>2,6–10</sup> Such logic functions are of elevated interest for

applications such as object coding,<sup>11</sup> intelligent materials,<sup>12–14</sup> pro-drug activation,<sup>15–17</sup> and diagnostics/actuation.<sup>18–20</sup> While these systems work independently of the order of input application (combinational logic), the molecular memorization of information is a precondition for applications which profit from a sequential application of input signals.<sup>21,22</sup> This behavior is reflected in the

<sup>†</sup> Universidad de Huelva.

<sup>‡</sup> Universidad de Málaga.

<sup>§</sup> Chalmers University of Technology.

(1) de Silva, A. P.; Uchiyama, S. *Nat. Nanotechnol.* **2007**, *2*, 399–410.

(2) Pischel, U. *Angew. Chem., Int. Ed.* **2007**, *46*, 4026–4040.

(3) Szaciłowski, K. *Chem. Rev.* **2008**, *108*, 3481–3548.

(4) Andréasson, J.; Pischel, U. *Chem. Soc. Rev.* **2010**, *39*, 174–188.

(5) Pischel, U. *Aust. J. Chem.* **2010**, *63*, 148–164.

(6) Margulies, D.; Melman, G.; Shanzer, A. *J. Am. Chem. Soc.* **2006**, *128*, 4865–4871.

(7) Amelia, M.; Baroncini, M.; Credi, A. *Angew. Chem., Int. Ed.* **2008**, *47*, 6240–6243.

(8) Pérez-Inestrosa, E.; Montenegro, J.-M.; Collado, D.; Suau, R. *Chem. Commun.* **2008**, 1085–1087.

(9) Ceroni, P.; Bergamini, G.; Balzani, V. *Angew. Chem., Int. Ed.* **2009**, *48*, 8516–8518.

(10) Andréasson, J.; Pischel, U.; Straight, S. D.; Moore, T. A.; Moore, A. L.; Gust, D. *J. Am. Chem. Soc.* **2011**, *133*, 11641–11648.

(11) de Silva, A. P.; James, M. R.; McKinney, B. O. F.; Pears, D. A.; Weir, S. M. *Nat. Mater.* **2006**, *5*, 787–790.

(12) Motornov, M.; Zhou, J.; Pita, M.; Gopishetty, V.; Tokarev, I.; Katz, E.; Minko, S. *Nano Lett.* **2008**, *8*, 2993–2997.

(13) Angelos, S.; Yang, Y.-W.; Khashab, N. M.; Stoddart, J. F.; Zink, J. I. *J. Am. Chem. Soc.* **2009**, *131*, 11344–11346.

(14) Tokarev, I.; Gopishetty, V.; Zhou, J.; Pita, M.; Motornov, M.; Katz, E.; Minko, S. *ACS Appl. Mater. Interfaces* **2009**, *1*, 532–536.

(15) Amir, R. J.; Popkov, M.; Lerner, R. A.; Barbas, C. F., III; Shabat, D. *Angew. Chem., Int. Ed.* **2005**, *44*, 4378–4381.

(16) Ozlem, S.; Akkaya, E. U. *J. Am. Chem. Soc.* **2009**, *131*, 48–49.

(17) Hammarson, M.; Andersson, J.; Li, S.; Lincoln, P.; Andréasson, J. *Chem. Commun.* **2010**, 46, 7130–7132.

(18) Konry, T.; Walt, D. R. *J. Am. Chem. Soc.* **2009**, *131*, 13232–13233.

(19) Margulies, D.; Hamilton, A. D. *J. Am. Chem. Soc.* **2009**, *131*, 9142–9143.

(20) Privman, M.; Tam, T. K.; Bocharova, V.; Halámek, J.; Wang, J.; Katz, E. *ACS Appl. Mater. Interfaces* **2011**, *3*, 1620–1623.

(21) Raymo, F. M.; Alvarado, R. J.; Giordani, S.; Cejas, M. A. *J. Am. Chem. Soc.* **2003**, *125*, 2361–2364.


(22) Pischel, U. *Angew. Chem., Int. Ed.* **2010**, *49*, 1356–1358.

(23) Guo, Z.; Zhu, W.; Shen, L.; Tian, H. *Angew. Chem., Int. Ed.* **2007**, *46*, 5549–5553.

function of molecular keypad locks<sup>10,23–28</sup> and memory devices.<sup>29–35</sup>

The set-reset (S-R) latch was one of the first memory devices that was implemented at the molecular level by using electrochemical, chemical, and photonic signaling.<sup>29,31–35</sup> The device is characterized by a high state (binary 1) whenever the set input is applied ( $S = 1$ ) and which upon reset ( $R = 1$ ) has a binary 0 (low) state. The herein described toggle-latch (T-latch) is a different logic switch with memory capacity. Its working principle is well illustrated with the function of a conventional light switch or of the push button of a ballpoint pen: every time the toggle input is activated, the state  $Q$  of the system changes (see Scheme 1). The device “remembers” if a 0 or a 1 state was memorized ( $Q_{\text{current}}$ ) and upon each T input application, the new state ( $Q_{\text{next}}$ ) has the opposite value ( $0 \rightarrow 1$  and  $1 \rightarrow 0$ ). The “do nothing” situation leaves the system state unchanged.


**Scheme 1.** Presentation of the T-Latch Function




We anticipated that a molecular OFF-ON-OFF fluorescent switch could integrate this function. In detail, we needed a switch which upon single application of an input changes to the ON state and is set back to the OFF state by

- (24) Margulies, D.; Felder, C. E.; Melman, G.; Shanzer, A. *J. Am. Chem. Soc.* **2007**, *129*, 347–354.  
 (25) Suresh, M.; Ghosh, A.; Das, A. *Chem. Commun.* **2008**, 3906–3908.  
 (26) Kumar, S.; Luxami, V.; Saini, R.; Kaur, D. *Chem. Commun.* **2009**, 3044–3046.  
 (27) Andréasson, J.; Straight, S. D.; Moore, T. A.; Moore, A. L.; Gust, D. *Chem.—Eur. J.* **2009**, *15*, 3936–3939.  
 (28) Remón, P.; Hammarson, M.; Li, S.; Kahnt, A.; Pischel, U.; Andréasson, J. *Chem.—Eur. J.* **2011**, *17*, 6492–6500.  
 (29) Baron, R.; Onopriyenko, A.; Katz, E.; Lioubashevski, O.; Willner, I.; Wang, S.; Tian, H. *Chem. Commun.* **2006**, 2147–2149.  
 (30) Elbaz, J.; Moshe, M.; Willner, I. *Angew. Chem., Int. Ed.* **2009**, *48*, 3834–3837.  
 (31) Periyasamy, G.; Collin, J.-P.; Sauvage, J.-P.; Levine, R. D.; Remacle, F. *Chem.—Eur. J.* **2009**, *15*, 1310–1313.  
 (32) Pita, M.; Strack, G.; MacVittie, K.; Zhou, J.; Katz, E. *J. Phys. Chem. B* **2009**, *113*, 16071–16076.  
 (33) de Ruiter, G.; Motiei, L.; Choudhury, J.; Oded, N.; van der Boom, M. E. *Angew. Chem., Int. Ed.* **2010**, *49*, 4780–4783.  
 (34) de Ruiter, G.; Tartakovsky, E.; Oded, N.; van der Boom, M. E. *Angew. Chem., Int. Ed.* **2010**, *49*, 169–172.  
 (35) Pischel, U.; Andréasson, J. *New J. Chem.* **2010**, *34*, 2701–2703.

**Scheme 2.** (a) Structures of Triad **2** and the Naphthalimide Models **3** and **5** with One Receptor Unit and (b) Synthesis of Triad **2** (for Compound **4**, an Intermediary Product, see Supporting Information)



a second equal input. Fluorescent systems, which change their emission properties upon application of chemical input information, have been often explored in the design of logic switches and chemical sensors.<sup>3,4,36–38</sup> The integrated receptor<sub>1</sub>-fluorophore-receptor<sub>2</sub> architecture **2** (Scheme 2a) was identified as an excellent candidate to put the molecular T-latch function into practice.



**Figure 1.** Relative absorption spectra (dashed lines) and normalized fluorescence spectra (solid lines) for **2** (red), **2H<sup>+</sup>** (blue), and **2H<sub>2</sub><sup>2+</sup>** (black). Note that the low fluorescence emissions of **2** and **2H<sub>2</sub><sup>2+</sup>** are hardly distinguishable.

- (36) de Silva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. *Chem. Rev.* **1997**, *97*, 1515–1566.  
 (37) de Silva, A. P.; Vance, T. P.; West, M. E. S.; Wright, G. D. *Org. Biomol. Chem.* **2008**, *6*, 2468–2480.  
 (38) Duke, R. M.; Veale, E. B.; Pfeffer, F. M.; Kruger, P. E.; Gunnlaugsson, T. *Chem. Soc. Rev.* **2010**, *39*, 3936–3953.

The synthesis of the new triad **2** is briefly sketched in Scheme 2b. The sequence started with the commercial 4-bromo-1,8-naphthalic anhydride, which was condensed with 8-aminoquinoline (74% yield). Further aromatic nucleophilic substitution of the intermediary 4-bromo-1,8-naphthalimide derivative with *N*-methylpiperazine resulted in the final product **2** with a yield of 52%. The synthesis of the naphthalimide derivatives **3** and **5** (Scheme 2a), which served herein as model structures, is described in the Supporting Information.


**Table 1.** Photophysical Properties of Compounds **2**, **3**, and **5** and Their Protonated Forms in Aerated Acetonitrile Solution

	$\lambda_{\text{abs,max}}/\text{nm}$	$\epsilon/\text{M}^{-1}\text{cm}^{-1}$	$\lambda_{\text{fluo,max}}/\text{nm}$	$\Phi_{\text{f}}$	$\tau_{\text{f}}/\text{ns}$
<b>2</b>	401	10100	504	0.017	<sup>a</sup>
<b>2H</b> <sup>+</sup>	377	10500	499	0.67	8.90
<b>2H</b> <sub>2</sub> <sup>2+</sup>	383	11100	499	0.017	<sup>a</sup>
<b>3</b>	433	13000	520	0.56	10.00
<b>3H</b> <sup>+</sup>	441	13400	519	0.006	<sup>a</sup>
<b>5</b>	398	9700	502	0.018	<sup>a</sup>
<b>5H</b> <sup>+</sup>	374	10100	498	0.62	9.06

<sup>a</sup> Not determined due to low signal intensity.

Triad **2** contains two proton receptors: a piperazinyl and a quinolinyl moiety. The receptors have sufficiently different  $\text{p}K_{\text{a}}$  values (7.78 for *N*-benzoylpiperazine versus 4.60 for 8-methylquinoline as models)<sup>39</sup> so that they can be stepwise protonated. In accordance with this assumption and as shown in Figures 1 and 2, the addition of 1 equiv of protons (triflic acid;  $\text{CF}_3\text{SO}_3\text{H}$ ) yielded a pronounced fluorescence enhancement (fluorescence quantum yield  $\Phi_{\text{f}} = 0.67$  versus 0.017 for **2H**<sup>+</sup> and **2**, respectively) of the 4-amino-1,8-naphthalimide chromophore ( $\lambda_{\text{fluo,max}} = 504$  nm for **2** and 499 nm for **2H**<sup>+</sup>). However, the subsequent addition of a second equivalent of  $\text{CF}_3\text{SO}_3\text{H}$  caused practically quantitative fluorescence quenching (98% quenching). The photophysical properties of all investigated compounds and their protonated forms are summarized in Table 1. The independent actuation of both receptors in **2** was supported by the observation of the same differential photophysical effects upon protonation of the model compounds **3** and **5**, which contain each only one of the two receptors (Scheme 2a). In accordance with the fluorescence response of **2** upon stepwise protonation, **3** showed quenching and **5** enhancement of the emission for the addition of 1 equiv of protons (Supporting Information). The superposition of the photophysical

trends of the model compounds in the triad was also noted for the absorption spectra.<sup>40</sup>



**Figure 2.** Fluorescence titration curve ( $\lambda_{\text{exc}} = 388$  nm,  $\lambda_{\text{obs}} = 499$  nm) of **2** (12.5  $\mu\text{M}$  in acetonitrile) upon  $\text{CF}_3\text{SO}_3\text{H}$  addition.

**Table 2.** Truth Table for the Implemented Molecular T-Latch

<i>T</i> input (1 equiv $\text{H}^+$ )	$Q_{\text{current}}$ (fluo)	$Q_{\text{next}}$ (fluo)	control channel (abs, 313 nm)
0	0	0	0
1	0	1	0
0	1	1	0
1	1	0	1

The fluorescence switching of triad **2** can be mechanistically rationalized as follows. The electron-donating methyl-substituted piperazinyl nitrogen atom is protonated upon the addition of the first equivalent of protons, which leads to blocking of photoinduced electron transfer (PET) and consequently fluorescence ON switching.<sup>41–43</sup> The second equivalent of protons serves to transform the quinolinyl residue into a quinolinium cation. The hydrogen-bonding interaction of  $\text{NH}^+$  with the imide carbonyl  $\text{C}=\text{O}$  is assumed to be at the origin of the fluorescence quenching of the 4-amino-1,8-naphthalimide derivative.<sup>41</sup> However, PET from the singlet-excited fluorophore to the electron-accepting quinolinium cation may also be involved in the observed fluorescence OFF switching.<sup>44,45</sup> Noteworthy, the control of 4-aminonaphthalimide fluorescence by a receptor linked to the “imide side” of the fluorophore has been rarely observed.<sup>41,42,46–48</sup>

(39) The  $\text{p}K_{\text{a}}$  data were taken from [http://research.chem.psu.edu/brpgrp/pKa\\_compilation.pdf](http://research.chem.psu.edu/brpgrp/pKa_compilation.pdf).

(40) The protonation of the piperazinyl residue leads to a blue shift (by 24 nm) of the long wavelength aminonaphthalimide absorption band (for **2** and **5**), which is indicative of the destabilization of the charge transfer state through the repulsive interaction between the protonated distant methyl-substituted N and the positive pole of the charge transfer state at the aromatic N (cf. ref 47). The protonation of the quinolinyl moiety has stabilizing effects on the charge transfer state, which is expressed by a red shift (by 6–8 nm) of the long wavelength absorption band (for **2** and **3**).

(41) de Silva, A. P.; Gunaratne, H. Q. N.; Habib-Jiwan, J.-L.; McCoy, C. P.; Rice, T. E.; Soumillon, J.-P. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1728–1731.

(42) de Silva, A. P.; Rice, T. E. *Chem. Commun.* **1999**, 163–164.


(43) Tian, H.; Xu, T.; Zhao, Y.; Chen, K. *J. Chem. Soc., Perkin Trans. 2* **1999**, 545–549.

(44) Greenfield, S. R.; Svec, W. A.; Gosztola, D.; Wasielewski, M. R. *J. Am. Chem. Soc.* **1996**, *118*, 6767–6777.

(45) Ferreira, R.; Remón, P.; Pischel, U. *J. Phys. Chem. C* **2009**, *113*, 5805–5811.

(46) Gao, Y. Q.; Marcus, R. A. *J. Phys. Chem. A* **2002**, *106*, 1956–1960.

The first three columns of the truth table (Table 2) describe the implementation of the T-latch function, which is mimicked by the above-discussed fluorescence switching. Starting with the triad in its unprotonated state (**2**), only low fluorescence is observed for  $T = 0$  (no addition of acid). This situation corresponds to  $Q_{\text{current}} = Q_{\text{next}} = 0$ . However, protonation of **2** with 1 equiv of acid ( $T = 1$ ) leads to  $2\text{H}^+$  and consequently a high fluorescence output (toggling from  $Q_{\text{current}} = 0$  to  $Q_{\text{next}} = 1$ ). Again, the “do nothing situation” ( $T = 0$ ) preserves the  $Q$  state (i.e.,  $Q_{\text{current}} = Q_{\text{next}} = 1$  in this case). The second addition of 1 equiv of acid ( $T = 1$ ) to  $2\text{H}^+$  yields  $2\text{H}_2^{2+}$  and concomitant fluorescence quenching, corresponding to a switching from  $Q_{\text{current}} = 1$  to  $Q_{\text{next}} = 0$ .



**Figure 3.** Recycling of fluorescence switching ( $\lambda_{\text{obs}} = 499 \text{ nm}$ ) of **2** ( $8.7 \mu\text{M}$  in acetonitrile) upon consecutive addition of 1 equiv of  $\text{CF}_3\text{SO}_3\text{H}$  followed by 1 equiv of  $\text{P}_2\text{-Et}$  phosphazene base. The dashed line marks the threshold. A conservative estimation yields that up to 10 cycles are possible, maintaining a dynamic switching range of  $I(\text{ON})/I(\text{OFF}) \geq 2$ .

The protonation state of the triad can be easily reset by application of a strong base ( $\text{P}_2\text{-Et}$  phosphazene), leading to the inverse titration curve (see Supporting Information). The consecutive protonation/deprotonation of **2** with

acid/base can be repeated for at least five cycles without significant loss of the dynamic fluorescence switching range (Figure 3). The correct functioning of the T-latch requires that the initial device state is represented by the unprotonated triad **2**. However, by solely reading the fluorescence output  $Q$ , it cannot be decided whether at a random point of operation  $Q_{\text{current}} = 0$  corresponds to **2** or  $2\text{H}_2^{2+}$ . The unambiguous assignment of an output to a concrete input situation can be resolved by reading a control channel, as has been shown previously for the implementation of reversible logic functionality.<sup>10,28,49,50</sup> This control signal is provided herein by the absorption of the quinolinium cation at ca. 313 nm (fourth column in Table 2).<sup>51</sup> This spectral signature only evolves when the quinoline unit becomes protonated (Supporting Information). Hence, when the fluorescence is low and the absorbance at 313 nm is high, 2 equiv of base is needed to reset the system to its initial state (unprotonated **2**). If the fluorescence output and the absorbance at 313 nm are both low, then the system is already in its initial state. Hence, the two  $Q = 0$  situations are now clearly distinguishable.

In summary, we have shown that an OFF-ON-OFF fluorescence switch with two degenerate proton inputs can integrate the function of a molecular T-latch. The photo-physical design of the switch is based on the control of electron transfer and hydrogen-bond interaction. Work on an all-optical version, exploring photoinduced proton transfer as relay mechanism, is underway.

**Acknowledgment.** Financial support by the Spanish MICINN (CTQ2008-06777-C02-02 for U.P., CTQ2010-20303 for E.P.-I.), the Consejería de Economía, Innovación y Ciencia—Junta de Andalucía (FQM-3685 for U.P.), FEDER, the Swedish Research Council (622-2010-280 for J.A.), and the European Research Council (FP7/2007-2013 No. 203952 for J.A.) is acknowledged.

**Supporting Information Available.** Details on the synthesis of **1–5**,  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra, additional spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(47) de Silva, A. P.; Goligher, A.; Gunaratne, H. Q. N.; Rice, T. E. *ARKIVOC* **2003**, (vii), 229–243.

(48) Veale, E. B.; Gunnlaugsson, T. *J. Org. Chem.* **2008**, *73*, 8073–8076.

(49) Remón, P.; Ferreira, R.; Montenegro, J.-M.; Suau, R.; Pérez-Inestrosa, E.; Pischel, U. *ChemPhysChem* **2009**, *10*, 2004–2007.

(50) Semeraro, M.; Credi, A. *J. Phys. Chem. C* **2010**, *114*, 3209–3214.

(51) Schulman, S. G.; Capomacchia, A. C. *J. Am. Chem. Soc.* **1973**, *95*, 2763–2766.