

Correction to “Tautomerism in 4-Hydroxypyrimidine, S-Methyl-2-thiouracil, and 2-Thiouracil” [*The Journal of Physical Chemistry A* **2010**, *114*, 12725–12730. DOI: 10.1021/jp106883s]. Barbara M. Giuliano, Vitaliy Feyer,* Kevin C. Prince, Marcello Coreno, Luca Evangelisti, Sonia Melandri, and Walther Caminati

The Gibbs free energy (ΔG) reported in Tables 1–3, calculated using the B3LYP and MP2 basis sets, and the corresponding populations (P) are corrected in the tables here.

Table 1. Relative Energies and Populations (%) of 4HP and 4PO, at the B3LYP and MP2 Levels of Approximation, with 6-311++G(d,p) Basis Sets^a


| | 4HP | | 4PO | |
|--|-------|------|-------|------|
| | B3LYP | MP2 | B3LYP | MP2 |
| $\Delta E/\text{kJ mol}^{-1}$ | 8.4 | 0.0 | 0.0 | 4.2 |
| $\Delta E_{\text{ZPE}}/\text{kJ mol}^{-1}$ | 8.3 | 0.0 | 0.0 | 3.6 |
| $\Delta G_{(348\text{ K})}/\text{kJ mol}^{-1}$ | 8.7 | 0.0 | 0.0 | 2.3 |
| $P_{(348\text{ K})}/\%$ | 4.7 | 69.1 | 95.3 | 30.9 |
| $\Delta G_{(365\text{ K})}/\text{kJ mol}^{-1}$ | 8.8 | 0.0 | 0.0 | 2.3 |
| $P_{(365\text{ K})}/\%$ | 5.3 | 68.3 | 94.7 | 31.7 |



^a ΔE = electronic energy difference. ΔE_{ZPE} = zero-point corrected energy difference. ΔG = Gibbs free energy difference calculated at 348 and 365 K. P = population in % of the tautomers, estimated using the calculated ΔG value.

Table 2. Relative Energies and Populations (%) of the Most Stable Species of 2TU, at the B3LYP and MP2 Levels of Approximation, with 6-311++G(d,p) Basis Sets^a





| | 2TU/diketo | | 2TU/keto | | 2TU/enol | |
|--|------------|------|----------|------|----------|------|
| | B3LYP | MP2 | B3LYP | MP2 | B3LYP | MP2 |
| $\Delta E/\text{kJ mol}^{-1}$ | 0.0 | 0.0 | 48.8 | 36.4 | 52.3 | 28.7 |
| $\Delta E_{\text{ZPE}}/\text{kJ mol}^{-1}$ | 0.0 | 0.0 | 38.2 | 23.8 | 42.1 | 17.4 |
| $\Delta G_{(339\text{ K})}/\text{kJ mol}^{-1}$ | 0.0 | 0.0 | 37.4 | 19.9 | 42.2 | 16.1 |
| $P_{(339\text{ K})}/\%$ | 100.0 | 99.4 | 0.0 | 0.1 | 0.0 | 0.5 |
| $\Delta G_{(435\text{ K})}/\text{kJ mol}^{-1}$ | 0.0 | 0.0 | 36.8 | 18.0 | 42.0 | 16.0 |
| $P_{(435\text{ K})}/\%$ | 100.0 | 98.1 | 0.0 | 0.7 | 0.0 | 1.2 |



^a ΔE = electronic energy difference. ΔE_{ZPE} = zero-point corrected energy difference. ΔG = Gibbs free energy difference calculated at 339 and 435 K. P = population in % of the tautomers, estimated using the calculated ΔG value.

Table 3. Relative Energies and Populations (%) of the Most Stable Species of SM2T, at the B3LYP and MP2 Levels of Approximation, with 6-311++G(d,p) Basis Sets^a

| | SM2T/hydroxyl-1 | | SM2T/oxo-1 | | SM2T/hydroxyl-2 | | SM2T/oxo-2 | |
|--|-----------------|------|------------|------|-----------------|------|------------|------|
| | B3LYP | MP2 | B3LYP | MP2 | B3LYP | MP2 | B3LYP | MP2 |
| $\Delta E/\text{kJ mol}^{-1}$ | 6.6 | 0.0 | 0.0 | 5.2 | 9.2 | 2.7 | 12.5 | 18.7 |
| $\Delta E_{\text{ZPE}}/\text{kJ mol}^{-1}$ | 6.7 | 0.0 | 0.0 | 4.2 | 9.1 | 2.6 | 12.4 | 18.7 |
| $\Delta G_{(343\text{ K})}/\text{kJ mol}^{-1}$ | 8.2 | 0.0 | 0.0 | 1.5 | 10.6 | 2.5 | 13.2 | 18.4 |
| $P_{(343\text{ K})}/\%$ | 4.9 | 48.9 | 92.2 | 29.9 | 2.1 | 21.1 | 0.8 | 0.1 |
| $\Delta G_{(360\text{ K})}/\text{kJ mol}^{-1}$ | 8.3 | 0.0 | 0.0 | 1.3 | 10.7 | 2.5 | 13.3 | 18.4 |
| $P_{(360\text{ K})}/\%$ | 5.4 | 48.0 | 91.2 | 31.1 | 2.4 | 20.8 | 1.0 | 0.1 |



^a ΔE = electronic energy difference. ΔE_{ZPE} = zero-point corrected energy difference. ΔG = Gibbs free energy difference calculated at 343 and 360 K. P = population in % of the tautomers, estimated using the calculated ΔG value.

DOI: 10.1021/jp204996u

Published on Web 06/22/2011