



# Iron(II)-Catalyzed Intermolecular Amino-Oxygenation of Olefins through the N–O Bond Cleavage of Functionalized Hydroxylamines

Deng-Fu Lu, Cheng-Liang Zhu, Zhen-Xin Jia, and Hao Xu\*

Department of Chemistry, Georgia State University, 100 Piedmont Avenue SE, Atlanta, Georgia 30303, United States

**S** Supporting Information

**ABSTRACT:** An iron-catalyzed diastereoselective *intermolecular* olefin amino-oxygenation reaction is reported, which proceeds via an iron-nitrenoid generated by the N–O bond cleavage of a functionalized hydroxylamine. In this reaction, a bench-stable hydroxylamine derivative is used as the amination reagent and oxidant. This method tolerates a range of synthetically valuable substrates that have been all incompatible with existing amino-oxygenation methods. It can also provide amino alcohol derivatives with regio- and stereochemical arrays complementary to known amino-oxygenation methods.


Selective olefin difunctionalization with an amino- and an oxygen-based group is an important transformation for organic synthesis because vicinal amino alcohol derivatives are widely present in synthetically valuable molecules. The osmium-based Sharpless aminohydroxylation continues to be a prevalent stereospecific method for olefin amino-oxygenation.<sup>1</sup> This pioneering method has also inspired extensive efforts for the development of alternative approaches for a broader substrate scope and better regioselectivity.<sup>2,3</sup> Among these approaches, nonprecious metal-catalyzed processes emerge with increasing interest: Chemler developed Cu-catalyzed methods for olefin amino-oxygenation and other difunctionalizations;<sup>2a–d</sup> Yoon developed Cu- and Fe-catalyzed sulfonyl oxaziridine based methods.<sup>2e–h</sup> Despite these and other excellent discoveries, new nonprecious metal-catalyzed olefin amino-oxygenation methods which achieve a broader substrate scope and regio- and stereoselectivity complementary to known methods are greatly desirable. In particular, the *intermolecular* olefin amino-oxygenation mediated by an iron nitrenoid has not been reported.

Unlike the N-atom transfer mediated by a rhodium nitrenoid,<sup>3,4</sup> the iron nitrenoid mediated process is more prone to proceed through radical pathways.<sup>5</sup> Therefore, new strategies are required to control an iron nitrenoid's reactivity in an olefin amino-oxygenation reaction. We have previously discovered iron-catalyzed *intramolecular* olefin amino-oxygenation and amino-fluorination reactions (Scheme 1A).<sup>6</sup> Our studies suggested that an iron nitrenoid is a possible intermediate in these stereoconvergent transformations and that the stereoselectivity can be modulated by N-based bidentate ligands.<sup>7</sup>


Our initial attempts to develop an *intermolecular* olefin amino-oxygenation with the catalyst previously identified to be effective for *intramolecular* amino-oxygenation failed due to the lack of reactivity. To modulate the reactivity of the iron nitrenoid to achieve a fine balance between reactivity, stability, and selectivity,

## Scheme 1. Iron-Catalyzed Olefin Amino-Oxygenation with Functionalized Hydroxylamines

A) previous work: intramolecular olefin amino-oxygenation



B) current work: intermolecular olefin amino-oxygenation



we explored a variety of new amination reagents, iron catalysts, and ligands. Herein, we disclose an iron-catalyzed intermolecular olefin amino-oxygenation that proceeds through the N–O bond cleavage of a functionalized hydroxylamine. In this transformation, a bench-stable hydroxylamine derivative is applied as the amination reagent and oxidant (Scheme 1B).

This method has a few unique features that complement the existing iron-catalyzed olefin amino-oxygenation method with sulfonyl oxaziridines.<sup>2g,h</sup> First, this method allows significant asymmetric induction with internal olefinic substrates, while the oxaziridine-based asymmetric approach is only effective for terminal olefins. Second, this method tolerates a broad range of synthetically valuable substrates, including allyl silanes, cyclopentadienes, enol ethers, glycals, indene, and silyl dienols, which are all incompatible with the iron-catalyzed olefin amino-oxygenation method with sulfonyl oxaziridines. Furthermore, this method can effectively afford amino alcohol derivatives with regio- and stereochemical arrays complementary to existing amino-oxygenation methods, especially osmium-based approaches. Therefore, we envision that this discovery will be a valuable tool for selective olefin amino-oxygenation.

Styrene **1** was selected as a model substrate for catalyst discovery (Table 1). Our initial attempts with Fe(OTf)<sub>2</sub>–N,N'-bidentate ligands failed due to the lack of reactivity. Inspection of a range of ligands revealed that the N-based tridentate ligands are necessary for the proposed reactivity and that an achiral bisoxazoline PyBOX ligand **L1** is uniquely effective:<sup>8</sup> the Fe(OTf)<sub>2</sub>–**L1** complex catalyzes the styrene amino-oxygenation with a range of functionalized hydroxyl amines (**2a–2d**, entries 1–4), affording both an alkoxyl oxazoline **3** and a protected amino alcohol **4** with good to excellent combined yields and regioselectivity complementary to the osmium-based methods.<sup>1a</sup>


**Received:** August 6, 2014

**Published:** August 28, 2014



**Table 2. Substrate Scope for the Iron-Catalyzed Olefin Amino-Oxygenation**

### Scheme 2. Iron-Catalyzed Asymmetric Olefin Amino-Oxygenation and Control Experiments To Probe Reaction Mechanisms



<sup>a</sup>LAH, THF,  $-20^{\circ}\text{C}$ , 85%. <sup>b</sup>  $\text{Fe}(\text{OTf})_2$  (10 mol %), **L1** (10 mol %),  $\text{CH}_2\text{Cl}_2/\text{MeCN}$  (15:1),  $-15^{\circ}\text{C}$ , 1 h. <sup>c</sup>  $\text{Fe}(\text{ClO}_4)_2$  (20 mol %), **L1** (20 mol %),  $\text{CH}_2\text{Cl}_2/\text{MeCN}$  (15:1),  $-15^{\circ}\text{C}$ , 2 h. <sup>d</sup>  $\text{R}^1$ :  $\text{CF}_3\text{CH}_2$ ;  $\text{R}^2$ : 2,4- $\text{Cl}_2$ -benzoyl.


with either aliphatic or aromatic substituents can also be efficiently transformed into protected 1,2-amino alcohols with excellent regioselectivity (entries 11–13).<sup>12</sup> To our delight, this method is also compatible with a silyl dienol with a labile C–H bond (entry 14, 63% yield). Also, a dienolate proves to be an acceptable substrate for this transformation (entry 15, 54% yield).

Additionally, we applied this method to isolated olefins. The  $\text{Fe}(\text{ClO}_4)_2$ –**L1** complex can catalyze the amino-oxygenation of a 1,1-disubstituted olefin with **2b**, affording an oxazolidinone (entry 16, 51% yield).<sup>13</sup> We further observed that the  $\text{Fe}(\text{OTf})_2$ –**L1** complex catalyzes the reaction of a monosubstituted olefin with **2c** to afford the oxazolidinone with a fair yield (entry 17, 48% yield).

The catalytic asymmetric amino-oxygenation of indene **6** has been a challenge in synthetic chemistry, and osmium-based protocols deliver a mixture of racemic 1- and 2-amino indanols.<sup>11</sup> In order to fill this gap, we have explored the asymmetric induction for the indene amino-oxygenation and discovered that an iron–chiral ligand **L4**<sup>14</sup> complex is uniquely effective to deliver a 2-amino indanol derivative **7** with a significant *ee* (Scheme 2A, 81% *ee*, *dr* > 20:1). Facile transformation converts **7** to **8** without erosion of its *ee* and *dr*. The asymmetric enol ether amino-oxygenation has also been unprecedented, and we observed that **L4** is effective for asymmetric induction with dihydrofuran **9** as well (57% *ee*, *dr* > 20:1).

In order to gather evidence for a mechanistic working hypothesis, we have carried out several control experiments (Scheme 2B). First, two analogues (**11** and **12**) of reagent **2c** were prepared, such that the N–H group was masked by either an acetyl or a methyl group. Both were evaluated for the model reaction and neither was found to be reactive (eq a). These experiments suggest that the N–H group in **2c** is critical for its activation. Next, we evaluated a cyclopropyl-substituted olefin **13**

### Scheme 3. Mechanistic Working Hypothesis for the Iron-Catalyzed Olefin Amino-Oxygenation



as a radical clock probe under the reaction conditions and observed the presence of both the ring-opening product **14** and the 1,2-amino-oxygenation product **15** (eq b). This result suggests that the reaction proceeds through a stepwise process that includes a radical amination step.

To probe the mechanism beyond the radical amination step, we further evaluated an isopropyl-substituted terminal olefin **16** (eq c). If a carbocation is generated after the radical amination, 1,2-hydride shift products may be observed. The amino-oxygenation with **16** afforded four products: in addition to the standard 1,2-amino-oxygenation product **17**, an 1,3-amino-oxygenation product **18**, aziridine **19**, and allylic amine **20** were isolated (eq c). Importantly, **19** cannot be converted to any of the three other products under the reaction conditions. These results suggest that a carbocation may be involved in the olefin amino-oxygenation and that the corresponding aziridine is unlikely an intermediate along this pathway.

Furthermore, we studied *cis/trans*  $\beta$ -methyl styrenes as mechanistic probes and the experimental results corroborate that the amino-oxygenation occurs in a stepwise fashion and they also suggest that the C–N bond formation is likely the rate-determining step.<sup>15</sup> Finally, we evaluated the electronic effect on styrene amino-oxygenation and concluded that amino alcohol formation is favored with substrates that can stabilize electrophilic radical species.<sup>15</sup>

Based upon the collective evidence, a mechanistic working hypothesis of olefin amino-oxygenation that best corroborates the experimental data is presented in Scheme 3. First, the iron–ligand complex may reductively cleave the N–O bond in **2c**, possibly converting it to an iron–nitrenoid **A**. **A** may then initiate radical amination with olefin **21** to afford radical species **B1** together with its conformer **B2** in equilibrium. Presumably, **B1** can be oxidized by the iron center to a carbocation **B3**,<sup>16,17</sup> which will be rapidly captured by the neighboring carbamate group, thereby affording **22**. Alternatively, oxidative carboxylate ligand transfer<sup>16</sup> may directly occur with **B2** to afford the protected amino alcohol **31**. We still cannot completely rule out the possibility that electron transfer from **B2** to the iron center occurs first and that the oxidation product **B4** will then be captured by a carboxylate to deliver **23**. When the substituent ( $\text{R}^3$ ) has a less significant radical-stabilizing effect, **B1** and **B2** are relatively short-lived high energy species; therefore, the oxidative neighboring group participation through **B1** may be favored to afford **22**. However, when the substituent has a strong radical-stabilizing effect and both species are relatively long-lived, the ligand transfer from the iron center through **B2** may become dominant to deliver **23**.



In conclusion, we have discovered a new iron-catalyzed stereoselective olefin amino-oxygenation method. This method tolerates a broad range of synthetically valuable olefins including those that are incompatible with existing amino-oxygenation methods. Our preliminary mechanistic studies revealed that an iron nitrenoid is a possible intermediate and its enantioselectivity can be controlled by chiral ligands. This discovery demonstrates the feasibility of developing a unique approach for iron-catalyzed selective olefin difunctionalization. Our ongoing efforts focus on understanding the mechanism of this new reaction and its applications in organic synthesis.

## ■ ASSOCIATED CONTENT

### ■ Supporting Information

Experimental procedure, characterization data for all new compounds, selected NMR spectra, and HPLC traces. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

[hxu@gsu.edu](mailto:hxu@gsu.edu)

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by the NIH (GM110382), Georgia State University, and the ACS Petroleum Research Fund (ACS PRF 51571-DN11). We thank Jeffrey Sears for synthesis of a few substrates.

## ■ REFERENCES

- (1) (a) Sharpless, K. B.; Chong, A. O.; Oshima, K. *J. Org. Chem.* **1976**, *41*, 177. (b) Li, G. G.; Chang, H. T.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **1996**, *35*, 451.
- (2) For selected references of olefin amino-oxygenation and related transformations, see: (a) Fuller, P. H.; Kim, J.-W.; Chemler, S. R. *J. Am. Chem. Soc.* **2008**, *130*, 17638. (b) Paderes, M. C.; Chemler, S. R. *Org. Lett.* **2009**, *11*, 1915. (c) Sequeira, F. C.; Chemler, S. R. *Org. Lett.* **2012**, *14*, 4482. (d) Turnpenny, B. W.; Chemler, S. R. *Chem. Sci.* **2014**, *5*, 1786. (e) Michaelis, D. J.; Shaffer, C. J.; Yoon, T. P. *J. Am. Chem. Soc.* **2007**, *129*, 1866. (f) Benkovics, T.; Guzei, I. A.; Yoon, T. P. *Angew. Chem., Int. Ed.* **2010**, *49*, 9153. (g) Williamson, K. S.; Yoon, T. P. *J. Am. Chem. Soc.* **2010**, *132*, 4570. (h) Williamson, K. S.; Yoon, T. P. *J. Am. Chem. Soc.* **2012**, *134*, 12370. (i) Alexanian, E. J.; Lee, C.; Sorensen, E. J. *J. Am. Chem. Soc.* **2005**, *127*, 7690. (j) Liu, G.; Stahl, S. S. *J. Am. Chem. Soc.* **2006**, *128*, 7179. (k) Desai, L. V.; Sanford, M. S. *Angew. Chem., Int. Ed.* **2007**, *46*, 5737. (l) Weinstein, A. B.; Schuman, D. P.; Tan, Z. X.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2013**, *52*, 11867. (m) Zhu, H.; Chen, P.; Liu, G. *J. Am. Chem. Soc.* **2014**, *136*, 1766. (n) Muñoz, K.; Iglesias, A.; Fang, Y. *Chem. Commun.* **2009**, 5591. (o) de Haro, T.; Nevado, C. *Angew. Chem., Int. Ed.* **2011**, *50*, 906. (p) Wardrop, D. J.; Bowen, E. G.; Forslund, R. E.; Sussman, A. D.; Weerasekera, S. L. *J. Am. Chem. Soc.* **2010**, *132*, 1188. (q) Lovick, H. M.; Michael, F. E. *J. Am. Chem. Soc.* **2010**, *132*, 1249. (r) Schmidt, V. A.; Alexanian, E. J. *J. Am. Chem. Soc.* **2011**, *133*, 11402. (s) Xu, H.-C.; Moeller, K. D. *J. Am. Chem. Soc.* **2008**, *130*, 13542. (t) Donohoe, T. J.; Johnson, P. D.; Cowley, A.; Keenan, M. *J. Am. Chem. Soc.* **2002**, *124*, 12934. (u) Harris, L.; Mee, S. P. H.; Furneaux, R. H.; Gainsford, G. J.; Luxemburger, A. *J. Org. Chem.* **2011**, *76*, 358. (v) Farid, U.; Wirth, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 3462. (w) Mahoney, J. M.; Smith, C. R.; Johnston, J. N. *J. Am. Chem. Soc.* **2005**, *127*, 1354. (x) Zhang, B.; Studer, A. *Org. Lett.* **2013**, *15*, 4548.
- (3) For selected references of rhodium nitrenoid-mediated amino-acetoxylation, see: (a) Levites-Agababa, E.; Menhaji, E.; Perlson, L. N.; Rojas, C. M. *Org. Lett.* **2002**, *4*, 863. (b) Padwa, A.; Stengel, T. *Org. Lett.*

**2002**, *4*, 2137. (c) Mulcahy, J. V.; Du Bois, J. *J. Am. Chem. Soc.* **2008**, *130*, 12630. (d) Beaumont, S.; Pons, V.; Retaillieu, P.; Dodd, R. H.; Dauban, P. *Angew. Chem., Int. Ed.* **2010**, *49*, 1634. (e) Gigant, N.; Dequierez, G.; Retaillieu, P.; Gillaizeau, I.; Dauban, P. *Chem.—Eur. J.* **2012**, *18*, 90. (f) Dequierez, G.; Ciesielski, J.; Retaillieu, P.; Dauban, P. *Chem.—Eur. J.* **2014**, *20*, 8929. For a manganese-mediated stoichiometric method, see: (g) Du Bois, J.; Tomooka, C. S.; Hong, J.; Carreira, E. M. *J. Am. Chem. Soc.* **1997**, *119*, 3179.

(4) For selected references of rhodium nitrenoid-mediated  $sp^3$  C—H amination, see: (a) Espino, C. G.; Du Bois, J. *Angew. Chem., Int. Ed.* **2001**, *40*, 598. (b) Zalatan, D. N.; Du Bois, J. *J. Am. Chem. Soc.* **2008**, *130*, 9220. (c) Nguyen, Q.; Sun, K.; Driver, T. G. *J. Am. Chem. Soc.* **2012**, *134*, 7262. For selected references of olefin aziridination through Cu and Rh metallonitrenes, see: (d) Li, Z.; Conser, K. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1993**, *115*, 5326. (e) Evans, D. A.; Faul, M. M.; Bilodeau, M. T.; Anderson, B. A.; Barnes, D. M. *J. Am. Chem. Soc.* **1993**, *115*, 5328. (f) Guthikonda, K.; Du Bois, J. *J. Am. Chem. Soc.* **2002**, *124*, 13672. (g) Lebel, H.; Lectard, S.; Parmentier, M. *Org. Lett.* **2007**, *9*, 4797. (h) Jat, J. L.; Paudyal, M. P.; Gao, H.; Xu, Q.-L.; Yousufuddin, M.; Devarajan, D.; Ess, D. H.; Kürti, L.; Falck, J. R. *Science* **2014**, *343*, 61. (5) For selected references of iron-nitrenoid mediated N-atom transfer, see: (a) Breslow, R.; Gellman, S. H. *J. Chem. Soc., Chem. Commun.* **1982**, 1400. (b) Mahy, J. P.; Battioni, P.; Mansuy, D. *J. Am. Chem. Soc.* **1986**, *108*, 1079. (c) Nakanishi, M.; Salit, A. F.; Bolm, C. *Adv. Synth. Catal.* **2008**, *350*, 1835. (d) Paradine, S. M.; White, M. C. *J. Am. Chem. Soc.* **2012**, *134*, 2036. (e) Hennessy, E. T.; Betley, T. A. *Science* **2013**, *340*, 591. (f) Liu, Y.; Guan, X.; Wong, E. L.-M.; Liu, P.; Huang, J.-S.; Che, C.-M. *J. Am. Chem. Soc.* **2013**, *135*, 7194. (g) Bach, T.; Schlummer, B.; Harms, K. *Chem. Commun.* **2000**, 287. (h) Churchill, D. G.; Rojas, C. M. *Tetrahedron Lett.* **2002**, *43*, 7225.

(6) (a) Liu, G.-S.; Zhang, Y.-Q.; Yuan, Y.-A.; Xu, H. *J. Am. Chem. Soc.* **2013**, *135*, 3343. (b) Zhang, Y.-Q.; Yuan, Y.-A.; Liu, G.-S.; Xu, H. *Org. Lett.* **2013**, *15*, 3910. (c) Lu, D.-F.; Liu, G.-S.; Zhu, C.-L.; Yuan, B.; Xu, H. *Org. Lett.* **2014**, *16*, 2912.

(7) For leading references of chiral BOX and relevant ligands, see: (a) Evans, D. A.; Woerpel, K. A.; Hinman, M. M.; Faul, M. M. *J. Am. Chem. Soc.* **1991**, *113*, 726. (b) Nishiyama, H.; Itoh, Y.; Matsumoto, H.; Park, S.-B.; Itoh, K. *J. Am. Chem. Soc.* **1994**, *116*, 2223. (c) Nishikawa, Y.; Yamamoto, H. *J. Am. Chem. Soc.* **2011**, *133*, 8432.

(8) For details of ligand synthesis, see Supporting Information (SI).

(9) For stereochemistry determination, see SI.

(10) For a manganese-mediated glycol aziridination for 2-amino sugar synthesis, see: Reference 3g. For a selected indirect method for 2-amino sugar synthesis, see: Griffith, D. A.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1990**, *112*, 5811.

(11) Osmium-based aminohydroxylation of indene affords racemic mixtures of 1- and 2-amino indanols; see: (a) Masruri; Willis, A. C.; McLeod, M. D. *J. Org. Chem.* **2012**, *77*, 8480. For an indirect method of 2-amino indanol synthesis, see: (b) Corey, E. J.; Roper, T. D.; Ishihara, K.; Sarakinos, G. *Tetrahedron Lett.* **1993**, *34*, 8399.

(12)  $\text{Fe}(\text{OTf})_2/\text{FeCl}_2$  mixed salts were applied as the catalyst in entry 11.  $\text{Fe}(\text{OTf})_2$  led to rapid decomposition of the diene;  $\text{FeCl}_2$  was inactive. See SI for details.

(13) Other Fe salts led to rapid allylic amination products. See SI for details.

(14) Davies, I. W.; Gerena, L.; Lu, N.; Larsen, R. D.; Reider, P. J. *J. Org. Chem.* **1996**, *61*, 9629.

(15) *Cis/trans* isomeric olefins have been observed to present different reactivity and selectivity in atom transfer reactions that proceed via a stepwise mechanism. For a selected reference, see: Lee, N. H.; Jacobsen, E. N. *Tetrahedron Lett.* **1991**, *32*, 6533. See SI for experimental details.

(16) For the oxidation of a radical species by a high-valent metal through ligand transfer or electron transfer, see: (a) Kharasch, M. S.; Sosnovsky, G. *J. Am. Chem. Soc.* **1958**, *80*, 756. (b) Kochi, J. K. *Science* **1967**, *155*, 415.

(17) For evidence for involvement of a possible carbocation intermediate, see eq c in Scheme 2B. For control experiments that exclude the aziridine as a possible intermediate, see SI.