


Solvent-Dependent Divergent Functions of $\text{Sc}(\text{OTf})_3$ in Stereoselective Epoxide-Opening Spiroketalizations

Indrajeet Sharma,[†] Jacqueline M. Wurst,[‡] and Derek S. Tan^{*,†,‡}

[†]Molecular Pharmacology & Chemistry Program, [‡]Tri-Institutional PhD Program in Chemical Biology, and Tri-Institutional Research Program, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, Box 422, New York, New York 10065, United States

S Supporting Information

ABSTRACT: A stereocontrolled synthesis of benzannulated spiroketals has been developed using solvent-dependent $\text{Sc}(\text{OTf})_3$ -mediated spirocyclizations of *exo*-glycal epoxides having alcohol side chains. In THF, the reaction proceeds via Lewis acid catalysis under kinetic control with inversion of configuration at the anomeric carbon. In contrast, in CH_2Cl_2 , Brønsted acid catalysis under thermodynamic control leads to retention of configuration. The reactions accommodate a variety of aryl substituents and ring sizes and provide stereochemically diverse spiroketals.



Benzannulated spiroketal natural products exhibit a broad array of biological activities.¹ Examples include the matrix metalloproteinase inhibitor berkeleyic acid,² the fungal cell wall glucan synthase inhibitory papulacandins,³ and the anti-inflammatory aquilarinoside A.⁴ Bisbenzannulated spiroketals include the rubromycin family of human telomerase and HIV reverse transcriptase inhibitors,⁵ the DNA helicase inhibitor heliquinomycin, and the antibiotic purpurumycin, which inhibits aminoacyl-tRNA synthesis by a novel mechanism involving direct binding to the tRNA substrate.⁶ Notably, the benzannulated spiroketal core is essential for telomerase inhibition in the rubromycin family.⁷ Numerous approaches to the synthesis of benzannulated spiroketals have been reported.^{8,9} Despite these notable advances, most strategies rely upon thermodynamically controlled reactions that often lead to stereoisomeric mixtures at the anomeric carbon.¹

We have previously developed stereocontrolled approaches to aliphatic spiroketals using stereocomplementary kinetic spirocyclization reactions of *endo*-glycal epoxides that proceed with either inversion or retention of configuration at the anomeric carbon, independent of thermodynamic preferences.¹⁰ We have also extended this approach to benzannulated spiroketals via incorporation of aromatic rings on the cyclizing side chain.¹¹ Unfortunately, this approach provides low diastereoselectivity in spirocyclization reactions with phenolic nucleophiles (45:55 to 58:42 dr).¹¹

To address this problem, we envisioned an alternative entry to phenolic spiroketals involving stereoselective spirocyclizations of benzannulated *exo*-glycal epoxides (dihydrobenzofuran spiroepoxides). Spirocyclizations of exocyclic enol ether epoxides have apparently not been explored previously, although classical acid-catalyzed spiroketalizations of the parent exocyclic enol ethers are well-known,¹² and the corresponding epoxides¹³ have been used in intermolecular alcohol additions.¹⁴

Thus, the requisite benzannulated *exo*-glycal epoxide substrates were synthesized from salicylaldehydes **1** via alkyne additions to form propargyl alcohols **2a–h** (Figure 1).¹⁵ Au(I)-mediated cycloisomerization, previously restricted to aromatic alkynes,^{15,16} then afforded *exo*-glycals **3a–h**. Diastereoselective *anti*-epoxidation with dimethyldioxirane (DMDO)¹⁷ provided *exo*-glycal epoxides **6a–h**. Interestingly, these epoxides were




Figure 1. Synthesis of *exo*-glycal epoxide substrates **6a–h**.


Received: March 21, 2014

Published: April 17, 2014

stable upon warming to rt, in stark contrast to the corresponding *endo*-glycal epoxides, which cyclize spontaneously at $-35\text{ }^{\circ}\text{C}$.¹⁸

We next explored spirocyclization reactions of benzannulated *exo*-glycal epoxide **6a** (Table 1). Notably, **6a** proved unreactive

Table 1. Spirocyclization Reactions of *exo*-Glycal Epoxide **6a^a**



entry	reagent (equiv)	solvent, temp ($^{\circ}\text{C}$)	7a:8a
1	MeOH (excess)	MeOH, rt	NR
2	Ti(O ^{<i>i</i>} Pr) ₄ (2.0)	CH ₂ Cl ₂ , rt	NR
3		toluene, 120	NR
4	Sc(OTf) ₃ (2.0)	CH ₂ Cl ₂ , $-78 \rightarrow 0$	75:25
5	Sc(OTf) ₃ (2.0)	THF, $-78 \rightarrow 0$	>98:2
6	Sc(OTf) ₃ (1.0)	THF, $-78 \rightarrow 0$	>98:2
7	Sc(OTf) ₃ (0.1)	THF, $-78 \rightarrow 0$	93:7
8	Sc(OTf) ₃ (1.0)	THF, -20	>98:2
9	Sc(OTf) ₃ (1.0)	THF, rt	90:10
10	Sc(OTf) ₃ (1.0)	CH ₂ Cl ₂ , $0 \rightarrow \text{rt}$	<2:98
11	Sc(OTf) ₃ (0.5)	CH ₂ Cl ₂ , $0 \rightarrow \text{rt}$	<2:98
12	Sc(OTf) ₃ + DTBMP (1.0 + 1.0)	THF, -20	>98:2
13	ScCl ₃ (1.0)	THF, rt	>98:2
14	TfOH (1.0)	THF, -20	30:70
15	Sc(OTf) ₃ + DTBMP (0.5 + 0.5)	CH ₂ Cl ₂ , $0 \rightarrow \text{rt}$	75:25
16	TfOH (0.5)	CH ₂ Cl ₂ , $0 \rightarrow \text{rt}$	<2:98
17	TfOH + DTBMP (0.5 + 0.5)	CH ₂ Cl ₂ , $0 \rightarrow \text{rt}$	51:49

^aProduct ratios determined by ¹H NMR analysis of crude reaction products. NR = no reaction; DTBMP = 2,6-di-*tert*-butyl-4-methylpyridine. See the Supporting Information for the complete table.

under our previously reported MeOH and Ti(O-*i*-Pr)₄ spirocyclization conditions, as well as upon heating to $120\text{ }^{\circ}\text{C}$ in toluene (entries 1–3). After investigating a wide range of Lewis acids,¹⁶ we were encouraged to find that Sc(OTf)₃ favored the inversion product **7a** (entry 4), which could be formed exclusively by changing the reaction solvent from CH₂Cl₂ to THF (entry 5). The diastereoselectivity decreased slightly with substoichiometric Sc(OTf)₃ (entries 6 and 7). Other Lewis acids gave lower or even reversed diastereoselectivity.¹⁶

In low-temperature ¹H NMR experiments, we found that the Sc(OTf)₃-mediated spirocyclization begins to occur at $-35\text{ }^{\circ}\text{C}$.¹⁶ Complete selectivity for spirocyclization with inversion of configuration was maintained when the reaction was run at $-20\text{ }^{\circ}\text{C}$ (entry 8), but selectivity decreased at higher temperatures (entry 9), suggesting that the reaction proceeds under kinetic control between -35 and $-20\text{ }^{\circ}\text{C}$.

Strikingly, when the room-temperature reaction was carried out in CH₂Cl₂ instead of THF, thermodynamic equilibration of an initially formed diastereomeric mixture afforded the retention product **8a** with complete stereoselectivity (entries 10, 11).¹⁹ A structural rationale for the observed thermodynamic preference is nonobvious, due to the conformational flexibility of 5-membered rings,²⁰ and remains a subject for further investigation. However, on the basis of these results, it is apparent that Sc(OTf)₃ plays divergent roles in the

spirocyclization reactions depending upon solvent selection (THF vs CH₂Cl₂, entry 9 vs 10).

It is known that metal triflates can serve as a mild source of triflic acid.²¹ Thus, we carried out mechanistic studies to differentiate between the Lewis and Brønsted acid activities of Sc(OTf)₃. Inclusion of the noncoordinating Brønsted base, 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP),^{21a} in the reaction in THF did not affect diastereoselectivity (entry 8 vs 12). Treatment with ScCl₃ at rt also led to complete stereoselectivity for the contrathermodynamic spiroketal **7a** (entry 13). In contrast, spirocyclization with TfOH afforded a diastereomeric mixture favoring the retention product **8a** (entry 14). Taken together, these results suggest that Sc(OTf)₃ acts as a Lewis acid in THF at reduced temperatures, catalyzing formation of the contrathermodynamic spiroketal **7a** under kinetic control.

We next carried out the analogous experiments in CH₂Cl₂ where, upon warming to rt, Sc(OTf)₃ favors the retention product **8a** (entry 11). In contrast, inclusion of DTBMP resulted in a diastereomeric mixture favoring the inversion product **7a** (entry 15). However, spirocyclization with TfOH provided the retention product **8a** exclusively (entry 16). Treatment with both TfOH and DTBMP afforded a diastereomeric mixture of spiroketals, similar to the result observed with Sc(OTf)₃ and DTBMP (entry 17 vs 15). Collectively, these results suggest that Sc(OTf)₃ acts as a mild source of Brønsted acid in CH₂Cl₂ at rt, catalyzing formation of the thermodynamically favored spiroketal **8a** under equilibrium control.

We then investigated the scope of these stereocomplementary Sc(OTf)₃-catalyzed spirocyclization reactions. Substrates with longer side chains (**6b**, **6c**) and various aryl substituents (**6d–h**) were synthesized from the corresponding alkyne and salicylaldehyde precursors (Figure 1). The bromide intermediate **4h** was also used to introduce other substituents (aryl, alkyne, azide, aldehyde, ester, imide) in **4i–n** to examine the functional group tolerance of the spirocyclization reactions (Figure S2, Supporting Information).¹⁶ The *exo*-glycals **4i–n** were then converted to the corresponding epoxide substrates **6i–n** (Figure S2, Supporting Information).¹⁶

In the spirocyclization reactions, both diastereomers of the larger 6- and 7-membered ring spiroketals (**7b,c** and **8b,c**) could be obtained with complete diastereoselectivity based on solvent selection (Figure 2). For **8b**, equilibration with Sc(OTf)₃ in CH₂Cl₂ required elevated temperature ($60\text{ }^{\circ}\text{C}$). The 7-membered ring spiroketal **7c** was obtained in somewhat lower yield due to an unexpected anti-Markovnikov 6-*exo* epoxide opening side reaction leading to a benzofuran product.¹⁶

Next, we investigated the electronic effects of various aryl substituents. A wide range of electron-withdrawing and -donating groups were tolerated (**7d–n**, **8d–n**), and high diastereoselectivities were maintained. Notably, the nitro-substituted substrate **6d** was less reactive and required more forcing conditions (**7d**: rt; **8d**: 6 h). Conversely, the methoxy-substituted substrate **6e** was highly reactive, providing slightly decreased diastereoselectivity in the THF reaction (**7e**: 93:7 dr) and rapid equilibration in the CH₂Cl₂ reaction (**8e**: 1 h). These results are consistent with the expected electronic influence of these *para* substituents upon the reactive anomeric spiroepoxide center.¹⁸

The reactions also tolerated other reactive functionalities including alkyne (**7j**, **8j**), azide (**7k**, **8k**), aldehyde (**7l**, **8l**), ester (**7m**, **8m**), and phthalimide (**7n**, **8n**) groups. In the case of




Figure 2. Scope of Sc(OTf)₃-mediated spirocyclization reactions. (a) THF: 1.0 equiv Sc(OTf)₃, THF, −20 °C, 2–3 h; CH₂Cl₂: 0.5 equiv Sc(OTf)₃, CH₂Cl₂, 0 °C to rt, 1–12 h; diastereomeric ratios determined by ¹H NMR analysis of crude reaction mixtures; stereochemistry assigned based on NOESY analysis except 8b, which was determined by X-ray crystallography;¹⁶ isolated yields after column chromatography shown in parentheses. (b) 60 °C. (c) 30% anti-Markovnikov 6-*exo*-cyclization side product also recovered.¹⁶ (d) rt.

azide **8k**, Sc(OTf)₃ equilibration in CH₂Cl₂ required elevated temperature (60 °C).

In conclusion, we have developed novel, solvent-dependent Sc(OTf)₃-mediated spirocyclizations of *exo*-glycal epoxides for the stereocontrolled synthesis of benzannulated spiroketals. This *exo*-glycal-based approach overcomes a key limitation of our previous *endo*-glycal-based approach and tolerates a wide range of functionalities. Applications to the diversity-oriented synthesis of stereochemically diverse spiroketal libraries are ongoing and will be reported in due course.

■ ASSOCIATED CONTENT

Supporting Information

Detailed experimental procedures and analytical data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: tand@mskcc.org.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Dr. George Sukenick, Rong Wang, Dr. Hui Liu, Hui Fang, and Dr. Sylvi Rusli (MSKCC) for expert NMR and mass spectral support, Emil Lobkovsky (Cornell University) for X-ray crystallographic analysis, and the NIH (P41 GM076267) and Lucille Castori Center for Microbes, Inflammation, and Cancer (postdoctoral fellowship to I.S.) for generous financial support.

■ REFERENCES

- (1) Reviewed in: (a) Sperry, J.; Wilson, Z. E.; Rathwell, D. C. K.; Brimble, M. A. *Nat. Prod. Rep.* **2010**, *27*, 1117–1137. For general reviews on spiroketals, see: (b) Aho, J. E.; Pihko, P. M.; Rissa, T. K. *Chem. Rev.* **2005**, *105*, 4406–4440. (c) Perron, F.; Albizzati, K. F. *Chem. Rev.* **1989**, *89*, 1617–1661.
- (2) (a) Stierle, A. A.; Stierle, D. B.; Kelly, K. J. *Org. Chem.* **2006**, *71*, 5357–5360. (b) Buchgraber, P.; Snaddon, T. N.; Wirtz, C.; Mynott, R.; Goddard, R.; Fürstner, A. *Angew. Chem., Int. Ed.* **2008**, *47*, 8450–8454.
- (3) (a) Traxler, P.; Fritz, H.; Fuhrer, H.; Richter, W. J. *J. Antibiot.* **1980**, *33*, 967–978. (b) Barrett, A. G. M.; Peña, M.; Willardsen, J. A. *J. Org. Chem.* **1996**, *61*, 1082–1100. (c) Varona, R.; Perez, P.; Duran, A. *FEMS Microbiol. Lett.* **1983**, *20*, 243–247.
- (4) Qi, J.; Lu, J.-J.; Liu, J.-H.; Yu, B.-Y. *Chem. Pharm. Bull.* **2009**, *57*, 134–137.

- (5) Reviewed in: Brasholz, M.; Soergel, S.; Azap, C.; Reissig, H.-U. *Eur. J. Org. Chem.* **2007**, 3801–3814.
- (6) (a) Bardone, M. R.; Martinelli, E.; Zerilli, L. F.; Coronelli, C. *Tetrahedron* **1974**, 30, 2747–2751. (b) Kirillov, S.; Vitali, L. A.; Goldstein, B. P.; Monti, F.; Semenov, Y.; Makhno, V.; Ripa, S.; Pon, C. L.; Gualerzi, C. O. *RNA* **1997**, 3, 905–913.
- (7) Ueno, T.; Takahashi, H.; Oda, M.; Mizunuma, M.; Yokoyama, A.; Goto, Y.; Mizushima, Y.; Sakaguchi, K.; Hayashi, H. *Biochemistry* **2000**, 39, 5995–6002.
- (8) For examples from benzannulated spiroketal natural product total synthesis, see the following. Berkelic acid: (a) Wu, X.; Zhou, J.; Snider, B. B. *Angew. Chem., Int. Ed.* **2009**, 48, 1283–1286. (b) Bender, C. F.; Yoshimoto, F. K.; Paradise, C. L.; De Brabander, J. K. *J. Am. Chem. Soc.* **2009**, 131, 11350–11352. (c) Snaddon, T. N.; Buchgraber, P.; Schulthoff, S.; Wirtz, C.; Mynott, R.; Fürstner, A. *Chem.—Eur. J.* **2010**, 16, 12133–12140. (d) McLeod, M. C.; Wilson, Z. E.; Brimble, M. A. *Org. Lett.* **2011**, 13, 5382–5385. (e) Wenderski, T. A.; Marsini, M. A.; Pettus, T. R. *Org. Lett.* **2011**, 13, 118–121. (f) Fañanás, F. J.; Mendoza, A.; Arto, T.; Temelli, B.; Rodríguez, F. *Angew. Chem., Int. Ed.* **2012**, 51, 4930–4933. Papulacandins: (g) Barrett, A. G. M.; Peña, M.; Willardsen, J. A. *J. Chem. Soc., Chem. Commun.* **1995**, 1147–1148. (h) Denmark, S. E.; Regens, C. S.; Kobayashi, T. *J. Am. Chem. Soc.* **2007**, 129, 2774–2776. Heliquinomycinone: (i) Qin, D.; Ren, R. X.; Siu, T.; Zheng, C.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2001**, 40, 4709–4713. (j) Siu, T.; Qin, D.; Danishefsky, S. J. *Angew. Chem., Int. Ed.* **2001**, 40, 4713–4716. Rubromycins: (k) Reviewed in refs 1 and 5. (l) Akai, S.; Kakiguchi, K.; Nakamura, Y.; Kuriwaki, I.; Dohi, T.; Harada, S.; Kubo, O.; Morita, N.; Kita, Y. *Angew. Chem., Int. Ed.* **2007**, 46, 7458–7461. (m) Rathwell, D. C. K.; Yang, S.-H.; Tsang, K. Y.; Brimble, M. A. *Angew. Chem., Int. Ed.* **2009**, 48, 7996–8000. (n) Wu, K.-L.; Mercado, E. V.; Pettus, T. R. *J. Am. Chem. Soc.* **2011**, 133, 6114–6117. (o) Wei, L.; Xue, J.; Liu, H.; Wang, W.; Li, Y. *Org. Lett.* **2012**, 14, 5302–5305. (p) Wang, W.; Xue, J.; Tian, T.; Zhang, J.; Wei, L.; Shao, J.; Xie, Z.; Li, Y. *Org. Lett.* **2013**, 15, 2402–2405.
- (9) For other selected approaches to benzannulated spiroketals, see the following. (a) Acid-catalyzed *endo*-glycol spirocyclization: Knabe, J.; Schaller, K. *Arch. Pharm.* **1968**, 301, 457–464. (b) Chromone intramolecular Michael addition and chromone epoxide spirocyclization: Cremins, P. J.; Wallace, T. W. *J. Chem. Soc., Chem. Commun.* **1986**, 1602–1603. (c) Dihydroxy ketone bicyclization: Tanaka, T.; Miyaguchi, M.; Mochisuki, R. K.; Tanaka, S.; Okamoto, M.; Kitajima, Y.; Miyazaki, T. *Heterocycles* **1987**, 25, 463–484. (d) *endo*-Glycol epoxide spirocyclization: Friesen, R. W.; Sturino, C. F. *J. Org. Chem.* **1990**, 55, 5808–5810. (e) *endo*-Glycol intramolecular alkoxysele- nation: Elsley, D. A.; MacLeod, D.; Miller, J. A.; Quayle, P.; Davies, G. M. *Tetrahedron Lett.* **1992**, 33, 409–412. (f) Glycoside dehydrative spirocyclization: Barrett, A. G. M.; Peña, M.; Willardsen, J. A. *J. Chem. Soc., Chem. Commun.* **1995**, 1147–1148. (g) Alkyne cyclotrimeriza- tion: McDonald, F. E.; Zhu, H. Y. H.; Holmquist, C. R. *J. Am. Chem. Soc.* **1995**, 117, 6605–6606. (h) C-Aryl glycoside ring rearrangement: Kumazawa, T.; Asahi, Matsuba, S.; Sato, S.; Furuhashi, K.; Onodera, J.-I. *Carbohydr. Res.* **1998**, 308, 213–216. (i) Arylfuran Achmatowicz reaction: Balachari, D.; O'Doherty, G. A. *Org. Lett.* **2000**, 2, 4033–4036. (j) C-Vinyl O-arylglycoside Claisen rearrangement–spirocyc- lization: Van Hooft, P. A. V.; Van Swieten, P. F.; Van der Marel, G. A.; Van Boeckel, C. A. A.; Van Boom, J. H. *Synlett* **2001**, 269–271. (k) *o*-Quinone methide–*exo*-glycol hetero-Diels–Alder: Zhou, G.; Zheng, D.; Da, S.; Xie, Z.; Li, Y. *Tetrahedron Lett.* **2006**, 47, 3349–3352. (l) β -Diketone–*exo*-glycol 1,3-dipolar cycloaddition: Lindsey, C. C.; Wu, K. L.; Pettus, T. R. *Org. Lett.* **2006**, 8, 2365–2367. (m) Alkyne diol hydroalkoxylation: Messerle, B. A.; Vuong, K. Q. *Organometallics* **2007**, 26, 3031–3040. (n) Oxidative radical spirocyclization: Liu, Y.-C.; Sperry, J.; Rathwell, D. C. K.; Brimble, M. A. *Synlett* **2009**, 793–797. (o) Intramolecular Ullmann ether coupling: Yoshioka, K.; Takaishi, I.; Shiozawa, K.; Fukushi, Y.; Tahara, S. *Biosci., Biotechnol., Biochem.* **2008**, 72, 2632–2639. (p) Multicomponent cascade reaction: Barluenga, J.; Mendoza, A.; Rodríguez, F.; Fananas, F. J. *Angew. Chem., Int. Ed.* **2009**, 48, 1644–1647. (q) Enantioselective *endo*-glycol spirocyclization: Sun, Z.; Winschel, G. A.; Borovika, A.; Nagorny, P. *J. Am. Chem. Soc.* **2012**, 134, 8074–8077.
- (10) (a) Potuzak, J. S.; Moilanen, S. B.; Tan, D. S. *J. Am. Chem. Soc.* **2005**, 127, 13796–13797. (b) Moilanen, S. B.; Potuzak, J. S.; Tan, D. S. *J. Am. Chem. Soc.* **2006**, 128, 1792–1793.
- (11) Liu, G.; Wurst, J. M.; Tan, D. S. *Org. Lett.* **2009**, 11, 3670–3673.
- (12) (a) Fittig, R.; Strom, K. T. *Justus Liebigs Ann. Chem.* **1892**, 267, 191–203. (b) Dedek, V.; Trska, P. *Collect. Czech. Chem. Commun.* **1970**, 35, 651–660. (c) Ley, S. V.; Lygo, B.; Organ, H. M.; Wonnacott, A. *Tetrahedron* **1985**, 41, 3825–3836.
- (13) (a) Ahmad, M. S.; Baddeley, G. J. *Chem. Soc.* **1961**, 4303–4306. (b) Ernest, I.; Fritz, H.; Rihs, G. *Helv. Chim. Acta* **1987**, 70, 203–209. (c) Adam, W.; Hadjirapoglou, L.; Jaeger, V.; Seidel, B. *Tetrahedron Lett.* **1989**, 30, 4223–4226. (d) Adam, W.; Hadjirapoglou, L.; Wang, X. *Tetrahedron Lett.* **1989**, 30, 6497–6500.
- (14) (a) Brady, B. A.; Geoghegan, M.; O'Sullivan, W. I. *J. Chem. Soc., Perkin Trans. 1* **1989**, 1557–1562. (b) Nicotra, F.; Panza, L.; Russo, G. *Tetrahedron Lett.* **1991**, 32, 4035–4038.
- (15) Harkat, H.; Blanc, A.; Weibel, J.-M.; Pale, P. *J. Org. Chem.* **2008**, 73, 1620–1623.
- (16) See the Supporting Information for full details.
- (17) Halcomb, R. L.; Danishefsky, S. J. *J. Am. Chem. Soc.* **1989**, 111, 6661–6666.
- (18) Wurst, J. M.; Liu, G.; Tan, D. S. *J. Am. Chem. Soc.* **2011**, 133, 7916–7925.
- (19) A benzyldenedihydrofuranone, presumed to arise from opening of the benzannulated ring followed by elimination of the C2-OTIPS group, was also recovered as a minor byproduct (Figure S1, Supporting Information).¹⁶
- (20) Tlais, S. F.; Dudley, G. B. *Org. Lett.* **2010**, 12, 4698–4701.
- (21) (a) Wabnitz, T. C.; Yu, J.-Q.; Spencer, J. B. *Chem.—Eur. J.* **2004**, 10, 484–493. (b) Rosenfeld, D. C.; Shekhar, S.; Takemiya, A.; Utsunomiya, M.; Hartwig, J. F. *Org. Lett.* **2006**, 8, 4179–4182. (c) Dang, T. T.; Boeck, F.; Hintermann, L. *J. Org. Chem.* **2011**, 76, 9353–9361.