

Correction to “Localized Electronic States from Surface Hydroxyls and Polarons in $\text{TiO}_2(110)$ ”, “Defining the Role of Excess Electrons in the Surface Chemistry of TiO_2 ”, and “Distribution of Ti^{3+} Surface Sites in Reduced TiO_2 ”

N. Aaron Deskins,* Roger Rousseau, and Michel Dupuis

J. Phys. Chem. C **2009**, *113* (33), 14583–14586. DOI: 10.1021/jp9037655

J. Phys. Chem. C **2010**, *114* (13), 5891–5897. DOI: 10.1021/jp101155t

J. Phys. Chem. C **2011**, *115* (15), 7562–7572. DOI: 10.1021/jp2001139

We have become aware that in a series of papers^{1–4} utilizing density functional theory (DFT) to model TiO_2 surfaces we reported erroneously the nature of the atomic basis set for Ti used in some of the calculations, a double- ζ basis set when in fact we used a single- ζ basis set when applying the DFT+U method.⁵ The purpose of this erratum is to correct our error. We also report data that suggest that the conclusions stated in these papers hold true.

In refs 1 and 2 dealing with excess electron localization associated with defects (surface hydroxyls, charged polarons, and oxygen vacancies) we utilized DFT+U for the entire studies, and the calculations were performed with a single- ζ basis set on Ti. Double- ζ basis sets were used for other elements as well as for Ti when using conventional DFT (non DFT+U calculations). In ref 3 dealing with substrate-to-adsorbate charge transfer we used conventional DFT for the majority of calculations. Those were carried out with a double- ζ basis set for Ti. A few calculations used DFT+U, and those were carried out with a single- ζ basis set for Ti.

For our electron localization work^{1,2} we predicted that electrons at Ti^{3+} centers preferentially localize at subsurface sites over surface sites. Use of both single- ζ and double- ζ basis sets leads to electron localization. The single- ζ basis set for Ti allowed control of the final location of the Ti^{3+} site. This was important as we were systematically comparing the stability of many Ti^{3+} centers (i.e., surface and subsurface sites). Localization of an excess electron to a designated Ti atom is possible with a double- ζ basis set but would have required many intricate attempts before converging to a wave function with the correct character. Ultimately we determined that a large number of states are nearly degenerate in energy, indicating that the electrons are in thermal equilibrium across an ensemble of electronic states at finite temperature and a large fraction of these states are subsurface. Similar conclusions were obtained by other authors^{6–8} using planewave basis sets, as well as from experiment.^{9–12} Our other work³ examined how surface reduction affected surface adsorption and was largely based on DFT calculations with a double- ζ basis set for Ti. We used DFT+U and a single- ζ basis set for Ti to assess how increased electron localization (due to the U potential) affects electron transfer to adsorbates. Other work¹³ examined surface reduction and adsorption over group 4 metal oxides and obtained similar conclusions using double- ζ basis sets with both DFT and DFT+U.

We have recently examined the effect of the Ti basis set on adsorption energies for several adsorbates using both single- ζ and double- ζ Ti basis sets (see Figure 1). For all of the




Figure 1. Comparison of adsorption energies for several adsorbates over the rutile (110) surface using single- ζ or double- ζ basis sets for Ti. All adsorbates are bound to a five-coordinated Ti atom, except for the one case of a H atom bound to a surface bridging O atom (O_b).

adsorbates, the difference in adsorption energy between the single- ζ and double- ζ Ti basis set calculations is ~ -0.5 eV, indicating that a double- ζ basis set binds molecules more strongly compared to a single- ζ basis set. The use of a single- ζ basis set introduces a systematic error (~ 0.5 eV) in adsorption energies. This error, however, does not change the qualitative conclusions of our previous study.³ Furthermore, in one of our papers¹⁴ we compared the effect of basis set choice for Ti on the H diffusion from a surface of TiO_2 into bulk. We found the diffusion energies to be nearly identical when using single- ζ and double- ζ basis sets for Ti.

In summary we believe that the findings reported in our papers hold qualitatively true in spite of the use of a single- ζ basis for Ti in the DFT+U calculations.

REFERENCES

- (1) Deskins, N. A.; Rousseau, R.; Dupuis, M. *J. Phys. Chem. C* **2009**, *113*, 14583.
- (2) Deskins, N. A.; Rousseau, R.; Dupuis, M. *J. Phys. Chem. C* **2011**, *115*, 7562.
- (3) Deskins, N. A.; Rousseau, R.; Dupuis, M. *J. Phys. Chem. C* **2010**, *114*, 5891.

Published: June 10, 2014

- (4) Du, Y.; Deskins, N. A.; Zhang, Z.; Dohnalek, Z.; Dupuis, M.; Lyubinetsky, I. *Phys. Chem. Chem. Phys.* **2010**, *12*, 6337.
- (5) Dudarev, S. L.; Botton, G. A.; Savrasov, S. Y.; Humphreys, C. J.; Sutton, A. P. *Phys. Rev. B* **1998**, *57*, 1505.
- (6) Kowalski, P. M.; Camellone, M. F.; Nair, N. N.; Meyer, B.; Marx, D. *Phys. Rev. Lett.* **2010**, *105*, 146405.
- (7) Chrétien, S.; Metiu, H. *J. Phys. Chem. C* **2011**, *115*, 4696.
- (8) Amore Bonapasta, A.; Filippone, F.; Mattioli, G.; Alippi, P. *Catal. Today* **2009**, *144*, 177.
- (9) Krüger, P.; Jupille, J.; Bourgeois, S.; Domenichini, B.; Verdini, A.; Floreano, L.; Morgante, A. *Phys. Rev. Lett.* **2012**, *108*, 126803.
- (10) Kruger, P.; Bourgeois, S.; Domenichini, B.; Magnan, H.; Chandresis, D.; Le Fevre, P.; Flank, A. M.; Jupille, J.; Floreano, L.; Cossaro, A.; Verdini, A.; Morgante, A. *Phys. Rev. Lett.* **2008**, *100*, 055501.
- (11) Kruger, P.; Bourgeois, S.; Domenichini, B.; Magnan, H.; Chandresis, D.; Le Fevre, P.; Floreano, L.; Cossaro, A.; Verdini, A.; Morgante, A. *Surf. Sci.* **2007**, *601*, 3952.
- (12) Minato, T.; Sainoo, Y.; Kim, Y.; Kato, H. S.; Aika, K.-i.; Kawai, M.; Zhao, J.; Petek, H.; Huang, T.; He, W.; Wang, B.; Wang, Z.; Zhao, Y.; Yang, J.; Hou, J. G. *J. Chem. Phys.* **2009**, *130*, 124502.
- (13) Garcia, J. C.; Deskins, N. A. *J. Phys. Chem. C* **2012**, *116*, 16573.
- (14) Du, Y.; Petrik, N. G.; Deskins, N. A.; Wang, Z.; Henderson, M. A.; Kimmel, G. A.; Lyubinetsky, I. *Phys. Chem. Chem. Phys.* **2012**, *14*, 3066.