

Nitrogen-Containing Dimeric *nor*-Multiflorane Triterpene from a *Turraea* sp.

Vincent E. Rasamison,^{†,⊥} L. Harinantenaina Rakotondraibe,^{‡,⊥,||} Carla Slebodnick,[‡] Peggy J. Brodie,[‡] Michel Ratsimbason,[†] Karen TenDyke,[§] Yongchun Shen,[§] Lucien M. Randrianjanaka,[▽] and David G. I. Kingston^{*,‡}

[†]Centre National d'Application des Recherches Pharmaceutiques B.P, 702, Antananarivo 101, Madagascar


[‡]Department of Chemistry and the Virginia Tech Center for Drug Discovery, Virginia Polytechnic Institute and State University, M/C 0212, Blacksburg, Virginia 24061, United States

[§]Next Generation Systems, Eisai Inc., 4 Corporate Drive, Andover, Massachusetts 01810, United States

[▽]Conservation International, Villa Hajaririna, Lot II W 27 B, Ankorahotra, Antananarivo 101, Madagascar

Supporting Information

ABSTRACT: The new triterpene turraoic acid (**1**) and the new N-containing *nor*-triterpene turraenine (**2**), along with triptocallic acid B (**3**) and esculentoic acid (**4**) were isolated from leaves of a *Turraea* sp. Compounds **1**–**3** showed weak to moderate in vitro antiparasmodial activity against the chloroquine-resistant *Plasmodium falciparum* strain FCM29. Compound **1** also displayed weak cytotoxic activity against the nonsmall lung cancer cell line H522-T1 with an IC₅₀ value of 16.4 μ M.



Photograph courtesy of Patrice Antilahimena, Missouri Botanical Garden

Plants from the genus *Turraea* (family Meliaceae, order Rutales) have been extensively investigated due to their high content of bioactive limonoids^{1–7} and triterpenoids such as turrapubesols^{8–10} and pregnanes.^{8,9,11,12} These compounds have been reported to have a wide range of biological activities including cytotoxic, insect antifeedant, and mosquito larvicidal activity. As part of a joint International Cooperative Biodiversity Group (ICBG) research program to search for new antimalarial and anticancer secondary metabolites from the natural resources of Madagascar, we selected a plant species from the genus *Turraea* for investigation. The genus *Turraea* contains approximately 85 species, about 60 of them found in tropical and southern Africa, one species in Australia, and 24 species in Madagascar.¹³ In African ethnobotany, species of this genus have been used as an aphrodisiac and to treat wounds, parasites (bilharzias), abscesses, and impotence, and the ripe fruit and the bitter bark of *Turraea* species are used in Madagascar to treat throat problems.

A crude EtOH extract made from leaves of a Madagascar species of *Turraea* was selected for investigation since it demonstrated in vitro antiparasmodial activity at 3.9 μ g/mL against the chloroquine-resistant strain FCM29 of *Plasmodium falciparum* during our preliminary screening. Bioassay-guided fractionation of this extract resulted in the isolation of turraoic acid (**1**), a new triterpenoid with a multiflorane skeleton, and turraenine (**2**), a new nitrogen-containing *nor*-multiflorane-type triterpene. Triptocallic acid B (**3**) and esculentoic acid (**4**) were also isolated (Figure 1). In this paper, we report the isolation

and the structural elucidation of the new compounds **1** and **2** and the biological activity of all four compounds.

The EtOH extract was subjected to liquid–liquid partitioning followed by column chromatography over silica gel, RP-C18




Figure 1. Structures of compounds **1**–**4**.

Received: March 13, 2014

Published: April 28, 2014

silica gel, and Sephadex LH-20. Final crystallization afforded the four compounds 1–4. Compounds 3 and 4 were identified as triptocalic acid B and esculentoic acid, respectively, based on the comparison of their spectroscopic data with those reported in the literature.^{14,15}

The positive-ion high-resolution electrospray ionization (HRESI) mass spectrum of compound 1 displayed a protonated molecular ion peak at m/z 455.3517 corresponding to the molecular formula $C_{30}H_{47}O_3$ (required for $[M + H]^+$: m/z 455.3520). The IR spectrum showed a typical absorption band at 3443 cm^{-1} and strong absorptions at 1709 and 1732 cm^{-1} which were suggestive of hydroxyl, ketone carbonyl, and carboxyl functions. The ^1H NMR spectroscopic data of 1 (Table 1) exhibited seven methyl singlets (δ 0.93, 3H; 0.98, 6H; 1.01, 3H; 1.03, 3H; 1.10, 3H; 1.23, 3H) and one olefin methine at δ 5.47 (dd, 6.4, 3.2 Hz) which was coupled to a methylene in a cyclohexene ring. The ^{13}C NMR data (Table 1) displayed 30 carbon resonances which were assigned to one ketone carbonyl (δ 217.2), one carboxylic acid (δ 184.5), one olefin methine (δ 117.3), one sp^2 -hybridized quaternary carbon (δ 145.6), 10

methylenes, three methines, six sp^3 -hybridized quaternary carbons, and seven methyls by DEPT 135 experiments. The above NMR data were very similar to those of triptocalic acid B (3, Table 1), a multiflorane-type triterpene acid which was first isolated from a callus cell culture of *Trypterygium wilfordii* (Celastraceae).¹⁴

Comparison of the ^1H and ^{13}C NMR data of 1 with those of 3 indicated that the chemical shifts arising from the methyl, the methylene, and the methine groups were essentially the same except for the signals due to the A-ring. The ^{13}C NMR spectrum of 1 showed the presence of a signal at δ 217.2 (C-3) instead of the oxygen-bearing methine (δ 79.3) in 3. The carbon chemical shifts of C-23 and C-24 shifted from 27.8 and 15.0 in 3 to 24.0 and 21.8, respectively, while the other signals remained almost the same.

In addition, the methylene signals at δ 2.25 dt ($J = 14.5, 3.7$ Hz, H-2a) and δ 2.73 dt ($J = 14.5, 5.5$ Hz, H-2b) in the proton spectrum of 1 corroborated the presence of a ketone carbonyl at C-3. Two-dimensional NMR data of 1 confirmed the assignment of the carbonyl to C-3, the olefin methine at C-7, the methyl groups at C-4, -10, -13, -14, -17, and -20, and the carboxyl group at C-29. The HSQC spectrum was used to assign the proton-bearing carbons (Table 1). The 2J and 3J couplings observed in the HMBC spectrum were then interpreted. The carbonyl carbon at C-3, the methyl group at C-10, and the location of the olefin methine (C-7) of the decalin were assigned by observing the long-range correlations from H-1 to C-3; CH_3 -23 and -24 to C-3, C-4, and C-5; CH_3 -25 to C-1, C-9, and C-5; and CH-7 to C-5, C-9, and C-14. The HMBC cross peaks between CH_3 -26 and C-12, C-18, and C-27 on one hand and between CH_3 -30 and C-29, C-19, and C-21 on the other hand indicated the presence of a carboxylic acid at C-29. In the same manner, the methyl groups at C-13 and C-14 were allocated. The relative and absolute configurations at C-20 and at other chiral centers were assigned by interpretation of the NOESY spectroscopic data and X-ray diffraction analysis of a single crystal of 1. The X-ray structure of 1 showing anisotropic displacement ellipsoids at the 50% probability level is shown in Figure 2.¹⁶ These findings confirmed the structure of 1 as multiflor-3-on-7-en-29-oic acid, named turranoic acid.

Table 1. ^1H NMR Data of Compound 1 and ^{13}C NMR Data of Compounds 1 and 3 in CDCl_3 (δ in ppm)^a

position	1	3	
	δ_{H} (multiplicity (m), J in Hz)	δ_{C}	δ_{C}
1	1.44, m 2.00, m	38.5	37.6
2	2.25, dt (14.5, 3.7) 2.73, td (14.5, 5.5)	35.0	27.6
3		217.2	79.3
4		47.8	39.2
5	1.64, dd (7.8, 6.5)	52.1	50.7
6	2.08, m	24.7	24.5
7	5.47, dd (6.4, 3.2)	117.3	117.6
8		145.6	145.7
9	2.12, m	48.1	48.3
10		35.3	35.5
11	1.52, m, 1.63, m	17.4	17.5
12	0.86 br d (13.8) 2.00, td (13.8, 4.3)	32.8	33.3
13		36.9	37.3
14		42.5	42.5
15	1.47–1.41 ^b	29.4	30.9
16	2.00, m ^b	36.9	37.3
17		31.2	31.6
18	1.48, m	47.2	45.8
19	1.67, m 2.37, br d (16.0)	30.4	30.9
20		40.4	40.6
21	1.37–1.41 ^b	29.4	29.5
22	1.37, m, 1.76, m	35.6	36.0
23	0.98, s	24.0	27.8
24	1.10, s	21.8	15.0
25	0.98, s	13.0	13.4
26	1.03, s	24.7	25.2
27	0.93, s	25.5	27.8
28	1.01, s	31.4	31.5
29		184.5	182.8
30	1.23, s	33.1	33.5

^a ^1H NMR recorded at 600 MHz; ^{13}C NMR recorded at 150 MHz.

^bSignals overlapped.




Figure 2. Displacement ellipsoid drawing (50% probability) of the single-crystal X-ray structure of 1. One of two crystallographically independent molecules is shown. Aliphatic H atoms are omitted for clarity.

Compound 2 had the molecular formula $C_{58}H_{83}NO_6$ as determined by positive high-resolution ESIMS (observed m/z 890.6311, required for $C_{58}H_{84}NO_6$ $[M + H]^+$, 890.6293). Its ^1H NMR spectrum displayed a deshielded triplet resonance due to a hydrogen-bonded amine proton (δ 14.0, t, $J = 11.4$ Hz, 1H), two 2H olefin methine signals (δ 6.46, d, $J = 11.4$, 2H, H-24 and -24' and δ 5.40, brs, 2H, H-7 and -7'), and two sets of signals superposable on those of 1. Excitation of the proton at δ

14.0 by 1D TOCSY collapsed the 2H olefin methine signal at δ 6.46, indicating the presence of the partial structure =CHNHCH= in **2**.

The ^{13}C NMR spectrum of **2** showed signals for conjugated ketone carbonyl (δ 200.0, C-3) and carboxyl (δ 185.9) groups, signals due to olefin methine carbons (δ 114.0, C; 115.8, CH; 141.5, CH and 146.5, C), one of which was assignable to a methine attached via a heteroatom, and signals ascribable to a multiflorane-type triterpene dimer.

Inspection of the ^1H and ^{13}C NMR data (Table 2) of **2** revealed the presence of two sets of signals which were very

Table 2. ^1H and ^{13}C NMR Data for **2** in CDCl_3 (δ in ppm)^a

position	δ_{H} (multiplicity (m), J in Hz)	δ_{C}
1,1'	2.49 m, 1.41–1.44 ^b	34.9
2, 2'	1.40 m, 1.98 m	33.7
3,3'		200.0
4,4'		114.0
5,5'	2.39 dd (10.1, 5.6)	42.4
6,6'	1.30–1.38 ^b	29.1
7,7'	5.40 br	115.8
8,8'		146.5
9,9'	2.11 brd (13.6)	44.8
10,10'		34.4
11,11'	1.48 m, 1.78 m	17.9
12,12'	0.90 m, 2.04 m	32.8
13,13'		36.7
14,14'		43.0
15,15'	1.68 m, 2.44 m	30.3
16,16'	1.36 m, 1.70 m	36.8
17,17'		31.2
18,18'	1.49 brd (7.7)	47.2
19,19'	1.35–1.39 ^b	29.2
20,20'		40.4
21,21'	1.35–1.39 ^b	29.3
22,22'	1.38, m, 1.82, m	35.5
24,24'	6.46, d (11.4)	141.5
25,25'	0.67, s	11.6
26,26'	0.96, s	24.4
27,27'	0.89, s	25.1
28,28'	1.01, s	31.4
29,29'		185.9
30,30'	1.25, s	33.1
NH	14.0, t (11.4)	

^a ^1H NMR recorded at 600 MHz; ^{13}C NMR recorded at 150 MHz.

^bSignals overlapped.

similar to those of **1**. In addition, the IR spectroscopic data of **2** showed absorption bands superposable to those of **1**, suggesting that compound **2** also had ketone carbonyl, carboxyl, and olefin methine functions in its skeleton. A comparison of the ^1H NMR and ^{13}C NMR spectroscopic data of **2** with those of **1** disclosed the absence of the two methyl signals at δ_{H} 0.98 (CH_3 -23) and 1.10 (CH_3 -24) of **1** in compound **2** and the presence instead of the olefin methine signal at δ 6.46. These data suggested that the CH_3 -23 and -24 methyls were replaced by an N-bearing olefin methine which could be the dimerization site of two multiflorane-type triterpenes.

In order to assign the allocations of all functionalities present in **2**, to confirm the site of dimerization, and to elucidate its planar structure, an HMBC experiment was carried out. ^2J and ^3J long-range correlations between the hydrogen-bonded

secondary amine proton at δ_{H} 14.0 and two olefin methine and quaternary carbons at δ_{C} 141.5 (C-24, C-24') and 114.0 (C-4, C-4') were observed. The ^3J long-range correlation between the two olefin protons at δ 6.46 (H-24, 24') and their bearing carbons confirmed that the C-24 (24') of the two units present in **2** were connected with the tertiary amine at δ_{H} 14.0. Moreover, the presence of a ketone carbonyl at C-3 was evidenced by the cross-peaks observed from H-24 (24') to the carbonyl at δ 200.0 and to the methine at C-5 (C-5', δ 42.4). The double bond at C-7 (C-8), the carboxyl group at C-20, and the locations of the methyl groups were evidenced by interpretation of the HMBC spectroscopic data. The key correlations observed to support the structure of **2** are shown in Figure S3 (Supporting Information). The molecular formula of **2** required 18 degrees of unsaturation, 9 of which could be assigned to 23-nor-multiflora-7(8),4(24)-dien-3-on-29-oic acid. The remaining 9 were thus due to the second occurrence of the same monomer.

The relative configuration of **2** was determined by a NOESY experiment and by X-ray diffraction analysis of a single crystal obtained from a chloroform/methanol solution of **2**. The absolute configuration of **2** was assumed to be the same as the absolute configuration of **1**. The X-ray structure of **2** showing anisotropic displacement ellipsoids at the 50% probability is depicted in Figure 3. The structure of turraenine (**2**) was thus determined to be as shown in Figure 1.




Figure 3. Displacement ellipsoid drawing (50% probability) of the single-crystal X-ray structure of **2**. Aliphatic H atoms and a methanol solvate are omitted for clarity.

The dimeric structure of turraenine immediately raised the question of whether it might be a simple artifact of the reaction of ammonia with a suitable aldehyde precursor, as observed in a study of acid hydrolysis of some 16-methylamino steroids.¹⁷ This is considered to be highly unlikely for three reasons. In the first place, the required aldehyde precursor, 13-methyl-3,23-dioxo-24,26-bisnorolean-7-en-29-oic acid, is not a known natural product and would be relatively unstable. Second, no ammonia was used in the initial extraction process or in any subsequent purification step. Finally, compound **2** was shown to be present in the crude extract by direct ^1H NMR analysis, which clearly showed the presence of the triplet due to the hydrogen-bonded secondary amine (Figure S4, Supporting Information). When this signal was excited, the doublet signal due to the olefin methine (δ_{H} 6.46, d, J = 11.4) was collapsed.

Compound **2** is the first example known of a naturally occurring N-linked triterpene dimer. The biogenesis of **2** can be plausibly traced from **1**. Successive oxidation and decarboxylation of **1** would lead to a C-23 demethylated compound which could be oxidized to afford C-23 aldehyde **S3**. Schiff base formation with ammonia, followed by tautomerism and reaction with another molecule of aldehyde **S3** and loss of

water, would give the dimerized compound **2** (Figure S5, Supporting Information).

All of the isolated compounds (**1–4**) were evaluated for antimalarial activity against the chloroquine-resistant strain FCM29 of *Plasmodium falciparum*. Turraoic acid (**1**) was the most active (IC₅₀ 5.2 μM) among all compounds tested. The activities of triptocallic acid **B** (**3**) and turraenine (**2**) are comparable (16.4 and 16.6 μM, respectively) while esculentoic acid (**4**) was not active. Among the four compounds isolated, only compound **1** exhibited antiproliferative activity, albeit weak, against the A2780 human ovarian cancer cell line (IC₅₀ 20 μM). Compound **1** was inactive against the H522-T1 nonsmall cell lung cancer cell line (IC₅₀ > 20 μM) and showed weak activity against the human A2058 melanoma cancer cell line (IC₅₀ = 16.4 μM).

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental details for the isolation of compounds **1–4**; characterization data for compounds **1** and **2**; key HMBC correlations of **1** and **2** and NOESY correlations of **1**; single-crystal X-ray data for compounds **1** and **2**; ¹H NMR and 1D-TOCSY spectra of the crude ethanol extract of *Turraea* sp.; plausible biosynthetic pathway for **2**; ¹H, ¹³C, DEPT 135, HSQC, and HMBC spectra of compounds **1** and **2**; 1D-TOCSY spectrum of **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: dkingston@vt.edu.

Present Address

^{||}Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, OH 43210.

Author Contributions

[⊥]These authors contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This project was supported by the Fogarty International Center, the National Cancer Institute, the National Science Foundation, the National Heart, Lung and Blood Institute, the National Institute of Mental Health, the Office of Dietary Supplements, and the Office of the Director of NIH, under Cooperative Agreement U01 TW000313 with the International Cooperative Biodiversity Groups. This project was also supported by the National Research Initiative of the Cooperative State Research, Education and Extension Service, USDA, Grant No. 2008-35621-04732. This work was also supported by the National Science Foundation under Grant No. CHE-0619382 for purchase of the Bruker Avance 600 NMR spectrometer and Grant No. CHE-0722638 for the purchase of the Agilent 6220 mass spectrometer. We thank Mr. B. Bebout (Virginia Tech) for obtaining the mass spectra. We gratefully acknowledge courtesies extended by the Government of Madagascar (Ministère des Eaux et Forêts). We thank The Ohio State University, College of Pharmacy, and the Campus Chemical Instrument Center (CCIC) NMR facility at The

Ohio State University for facilitating the acquisition of a part of the 2D NMR spectra.

■ REFERENCES

- (1) Akinniyi, J. A.; Connolly, J. D.; Mulholland, D. A.; Rycroft, D. S.; Taylor, D. A. H. *Phytochemistry* **1986**, *25*, 2187–2189.
- (2) Bentley, M. D.; Adul, G. O.; Alford, A. R.; Huang, F.-Y.; Gelbaum, L.; Hassanali, A. *J. Nat. Prod.* **1995**, *58*, 748–750.
- (3) Cheplogoi, P. K.; Mulholland, D. A. *Biochem. Syst. Ecol.* **2003**, *31*, 799–803.
- (4) McFarland, K.; Mulholland, D. A.; Fraser, L.-A. *Phytochemistry* **2004**, *65*, 2031–2037.
- (5) Mulholland, D. A.; Monkhe, T. V.; Coombes, P. H.; Rajab, M. S. *Phytochemistry* **1998**, *49*, 2585–2590.
- (6) Rajab, M. S.; Bentley, M. D.; Hassanali, A.; Chapya, A. *Phytochemistry* **1988**, *27*, 2353–2355.
- (7) Sarker, S. D.; Savchenko, T.; Whiting, P.; Sik, V.; Dinan, L. *Arch. Insect Biochem. Physiol.* **1997**, *35*, 211–217.
- (8) Bentley, M. D.; Gaul, F.; Rajab, M. S.; Hassanali, A. *J. Nat. Prod.* **1992**, *55*, 84–87.
- (9) Mulholland, D. A.; Monkhe, T. V.; Taylor, D. A. H.; Rajab, M. S. *Phytochemistry* **1999**, *52*, 123–126.
- (10) Wang, X.-N.; Fan, C.-Q.; Yin, S.; Lin, L.-P.; Ding, J.; Yue, J.-M. *Helv. Chim. Acta* **2008**, *91*, 510–519.
- (11) Chiplunkar, Y. G.; Nagasampagi, B. A.; Tavale, S. S.; Puranik, V. G. *Phytochemistry* **1993**, *33*, 901–903.
- (12) Wang, X.-N.; Fan, C.-Q.; Yue, J.-M. *Steroids* **2006**, *71*, 720–724.
- (13) Mabberly, D. J. *The Plant Book*, 2nd ed.; Cambridge University Press: Cambridge, UK, 1997; p 733.
- (14) Kakano, K.; Oose, Y.; Masuda, Y.; Kamada, H.; Takaishi, Y. *Phytochemistry* **1997**, *45*, 293–296.
- (15) Chaturvedula, V. S. P.; Schilling, J. K.; Malone, S.; Wisse, J. H.; Werkhoven, M. C. M.; Kingston, D. G. I. *Planta Med.* **2003**, *69*, 271–274.
- (16) Crystallographic data have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC 980410 (**1**) and no. CCDC 980561 (**2**). These data can be obtained free of charge on application to Cambridge Crystallographic Data Centre, 2 Union Road, Cambridge CB2 1EZ, UK [fax: (+44)1223-336-033; e-mail: deposit@ccdc.cam.ac.uk].
- (17) Gerali, G.; Cecchi, L.; Parini, C.; Saglia, G. *Farmaco, Ed. Sci.* **1971**, *26*, 1089–1099.

■ NOTE ADDED AFTER ASAP PUBLICATION

In the Supporting Information, Figure S5 was submitted in its uncorrected form. Figure S5 was replaced on April 29, 2014.