



Flavone-Based ESIPT Ratiometric Chemodosimeter for Detection of Cysteine in Living Cells

Bin Liu,^{†,‡} Junfeng Wang,[†] Ge Zhang,[§] Ruke Bai,^{*,‡} and Yi Pang^{*,†}


[†]Department of Chemistry & Maurice Morton Institute of Polymer Science, and [§]Department of Biomedical Engineering, The University of Akron, Akron, Ohio 44325, United States

[‡]CAS Key Laboratory of Soft Matter Chemistry, Department of Polymer Science and Engineering, University of Science and Technology of China, Hefei, P. R. China 230026

Supporting Information

ABSTRACT: We have designed and synthesized a novel ratiometric fluorescent chemodosimeter **MHF**-based ESIPT process for specific detection of cysteine among the biological thiols. The probe **MHF** shows very weak blue fluorescence under UV excitation. Upon addition of cysteine (Cys), the reaction of Cys with **MHF** induces acrylate hydrolysis, thereby enabling the ESIPT process to shift the weak blue emission to a strong green emission with about 20-fold enhancement. We utilized ¹H NMR spectra to elucidate the fluorescence sensing mechanism. Moreover, the cellular imaging experiment indicated the **MHF** possessed excellent selectivity, low cytotoxicity, and desirable cell permeability for biological applications.

KEYWORDS: chemodosimeter, fluorescence, cysteine, ratiometric, selectivity, bioimaging



■ INTRODUCTION

As one of the most important biological thiols, cysteine (Cys) plays a pivotal role in many biological processes such as reversible redox reaction and cellular detoxification and metabolism.^{1,2} Low Cys levels could be related to many health issues such as hematopoiesis reduction, hair depigmentation, skin lesion development, and cancer,^{3–5} as Cys is involved in the chemical regulation of many biological processes. A number of analytical methods for the detection of Cys have been developed using high-performance liquid chromatography (HPLC),⁶ capillary electrophoresis,⁷ electrochemical assay,⁸ UV/Vis,⁹ FTIR,¹⁰ mass,¹¹ and fluorescence spectroscopy.^{12,13} Among these methods, fluorescence probes are more desirable due to its high selectivity, low detection limit, fast response and great potential for bioimaging.^{14–16}

Current fluorescent probes for Cys often utilize the nucleophilicity of thiol group in the sensing scheme.^{17–35} However, lots of reported probes suffer from low selectivity, poor cellular uptake, interference of autofluorescence, and high cytotoxicity, which significantly limits their biological applications. Therefore, developing highly biocompatible and selective fluorescent probes to monitor the Cys levels is of great scientific interest, which requires the integration of the Cys' unique reactivity with a selective and reliable chemical event.

Recently, the fluorescent dyes based on excited-state intramolecular proton transfer (ESIPT) process, as seen from 2-(2'-hydroxyphenyl)benzoxazole, 1-aminoanthraquinone, and flavone, have been used as an attractive fluorescent signal transducer in sensors.^{36–44} In comparison with the other fluorescent processes, such as electron transfer, ESIPT process

can occur at a much faster rate ranging from fractions of picoseconds to tens of picoseconds.⁴⁵ Moreover, ESIPT dyes generally have large Stokes' shift (>150 nm), which minimizes the self-absorption and reduce the interference from autofluorescence for in vivo application.⁴⁶ Although the unique photophysical properties are known for decades, only one ESIPT sensor has been designed for Cys detection.²⁶

Among ESIPT dyes, flavone dyes are a broad class of natural products, and have been extensively studied for their antioxidant properties and anticancer activities in the food and health sciences.⁴⁷ However, few flavone-based biosensors have been studied for bioimaging application (Scheme S2 in the Supporting Information). Herein, we present a novel flavone-based ratiometric fluorescence probe, 4-oxo-2-phenyl-4H-chromen-3-yl acrylate (**MHF**), which gives ESIPT emission upon binding cysteine in living cells (Scheme 1a). The sensor design utilizes both thiol and amino groups of Cys in a nucleophilic addition and subsequent cyclization reaction, in order to achieve specific recognition of Cys. Other prominent features of **MHF** include: (1) large emission spectral shift (from weak 380 nm to strong 510 nm) in responding to Cys, as a consequence of the ESIPT turn-on;⁴⁸ (2) linear response to Cys; (3) the flavone-based dye is of low cytotoxicity, good cellular uptake, which are desirable for medicinal biology and diagnostic applications.

Received: January 6, 2014

Accepted: February 27, 2014

Published: February 27, 2014

Scheme 1. (a) Chemical Structures of MHF along with the Proposed Sensing Mechanism; (b) Schematic Representation of ESIPT Process of HF





Figure 1. Time-dependent (a) absorption spectral changes and (b) absorbance changes ($\lambda = 285$ nm and 350 nm) of MHF (10 μ M) in the presence of 100 μ M Cys in MeCN- H_2O (1:1, v/v) solution with 10 mM HEPES buffer.

EXPERIMENTAL SECTION

Reagents and Instrumentation. 1H NMR and ^{13}C NMR spectra were obtained using a Bruker AVANCE II. UV-vis spectra were acquired on a Hewlett-Packard 8453 diode-array spectrometer. Fluorescence spectra were measured by RF-5301PC spectrometer. The fluorescence quantum yields were obtained using quinine sulfate as the standard ($\Phi_f = 0.53$, 0.1 M H_2SO_4). Electrospray ionization (ESI) mass spectra were acquired with a Waters Synapt HDMS quadrupole/time-of-flight (Q/ToF) mass spectrometer. All the solvents for the fluorescence experiments were analytic grade, which were purchased from Fisher Scientific and used without further purification. 1 mmol/L HF and MHF were dissolved in MeCN as stock solutions and 10 mM biologically relevant analytes (Cys, Hcy, GSH, NaSH, Ala, Arg, Asn, Asp, Gln, Gluc, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Tau, Thr, Trp, Tyr, Val) were separately dissolved in distilled water. 10 mM HEPES solution was prepared as buffer solution. All UV/Vis and fluorescence titration experiments were

performed using 10 μ M of HF or MHF in 50% MeCN aqueous solution (pH 7.4, 10 mM PBS buffer) with varying concentrations of analytes at room temperature with 1 h reaction time. The cell imaging was obtained by X-Cite Series 120Q fluorescence microscopy. The blue channel filter: excitation 365 nm, beam splitter FT 395 nm, emission 445/50 nm. The green channel filter: excitation 450–490 nm, beam splitter FT 510 nm, emission 515–565 nm.

Synthesis of 4-Oxo-2-phenyl-4H-chromen-3-yl acrylate (MHF). 1 mmol HF and 1.2 mmol K_2CO_3 was dissolved in 20 mL of dry acetone in ice-water bath then 1.2 mmol acryloyl chloride in 10 mL dry acetone was slowly added into the solution. The mixture was stirred for 12 h then the solvent was removed in reduced pressure. The crude product was purified by column chromatography on silica gel. Yield = 52%. 1H NMR ($CDCl_3$, 300 MHz): $\delta = 6.04$ – 6.07 (d, 1H), 6.33– 6.42 (m, 1H), 6.61– 6.67 (d, 1H), 7.26– 7.45 (t, 1H), 7.47– 7.56 (m, 3H), 7.59– 7.61 (d, 1H), 7.71– 7.76 (t, 1H), 7.87– 7.90 (dd, 2H), 8.27– 8.30 (d, 1H). ^{13}C NMR (75 MHz, $CDCl_3$): $\delta = 118.1$, 123.7, 125.3, 126.2, 126.9, 128.3, 128.7, 130.0, 131.3, 133.7, 134.0, 155.7,




Figure 2. Time-dependent (a) fluorescence spectral changes and (b) fluorescence intensities ($\lambda=510$ nm) of MHF (10 μ M) in the presence of 100 μ M Cys in MeCN- H_2O (1:1, v/v) solution with 10 mM HEPES buffer. (c) Fluorescence spectra changes and (d) fluorescence intensity changes ($\lambda=510$ nm) of 10 μ M MHF in the presence of increasing concentrations of Cys (final concentration: 0, 0.001, 0.0025, 0.005, 0.0075, 0.01, 0.02, 0.04, 0.06, 0.08, 0.1 mM) in MeCN- H_2O (1:1, v/v) solution with 10 mM HEPES buffer. Each spectrum was recorded after 60 min.

156.4, 163.1, 172.1. HRMS: m/z calcd for $C_{18}H_{14}O_5$ ($M + Na$) $^+$, 315.0633; found, 315.0616.

Cell Culture. Human mesenchymal stem cells (hMSCs) (Lonza, Walkersville, MD) were cultured in serum-containing MSCBM medium (Lonza) supplemented with MSCGM SingleQuots (Lonza) according to manufacturer's specifications. hMSCs (Passage 5) were seeded at a density of 5.0×10^4 cell/ cm^2 . Before treatment of MHF, the control cells were incubated with media containing 100 μ M NEM for 30 min at 37 $^\circ$ C to react with cellular thiols. The cells were then briefly washed with 1 mL of PBS. After incubation with 20 μ M MHF (1% DMSO) for 1 h at 37 $^\circ$ C, fluorescence images were taken using a fluorescence microscope. The cytotoxicity of the MHF towards stem cells was determined by conventional MTT assays.

RESULTS AND DISCUSSION

MHF was conveniently synthesized from acylation of 3-hydroxyflavone (HF) with acryloyl chloride. MHF exhibited one absorption peak at 290 nm, while HF had two absorption peaks at 310 and 340 nm (see Figure S1 in Supporting Information). When being excited at 350 nm, MHF gave a weak emission peak at \sim 380 nm. Addition of Cys to MHF, however, gave two emission bands at 380 and 510 nm, which can be attributed to the normal isomer (N^* emission) and tautomer (T^* emission) of HF, respectively (see Scheme 1b). Observation of the intense green emission from "MHF + Cys"




Figure 3. 1H NMR spectrum of MHF in d_6 -DMSO, and the resulting spectrum after addition of 1 equiv. Cys in D_2O for 5 min, 1 h, and 5 h. The starred "*" signals are attributed to DMSO and water solvents.

sample indicated the formation of HF, as the reaction of Cys with MHF released the hydroxyl group in flavone, thereby enabling the ESIPT process to shift the emission signal to a longer wavelength. The new emission peak can be used for the ratiometric fluorescent measurement, as the ratio of two fluorescent bands (instead of the absolute intensity of one band) can determine the analytes more accurately with the minimization of the background signal.⁴⁹




Figure 4. The fluorescence intensities ($\lambda = 510$ nm) of $10 \mu\text{M}$ MHF upon addition of $100 \mu\text{M}$ physiological important amino acids (Glu, Asp, His, Arg, Lys, Gln, Asn, Tyr, Thr, Ser, Cys, Gly, Met, Trp, Phe, Pro, Ile, Leu, Val, and Ala) in MeCN- H_2O (1:1, v/v) solution with 10 mM HEPES buffer.




Figure 5. The fluorescence intensities ($\lambda = 510$ nm) of $10 \mu\text{M}$ MHF upon addition of $100 \mu\text{M}$ biologically important thiols (Cys, GSH, Hcy, and NaHS) in MeCN- H_2O (1:1, v/v) solution with 10 mM HEPES buffer.

The optical sensing behavior of MHF toward Cys was investigated by using a $10 \mu\text{M}$ MHF in MeCN- H_2O (1:1, v/v) solution (pH 7.4, 10 mM PBS buffer). Upon addition of $100 \mu\text{M}$ Cys to the solution of MHF, the absorption band at around 350 nm increased gradually over time, meanwhile the band at 290 nm decreased with a 10 nm red shift, shown in Figure 1. For the fluorescence spectra, the addition of Cys caused an apparent ratiometric fluorescence response. The N^* emission slowly increased and became doubled after one hour reaction, whereas a significantly higher fluorescence (>20 -fold) was observed from the tautomer (T^* emission), as seen in Figure 2a, b. To verify this mechanism of the Cys-induced acrylate cyclization,⁵⁰ we examined the ^1H NMR of MHF in d_6 -DMSO at room temperature. As seen in Figure 3, after addition of 1 equiv. of Cys in D_2O for 5 min, the characteristic alkenyl proton H_a (labeled in the structure in scheme 1a) from 6 to 7 ppm disappeared completely, suggesting a very fast reaction between thiol and alkene, which produced intermediate **1**. However, the lactam proton H_c at ~ 4.3 ppm in **2** wasn't found, which indicated the cyclization was relatively slow. As the reaction proceeds, the intensity of H_c became relatively higher than the intensity of methine proton H_b in Cys and **1** at ~ 3.7 ppm. The proton signal $H_{d'}$ in **HF** was gradually increased, along with the decrease in proton signals H_b and $H_{d''}$ in **1**. The reaction over 10 h at room temperature (see Figure S2 in the Supporting Information) showed that the reaction sequence in Scheme 1a proceeded cleanly, making the process reliable for Cys detection.

The sensitivity of MHF was studied by fluorescence response towards various concentrations of cysteine. Panels c and d in Figure 2 showed that with the increase of Cys concentration, the fluorescence intensity at 510 nm was enhanced dramatically. The fluorescence intensity at $\lambda = 510$ nm was linearly proportional to the amount of Cys ranging from $10 \mu\text{M}$ to $100 \mu\text{M}$ with a detection limit of lower than $1 \mu\text{M}$. Cys concentration normally range from 16.5 to $33.0 \mu\text{M}$ in healthy individual urine.⁵¹ Because $10 \mu\text{M}$ of Cys could enhance T^* emission intensity of MHF for about 2-fold in pure water, this probe could be used for early detection of Cys-related metabolism disease (see Figure S3 in the Supporting Information). The selectivity of chemodosimeter MHF towards various physiological important amino acids and biological




Figure 6. Fluorescence microscopy images of hMSCs. (a) Bright-field image, and fluorescence images in (c) blue and (e) green channel after hMSCs were pre-treated with $100 \mu\text{M}$ NEM for 30 min, and then incubated with $20 \mu\text{M}$ MHF for 1 h. (b) Bright-field image, fluorescence images in (d) blue and (f) green channel after hMSCs being incubated with $20 \mu\text{M}$ MHF for 1 h at 37°C .

thiols was also investigated. As showed in Figure 4 and Figure 5, **MHF** showed highly selective for Cys with remarkable fluorescence intensity enhancement, only mercapto species such as Hcy and GSH showed slight interference. Therefore, **MHF** could be a practically useful probe for effective recognition of Cys.

To further investigate the biological application of **MHF**, the fluorescence microscopy experiment was carried out. When human mesenchymal stem cells (hMSCs) were incubated with 20 μM **MHF** in culture medium at 37 $^{\circ}\text{C}$ for 1 h, relatively weak blue N^* emission (Figure 6d) but strong green T^* emission (Figure 6f) were observed, which was attributed to the formation of **HF** via Cys-induced acrylate hydrolysis and indicated a very good cellular uptake. A control experiment was performed to verify that the sensor's green fluorescence was attributed to the reaction with Cys. Thus, when hMSCs were pretreated with 100 μM NEM (N-ethylmaleimide, an efficient thio-reactive compound) for 30 min, and then incubated with 20 μM **MHF** in culture medium at 37 $^{\circ}\text{C}$ for 1 h, the green emission was very weak (Figure 6e), because Cys were consumed by NEM. Moreover, **MHF** exhibited very low cytotoxicity towards hMSCs, which was evaluated by means of MTT assays,⁵² after the cells were incubated for 24 h in the presence of 50 μM **MHF** (see Figure S4 in the Supporting Information).

CONCLUSION

In summary, we have developed a novel ratiometric chemodosimeter **MHF** by using the low-cytotoxic flavonoid dye and ESPT turn-on. The **MHF** exhibited high sensitivity for Cys (detection limit 1 μM) whose excellent selectivity differentiates it not only from the essential amino acids but also from the biologically important thiols. The probe was successfully used for fluorescent imaging of intracellular Cys, demonstrating its potential for a broader range of biological sample analysis.

ASSOCIATED CONTENT

Supporting Information

Experimental details for synthesis of **MHF**, its ^1H and ^{13}C NMR data, and additional absorption and fluorescence spectra for **MHF** and **HF**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

*E-mail: bairk@ustc.edu.cn.

*E-mail: yp5@uakron.edu.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by NIH (Grant 1R15EB014546-01A1). We also thank the Coleman endowment from the University of Akron for partial support.

REFERENCES

- (1) Chen, X.; Zhou, Y.; Peng, X.; Yoon, J. Fluorescent and Colorimetric Probes for Detection of Thiols. *Chem. Soc. Rev.* **2010**, *39*, 2120–2135.
- (2) Yang, Y.; Zhao, Q.; Feng, W.; Li, F. Luminescent Chemodosimeters for Bioimaging. *Chem. Rev.* **2013**, *113*, 192–270.

- (3) Reddie, K. G.; Carroll, K. S. Expanding the Functional Diversity of Proteins through Cysteine Oxidation. *Curr. Opin. Chem. Biol.* **2008**, *12*, 746–754.
- (4) Weerapana, E.; Wang, C.; Simon, G. M.; Richter, F.; Khare, S.; Dillon, M. B. D.; Bachovchin, D. A.; Mowen, K.; Baker, D.; Cravatt, B. F. Quantitative Reactivity Profiling Predicts Functional Cysteines in Proteomes. *Nature* **2010**, *468*, 790–795.
- (5) Townsend, D. M.; Tew, K. D.; Tapiero, H. The Importance of Glutathione in Human Disease. *Biomed. Pharmacother.* **2003**, *57*, 145–155.
- (6) Ercal, N.; Yang, P.; Aykin, N. Determination of Biological Thiols by High-Performance Liquid Chromatography Following Derivatization by ThioGlo Maleimide Reagents. *J. Chromatogr. B* **2001**, *753*, 287–292.
- (7) Ryant, P.; Dolezelova, E.; Fabrik, I.; Baloun, J.; Adam, V.; Babula, P.; Kizek, R. Electrochemical Determination of Low Molecular Mass Thiols Content in Potatoes (*Solanum Tuberosum*) Cultivated in the Presence of Various Sulphur Forms and Infected by Late Blight (*Phytophthora Infestans*). *Sensors* **2008**, *8*, 3165–3182.
- (8) Potesil, D.; Petrova, J.; Adam, V.; Vacek, J.; Klejdus, B.; Zehnalek, J.; Trnkova, L.; Havel, L.; Kizek, R. Simultaneous Femtomole Determination of Cysteine, Reduced and Oxidized Glutathione, and Phytochelatin in Maize (*Zea mays* L.) Kernels Using High-Performance Liquid Chromatography with Electrochemical Detection. *J. Chromatogr. A* **2005**, *1084*, 134–144.
- (9) Rusin, O.; Luce, N. N. S.; Agbaria, R. A.; Escobedo, J. O.; Jiang, S.; Warner, I. M.; Dawan, F. B.; Lian, K.; Strongin, R. M. Visual Detection of Cysteine and Homocysteine. *J. Am. Chem. Soc.* **2004**, *126*, 438–439.
- (10) Sato, Y.; Iwata, T.; Tokutomi, S.; Kandori, H. Reactive Cysteine is Protonated in the Triplet Excited State of the LOV2 Domain in *Adiantum Phytochrome3*. *J. Am. Chem. Soc.* **2005**, *127*, 1088–1089.
- (11) Guan, X.; Hoffman, B.; Dwivedi, C.; Matthees, D. P. A Simultaneous Liquid Chromatography/Mass Spectrometric Assay of Glutathione, Cysteine, Homocysteine and Their Disulfides in Biological Samples. *J. Pharm. Biomed. Anal.* **2003**, *31*, 251–261.
- (12) Zhou, Y.; Yoon, J. Recent Progress in Fluorescent and Colorimetric Chemosensors for Detection of Amino Acids. *Chem. Soc. Rev.* **2012**, *41*, 52–67.
- (13) Chen, X.; Pradhan, T.; Wang, F.; Kim, J. S.; Yoon, J. Fluorescent Chemosensors Based on Spiroring-Opening of Xanthenes and Related Derivatives. *Chem. Rev.* **2012**, *112*, 1910–1956.
- (14) Wright, A. T.; Anslyn, E. V. Differential Receptor Arrays and Assays for Solution-Based Molecular Recognition. *Chem. Soc. Rev.* **2006**, *35*, 14–28.
- (15) Yoon, J.; Kim, S. K.; Singh, N. J.; Kim, K. S. Imidazolium Receptors for the Recognition of Anions. *Chem. Soc. Rev.* **2006**, *35*, 355–360.
- (16) Wu, J.; Liu, W.; Ge, J.; Zhang, H.; Wang, P. F. New Sensing Mechanisms for Design of Fluorescent Chemosensors Emerging in Recent Years. *Chem. Soc. Rev.* **2011**, *40*, 3483–3495.
- (17) Yi, L.; Li, H.; Sun, L.; Liu, L.; Zhang, C.; Xi, Z. A Highly Sensitive Fluorescence Probe for Fast Thiol-Quantification Assay of Glutathione Reductase. *Angew. Chem., Int. Ed.* **2009**, *48*, 4034–4037.
- (18) Hewage, H. S.; Anslyn, E. V. Pattern-Based Recognition of Thiols and Metals Using a Single Squaraine Indicator. *J. Am. Chem. Soc.* **2009**, *131*, 13099–13106.
- (19) Hong, V.; Kislukhin, A. A.; Finn, M. G. Thiol-Selective Fluorogenic Probes for Labeling and Release. *J. Am. Chem. Soc.* **2009**, *131*, 9986–9994.
- (20) Li, H.; Fan, J.; Wang, J.; Tian, M.; Du, J.; Sun, S.; Peng, X. A Fluorescent Chemodosimeter Specific for Cysteine: Effective Discrimination of Cysteine from Homocysteine. *Chem. Commun.* **2009**, *45*, 5904–5906.
- (21) Kwon, H.; Lee, K.; Kim, H.-J. Coumarin-Malonitrile Conjugate as a Fluorescence Turn-On Probe for Biothiols and Its Cellular Expression. *Chem. Commun.* **2011**, *47*, 1773–1775.

- (22) Yang, Y.-K.; Shim, S.; Tae, J. Rhodamine-Sugar Based Turn-On Fluorescent Probe for the Detection of Cysteine and Homocysteine in Water. *Chem. Commun.* **2010**, 46, 7766–7768.
- (23) Ruan, Y.-B.; Li, A.-F.; Zhao, J.-S.; Shen, J.-S.; Jiang, Y.-B. Specific Hg(2+)-Mediated Perylene Bisimide Aggregation for Highly Sensitive Detection of Cysteine. *Chem. Commun.* **2010**, 46, 4938–4940.
- (24) Kim, G.-J.; Lee, K.; Kwon, H.; Kim, H. -J. Ratiometric Fluorescence Imaging of Cellular Glutathione. *Org. Lett.* **2011**, 13, 2799–2801.
- (25) Lin, W.; Yuan, L.; Cao, Z.; Feng, Y.; Long, L. A Sensitive and Selective Fluorescent Thiol Probe in Water Based on the Conjugate 1,4-Addition of Thiols to α , β -Unsaturated Ketones. *Chem.—Eur. J.* **2009**, 15, 5096–5103.
- (26) Yang, X.; Guo, Y.; Strongin, R. M. Conjugate Addition/Cyclization Sequence Enables Selective and Simultaneous Fluorescence Detection of Cysteine and Homocysteine. *Angew. Chem., Int. Ed.* **2011**, 50, 10690–10693.
- (27) Yuan, L.; Lin, W.; Yang, Y. A Ratiometric Fluorescent Probe for Specific Detection of Cysteine over Homocysteine and Glutathione Based on the Drastic Distinction in the Kinetic Profiles. *Chem. Commun.* **2011**, 47, 6275–6277.
- (28) Zhang, H.; Wang, P.; Yang, Y.; Sun, H. A Selective Fluorescent Probe for Thiols Based on α , β -Unsaturated Acyl Sulfonamide. *Chem. Commun.* **2012**, 48, 10672–10674.
- (29) Wang, H.; Zhou, G.; Gai, H.; Chen, X. A Fluorescein-Based Probe With High Selectivity to Cysteine over Homocysteine and Glutathione. *Chem. Commun.* **2012**, 48, 8341–8343.
- (30) Bao, Y.; Li, Q.; Liu, B.; Du, F.; Tian, J.; Wang, H.; Wang, Y.; Bai, R. Conjugated Polymers Containing a 2,2'-Biimidazole Moiety—A Novel Fluorescent Sensing Platform. *Chem. Commun.* **2012**, 48, 118–120.
- (31) Shao, J.; Guo, H.; Ji, S.; Zhao, J. Styryl-BODIPY Based Red-Emitting Fluorescent OFF-ON Molecular Probe For Specific Detection of Cysteine. *Biosens. Bioelectron.* **2011**, 26, 3012–3017.
- (32) Kwon, N. Y.; Kim, D.; Jang, G.; Lee, J. H.; So, J.-H.; Kim, C. -H.; Kim, T. H.; Lee, T. S. Highly Selective Cysteine Detection and Bioimaging in Zebrafish through Emission Color Change of Water-Soluble Conjugated Polymer-Based Assay Complex. *ACS Appl. Mater. Interfaces* **2012**, 4, 1429–1433.
- (33) Zhang, M.; Yu, M.; Li, F.; Zhu, M.; Li, M.; Gao, Y.; Li, L.; Liu, Z.; Zhang, J.; Zhang, D.; Yi, T.; Huang, C. A Highly Selective Fluorescence Turn-On Sensor for Cysteine/Homocysteine and Its Application in Bioimaging. *J. Am. Chem. Soc.* **2007**, 129, 10322–10323.
- (34) Xiong, L.; Zhao, Q.; Chen, H.; Wu, Y.; Dong, Z.; Zhou, Z.; Li, F. Phosphorescence Imaging of Homocysteine and Cysteine in Living Cells Based on a Cationic Iridium(III) Complex. *Inorg. Chem.* **2010**, 49, 6402–6408.
- (35) Li, Z.; Lou, X.; Li, Z.; Qin, J. A New Approach to Fluorescence “Turn-On” Sensing of α -Amino Acids. *ACS Appl. Mater. Interfaces* **2009**, 1, 232–234.
- (36) Wu, Y. K.; Peng, X. J.; Fan, J. L.; Gao, S.; Tian, M. Z.; Zhao, J. Z.; Sun, S. G. Fluorescence Sensing of Anions Based on Inhibition of Excited-State Intramolecular Proton Transfer. *J. Org. Chem.* **2007**, 72, 62–70.
- (37) Xu, Y. Q.; Pang, Y. Zinc Binding-Induced Near-IR Emission From Excited-State Intramolecular Proton Transfer of a Bis-(benzoxazole) Derivative. *Chem. Commun.* **2010**, 46, 4070–4072.
- (38) Kim, T.; Kang, H. J.; Han, G.; Chung, S. J.; Kim, Y. A Highly Selective Fluorescent ESIPT Probe for the Dual Specificity Phosphatase MKP-6. *Chem. Commun.* **2009**, 45, 5895–5897.
- (39) Hu, R.; Feng, J.; Hu, D. H.; Wang, S. Q.; Li, S. Y.; Li, Y.; Yang, G. Q. A Rapid Aqueous Fluoride Ion Sensor with Dual Output Modes. *Angew. Chem., Int. Ed.* **2010**, 49, 4915–4918.
- (40) Xu, Z.; Xu, L.; Zhou, J.; Xu, Y.; Zhu, W.; Qian, X. A Highly Selective Fluorescent Probe for Fast Detection of Hydrogen Sulfide in Aqueous Solution and Living Cells. *Chem. Commun.* **2012**, 48, 10871–10873.
- (41) Murale, D. P.; Kim, H.; Choi, W. S.; Churchill, D. G. Highly Selective Excited State Intramolecular Proton Transfer (ESIPT)-Based Superoxide Probing. *Org. Lett.* **2013**, 15, 3946–3949.
- (42) Klymchenko, A. S.; Duportail, G.; Mely, Y.; Demchenko, A. P. Ultrasensitive Two-Color Fluorescence Probes For Dipole Potential in Phospholipid Membranes. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, 100, 11219–11224.
- (43) Yushchenko, D. A.; Fauerbach, J. A.; Thirunavukkuarasu, S.; Jares-Erijman, E. A.; Jovin, T. M. Fluorescent Ratiometric MFC Probe Sensitive to Early Stages of Alpha-Synuclein Aggregation. *J. Am. Chem. Soc.* **2010**, 132, 7860–7861.
- (44) Liu, B.; Wang, H.; Wang, T.; Bao, Y.; Du, F.; Tian, J.; Li, Q.; Bai, R. A New Ratiometric ESIPT Sensor for Detection of Palladium Species in Aqueous Solution. *Chem. Commun.* **2012**, 48, 2867–2869.
- (45) Berezin, M. Y.; Achilefu, S. Fluorescence Lifetime Measurements and Biological Imaging. *Chem. Rev.* **2010**, 110, 2641–2684.
- (46) Hanaoka, K.; Kikuchi, K.; Kojima, H.; Urano, Y.; Nagano, T. Development of a Zinc Ion-Selective Luminescent Lanthanide Chemosensor for Biological Applications. *J. Am. Chem. Soc.* **2004**, 126, 12470–12476.
- (47) Nijveldt, R. J.; Nood, E. van; Hoorn, D. E. C. van; Boelens, P. G.; Norren, K. van; Leeuwen, P. A. M. van. Flavonoids: A Review of Probable Mechanisms of Action and Potential Applications. *Am. J. Clin. Nutr.* **2001**, 74, 418–425.
- (48) Qian, F.; Zhang, C.; Zhang, Y.; He, W.; Gao, X.; Hu, P.; Guo, Z. Visible Light Excitable Zn²⁺ Fluorescent Sensor Derived From an Intramolecular Charge Transfer Fluorophore and Its in Vitro and in Vivo Application. *J. Am. Chem. Soc.* **2009**, 131, 1460–1468.
- (49) Domaille, D.; Zeng, W. L.; Chang, C. J. Visualizing Ascorbate-Triggered Release of Labile Copper within Living Cells Using a Ratiometric Fluorescent Sensor. *J. Am. Chem. Soc.* **2010**, 132, 1194–1195.
- (50) Blondeau, P.; Gauthier, R.; Berse, C.; Gravel, D. Synthesis of Some Stable 7-Halo-1,4-Thiazepines. Potential Substituted Penam Precursors. *Can. J. Chem.* **1971**, 49, 3866–3876.
- (51) Brigham, M. P.; Stein, W. H.; Moore, S. The Concentrations of Cysteine and Cysteine in Human Blood Plasma. *J. Clin. Invest.* **1960**, 39, 1633–1638.
- (52) Rubinstein, L. V.; Shomaker, R. H.; Paull, K. D.; Simon, R. M.; Tosini, S.; Skehan, P.; Scudiero, D. A.; Monko, A.; Boyd, M. R. Comparison of in Vitro Anticancer-Drug-Screening Data Generated with a Tetrazolium Assay Versus a Protein Assay Against a Diverse Panel of Human Tumor Cell Lines. *J. Natl. Cancer Inst.* **1990**, 82, 1113–1117.