Correction to New Heterometallic Hybrid Polymers Constructed with Aromatic Sulfonate-Carboxylate Ligands: Synthesis, Layered Structures, and Properties

Teresa Kurc, Jan Janczak, Józef Hoffmann, and Veneta Videnova-Adrabinska*

In the original paper, bond lengths in Table 2 were omitted. The corrected table appears below.

Table 2. Selected Bond Lengths (Å) 1−3

Crystal 1[a]			
Cd1−O1	2.2745(18)	K1−O21	2.7507(19)
Cd1−O13	2.2770(15)	K1−O12[(ii)]	2.779(2)
Cd1−O23	2.2970(18)	K1−O21[(iii)]	2.970(2)
K1−O2	2.849(2)	K1−O24[(iv)]	2.8165(17)
K1−O3	2.871(2)	K1···K1[(iii)]	3.7915(16)
K1−O11	2.967(2)		
Crystal 2[b]			
Cd−O4	2.1976(12)	Na−O5	2.3849(14)
Cd−O5	2.7610(14)	Na−O9	2.3584(15)
Cd−O8	2.3128(15)	Na−O3[(ii)]	2.3839(14)
Cd−O10	2.3198(14)	Na−O3[(iii)]	2.4406(14)
Cd−O11	2.2757(15)	Na−O6[(iv)]	2.3555(16)
Cd−O6[(i)]	2.6139(13)	Na−Na[(iv)]	3.7587(12)
Cd−O7[(i)]	2.2927(12)	Cd−Na	4.5629(11)
Cd−Cd[(v)]	4.8495(x)	Cd−Na[(v)]	4.1167(10)
Crystal 3[c]			
Cd1−O6	2.401(6)	Cd2−O26	2.285(5)
Cd1−O7	2.306(5)	Cd2−O27	2.502(5)
Cd1−O22	2.335(6)	Cd2−O31	2.345(6)
Cd1−O1[(i)]	2.319(5)	Cd2−O41	2.287(5)
Cd1−O24[(i)]	2.424(5)	Cd2−O51	2.307(6)
Cd1−O25[(i)]	2.315(5)	Cd2−O4[(iii)]	2.502(5)
Cd1−O32[(i)]	2.400(5)	Cd2−O5[(iii)]	2.267(5)
Na1−O2	2.355(5)	Na2−O4	2.441(6)
Na1−O25	2.334(5)	Na2−O27	2.276(5)
Na1−O32	2.592(5)	Na2−O31	2.597(6)
Cd1···Na1[(ii)]	3.7119(9)	Cd2···Na2	3.2892(8)
Na1···Cd1[(i)]	3.7119(9)	Na2···Cd2	3.2892(8)
Na1···Cd1[(v)]	3.7119(9)	Na2···Cd2[(iii)]	3.2892(8)

[a]Symmetry codes: (ii) −x, 1 − y, −z; (iii) 1 − x, 1 − y, −z; (iv) 1 − x, 1 − y, 1 − z. [b]Symmetry code: (i) x, 1 + y, z; (ii) 1 + x, y, z; (iii) −x, 1 − y, 1 − z; (iv) 1 − x, 1 − y, 1 − z; (v) 1 − x, 2 − y, 1 − z. [c]Symmetry code: (i) 1 + x, y, z; (ii) −1 + x, y, z; (iii) −x, 1 − y, −z; (v) 1 − x, 2 − y, 1 − z.