

Correction to Optical and Chemical Characterization of Aerosols Emitted from Coal, Heavy and Light Fuel Oil and Small-Scale Wood Combustion

Anna K. Frey,* Karri Saarnio, Heikki Lamberg, Fanni Mylläri, Panu Karjalainen, Kimmo Teinilä, Samara Carbone, Jarkko Tissari, Ville Niemelä, Anna Häyrinen, Jani Rautiainen, Jorma Kytömäki, Paulo Artaxo, Aki Virkkula, Liisa Pirjola, Topi Rönkkö, Jorma Keskinen, Jorma Jokiniemi, and Risto Hillamo

Environ. Sci. Technol., **2014**, 48, 827–836. DOI: 10.1021/es4028698

Unfortunately, in the original manuscript there was an erratum in the description of the aerosol forcing efficiency calculation.

Radiative forcing influence of emission particles was assessed by eq 1 (page 830, last paragraph of the left column before Results and Discussion section) that included a formula of upscatter fraction $\beta = 0.817 + 1.8495b - 2.9682b^2$. In this formula, the value 0.817 is erroneous. Based on the original formula presented, e.g., by Delene and Ogren (2002)¹ this value should be 0.0817. The right formula is then $\beta = 0.0817 + 1.8495b - 2.9682b^2$.¹ Despite of the erratum in the text, the results in this original study were calculated based on the correct equations and thus, results and conclusions remained unchanged.

■ REFERENCES

(1) Delene, D. J.; Ogren, J. A. Variability of aerosol optical properties at four North American surface monitoring sites. *J. Atmos. Sci.* **2002**, 59, 1135–1150.