

**Correction to Analysis of the Molecular Weight Distribution of Petroleum Asphaltenes Using Laser Desorption–Mass Spectrometry** [*Energy Fuels* **2004**, *18*, 1405. DOI: ef034083r].  
Ryuzo Tanaka,<sup>\*,†</sup> Shinya Sato,<sup>‡</sup> Toshimasa Takanohashi,<sup>‡</sup>  
Jerry E. Hunt,<sup>§</sup> and Randall E. Winans<sup>§</sup>

<sup>†</sup>Central Research Laboratories, Idemitsu Kosan Co., Ltd.,  
1280 Kamiizumi, Sodegaura 299-0293, Japan, <sup>‡</sup>Institute for  
Energy Utilization, National Institute of Advanced Industrial  
Science and Technology, 16-1 Onogawa, Tsukuba 305-8569,  
Japan, and <sup>§</sup>Chemistry Division, Argonne National Laboratory,  
9700 South Cass Avenue, Argonne, Illinois 60439

Page 1406. Table 1 should be and not as originally published.

**Table 1. Properties of Intact Asphaltenes and Their Sub-fractions' Yields**

sample	MY	KF	IL
yields in VR (wt %)	24.9	14.2	6.3
elemental analysis (wt %)			
C	82.0	82.2	83.2
H	7.5	7.6	6.8
N	1.3	0.9	1.4
S	7.1	7.6	5.9
O	1.2	1.1	1.5
metals (wt ppm)			
Ni	390	200	390
V	1800	550	1200
atomic ratio			
H/C	1.09	1.10	0.97
N/C	0.01	0.00	0.01
S/C	0.07	0.08	0.06
O/C	0.01	0.01	0.01
$M_n$ (VPO)	4000	4000	2400
$f_a$ ( $^{13}\text{C}$ NMR)	0.53	0.51	0.54
subfraction yields (wt %)			
Fr-1	17.5	16.7	16.9
Fr-2	12.7	10.8	17.2
Fr-3	13.1	10.4	17.3
Fr-4	13.4	10.3	16.2
Fr-5	12.3	12.9	14.5
Fr-6	31.0	38.8	18.0

DOI: 10.1021/ef9015679  
Published on Web 01/25/2010