

Correction to Nanoplastic Affects Growth of *S. obliquus* and Reproduction of *D. magna*

Ellen Besseling,* Bo Wang, Miquel Lürling, and Albert A. Koelmans

Environ. Sci. Technol. 2014, 48, 12336-12343; DOI: 10.1021/es503001d

Two corrections to the Implication Section of our paper are needed. The corrections do not change the conclusions of our paper. In the original paper, a range of 0.04–34 ng/L was mentioned for the microplastic concentrations found in fresh water in the U.S.² and Europe.³ This range should have been 0.4–34 ng/L. Furthermore, a trawling depth of 0.01 m was mentioned, which should have been 0.1 m. Correction of these errors results in the conclusion that our observed effect concentrations are a factor 10^6-10^8 higher (previously: 10^6) than the 0.4–34 ng/L microplastic concentrations found in fresh water in the U.S.² and Europe,³ and about 2 orders of magnitude higher than the highest reported microplastic concentration in marine water.⁴ The implications and conclusions based on these comparisons remain the same.

■ REFERENCES

- (1) Besseling, E.; Wang, B.; Lürling, M.; Koelmans, A. A. Nanoplastic affects growth of *S. obliquus* and reproduction of *D. magna. Environ. Sci. Technol.* **2014**, *48*, 12336–12343.
- (2) Eriksen, M.; Mason, S.; Wilson, S.; Box, C.; Zellers, A.; Edwards, W.; Farley, H.; Amato, S. Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar. Pollut. Bull.* **2013**, *77*, 177–182.
- (3) Besseling, E.; Foekema, E. M.; Koelmans, A. A. Verkennend onderzoek microplastic in het beheersgebied van waterschap rivierenland; Wageningen University; Wageningen, The Netherlands, 2014; pp 1–18
- (4) Lopez Lozano, R.; Mouat, J. Marine litter in the North-East Atlantic Region; OSPAR Commission: London, United Kingdom, 2009; pp 1–120.

