

Noncryogenic Preparation of Functionalized Arylboronic Esters through a Magnesium–Iodine Exchange with in Situ Quench


Emilien Demory, Véronique Blandin, Jacques Einhorn, and Pierre Y. Chavant*

Département de Chimie Moléculaire, UMR-5250, ICMG FR-2607, CNRS, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 09, France

Org. Process Res. Dev., **2011**, 15(3), 710–716; 10.1021/op2000089**S** Supporting Information

We recently found out that compound **3j** (Table 1, entry 10) had not been prepared from 2-bromo-6-iodopyridine, but from 2-bromo-5-iodopyridine. Thus, **3j** is actually 2-bromo-5-(4,4,6-trimethyl-1,3,2-dioxaborinan-2-yl)pyridine. This prompted us to check the reaction of 2-bromo-6-iodopyridine, as well as that of 2-iodopyridine, in our standard conditions; we were unable to isolate any boronic ester from these attempts. We apologize for any inconvenience caused by this error.

Thus the structure in Table 1, entry 10 should be:



Page 711, column 2: “2-Pyridyl boronic acids are notoriously unstable, and ester derivatives are therefore attractive compounds;²⁷ we were thus pleased to obtain 6-bromopyridin-2-yl boronic ester **3j** in excellent yield.” should be:

“2-Pyridyl boronic acids are notoriously unstable, and ester derivatives are therefore attractive compounds;²⁷ unfortunately, all our attempts to isolate (2-pyridyl)boronic esters by the present method failed. Conversely, the (6-bromo-3-pyridyl)-boronic ester **3j** could be obtained in excellent yield (Table 1, entry 10).”

Experimental section (page 715, column 1): **3j** should be 2-bromo-5-(4,4,6-trimethyl-1,3,2-dioxaborinan-2-yl)pyridine.

■ ASSOCIATED CONTENT**S** Supporting Information

The skeletal structure of **3j** was corrected. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION**Corresponding Author**

*Telephone: (+)33 476635796. Fax: (+)33 476635983. E-mail: Pierre-Yves.Chavant@ujf-grenoble.fr.

Published: July 3, 2012