

Correction to iTRAQ-based Quantitative Protein Expression Profiling and MRM Verification of Markers in Type 2 Diabetes

Prabhjit Kaur, Nasser M. Rizk, Sereen Ibrahim, Noura Younes, Arushi Uppal, Kevin Dennis, Tejaswita Karve, Kenneth Blakeslee, John Kwagyan, Mahmoud Zirie, Habtom W. Resson, and Amrita K. Cheema*

J. Proteome Res. **2012**, *11* (11), 5527–5537. 10.1021/pr300798z

On page 5531 of the original article, Table 1 was formatted incorrectly. The correct Table 1 is shown here.

Table 1. Characteristics of T2DM ($n = 106$) and Nondiabetic Control ($n = 76$) Volunteers^a

	control	T2DM
Clinical characteristics		
Age	32.2 \pm 1.45	53.6 \pm 1.43
BMI(kg/m ²)	28.9 \pm 0.90	32.37 \pm 0.97
HbA _{1c} (%)	5.57 \pm 0.05	7.66 \pm 0.06
Vitamin D (ng/mL)	17.5 \pm 1.40	12.90* \pm 1.34
Glucose (mM)	4.88 \pm 0.05	8.71** \pm 0.10
Triglycerides (mM)	1.16 \pm 0.77	1.72 \pm 0.90
LDL (mM)	2.80 \pm 0.05	2.54 \pm 0.07
HDL (mM)	4.00 \pm 1.23	1.23** \pm 0.01
Insulin (pM)	72.2 \pm 3.69	91.43* \pm 4.52

^aSerum samples obtained from individual subjects were analyzed to determine the levels of glucose, Vitamin D among others (as described in methods). The value spreads within the group for each parameter are indicated as mean \pm standard error. The clinical parameters that had a significant change ($P \leq 0.05$) between the control and T2DM groups are indicated with an asterix. As expected, the diabetic cohort showed an elevation in serum glucose, HbA_{1c}, and insulin levels and a decrease in the serum levels of Vitamin D and HDL.