

Copper-Catalyzed Synthesis of Phenanthridine Derivatives under an Oxygen Atmosphere Starting from Biaryl-2-carbonitriles and Grignard Reagents


Line Zhang, Gim Yean Ang, and Shunsuke Chiba*

Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore

shunsuke@ntu.edu.sg

Received June 29, 2010

ABSTRACT



A copper-catalyzed synthesis of phenanthridine derivatives was developed starting from biaryl-2-carbonitriles and Grignard reagents. The present transformation is carried out by a sequence of nucleophilic addition of Grignard reagents to biaryl-2-carbonitriles to form *N*-H imines and their Cu-catalyzed C–N bond formation on the aromatic C–H bond, where molecular oxygen is a prerequisite to achieve the catalytic process.

Phenanthridines and their derivatives are of great interest in medicinal chemistry and material science due to their potent biological activities¹ and optoelectronic properties.² Although diverse approaches toward the construction of a phenanthridine skeleton have been reported so far,³ versatile and efficient methodologies to synthesize phenanthridines with selective control of substitution patterns using readily accessible building blocks are still needed. Herein we wish to

report a copper-catalyzed synthesis of phenanthridine derivatives under an oxygen atmosphere starting from biaryl-2-

(1) (a) Bernardo, P. H.; Wan, K. F.; Sivaraman, T.; Xu, J.; Moore, F. K.; Hung, A. W.; Mok, H. Y. K.; Yu, V. C.; Chai, C. L. *J. Med. Chem.* **2008**, *51*, 6699. (b) Zhu, S.; Ruchelman, A. L.; Zhou, N.; Liu, A.; Liu, L. F.; LaVoie, E. J. *Bioorg. Med. Chem.* **2005**, *13*, 6782. (c) Lewis, W. G.; Green, L. G.; Grynszpan, F.; Radic, Z.; Carlier, P. R.; Taylor, P.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 1053. (d) Lynch, M. A.; Duval, O.; Sukhanova, A.; Devy, J.; MacKay, S. P.; Waigh, R. D.; Nabiev, I. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2643. (e) Cappelli, A.; Anzini, M.; Vomero, S.; Mannuni, L.; Makovec, F.; Doucet, E.; Hamon, M.; Bruni, G.; Romeo, M. R.; Menziani, M. C.; Benedetti, P. G.; Langer, T. *J. Med. Chem.* **1998**, *41*, 728.


(2) (a) Stevens, N.; O'Connor, N.; Vishwasrao, H.; Samaroo, D.; Kandel, E. R.; Akins, D. L.; Drain, C. M.; Turro, N. J. *J. Am. Chem. Soc.* **2008**, *130*, 7182. (b) Bondarev, S. L.; Knyukshto, V. N.; Tikhomirov, S. A.; Pyrko, A. N. *Opt. Spectrosc.* **2006**, *100*, 386. (c) Zhang, J.; Lakowicz, J. R. *J. Phys. Chem. B* **2005**, *109*, 8701.

(3) For recent reports on the synthesis of substituted phenanthridines and their derivatives, see: (a) Budén, M. E.; Dorn, V. B.; Gamba, M.; Pierini, A. B.; Rossi, R. A. *J. Org. Chem.* **2010**, *75*, 2206. (b) Youn, S. W.; Bihn, J. H. *Tetrahedron Lett.* **2009**, *50*, 4598. (c) Kohno, K.; Azuma, S.; Choshi, T.; Nobuhiro, J.; Hibino, S. *Tetrahedron Lett.* **2009**, *50*, 590. (d) Candito, D. A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 6713. (e) Gerfaud, T.; Neuville, L.; Zhu, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 572. (f) Portela-Cubillo, F.; Scanlan, E. M.; Scott, J. S.; Walton, J. C. *Chem. Commun.* **2008**, 4189. (g) Mandadapu, A. K.; Saifuddin, M.; Agarwal, P. K.; Kundu, B. *Org. Biomol. Chem.* **2009**, *7*, 2796. (h) Yanada, R.; Hashimoto, K.; Tokizane, R.; Miwa, Y.; Minami, H.; Yanada, K.; Ishikura, M.; Takemoto, Y. *J. Org. Chem.* **2008**, *73*, 5135. (i) Bowman, W. R.; Lyon, J. E.; Gareth, J. P. *Synlett* **2008**, *14*, 2169. (j) Sripada, L.; Teske, J. A.; Deiters, A. *Org. Biomol. Chem.* **2008**, *6*, 263. (ii) Sanz, R.; Fernández, Y.; Castroviejo, M. P.; Pérez, A.; Fañanás, F. J. *Eur. J. Org. Chem.* **2007**, *8*, 62. (k) Shabashov, D.; Daugulis, O. *J. Org. Chem.* **2007**, *72*, 7720. (l) Wang-Ge, S.; Yun-Yun, Y.; Yan-Guang, W. *J. Org. Chem.* **2006**, *71*, 9241. (m) Alonso, R.; Campos, P. J.; García, B.; Rodríguez, M. A. *Org. Lett.* **2006**, *8*, 3521. (n) Banwell, M. G.; Lupton, D. W.; Ma, X.; Renner, J.; Sydnese, M. O. *Org. Lett.* **2004**, *6*, 2741. (o) Mehta, B. K.; Yanagisawa, K.; Shiro, M.; Kotsuki, H. *Org. Lett.* **2003**, *5*, 1605. (p) Pawlas, J.; Begtrup, M. *Org. Lett.* **2002**, *4*, 2687. (q) Lysén, M.; Kristensen, J. L.; Vedsø, P.; Begtrup, M. *Org. Lett.* **2002**, *4*, 257. (r) Moreno, I.; Tellitu, I.; Etayo, J.; San Martín, R.; Dominguez, E. *Tetrahedron* **2001**, *57*, 5403.

carbonitriles and Grignard reagents. The present transformation is carried out by a sequence of addition of Grignard reagents to biaryl-2-carbonitriles to form *N*-H imines and their Cu-catalyzed intramolecular cyclization including C–N bond formation on the aromatic C–H bond, where molecular oxygen is a prerequisite to achieve the catalytic process.


We have recently reported generation of iminyl copper species from α -azido carbonyl compounds and their copper-catalyzed C–C bond cleavage under an oxygen atmosphere, where nitriles were synthesized.^{4,5} During the course of the study, it was found that a reaction of ethyl 2-azido-2-(biphenyl-2-yl)acetate (**1**) provided the desired biphenyl-2-carbonitrile (**2a**) in 46% yield along with 50% yield of phenanthridine **3**, which might be formed via aromatic C–H bond functionalization/C–N bond formation of the iminyl copper species (Scheme 1).

Scheme 1. Copper-Catalyzed Reaction of Ethyl 2-Azido-2-(biphenyl-2-yl)acetate (**1**)



To explore an efficient synthetic method of phenanthridines via the iminyl copper species, we planned to use biaryl-2-carbonitriles and organometallic reagents as shown in Scheme 2, which commences with nucleophilic addition

Scheme 2. Synthetic Plan of Phenanthridines from Biaryl-2-carbonitriles and Organometallic Reagents




of R-[M] to biaryl-2-carbonitriles to afford *N*-H imines **A** after proper protonation. Consecutive treatment of resulting

N-H imines with a catalytic amount of Cu salts under an oxygen atmosphere would give iminyl copper species **B**,⁶ which could lead to formation of phenanthridines **4**.

On the basis of this hypothesis, the formation of phenanthridines was investigated using biphenyl-2-carbonitrile (**2a**)⁷ and *p*-tolylmagnesium bromide, and Table 1 lists the represen-

Table 1. Optimization of Reaction Conditions



entry	metal salts	time (h)	yield (%) ^a	
			4a	5a
1	Cu(OAc) ₂	26	91	0
2	CuCl ₂	19	88	6
3	Cu(OTf) ₂	19	93	0
4	CuCl	18	89	0
5	CuTC	18	90	2
6 ^b	Cu(OAc) ₂	65	21	36
7	Pd(OAc) ₂	24	2	72
8	Co(OAc) ₂	24	1	99
9	Mn(OAc) ₃	24	2	83
10	FeCl ₃	24	0	71

^a Isolated yields. ^b The reaction was carried out using 1 equiv of Cu(OAc)₂ under an N₂ atmosphere using degassed DMF. Tol = 4-methylphenyl; CuTC = copper(I) thiophene-2-carboxylate.

tative data. Addition of *p*-tolylmagnesium bromide to nitrile **2a** occurred smoothly in Et₂O at 60 °C (in sealed tube). After protonation with MeOH,⁸ DMF (diluted to 0.1 M) and metal salts (10 mol %) were subsequently added, and the reaction mixture was stirred at 80 °C under an oxygen atmosphere (1 atm). It was found that several copper salts, either Cu(I) or Cu(II), exhibited good catalytic activity toward the formation of phenanthridine **4a** (entries 1–5). In contrast, the reaction using a stoichiometric amount of Cu(OAc)₂ without oxygen (under a N₂ atmosphere) provided phenanthridine **4a** in only 21% yield along with 36% yield of *N*-H imine **5a**, suggesting that molecular oxygen plays a vital role to achieve the catalytic C–N bond formation process (entry 6).⁹ Other metal complexes such as Pd(II), Co(II), Mn(III), and Fe(III) were not viable catalysts for this transformation (entries 7–10).

By utilizing Cu(OAc)₂ as the catalyst (entry 1 in Table 1), we examined the generality of this catalytic method for synthesis of substituted phenanthridines. First, scope of Grignard reagents were examined using biphenyl-2-carbonitrile (**2a**) (Table 2). Sterically hindered 2-methyl- and 2,6-dimethylphenyl moieties as well as an electron-deficient 4-chlorophenyl part could be installed with good to excellent

(4) Chiba, S.; Zhang, L.; Ang, G. Y.; Hui, B. W.-Q. *Org. Lett.* **2010**, *12*, 2052.

(5) For a report on copper-catalyzed synthesis of azaspirocyclohexadienones from α -azido-*N*-arylamides under an oxygen atmosphere via iminyl peroxycopper(III) intermediates, see: Chiba, S.; Zhang, L.; Lee, J.-Y. *J. Am. Chem. Soc.* **2010**, *132*, 7266.


(6) For a report on formation of iminyl rhodium(I) complexes from *N*-H diarylimines, see: Zhao, P.; Hartwig, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 11618.

Table 2. Reaction Scope of Grignard Reagents^a

entry	R ¹	time (h)	yield (%) ^b
1	2-Me-C ₆ H ₄	19	4b , 99
2	2,6-Me ₂ -C ₆ H ₃	20	4c , 88
3	4-Cl-C ₆ H ₄	32	4d , 91
4	4-MeO-C ₆ H ₄	36	4e , 5 (5e , 90)
5	2-thienyl	60	4f , 54 (2a , 41)
6 ^c	CH ₂ CH ₂ Ph	31	4g , 76
7 ^d	<i>i</i> -Pr	22	4h , 67

^a Unless otherwise noted, reactions were carried out using 0.5 mmol of biaryl-2-carbonitriles **2a** with 1.3 equiv of Grignard reagents in Et₂O (0.5 mL) at 60 °C (sealed tube) for 2 h followed by addition of MeOH (60 μL), DMF (4 mL), and Cu(OAc)₂ (10 mol %); the mixture was stirred at 80 °C under an O₂ atmosphere. ^b Isolated yields. ^c The reaction of **2a** with Ph(CH₂)₂MgBr was completed in 24 h at 60 °C. ^d The reaction of **2a** with *i*-PrMgBr (2 equiv) was carried out at 80 °C for 24 h.

yields (entries 1–3). An electron-donating group such as methoxy on the benzene ring retarded the present cyclization, affording the corresponding phenanthridine **4e** in only 5% yield along with 90% yield of *N*-H imine **5e** even after

Scheme 3. Reaction Scope on Substituents of Biaryl-2-carbonitriles **2**^{a,b}

^a Unless otherwise noted, reactions were carried out using 0.5 mmol of biaryl-2-carbonitriles **2** with 1.3 equiv of *p*-tolylmagnesium bromide in Et₂O (0.5 mL) at 60 °C (sealed tube) for 2 h followed by addition of MeOH (60 μL), DMF (4 mL), and Cu(OAc)₂ (10 mol %), and the mixture was stirred at 80 °C under an O₂ atmosphere. ^b Isolated yields and reaction times on copper-catalyzed cyclization are recorded in parentheses. ^c 20 mol % of Cu(OAc)₂ was used.

stirring for 36 h (entry 4). When 2-thienylmagnesium bromide was utilized, the desired phenanthridine **4d** and biphenyl-2-carbonitrile (**2a**) were isolated in 54% and 41% yields, respectively, although nitrile **2a** was once consumed by the reaction with the Grignard reagent (entry 5). It was speculated that regeneration of nitrile **2a** might proceed via C–C bond cleavage of 2-thienyliminocopper species (see Supporting Information for more detail). Alkyl Grignard reagents also could be used, affording phenanthridines **4** in good yields (entries 6 and 7).

Next, various biaryl-2-carbonitriles **2** were utilized to prepare substituted phenanthridines (Scheme 3). By varying substituent R² on C(3) of phenanthridine **4**, both electron-donating and -withdrawing groups could be installed (for **4i–4m**). Several substituents such as F, CF₃, and Me were also successfully introduced at the C(7), C(8), and C(9) of phenanthridines (for **4n–4q**).

This catalytic method allowed accessing polycyclic aza-aromatic hydrocarbons (aza-PAHs)¹⁰ (Table 3). The reaction

Table 3. Synthesis of Polycyclic Aza-Aromatic Compounds (aza-PAHs)^a

entry	nitriles 6	time (h) ^b	aza-PAHs 7	yield (%) ^c
1		46		7a : 86%
2		21		7b : 86%
3		64		7c : 81%
4 ^d		70		7d : 68%

^a Reactions were carried out using 0.5 mmol of biaryl-2-carbonitriles **6** with 1.3 equiv of *p*-tolylmagnesium bromide in Et₂O (0.5 mL) at 60 °C (sealed tube) for 2 h followed by addition of MeOH (60 μL), DMF (4 mL), and Cu(OAc)₂ (10 mol %); the mixture was stirred at 80 °C under an O₂ atmosphere. ^b Reaction times on copper-catalyzed cyclization. ^c Isolated yields. ^d The reaction of **6d** (0.3 mmol) with *p*-tolylmagnesium bromide (2 equiv) was carried out in toluene (1 mL) at 80 °C (sealed tube) for 2 h followed by addition of MeOH (60 μL), DMF (4 mL), and Cu(OAc)₂ (20 mol %); the mixture was stirred at 120 °C under an O₂ atmosphere.

of 1-phenyl-2-naphthonitrile (**6a**) with *p*-tolylmagnesium bromide provided tetracyclic benzo[*k*]phenanthridine **7a** in good yield (entry 1). Aza-chrysene (benzo[*c*]phenan-

thridine) **7b** could be accessed selectively from 2-(naphthalen-2-yl)benzonitrile (**6b**) (entry 2). Pentanuclear aza-aromatic hydrocarbons, dibenzo[*c,k*]phenanthridine **7c** and dibenzo[*c,i*]phenanthridine (aza-picene) **7d**, were also synthesized starting from binaphthyl-2-carbonitriles **6c** and **6d**, respectively, although longer reaction time and higher temperature (120 °C for **7d**) were required (entries 3 and 4). In the cases of **6b**, **6c**, and **6d**, C–H functionalization occurred exclusively on the α -carbon (marked in blue) of the naphthalene ring, which suggested that an electrophilic aromatic substitution pathway¹¹ might be involved in the mechanism of the copper-catalyzed C–N bond formation (see Supporting Information for putative reaction mechanisms).^{12–14}

(7) (a) Lysén, M.; Köhler, K. *Synlett* **2005**, 1671. (b) Cahiez, G.; Lepifre, F.; Ramiandrasoa, P. *Synthesis* **1999**, 2138.

(8) Pickard, P. L.; Tolbert, T. L. *J. Org. Chem.* **1961**, 26, 4886.

(9) No C–N bond formation was observed without Cu catalysts.

(10) For recent selected reports on mutagenic and carcinogenic activity of aza-PAHs, see: (a) Xue, W.; Warshawsky, D. *Toxicol. Appl. Pharmacol.* **2005**, 206, 73. (b) Yamada, K.; Suzuki, T.; Kohara, A.; Kato, T.; Hayashi, M.; Mizutani, T.; Saeki, K. *Mutat. Res.* **2005**, 586, 1. (c) Yuan, Z.-X.; Kumar, S.; Sikka, H. C. *Chem. Res. Toxicol.* **2004**, 17, 672.

(11) (a) Field, L. D.; Sternhell, S.; Wilton, H. V. *J. Chem. Educ.* **1999**, 76, 1246. (b) Gore, P. H. *Chem. Rev.* **1955**, 55, 229.

(12) For a review on metal-catalyzed C–H amination, see: Collet, F.; Dodd, R. H.; Dauban, P. *Chem. Commun.* **2009**, 5061. For a review on copper-catalyzed aromatic C–H functionalization, see: Zhang, M. *Appl. Organometal. Chem.* **2010**, 24, 269.

In conclusion, a facile synthetic method of phenanthridine derivatives has been developed starting from biaryl-2-carbonitriles and Grignard reagents via Cu-catalyzed C–N bond formation on the aromatic C–H bond under an O₂ atmosphere. Continuous studies on the scope, mechanistic evaluation, and synthetic applications of the present method toward various polynuclear aza-aromatic hydrocarbons are in progress.

Acknowledgment. This work was supported by funding from Nanyang Technological University and Singapore Ministry of Education.

Supporting Information Available: Experimental procedures and characterization of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL101490N

(13) For a report on copper-catalyzed functionalization of aryl C–H bonds/C–N bond formation of amidines for synthesis of benzimidazoles, see: Brasche, G.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2008**, 47, 1932.

(14) For reports on copper-catalyzed-aromatic C–H functionalization, see: (a) Bernini, R.; Fabrizi, G.; Sferrazza, A.; Cacchi, S. *Angew. Chem., Int. Ed.* **2009**, 48, 8078. (b) Yang, L.; Lu, Z.; Stahl, S. S. *Chem. Commun.* **2009**, 6460. (c) Ueda, S.; Nagasawa, H. *Angew. Chem., Int. Ed.* **2008**, 47, 6411. (d) Ban, I.; Sudo, T.; Taniguchi, T.; Itami, K. *Org. Lett.* **2008**, 10, 3607. (e) Chen, X.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, 128, 6790.