

## STRUCTURE NOTE

# Crystal structure of the proline iminopeptidase-related protein TTHA1809 from *Thermus thermophilus* HB8

Masahiko Okai,<sup>1</sup> Yumiko Miyauchi,<sup>1</sup> Akio Ebihara,<sup>2</sup> Woo Cheol Lee,<sup>1</sup> Koji Nagata,<sup>1,3</sup> and Masaru Tanokura<sup>1,3\*</sup>

<sup>1</sup> Department of Applied Biological Chemistry, Graduate School of Agricultural and Life Sciences, University of Tokyo, Tokyo 113-8657, Japan

<sup>2</sup> RIKEN SPring-8 Center, Harima Institute, Hyogo 679-5148, Japan

<sup>3</sup> Theoretical Structural Biology Laboratory, RIKEN Harima Institute at SPring-8, Sayo-gun, Hyogo 679-5148, Japan

**Key words:** hydrolase; X-ray; thermophilic bacterium; structural comparison; catalytic triad.

## INTRODUCTION

*Thermus thermophilus* HB8, an extremely thermophilic bacterium, is a model organism in structural biology. The complete sequence of the genome has been determined ([www.thermus.org](http://www.thermus.org)), and the sequence information provided us with the presence of a large number of hypothetical proteins.

The open reading frame of TTHA1809 from *Thermus thermophilus* HB8 was annotated as a proline iminopeptidase-related protein. Proline iminopeptidase (PIP) is a serine peptidase that catalyzes the removal of N-terminal proline from peptides with high specificity. The enzyme was reported for the first time by Sarid *et al.*,<sup>1</sup> and several genes have been cloned.<sup>2–7</sup> The crystal structure of proline iminopeptidase from *Xanthomonas campestris* revealed that the enzyme has an  $\alpha/\beta$ -hydrolase fold and a Ser-His-Asp catalytic triad.<sup>8</sup> The  $\alpha/\beta$ -hydrolase fold family is one of the largest families with a variety of catalytic functions and substrate specificities. Members in the family include proline iminopeptidase, acetylcholinesterase, proline oligopeptidase, haloalkane dehalogenase, haloperoxidase, epoxide hydrolase, and hydroxynitrile lyase.<sup>9</sup>

In this study, we report the crystal structure of the proline iminopeptidase-related protein TTHA1809 from *Thermus thermophilus* HB8, and compare the active site of the tricorner-interacting aminopeptidase F1, which is the best homolog found using the Dali program, with the corresponding region of TTHA1809.

## MATERIALS AND METHODS

The coding gene for TTHA1809 was amplified by polymerase chain reaction (PCR) using the genomic DNA of *Thermus thermophilus* HB8 as the template. The PCR product was cloned into pET-11a (Novagen). The *Escherichia coli* strain BL21(DE3) was used as the host for protein expression. The cells were suspended in 20 mM Tris-HCl buffer (pH 8.0) containing 50 mM NaCl and 5 mM  $\beta$ -mercaptoethanol. The cell lysate was heated at 343 K for 10 min, and after centrifugation, the supernatant was loaded onto a Resource ISO column. The protein was eluted with a linear gradient of 1.5 to 0M ammonium sulfate. The fractions containing TTHA1809 protein were dialyzed against 20 mM Tris-HCl (pH 8.0). The dialyzed protein was loaded onto a Resource Q column. The protein was eluted with a linear gradient of 0 to 0.5M NaCl. The dialyzed protein was applied to a hydroxyapatite CHT5 column equilibrated with 10 mM sodium phosphate buffer (pH 7.0), and the flow-through fractions were collected. The protein was loaded onto a HiLoad

Grant sponsor: Ministry of Education, Culture, Sports, Science, and Technology of Japan (National Project on Protein Structural and Functional Analyses).

\*Correspondence to: Masaru Tanokura, Department of Applied Biological Chemistry, Graduate School of Agricultural and Life Sciences, University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan. Tel: 81-3-5841-5165; Fax: 81-3-5841-8023. E-mail: [amtanok@mail.ecc.u-tokyo.ac.jp](mailto:amtanok@mail.ecc.u-tokyo.ac.jp)

Received 8 June 2007; Accepted 9 July 2007

Published online 3 January 2008 in Wiley InterScience ([www.interscience.wiley.com](http://www.interscience.wiley.com)). DOI: 10.1002/prot.21892

**Table 1**

Data Collection and Refinement Statistics for Native and Se-Met Labeled Crystals

Diffraction data	Native	Se-Met
Beamline	SPring-8 BL41XU	Photon Factory NW12
Wavelength (Å)	1.0000	0.9790
Space group	$P3_221$	$P3_221$
Unit-cell parameters (Å)	$a = b = 126.9$ $c = 114.3$	$a = b = 127.1$ $c = 114.3$
Resolution (Å)	20–2.20 (2.28–2.20)	20–2.10 (2.17–2.10)
No. of measurements	366887	687801
No. unique reflections	54254	62447
Completeness (%)	99.6 (100.0)	99.9 (99.9)
$R_{\text{merge}}^a$	0.063 (0.226)	0.062 (0.144)
$\langle I \rangle / \langle \sigma \rangle$	10.6	27.0
<b>Refinement statistics</b>		
Resolution range (Å)	20–2.2	
$R_{\text{cryst}}^b$ (%)	17.7	
$R_{\text{free}}^c$ (%)	20.8	
<b>RMS deviations</b>		
Bonds (Å)	0.009	
Angles (°)	1.229	
<b>Average B factors (Å<sup>2</sup>)</b>		
Protein	28.1	
Water	32.5	
Glycerol	67.4	
<b>Ramachandran plot</b>		
Most favored regions (%)	92.5	
Additionally favored regions (%)	7.5	
PDB ID	2YYS	

<sup>a</sup> $R_{\text{merge}} = \sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle| / \sum_{hkl} \sum_i I_i(hkl)$ , where  $\langle I(hkl) \rangle$  is the average of individual measurements of  $I_i(hkl)$ .

<sup>b</sup> $R_{\text{cryst}} = \sum_{hkl} \|F_{\text{obs}} - |F_{\text{calc}}|\| / \sum_{hkl} |F_{\text{obs}}|$ .

<sup>c</sup> $R_{\text{free}}$  was calculated using 5% of reflections excluded in the refinement.

16/60 Superdex 75 column and eluted with 20 mM Tris-HCl buffer (pH 8.0) containing 150 mM NaCl. The purified protein containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, and 1 mM dithiothreitol (DTT) was concentrated to 15 mg/mL. Se-Met labeled protein was overexpressed in *Escherichia coli* B834(DE3) and purified using the same steps as in the native protein.

Crystallization was performed using the sitting-drop vapor diffusion method at 293 K. Crystals of TTHA1809 (0.2 mm × 0.15 mm × 0.07 mm) were obtained in 2 days by mixing 1 μL of the protein solution and 1 μL of the reservoir solution containing 100 mM HEPES-Na (pH 7.2) and 0.85M potassium sodium tartrate tetrahydrate. Se-Met labeled crystals were obtained under the same conditions as the native protein crystals. Native and Se-Met labeled crystals were transferred to the reservoir solution containing 25% (v/v) glycerol as the cryoprotectant and flash-cooled at 100 K in liquid nitrogen. Diffraction data were collected in a nitrogen cryostream at the beamline BL41XU of SPring-8 and at the beamline NW12 of the Photon Factory Advanced Ring (PF-AR). The diffraction data were indexed and scaled with

HKL2000.<sup>10</sup> The crystal contained two monomers per asymmetric unit according to the Matthews coefficient.<sup>11</sup>

The structure of TTHA1809 was determined by the single-wavelength anomalous diffraction (SAD) method. Five of the six expected Se atoms in the asymmetric unit were found using the program SOLVE.<sup>12</sup> The program RESOLVE<sup>12</sup> was used to improve phases. The initial model was built with ARP/wARP.<sup>13</sup> The remainder were built manually with XtalView<sup>14</sup> and refined with REFMAC5.<sup>15</sup> The native structure was then determined by the molecular replacement program MOLREP.<sup>16</sup> Then, simulated annealing and B-factor refinements were performed using CNS,<sup>17</sup> and further refinements were performed using XtalView and Refmac5. The final model consisted of 564 amino acid residues, 6 glycerol molecules, and 413 water molecules in the asymmetric unit. Residues 285–286 of molecule A and residues 281–286 of molecule B were not visible due to disorder. The quality of the final model was checked by PROCHECK.<sup>18</sup>


## RESULTS AND DISCUSSION

### Overall structure

The TTHA1809 protein from *Thermus thermophilus* HB8 consists of 286 amino acid residues with a predicted molecular weight of 31,550. The crystal structure of the TTHA1809 protein was solved by the SAD method and refined with 20–2.2 Å resolution diffraction data (Table 1). The quality of the structure was assessed by Ramachandran plot, and no residue fell in the generously allowed and disallowed regions.

TTHA1809 exists as a monomer with approximate dimensions of 40 × 45 × 55 Å<sup>3</sup> and contains the two domains shown in Figure 1(A). The large domain (core domain) is composed of the N-terminal residues 1–128 and the C-terminal residues 210–286, and shows an α/β-hydrolase fold which consists of an eight-stranded mainly parallel β-sheet flanked on both sides by α-helices. The small domain (residues 129–209) contains five α-helices (α4(α8) and one <sub>3</sub><sub>10</sub>-helix (η3)). The two molecules of TTHA1809 in the asymmetric unit are quite similar, with a root mean square deviation (RMSD) of 0.52 Å (1–275 C<sup>α</sup> atoms).

A structural similarity search using DALI<sup>19</sup> showed that TTHA1809 resembles the following proteins: the tri-corn-interacting aminopeptidase F1 (proline iminopeptidase) from *Thermoplasma acidophilum* (Z-score = 24.8; PDB code 1MT3)<sup>20</sup> with an RMSD of 2.8 Å for 250 C<sup>α</sup>, 2-hydroxyl-6-oxo-6-phenylhexa-2,4-dienoate hydrolase BphD from *Rhodococcus* sp. strain RHA1 (Z-score = 24.8; PDB code 1C4X)<sup>21</sup> with an RMSD of 2.9 Å for 247 C<sup>α</sup>, chloroperoxidase A2 from *Streptomyces aureofaciens* (Z-score = 24.1; PDB code 1BRT)<sup>22</sup> with an RMSD of 2.6 Å for 242 C<sup>α</sup>, prolyl aminopeptidase from *Serratia marcescens* (Z-score = 23.6; PDB code 1QTR)<sup>23</sup> with an

**Figure 1**

(A) The ribbon representation of the TTHA1809 monomer. The helices,  $\beta$ -strands, and random coils are colored red, yellow, and green, respectively. (B) The superimposition between the TTHA1809 protein (green) and the tricorm-interacting aminopeptidase F1 (magenta) from *Thermoplasma acidophilum*. The catalytic triad residues of F1 and the corresponding regions of TTHA1809 are represented in stick models.

RMSD of 3.0 Å for 250 C $\alpha$ , and aclacinomycin methyl-esterase RdmC from *Streptomyces purpurascens* (Z-score = 23.2; PDB code 1Q0R)<sup>24</sup> with an RMSD of 3.0 Å for 248 C $\alpha$ . These proteins are members of the  $\alpha/\beta$ -hydrolase family. TTHA1809 differed from these proteins in orientation of the helices of the small domain.

#### Comparison with the tricorm-interacting aminopeptidase F1

The tricorm-interacting aminopeptidase F1<sup>20</sup> from *Thermoplasma acidophilum* has a Ser-His-Asp catalytic

triad in the active site. The superimposition between TTHA1809 and F1 revealed that the residue corresponding to the catalytic Ser105 in F1 is replaced by a Gly at TTHA1809 [Fig. 1(B)]. Asp229 and His255 of TTHA1809 were located at the same position as Asp244 and His271 of the catalytic triad of F1. His255 of TTHA1809 was located in a loop between the  $\beta$ 8-strand and the  $\alpha$ 10-helix. The ND-1 atom of His255 was hydrogen-bonded to Asp229 in a loop located between the  $\beta$ 7-strand and the  $\alpha$ 9-helix, whereas the NE-2 atom of His255 formed a hydrogen bond with a water molecule because of the lack of catalytic Ser residue. Thus, TTHA1809 and F1 would have different functions. A BLAST search using TTHA1809 revealed the possibility of the existence of enzymes lacking a catalytic Ser residue in several microorganisms (*Thermus thermophilus* HB27, *Deinococcus geothermalis* DSM 11300, *Legionella pneumophila* subsp. *pneumophila* str. Philadelphia 1, *Dechloromonas aromatica* RCB, and *Hahella chejuensis* KCTC 2396), but their functions have not yet been revealed.

#### REFERENCES

1. Sarid S, Berger A, Katchalski E. Proline iminopeptidase. *J Biol Chem* 1959;234:1740–1746.
2. Kitazono A, Yoshimoto T, Tsuru D. Cloning, sequencing, and high expression of the proline iminopeptidase gene from *Bacillus coagulans*. *J Bacteriol* 1992;174:7919–7925.
3. Albertson NH, Koomey M. Molecular cloning and characterization of a proline iminopeptidase gene from *Neisseria gonorrhoeae*. *Mol Microbiol* 1993;9:1203–1211.
4. Atlan D, Gilbert C, Blanc B, Portalier R. Cloning, sequencing and characterization of the pepIP gene encoding a proline iminopeptidase from *Lactobacillus delbrueckii* subsp. *bulgaricus* CNRZ 397. *Microbiology* 1994;140 (Part 3):527–535.
5. Kitazono A, Kitano A, Tsuru D, Yoshimoto T. Isolation and characterization of the prolyl aminopeptidase gene (pap) from *Aeromonas sobria*: comparison with the *Bacillus coagulans* enzyme. *J Biochem (Tokyo)* 1994;116:818–825.
6. Tamura T, Tamura N, Lottspeich F, Baumeister W. Tricorn protease (TRI) interacting factor 1 from *Thermoplasma acidophilum* is a proline iminopeptidase. *FEBS Lett* 1996;398:101–105.
7. Kabashima T, Kitazono A, Kitano A, Ito K, Yoshimoto T. Prolyl aminopeptidase from *Serratia marcescens*: cloning of the enzyme gene and crystallization of the expressed enzyme. *J Biochem (Tokyo)* 1997;122:601–605.
8. Medrano FJ, Alonso J, García JL, Romero A, Bode W, Gomis-Ruth FX. Structure of proline iminopeptidase from *Xanthomonas campestris* pv. *citri*: a prototype for the prolyl oligopeptidase family. *EMBO J* 1998;17:1–9.
9. Holmquist M. Alpha/beta-hydrolase fold enzymes: structures, functions and mechanisms. *Curr Protein Pept Sci* 2000;1:209–235.
10. Otwinowski Z, Minor W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol* 1997;276:307–326.
11. Matthews BW. Solvent content of protein crystals. *J Mol Biol* 1968;33:491–497.
12. Terwilliger TC, Berendzen J. Automated MAD and MIR structure solution. *Acta Crystallogr D* 1999;55:849–861.
13. Perrakis A, Morris R, Lamzin VS. Automated protein model building combined with iterative structure refinement. *Nat Struct Biol* 1999;6:458–463.

14. McRee DE. XtalView/Xfit—A versatile program for manipulating atomic coordinates and electron density. *J Struct Biol* 1999;125:156–165.
15. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D* 1997;53:240–255.
16. Vagin AA, Teplyakov A. MOLREP: an automated program for molecular replacement. *J Appl Cryst* 1997;30:1022–1025.
17. Brünger AT, Adams PD, Clore GM, DeLano WL, Gros P, Grosse-Kunstleve RW, Jiang JS, Kuszewski J, Nilges M, Pannu NS, Read RJ, Rice LM, Simonson T, Warren GL. Crystallography & NMR system: a new software suite for macromolecular structure determination. *Acta Crystallogr* 1998;D54:905–921.
18. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to check the stereochemical quality of protein structures. *J Appl Crystallogr* 1993;26:283–291.
19. Holm L, Sander C. Protein structure comparison by alignment of distance matrices. *J Mol Biol* 1993;233:123–138.
20. Goettig P, Groll M, Kim JS, Huber R, Brandstetter H. Structures of the tricorn-interacting aminopeptidase F1 with different ligands explain its catalytic mechanism. *EMBO J* 2002;21:5343–5352.
21. Nandhagopal N, Senda T, Hatta T, Yamada A, Masai E, Fukuda M, Mitsui Y. Three-dimensional structure of microbial 2-hydroxyl-6-oxo-6-phenylhexa-2,4-dienoic acid (HPDA) hydrolase (BphD enzyme) from *Rhodococcus* sp. strain RHA1, in the PCB degradation pathway. *Proc Jpn Acad Ser B* 1997;73:154–157.
22. Hofmann B, Tölzer S, Pelletier I, Altenbuchner J, van Pée KH, Hecht HJ. Structural investigation of the cofactor-free chloroperoxidases. *J Mol Biol* 1998;279:889–900.
23. Yoshimoto T, Kabashima T, Uchikawa K, Inoue T, Tanaka N, Nakamura KT, Tsuru M, Ito K. Crystal structure of prolyl aminopeptidase from *Serratia marcescens*. *J Biochem (Tokyo)* 1999;126:559–565.
24. Jansson A, Niemi J, Mantsala P, Schneider G. Crystal structure of aclacinomycin methylesterase with bound product analogues: implications for anthracycline recognition and mechanism. *J Biol Chem* 2003;278:39006–39013.