

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/260840472>

Synthesis and tripanocidal activity of ferrocenyl and benzyl diamines against *Trypanosoma brucei* and *Trypanosoma cruzi*

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY LETTERS · FEBRUARY 2014

Impact Factor: 2.42 · DOI: 10.1016/j.bmcl.2014.02.046 · Source: PubMed

CITATIONS

2

READS

56

11 AUTHORS, INCLUDING:



Andréa Kohatsu

Universidade Estadual Paulista "Júlio de M...

1 PUBLICATION 2 CITATIONS

SEE PROFILE



Vera Isaac

São Paulo State University

28 PUBLICATIONS 67 CITATIONS

SEE PROFILE



Maria D. Vargas

Universidade Federal Fluminense

112 PUBLICATIONS 1,395 CITATIONS

SEE PROFILE



Regina M B Cicarelli

São Paulo State University

79 PUBLICATIONS 1,263 CITATIONS

SEE PROFILE

Accepted Manuscript

Synthesis and tripanocidal activity of ferrocenyl and benzyl diamines against *Trypanosoma brucei* and *Trypanosoma cruzi*

Angela Maria Arenas Velásquez, Acácio Ivo Francisco, Andréa Akiko Nakaima Kohatsu, Flavia Alves de Jesus Silva, Danilo Fernando Rodrigues, Rafaela Gomes da Silva Teixeira, Bruna Galdorfini Chiari, Maria Gabriela José de Almeida, Vera Lucia Borges Isaac, Maria D. Vargas, Regina Maria Barretto Cicarelli


PII: S0960-894X(14)00178-4
DOI: <http://dx.doi.org/10.1016/j.bmcl.2014.02.046>
Reference: BMCL 21364

To appear in: *Bioorganic & Medicinal Chemistry Letters*

Received Date: 12 December 2013
Revised Date: 13 February 2014
Accepted Date: 18 February 2014

Please cite this article as: Velásquez, A.M.A., Francisco, A.I., Kohatsu, a.A.N., Silva, F.A.d., Rodrigues, D.F., Teixeira, R.G.d., Chiari, B.G., Almeida, M.G.J., Isaac, V.L.B., Vargas, M.D., Cicarelli, R.M.B., Synthesis and tripanocidal activity of ferrocenyl and benzyl diamines against *Trypanosoma brucei* and *Trypanosoma cruzi*, *Bioorganic & Medicinal Chemistry Letters* (2014), doi: <http://dx.doi.org/10.1016/j.bmcl.2014.02.046>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.




Graphical Abstract

To create your abstract, type over the instructions in the template box below.

Synthesis and trypanocidal activity of ferrocenyl and benzyl diamines against *Trypanosoma cruzi* and *Trypanosoma brucei*

Velásquez, A. M. A.^{a,b}; Francisco, A. I.^c; Kohatsu, A. A. N.^a; Silva, F. J. A.^a; Rodrigues, D. F.^{a,b}; Teixeira, R. G. S.^c; Chiari, B. G.^d; De Almeida, M. G. J.^d; Isaac, V. L.^d; Vargas, M. D.^c; Cicarelli, R. M. B.^a

Leave this area blank for abstract info.



Fonts or abstract dimensions should not be changed or altered.



Synthesis and tripanocidal activity of ferrocenyl and benzyl diamines against *Trypanosoma brucei* and *Trypanosoma cruzi*

Angela Maria Arenas Velásquez^{a,b,*}, Acácio Ivo Francisco^c, Andréa Akiko Nakaima Kohatsu^a, Flavia Alves de Jesus Silva^a, Danilo Fernando Rodrigues^{a,b}, Rafaela Gomes da Silva Teixeira^c, Bruna Galdorfini Chiari^d, Maria Gabriela José de Almeida^d, Vera Lucia Borges Isaac^d, Maria D. Vargas^c and Regina Maria Barretto Cicarelli^{a,*}.

^aDepartamento de Ciências Biológicas, Faculdade de Ciências Farmacêuticas, Universidade Estadual Paulista "Júlio de Mesquita Filho", Rodovia Araraquara-Jaú Km 01-Campus, Araraquara, 14.801-902, Brazil.

^bInstituto de Química, Universidade Estadual Paulista "Júlio de Mesquita Filho", Rua Prof. Francisco Degni, 55, Araraquara, 14.800-900, Brazil

^cInstituto de Química, Universidade Federal Fluminense, Campus do Valonguinho - Centro, Niterói, 24.020-150, Brazil.

^dDepartamento de Fármacos e Medicamentos, Faculdade de Ciências Farmacêuticas, Universidade Estadual Paulista "Júlio de Mesquita Filho", Rodovia Araraquara-Jaú Km 01-Campus, Araraquara, 14.801-902, Brazil.

ARTICLE INFO

Article history:

Received

Revised

Accepted

Available online

Keywords:

Sleeping sickness

Chagas disease

Benzyl diamines

Ferrocenyl diamines

ABSTRACT

Trypanosoma brucei and *Trypanosoma cruzi* are the etiologic agents of sleeping sickness and Chagas disease, respectively, two of the 17 preventable tropical infectious diseases (NTD) which have been neglected by governments and organizations working in the health sector, as well as pharmaceutical industries. High toxicity and resistance are problems of the conventional drugs employed against trypanosomiasis, hence the need for the development of new drugs with trypanocidal activity. In this work we have evaluated the trypanocidal activity of a series of *N1,N2*-dibenzylethane-1,2-diamine hydrochlorides (benzyl diamines) and *N1*-benzyl,*N2*-methylferrocenylethane-1,2-diamine hydrochlorides (ferrocenyl diamines) against *T. brucei* and *T. cruzi* parasite strains. We show that incorporation of the ferrocenyl group into the benzyl diamines increases the trypanocidal activity. The molecules exhibit potential trypanocidal activity *in vitro* against all parasite strains. Cytotoxicity assay was also carried out to evaluate the toxicity in HepG2 cells.

2009 Elsevier Ltd. All rights reserved.

* Corresponding author. Tel.: +55-16-33016946; fax: +55-16-3301-6940; e-mail: avellangel@gmail.com (Velásquez, A. M. A.), reginacicarelli@gmail.com (Cicarelli, R. M. B.).

The *Trypanosomatidae* family, flagellated parasitic protozoa, is responsible for important infectious diseases in humans: sleeping sickness, Chagas diseases and leishmaniasis. Cost, toxicity and resistance problems of conventional drugs point to an urgent need to identify and develop new therapeutic alternatives. Chagas disease is a zoonosis that spans the Great Lakes in North America to the southeast of Patagonia^{1,2} and occurs widely in Latin America¹. The infection can be readily transmitted by blood transfusion and migration from endemic areas. The causative agent is *T. cruzi*³. According to the World Health Organization, WHO, an estimated 10 million people are infected worldwide, mostly in Latin America and more than 25 million people are at risk of the disease. It is estimated that in 2008 Chagas disease killed over 10.000 people⁴. The only prescription drugs in medical practice to treat Chagas disease are nifurtimox and benznidazole^{5,6}. These medicines are curative only in acute or early chronic infections, and in congenital cases⁷. Benznidazole is more effective than nifurtimox in suppressing parasitemia during the chronic phase⁸, but neither is effective in curing the disease at this stage⁹. Both drugs have significant side effects, probably as a consequence of reducing oxidative damage in tissues or hosts¹⁰.

Sleeping sickness is caused by *T. b. gambiense* and *T. b. rhodesiense*, subspecies related to *T. b. brucei*, which is found in domestic animals and related to the disease called nagana¹¹. WHO reports that *T. b. gambiense* accounts for 95% of reported cases of sleeping sickness¹². The estimated number of actual cases is currently 30.000. The symptoms of sleeping sickness can be varied, but two clinical phases can be recognized during infection: an early hemolymphatic stage and a late meningoencephalitic stage. Treatment of the first phase of sleeping sickness relies on suramin and pentamidine and in the late phase, melarsoprol is active against *T. b. gambiense* and *T. b. rhodesiense*, whereas eflornithine is useful only against *T. b. gambiense*. Melarsoprol is a highly toxic arsenic-based drug, which can lead to patient death. A new combination therapy with the oral drugs nifurtimox and eflornithine is more effective when compared to nifurtimox and eflornithine monotherapies, but causes several side effects¹³. In addition, eflornithine reduces the number of leukocytes, erythrocytes and platelets¹⁴. These drugs are mostly antiquated, scarce, highly toxic and subject of parasite resistance¹³⁻¹⁷. Clearly there is an urgent need to develop new drugs to treat these diseases.

Polyamines are present in all parasitic protozoa, and in recent years, these compounds and their associated enzymes have been investigated as drug targets in search of novel parasitic therapy options¹⁸. A large number of *N*-alkylated polyamine analogs have been described which selectively interfere with *T. cruzi* polyamine metabolism¹⁹. Furthermore, in a series of recent reports, Yamanaka et al.²⁰ (2013) provided evidence that simpler lipophilic alkyl diamines can be leading molecules for the development of new antiparasitic drugs. They described the *in vitro* activity of several 1,2-ethane, 1,4-butane and 1,6-hexanediamine hydrochlorides against *T. cruzi*, *L. braziliensis* and *L. chagasi* and showed that in the case of *T. cruzi* the mitochondria is a target for the active compounds²¹. Differently, diaryl diamines [2-(3-(substituted-benzylamino)propylamino)-quinolin-4(1H)-ones] were shown to inhibit *T. brucei* and *T. cruzi* growth by selectively inhibiting their methionyl-tRNA synthetase²¹.

Incorporation of transition metal fragments to the structure of a drug often enhances its activity²²⁻²⁴. Of special interest in this work is the ferrocenyl (Fc) group²⁵, which is derived from ferrocene [bis(η^5 -cyclopentadienyl)iron(II)], a non-toxic and

stable organometallic compound²⁶. Incorporation of the Fc group into standard drugs offers new possibilities in therapeutic applications and reversal of drug resistances²⁷. E.g. introduction of Fc into the lateral chain of chloroquine yielded ferroquine²⁸, with excellent *in vitro* and *in vivo* activity against malaria parasites - *Plasmodium spp.* and particularly good activity against chloroquine-resistant ones^{29,30}. Furthermore substitution of a phenyl group in tamoxifen with Fc resulted in improved cytotoxic activity in breast cancer^{27,31,32}, and modification of ethambutol with Fc groups led to improved antimicrobial activity (*Mycobacterium tuberculosis*)²⁸.

Herein we describe the synthesis of several *N1,N2*-dibenzylethane-1,2-diamine hydrochlorides (benzyl diamines) and *N1*-benzyl,*N2*-methylferrocenylethane-1,2-diamine hydrochlorides (ferrocenyl diamines) and the evaluation of their trypanocidal activity (Table 1). Compounds **2-10** (Figure 1) have been obtained as stable solids (yields ranging from 66 to 80%); their structures are fully supported by ¹H and ¹³C NMR spectroscopy and elemental analysis. Compounds **2a-10a** (Figure 1), described previously, have been synthesized according to the literature³³⁻³⁵. Synthesis and full spectroscopic data are presented in the Supplementary Information.

The compounds were tested against *T. brucei* and *T. cruzi* strains using MTT colorimetric assay to evaluate the trypanocidal activity^{36,37}. Pentamidine and benznidazole were included in the assays as positive controls for *T. brucei* and *T. cruzi*, respectively. MTT assays were carried out after 24 h (*T. brucei*) and 72 h (*T. cruzi*) incubation. The methodology is detailed in the supplementary information. A cytotoxicity assay was also carried out to evaluate the toxicity of different concentrations of the compounds in HepG2 cells. This hepatoma cell line is used as a model to simulate the hepatic function of the human organism *in vitro*, as it exhibits similar metabolic functions to the human liver^{38,39}.

The benzyl diamines showed higher trypanocidal effect in *T. brucei* strains than in *T. cruzi* Y (see Table 1), in accord with a previous report on *T. cruzi* strains being less responsive than *T. brucei* to this type of compounds²⁰. Especially for *T. cruzi* Y the nature of the substituent strongly influenced the IC₅₀ values. Compound **3a** bearing a 4-OMe group on the phenyl rings is by far the most active compound against *T. cruzi* Y (IC₅₀ = 10.9 μ M) and also the least toxic one of the series, with an S.I. value of about 90, higher than both pentamidine and benznidazole. It was also the most active compound of the series against both strains of *T. brucei*, with IC₅₀ values comparable to that of pentamidine (8.25 and 8.63 vs. 6.43 μ M) and lower mammalian toxicity.

Upon substitution of one of the benzyl or pyridyl groups in the structures of the benzyl diamines **2a**, **3a**, **5a**, **7a**, **8a** and **10a** for an Fc group to generate the corresponding ferrocenyl diamines **2**, **3**, **5**, **7**, **8** and **10**, respectively (Figure 1), IC₅₀ values decreased considerably in *T. cruzi* (Table 1), demonstrating that the presence of the Fc group leads to increased trypanocidal activity to *T. cruzi*. In fact ferrocenyl diamines exhibited higher trypanocidal activity than the corresponding benzyl diamines on both parasites (*T. brucei* 427, *T. brucei* 29-13 and *T. cruzi* Y) which might be associated to increased lipophilicity and/or differences in electronic (redox properties) and geometric structure^{29,30}. Compounds **3** (IC₅₀ = 0.36 μ M) and **6** (IC₅₀ = 0.35 μ M) were especially active against *T. brucei* 427; compound **3** (IC₅₀ = 0.82 μ M), against *T. brucei* 29-13 and compound **4** (IC₅₀ = 2.21 μ M), against *T. cruzi* Y. All ferrocenyl diamines exhibited higher mammalian toxicity compared to the corresponding benzyl diamines.




Figure 1. Synthesis of benzyl and ferrocenyl diamines.

Table 1. Antiparasitic activities and safety index of benzyl diamines and ferrocenyl diamines.

Table 1. Antiparasitic activities and safety index of benzyl diamines and ferrocenyl diamines.									
	Compound	R	HepG2 cells ^a	<i>T. brucei</i> 427 ^a	S.I. ^b		S.I. ^b		S.I. ^b
					<i>T. brucei</i> 427	<i>T. brucei</i> 29-13 ^a	<i>T. brucei</i> 29-13	<i>T. cruzi</i> Y ^a	
Ferrocenyl diamines	2	C ₆ H ₅	30.99	0.51	<u>60.76</u>	4.04	7.67	6.03	5.14
	3	4-OMeC ₆ H ₄	18.60	0.36	<u>51.67</u>	0.82	<u>22.68</u>	4.23	4.40
	4	2-OMeC ₆ H ₄	32.26	1.78	<u>18.12</u>	7.69	4.20	2.21	<u>14.60</u>
	5	2-OHC ₆ H ₄	34.49	1.27	<u>27.16</u>	5.33	6.47	9.58	3.60
	6	Fc	9.20	0.35	<u>26.29</u>	4.81	1.91	5.59	1.65
	7	Py(3Cl ⁻)	85.45	7.46	<u>11.45</u>	10	8.55	3.74	<u>22.85</u>
	8	2,4-ClC ₆ H ₃	21.94	0.45	<u>48.76</u>	4.58	4.79	3.98	5.51
	9	2-BrC ₆ H ₄	26.20	0.41	<u>63.90</u>	1.13	<u>23.19</u>	3.37	7.77
	10	4-NO ₂ C ₆ H ₄	22.68	2.28	9.95	3.56	6.37	6.07	3.74
	11	4(R)-7-chloroquinoline	20.31	10.42	1.95	6.48	3.13	12.84	1.58
Benzyl diamines	2a	C ₆ H ₅	223.74	11.08	<u>20.19</u>	9.99	<u>22.40</u>	159.61	1.40
	3a	4-OMeC ₆ H ₄	990.62	8.25	<u>120.08</u>	8.63	<u>114.79</u>	10.9	<u>90.88</u>
	5a	2-OMeC ₆ H ₄	148.21	4.61	<u>32.15</u>	5.16	<u>28.72</u>	108.9	1.36
	7a	py	86.91	13.88	6.26	14.11	6.16	258.97	0.34
	8a	2,4-ClC ₆ H ₃	280.57	12.35	<u>22.72</u>	16.78	<u>16.72</u>	51.44	5.45
	10a	4-NO ₂ C ₆ H ₄	590.18	36.58	<u>16.13</u>	19.22	<u>30.71</u>	122.01	4.84
	Pentamidine	---	159.71	6.43	24.84	6.43	24.84	4.58	34.87
	Benznidazole	---	2126.16	207.57	10.24	149.58	14.21	34.00	62.53

^aIC₅₀ (μM): half maximal inhibitory concentration

^bS.I.: Safety index. OBS: The underlined S.I. values indicate promising compounds, i.e. those with low mammalian toxicity and high toxicity to parasites.

Although pentamidine and other diamidines, such as furamidine, exhibit high potency, they display poor oral bioavailability and unfavorable side effects⁴⁰. These dicationic compounds have been studied primarily against sleeping sickness and few have been assayed as anti-*T. cruzi* candidates⁴¹. Nevertheless, in this work, pentamidine was active against the epimastigote forms of *T. cruzi* Y strain (IC₅₀ = 4.58 μM).

In their study, de Almeida et al.²⁰ evaluated the *in vitro* trypanocidal activity of several lipophilic diamines against *T. cruzi* epimastigote (IC₅₀ = 11.9–15.1 μM) and amastigote forms (IC₅₀ = 1.6–23.6 μM), and observed mitochondrial depolarization in *T. cruzi* epimastigote forms treated with the most active compounds of the series, which suggested that this mechanism may be involved in the trypanocidal effect.

In biological systems ferrocene is oxidized by H_2O_2 in the presence of peroxidases to the ferrocenium cation, which forms charge-transfer complexes with protein donor groups⁴². For parasite survival and growth, trypanosomatids possess a defense mechanism against oxidative stress. Superoxide dismutase (SOD), old yellow enzyme (OYE) and peroxiredoxine (Prx) are involved in various mitochondrial activities in certain life cycle stages of trypanosomatids¹⁶. The ferrocenyl group could therefore interfere with the defense mechanisms of the parasites.

According to the IC₅₀ values in HepG2 cells (Table 1) the ferrocenyl amines showed lower toxic activity on mammalian cells, as compared with their toxicity against parasites, which is very interesting for new drugs evaluation.

These ferrocenyl diamines could therefore be potential trypanocidal drug candidates. Studies on the mechanism of action of the ferrocenyl diamines against both *T. cruzi* and *T. brucei* are underway, and so are the tests with other *T. cruzi* strains, due to genetic variability, and with the *T. cruzi* amastigote form and the bloodstream forms of *T. brucei*.

Acknowledgements

The authors thank FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), FUNDUNESP (Fundação para o Desenvolvimento da Unesp) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for research funding and financial support.

Supplementary Information

Supplementary information associated with this article can be found, in the online version, at doi:

References

- Jannin, J.; Villa, L. Mem. Inst. Oswaldo Cruz 2007, 102, 95.
- Schmunis, G. A. Mem. Inst. Oswaldo Cruz 2007, 102, 75.
- Teixeira, A. R. L.; Nitz, N.; Guimaro, M. C.; Gomes, C.; Santos-Buch, C. A. Postgrad. Med. J. 2006, 82, 788.
- World Health Organization. Chagas disease (American trypanosomiasis) Fact sheet N°340. Available at <http://www.who.int/mediacentre/factsheets/fs340/en/>, 2010 (accessed 05.16.12).
- Coura, J. R. Mem. Inst. Oswaldo Cruz 2009, 104, 549.
- Wilkinson, S. R.; Kelly, J. M. Expert Rev. Mol. Med. 2009, 11, 1.
- Rojo, G.; Castillo, C.; Duaso, J.; Liempi, A.; Droguett, D.; Galanti, N.; Maya, J.D.; López-Muñoz, R.; Kemmerling, U. Acta Trop. 2014, in press, Toxic and therapeutic effects of Nifurtimox and Benznidazol on *Trypanosoma cruzi* ex vivo infection of human placental chorionic villi explants, <http://dx.doi.org/10.1016/j.actatropica.2014.01.002>.
- Coura, J. R.; Borges-Pereira, J. Rev. Soc. Bras. Med. Trop. 2012, 45, 286.
- Pereira, P. C. M.; Navarro, E. C. J. Venom. Anim. Toxins Incl. Trop. Dis. 2013, 19, 34.
- Urbina, J. A.; Docampo, R. Trends Parasitol. 2003, 19, 495.
- Cox, F. E. G. Infect. Dis. Clin. North Am. 2004, 18, 231.
- World Health Organization. Human African trypanosomiasis (sleeping sickness) Fact sheet N°259. Available at

- <http://www.who.int/mediacentre/factsheets/fs259/en/>, 2012 (accessed 05.16.12).
- Babokhov, P.; Sanyaolu, A. O.; Oyibo, W. A.; Fagbenro-Beyioku, A. F.; Iriemenam, N. C. Pathog. Glob. Health 2013, 107, 242.
- Campbell, R. A.; Kurtz, G. L.; Norton-Reel, D. J. Life Sci. 1991, 48, 225.
- Nok, A. J. Parasitol. Res. 2003, 90, 71.
- Fidalgo, L. M.; Gille, L. Pharm. Res. 2011, 28, 2758.
- Legros D.; Ollivier G.; Gastellu-Etcheberry M.; Paquet C.; Burri C.; Jannin J.; Büscher P. Lancet Infect. Dis. 2002, 2, 437.
- Müller, S.; Coombs, G. H.; Walter, R. D. Trends Parasitol. 2001, 17, 242.
- Casero, R.A. Jr.; Woster, P.M. J. Med. Chem. 2001, 44, 1.
- Yamanaka, C. N.; Giordani, R. B.; Rezende Jr, C. O.; Eger, I.; Kessler, R. L.; Tonini, M. L.; Moraes, M. H.; Araújo, D. P.; Zuanazzi, J. A.; De Almeida, M. V.; Steidel, M. Chem. Biol. Drug Des. 2013, 82, 697 and references therein.
- Shibata, S.; Gillespie, J. R.; Kelley, A. M.; Napuli, A. J.; Zhang, Z.; Kovzun, K. V.; Pefley, R. M.; Lam, J.; Zucker, F. H.; Van Voorhis, W. C.; Merritt, E. A.; Hol, W. G. J.; Verlinde, C. L. M. J.; Fan, E.; Buckner, F. S. Antimicrob. Agents Chemother. 2011, 55, 1982.
- Jaouen, G.; Vessières, A.; Butler, I. S. Acc. Chem. Res. 1993, 26, 361.
- Sharma, V.; Piwnicka-Worms, D. Chem. Rev. 1999, 9, 2545.
- Schatzschneider, U.; Metzler-Nolte, N. Angew. Chem. Int. Ed. 2006, 45, 1504.
- Van Staveren, D. R.; Metzler-Nolte, N. Chem Rev. 2004, 104, 5931.
- Biot, C.; Glorian, G.; Maciejewski, L. A.; Brocard, J. S. J. Med. Chem. 1997, 40, 3715.
- Neuse, E. W. J. Inorg. Organomet. Polym. 2005, 15, 3.
- Ralambomanana, D. A.; Razafigahefa-Ramilison, D.; Rakotohova, A. C.; Maugein, J. Péliniski, L. Bioorg. Med. Chem. 2008, 16, 9546.
- Biot, C.; Nosten, F.; Fraisse, L.; Ter-Minassian, D.; Khalife, J.; Dive, D. Parasite, 2011, 18, 207.
- Domarle, O.; Blampain, G.; Agnani, H.; Nzadiyabi, T.; Lebibi, J.; Brocard, J.; Maciejewski, L.; Biot, C.; Georges, A. J.; Millet, P. Antimicrob. Agents Chemother. 1998, 42, 540.
- Hillard, E.; Vessières, A.; Thouin, L.; Jaouen, G.; Amatore, C. Angew. Chem. Int. Ed. Engl. 2006, 45, 285.
- Heilmann, J. B.; Hillard, E. A.; Plamont, M.; Pigeon, P.; Bolte, M.; Jaouen, G.; Vessières, A. J. Organomet. Chem. 2008, 693, 1716.
- Khan, M. S. Y.; Chawla, G. Indian J. Chem., Sect B 2002, 41B, 653.
- Sharma, V.; Khan, M. S. Y. Eur. J. Med. Chem. 2001, 36, 651.
- Khan, M. S. Y.; Husain, A.; Sharma, S.; Rashid, M. Indian J. Pharm. Sci. 2012, 74, 80.
- Mosmann, T. J. Immunol. Methods 1983, 65, 55.
- Cotinguiba, F.; Regasini, L. O.; Bolzani, V. S.; Debonsi, H. M.; Passerini, G. D.; Cicarelli, R. M. B.; Kato, M. J.; Furlan, M. Med. Chem. Res. 2009, 18, 703.
- Sassa, S.; Sugita, O.; Galbraith, R.A.; Kappas, A. Biochem. Biophys. Res. Commun. 1987, 143, 52.
- Natarajan, A.T.; Darroudi, F. Mutagenesis 1991, 6, 399.
- Werbovetz, K. Curr. Opin. Invest. Drugs 2006, 7, 147.

41. Wilson, W. D.; Tanious, F. A.; Mathis, A.; Tevis, D.; Hall, J. E.; Boykin, D. W. *Biochimie* 2008, 90, 999.
42. Sai-Krishna, A.D.; Panda, G.; Kondapi, A. K. *Arch. Biochem. Biophys.* 2005, 438, 206.