

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/265688602>

Synthesis of new 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles: A search for novel nitric oxide donor anti-inflammatory agen...

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY LETTERS · SEPTEMBER 2014

Impact Factor: 2.42 · DOI: 10.1016/j.bmcl.2014.09.024

CITATION

1

READS

26

3 AUTHORS:



Khaled Abdellatif

Beni Suef University

51 PUBLICATIONS 401 CITATIONS

SEE PROFILE



Abeer Moawad

Beni Suef University

8 PUBLICATIONS 19 CITATIONS

SEE PROFILE



Ed Knaus

University of Alberta

187 PUBLICATIONS 1,832 CITATIONS

SEE PROFILE



Synthesis of new 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles: A search for novel nitric oxide donor anti-inflammatory agents

Khaled R. A. Abdellatif^{a,*}, Abeer Moawad^b, Edward E. Knaus^c

^a Pharmaceutical Organic Chemistry Department, Faculty of Pharmacy, Beni-Suef University, Beni-Suef, Egypt

^b Pharmacognosy Department, Faculty of Pharmacy, Beni-Suef University, Beni-Suef, Egypt

^c Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta T6G 2N8, Canada

ARTICLE INFO

Article history:

Received 5 August 2014

Revised 5 September 2014

Accepted 8 September 2014

Available online 16 September 2014

Keywords:

Pyrazoles

Cyclooxygenase

Diazen-1-ium-1,2-diolate nitric oxide donors

Anti-inflammatory activity

ABSTRACT

A group of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (**12a–f**) was synthesized and evaluated as anti-inflammatory agents. While all the compounds (20 mg/kg) showed significant anti-inflammatory activity after 3 h of inflammation induction (69–89%) as compared to celecoxib (80%), 1-(4-methanesulfonylphenyl)-5-(4-methylaminomethylphenyl)-3-trifluoromethyl-1H-pyrazole (**12a**) was found to be the most effective one (89%). The synthesis of model hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives of 1-(4-methanesulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (**10a–f**) requires further investigation since the reaction of *N*-(4-(1-(4-(methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)ethanamine (**12b**) or 1-(4-(1-(4-(methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)piperazine (**12c**) with nitric oxide furnished *N*-nitroso derivatives (**13** and **14**), respectively, rather than the desired *N*-diazen-1-ium-1,2-diolate derivatives (**10b** and **10c**).

© 2014 Elsevier Ltd. All rights reserved.

The anti-inflammatory activity (AI) of non-steroidal anti-inflammatory drugs (NSAIDs) results from enzymatic inhibition of cyclooxygenase (COX) mediated production of pro-inflammatory prostaglandins (PGs) and thromboxanes (TXs).^{1,2} The cyclooxygenase enzyme exists in two distinct isoforms, a constitutive form (COX-1) and an inducible form (COX-2). The constitutive COX-1 plays an important role in the maintenance of physiological functions such as protection of gastric mucosa, vascular homeostasis, and platelet aggregation, whereas the inducible COX-2 is over expressed during acute and chronic inflammation.³ Traditional NSAIDs interact with both forms (COX-1 and COX-2) and therefore their long term administration often causes gastro intestinal side effects that encompass gastric ulceration and bleeding. For this reason, synthesis of selective COX-2 inhibitor drugs (coxibs) attracted much interest in recent years that achieve the same anti-inflammatory activity as traditional NSAIDs but with minimal risk of undesirable gastrointestinal side effects.⁴ Celecoxib (**1**), rofecoxib (**2**), valdecoxib (**3**), and etoricoxib (**4**) were the most common coxibs approved for therapeutic use. Unfortunately, some drugs within this group such as rofecoxib (**1**) and valdecoxib (**3**)

alter the natural balance in the COX pathway wherein the amount of the desirable vasodilatory and anti-aggregatory prostacyclin (PGI₂) produced is decreased in conjunction with a concomitant increase in the level of the undesirable prothrombotic thromboxane A₂ (TxA₂).^{5–7} This combination of adverse biochemical changes in the COX pathway are responsible for increased prevalence of high blood pressure and myocardial infarction that subsequently caused the withdrawal of rofecoxib (Vioxx®) and valdecoxib (Bextra®).^{8,9}

Nitric oxide (NO) exhibits a number of useful pharmacological actions that include vascular relaxation (vasodilation), and inhibition of platelet aggregation and adhesion.¹⁰ Hybrid COX-2 inhibitors possessing a NO-donor moiety (NO-coxibs) have been investigated as a method to increase the clinical safety of COX-2 inhibitors. In this regard, we previously reported NO-coxib ester prodrugs (**5–9**) having a NO-donor diazen-1-ium-1,2-diolate moiety that are effectively cleaved by esterases to release the parent anti-inflammatory drug and NO.^{11–15} The spontaneous decomposition reaction of *N*-diazoniumdiolates (**5–9**, Fig. 1) in physiological medium would release (upon esterase-mediated hydrolysis) not only the active components (coxib and NO) but also 1 equiv of the corresponding amine and 1 equiv formaldehyde for **5**, in addition to 1 equiv acetic acid for **6–9**.^{11–15} Considering the toxicity of

* Corresponding author. Tel.: +20 1002535444; fax: +20 822317958.

E-mail address: khaled.ahmed@pharm.bsu.edu.eg (K.R.A. Abdellatif).





Figure 1. Chemical structures of the selective cyclooxygenase-2 (COX-2) inhibitors rofecoxib (1), celecoxib (2), valdecoxib (3), etoricoxib (4), the NO coxibs (5–9), and the putative hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives of 1-(4-methanesulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (10a–f).

these byproducts we initiated a study to address this issue by developing strategies to minimize or eliminate the risk of exposure to toxic compounds in patients who may be treated with NO-coxibs. Accordingly, we now describe an investigation directed toward the design and synthesis of model hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives of 1-(4-methanesulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (10a–f) that upon decomposition releases only the active components (coxib and NO) (see structure 10a–f in Fig. 1).

A group of 1-(4-methanesulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (12a–f) were synthesized in good yield (57–71%) via reaction of 1-(4-methanesulfonylphenyl)-5-(4-bromomethylphenyl)-

3-trifluoromethyl-1H-pyrazole (11a) or (11b) with methylamine, ethylamine or piperazine in dry THF at 25 °C. An initial study was directed toward synthesis of the target hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives 10b and 10c from *N*-(4-(1-(4-(methanesulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)ethanamine (12b) and 1-(4-(1-(4-(methanesulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)piperazine (12c) respectively. However, reaction of the substituted amino derivatives 12b and 12c with nitric oxide gas (40 psi) under a variety of experimental conditions (aprotic and protic solvent systems, various temperatures) afforded the *N*-nitroso derivatives 13 and 14, respectively, as the sole isolable products rather than the desired products 10b and 10c (see Scheme 1). The formation of the *N*-nitroso products 13 and 14 upon reaction of 12b and 12c





Scheme 1. Reagents and conditions: (a) THF, 25 °C, 1 h; (b) THF, NaOMe (95%), NO (40 psi), –65 °C, 1 h or MeCN, NaOMe (25% in MeOH), NO (40 psi), 25 °C, 24 h.

with nitric oxide indicates that the initially formed unstable intermediates (*N*-amino-*N*-diazen-1-ium-1,2-diolate products of **12b** and **12c**) must undergo protonation of the diazen-1-ium-1,2-diolate *N*²-nitrogen which subsequently eliminates HNO (see mechanism shown in Scheme 1).^{16,17}

N-Ethyl-*N*-(4-(1-(4-(methylsulfonyl)phenyl)-3-(trifluoromethyl)-1*H*-pyrazol-5-yl)benzyl)nitrous amide (**13**) shows duplicate signals in the ¹H NMR and ¹³C NMR spectra (Figs. 2 and 3) which suggest the presence of a pair of rotamers. Rotamers are conformational isomers where interconversion by rotation around a single

bond is restricted. The phenomenon of hindered rotation around C–C bond due to the presence of a bulky group in the vicinity of this bond is well known in many organic compounds. For example, methoxy group can hinder the rotation around the biflavanyl bond in biflavonoids.¹⁸ HPLC analysis for compound **13**, using a variety of solvent systems, showed only one peak. The two rotamers existed in a ratio of 1:0.6 according to the integration for the CH₃ resonance of the ethyl group at δ 0.89 and δ 1.31 in CDCl₃ and in a ratio of 1:0.76 in DMSO-*d*₆. The restricted rotation appears to be around the benzyl (CH₂–N) bond. It can be seen that the signals

Figure 2. ^1H NMR for compound (13) (400 MHz, CDCl_3).




Figure 4. NOESY spectrum for compound (13).


aromatic protons of the methanesulfonylphenyl ring showed two overlapped doublets, and the $\text{SO}_2\text{-CH}_3$ protons appeared as one singlet that integrated as 5.6 protons at δ 3.00. Complete assignment of the two rotamers was performed using 1D and 2D NMR techniques (COSY, NOESY, HSQC, and HMBC). Careful examination of HMBC spectrum showed no long range coupling between the protons of one rotamer and the carbons of the other indicating that compound **13** is not a dimer. The possibility of having a dimer is also unlikely due to the presence of the two identical compounds in different ratios, and the fact that their ratio is altered depending upon the NMR solvent used. The NMR solvent is a determinant of hydrogen bonding capability which in turn affects the stability (ratio) of the rotamers.¹⁹

Hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1*H*-pyrazoles (see structure **10a–f** in Fig. 1) constitute a potential class of selective COX-2 inhibitor compounds that are devoid of adverse cardiovascular properties. The design of **10a–f** was based on structure-activity data showing that (i) a COX-2 pharmacophore such as MeSO_2 or H_2NSO_2 at the *para*-position of a *N*¹-phenyl ring on a pyrazole ring template confers potent and selective COX-2 inhibitory activity, (ii) attachment of a substituted-aminomethylphenyl ring substituent via its C-4 sp^2 hybridized carbon atom is consistent with the observation that two aryl rings on adjacent positions of a five-membered heterocyclic ring template (scaffold) generally provide optimum COX-2 inhibitory activity, and (iii) the *N,N*-disubstituted-aminomethylphenyl secondary amino group that is present in compounds **12a–f**, provides a logical synthon for elaboration to the corresponding nitric oxide donor *N*-diazen-1-ium-1,2-diolates.²⁰ Unfortunately, reaction of **12b** and **12c** with nitric oxide afforded the corresponding *N*-nitroso products **13** and **14** rather than the desired unisolable intermediate *N*-diazen-1-ium-1,2-diolate product **10b** and **10c**, respectively (see Scheme 1).

The anti-inflammatory (AI) activities for a group of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1*H*-pyrazoles (**12a–f**) and *N*-nitroso derivatives (**13** and **14**) were determined using a carrageenan-induced rat foot paw edema model (see data in Table 1). Structure-activity relationships acquired for **12a–f** showed that the relative AI potency profile (i) with respect to the COX-2 pharmacophore moiety was $\text{SO}_2\text{Me} > \text{SO}_2\text{NH}_2$ in all derivatives ($\text{R}^1 = \text{NHMe}$, NHEt , piperazino) and at all time intervals (1, 3 and 5 h), (ii) within the SO_2Me group of compounds, the relative potency profile was $\text{NHMe} > \text{piperazino} > \text{NHEt}$ with respect to the R^1 substituent, (iii) within the SO_2NH_2 group of compounds, the three derivatives showed similar AI activity (54%, 53%, 58% at the 1 h interval) and (69%, 71%, 70% at the 3 h interval). In comparison at the 5 h time interval, the relative potency order was NHMe (57%) $>$ NHEt (39%) $>$ piperazino (35%) with respect to the R^1 substituent, and (iv) the most potent compound **12a** ($\text{R} = \text{Me}$, $\text{R}^1 = \text{NHMe}$, % AI inhibition = 72%, 89%, 67% at 1, 3, 5 h intervals, respectively) exhibited anti-inflammatory activities higher than that of the reference drug celecoxib (% AI inhibition = 40%, 80%, 48% at 1, 3, 5 h intervals, respectively). In contrast, the *N*-nitroso derivatives **13** and **14** showed poor activity at all time intervals.

In conclusion, a group of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1*H*-pyrazoles (**12a–f**) were synthesized²¹ and evaluated as anti-inflammatory agents²². All compounds showed significant anti-inflammatory activity after 3 h of inflammation induction (69–89%) as compared to celecoxib (80%). The *N,N*-disubstituted amino moiety present in compounds **12a–f** is not a suitable precursor to prepare putative hybrid nitric oxide donor *N*-diazen-1-ium-1,2-diolate derivatives of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1*H*-pyrazoles **10a–f** since the reaction of **12b** or **12c** with NO afforded the unstable *N*-diazen-1-ium-1,2-diolate products that undergo protonation

Table 1
Anti-inflammatory activities for the 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (**12a–f**), nitroso derivatives (**13** and **14**) and reference drug (celecoxib)



| Compound | R | R ¹ | % of anti-inflammatory activity (AI) | | |
|------------|-----------------|----------------------------------|--------------------------------------|-----|-----|
| | | | 1 h | 3 h | 5 h |
| 12a | Me | NHCH ₃ | 72 | 89 | 67 |
| 12b | Me | NHC ₂ H ₅ | 66 | 72 | 66 |
| 12c | Me | | 67 | 73 | 66 |
| 12d | NH ₂ | NHCH ₃ | 54 | 69 | 57 |
| 12e | NH ₂ | NHC ₂ H ₅ | 53 | 71 | 39 |
| 12f | NH ₂ | | 58 | 70 | 35 |
| 13 | Me | NNOC ₂ H ₅ | 24 | 39 | 25 |
| 14 | Me | | 19 | 43 | 29 |
| Celecoxib | — | — | 40 | 80 | 48 |

All test compounds (**12a–f**, celecoxib) were administrated (20 mg/kg) 30 min prior to testing. The anti-inflammatory activity % is expressed according to the following equation:

AI (%) = $(V_c - V_t/V_c) \times 100$ where V_c represents the paw volume of control group of animals and V_t represents the paw volume in drug treated animals.

and elimination of HNO to furnish the *N*-nitroso derivatives (**13** and **14**), respectively.

References and notes

- Zarghi, A.; Najafnia, L.; Daraee, B.; Dadrass, O. G.; Hedayati, M. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5634.
- Zebardast, T.; Zarghi, A.; Daraie, B.; Hedayati, M.; Dadrass, O. G. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 3162.
- Al-Hourani, B. J.; Sharma, S. K.; Mane, J. Y.; Tuszyński, J.; Baracos, V.; Kniess, T.; Suresh, M.; Pietzsch, J.; Wuest, F. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 1823.
- Rathish, I. G.; Javed, K.; Ahmad, S.; Bano, S.; Alam, M. S.; Pillai, K. K.; Singh, S.; Bagchi, V. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 255.
- Hinz, B.; Brune, K. *J. Pharmacol. Exp. Ther.* **2002**, *300*, 367.
- Patel, H. H.; Gross, G. J. *J. Mol. Cell Cardiol.* **2002**, *34*, 1.
- Mukherjee, D. *Biochem. Pharmacol.* **2002**, *63*, 817.
- Scheen, A. J. *Rev. Med. Liege* **2004**, *59*, 565.
- Dogné, J.-M.; Supuran, C. T.; Pratico, D. *J. Med. Chem.* **2005**, *48*, 2251.
- Butler, A. R.; Williams, D. L. *H. Chem. Soc. Rev.* **1993**, *22*, 233.
- Abdellatif, K. R. A.; Dong, Y.; Chen, Q.-H.; Chowdhury, M. A.; Knaus, E. E. *Bioorg. Med. Chem.* **2007**, *15*, 6796.
- Abdellatif, K. R. A.; Chowdhury, M. A.; Dong, Y.; Chen, Q.-H.; Knaus, E. E. *Bioorg. Med. Chem.* **2008**, *16*, 3302.
- Abdellatif, K. R. A.; Chowdhury, M. A.; Dong, Y.; Knaus, E. E. *Bioorg. Med. Chem.* **2008**, *16*, 6528.
- Abdellatif, K. R. A.; Chowdhury, M. A.; Dong, Y.; Das, D.; Yu, G.; Knaus, E. E. *Bioorg. Med. Chem.* **2009**, *17*, 5182.
- Abdellatif, K. R. A.; Huang, Z.; Chowdhury, M. A.; Kaufman, S.; Knaus, E. E. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 3951.
- Toscano, J. P.; Pavlos, C. M.; Boppana, K. Int. PCT Patent WO 2005/074598 A2, Issued August 18, 2005.
- Abdellatif, K. R. A.; Chowdhury, M. A.; Knaus, E. E. *J. Heterocycl. Chem.* **2008**, *45*, 1707.
- Moawad, A.; Hetta, M.; Zjawiony, J. K.; Jacob, M. R.; Hifnawy, M.; Ferreira, D. *Planta Med.* **2010**, *76*, 796.
- Deet, M. J.; Fahey, J. E.; Smith, B. D. *J. Phys. Org. Chem.* **2001**, *14*, 463.
- Abdellatif, K. R. A.; Chowdhury, M. A.; Dong, Y.; Das, D.; Yu, G.; Velázquez, C.; Suresh, M. R.; Knaus, E. E. *Bioorg. Med. Chem.* **2008**, *16*, 8882.
- Experimental procedures and spectral data for compounds **12a–f**, **13** and **14**. General. Melting points were determined on a Thomas-Hoover capillary

apparatus and are uncorrected. Infrared (IR) spectra were recorded as films on NaCl plates using a Nicolet 550 Series II Magna FT-IR spectrometer. ¹H NMR and ¹³C NMR spectra were measured on a Bruker Avance III 400 MHz Spectrophotometer, Faculty of Pharmacy, Beni-Suef University, Egypt in CDCl₃ or DMSO-*d*₆, where *J* (coupling constant) values are estimated in Hertz (Hz). Microanalyses were performed for C, H and N were carried out on Perkin–Elmer 2400 analyzer (Perkin–Elmer, Norwalk, CT, USA) at the Micro analytical unit of Cairo University, Egypt. All compounds were within ±0.4% of the theoretical values. Silica gel column chromatography was performed using Merck silica gel 60 ASTM (70–230 mesh). 1-(4-Methane(amino)sulfonylphenyl)-5-(4-bromomethylphenyl)-3-trifluoromethyl-1H-pyrazole (**11a** and **11b**) were prepared according to our previously reported procedure (Abdellatif, K. R. A.; Chowdhury, M. A.; Dong, Y.; Velázquez, C.; Das, D.; Suresh, M. R.; Knaus, E. E. *Bioorg. Med. Chem.* **2008**, *16*, 9694). All other reagents, purchased from the Aldrich Chemical Company (Milwaukee, WI), were used without further purification.

General procedure for the synthesis of 1-(4-methane(amino)sulfonylphenyl)-5-(4-substituted-aminomethylphenyl)-3-trifluoromethyl-1H-pyrazoles (12a–f): The corresponding amine (2.2 mmol) was added drop wise with stirring to a solution of 1-(4-methane(amino)sulfonylphenyl)-5-(4-bromomethylphenyl)-3-trifluoromethyl-1H-pyrazole (**11a**) or (**11b**) (1.0 mmol) in dry THF (10 mL). The reaction mixture was stirred at 25 °C for 1 h, and the solvent was removed in vacuo. Potassium carbonate (0.138 g, 1.0 mmol) in water (5 mL) was added to the residue prior to extraction with EtOAc (3 × 15 mL). The combined organic fractions were washed successively with water and brine, and the organic fraction was dried (MgSO₄). Filtration and removal of the solvent from the organic fraction in vacuo afforded the crude product which was purified by silica gel column chromatography using EtOAc/MeOH (3:2, v/v) as eluent to furnish the respective title compounds **12a–f**. Physical and spectroscopic data for **12a–f** are listed below. *N*-Methyl-1-(4-(1-(4-(methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)phenyl)methan-amine (**12a**): The product was obtained as a pale yellow powder in 68% yield; mp 112–114 °C; IR (film) 3340 (broad NH), 2975 (C–H aromatic), 2932 (C–H aliphatic), 1319, 1155 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 1.86 (s, 1H, NH, D₂O exchangeable), 2.49 (s, 3H, NCH₃), 3.08 (s, 3H, SO₂CH₃), 3.80 (s, 2H, CH₂), 6.78 (s, 1H, pyrazole H-4), 7.21 (d, *J* = 8.0 Hz, 2H, methylaminomethanophenyl H-3, H-5), 7.36 (d, *J* = 8.0 Hz, 2H, methylaminomethanophenyl H-2, H-6), 7.55 (d, *J* = 8.8 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.94 (d, *J* = 8.8 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 36.17, 44.50, 55.49, 106.69, 120.98 (q, *J* = 268 Hz, CF₃), 125.69, 127.17, 128.57, 128.83, 128.90, 139.86, 142.03, 143.41, 144.35 (q, *J* = 38 Hz, C–CF₃), 145.11; MS 410.1 (M+1). Anal. Calcd for C₁₉H₁₈F₃N₃O₂S·1/5H₂O: C, 55.25; H, 4.49; N, 10.17. Found: C, 55.59; H, 4.79; N, 9.79.

N-(4-(1-(4-(Methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)ethanamine (**12b**): The product was obtained as a pale yellow powder in 62% yield; mp 83–85 °C; IR (film) 3335 (broad NH), 2966 (C–H aromatic), 2932 (C–H aliphatic), 1323, 1157 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 1.15 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 1.96 (s, 1H, NH, D₂O exchangeable), 2.71 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 3.06 (s, 3H, SO₂CH₃), 3.83 (s, 2H, CH₂), 6.77 (s, 1H, pyrazole H-4), 7.19 (d, *J* = 8.1 Hz, 2H, ethylaminomethanephenyl H-3, H-5), 7.36 (d, *J* = 8.1 Hz, 2H, ethylaminomethanephenyl H-2, H-6), 7.53 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.93 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 15.27, 43.91, 44.50, 53.39, 106.69, 120.98 (q, *J* = 268 Hz, CF₃), 125.70, 127.06, 128.72, 128.77, 128.88, 139.84, 142.36, 143.41, 144.33 (q, *J* = 38 Hz, C–CF₃), 145.13; MS 424.1 (M+1). Anal. Calcd for C₂₀H₂₀F₃N₃O₂S·1/3H₂O: C, 55.94; H, 4.85; N, 9.78. Found: C, 55.95; H, 4.82; N, 9.41.

N-(4-(1-(4-(Methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)piperazine (**12c**): The product was obtained as a yellow powder in 71% yield; mp 102–104 °C; IR (film) 3323 (broad NH), 2969 (C–H aromatic), 2937 (C–H aliphatic), 1321, 1156 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 2.33 (s, 1H, NH, D₂O exchangeable), 2.50 (t, *J* = 5.0 Hz, 4H, piperazinyl H-3, H-5), 2.97 (t, *J* = 5.0 Hz, 4H, piperazinyl H-2, H-6), 3.07 (s, 3H, SO₂CH₃), 3.49 (s, 2H, CH₂), 6.70 (s, 1H, pyrazole H-4), 7.10 (d, *J* = 8.0 Hz, 2H, piperazinylmethanephenyl H-3, H-5), 7.28 (d, *J* = 8.0 Hz, 2H, piperazinylmethanephenyl H-2, H-6), 7.47 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.86 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 44.50, 45.96, 54.35, 63.00, 106.69, 120.96 (q, *J* = 268 Hz, CF₃), 125.70, 127.18, 128.56, 128.71, 129.66, 139.86, 140.26, 143.44, 144.34 (q, *J* = 38 Hz, C–CF₃), 145.13; MS 465.2 (M+1). Anal. Calcd for C₂₂H₂₃F₃N₄O₂S: C, 56.89; H, 4.99; F, 12.27; N, 12.06. Found: C, 56.62; H, 4.87; N, 12.31.

N-(4-(1-(4-(Methylamino)methyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)benzenesulfonamide (**12d**): The product was obtained as a pale yellow powder in 57% yield; mp 121–123 °C; IR (film) 3247 (broad NH), 2968 (C–H aromatic), 2923 (C–H aliphatic), 1338, 1165 (SO₂) cm⁻¹; ¹H NMR (CDCl₃+DMSO-*d*₆) δ 2.36 (s, 3H, NCH₃), 3.01 (s, 2H, SO₂NH₂, D₂O exchangeable), 3.68 (s, 2H, CH₂), 6.69 (s, 1H, pyrazole H-4), 7.11 (d, *J* = 8.0 Hz, 2H, methylaminomethanephenyl H-3, H-5), 7.25 (d, *J* = 8.0 Hz, 2H, methylaminomethanephenyl H-2, H-6), 7.34 (d, *J* = 8.5 Hz, 2H, sulfamoylphenyl H-2, H-6), 7.83 (d, *J* = 8.5 Hz, 2H, sulfamoylphenyl H-3, H-5); ¹³C NMR (DMSO-*d*₆) δ 37.23, 54.13, 106.47, 121.01 (q, *J* = 268 Hz, CF₃), 125.53, 127.36, 127.45, 129.01, 129.53, 140.31, 141.74, 142.28, 144.13 (q, *J* = 38 Hz, C–CF₃), 145.07; MS 411.1 (M+1). Anal. Calcd for C₁₈H₁₇F₃N₄O₂S: C, 52.68; H, 4.18; N, 13.65. Found: C, 52.45; H, 4.22; N, 13.68.

N-(4-(1-(4-(Ethylamino)methyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)benzenesulfonamide (**12e**): The product was obtained as a pale yellow powder in 59% yield; mp 170–172 °C; IR (film) 3341 (broad NH), 2962 (C–H aromatic), 2924 (C–H aliphatic), 1338, 1163 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 1.16 (t, *J* = 7.1 Hz, 3H, CH₂CH₃), 2.73 (q, *J* = 7.1 Hz, 2H, CH₂CH₃), 2.92 (s, 2H, SO₂NH₂, D₂O exchangeable), 3.83 (s, 2H, CH₂), 6.77 (s, 1H, pyrazole H-4), 7.19 (d, *J* = 8.1 Hz, 2H, ethylaminomethanephenyl H-3, H-5), 7.35 (d, *J* = 8.1 Hz, 2H, ethylaminomethanephenyl H-2, H-6), 7.47 (d, *J* = 8.7 Hz, 2H, sulfamoylphenyl H-2, H-6), 7.91 (d, *J* = 8.7 Hz, 2H, sulfamoylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 13.75, 41.93, 55.03, 106.47, 121.00 (q, *J* = 268 Hz, CF₃), 125.50, 127.46, 127.57, 129.01, 129.11, 140.61, 141.75, 142.25, 144.11 (q, *J* = 38 Hz, C–CF₃), 144.88; MS 425.1 (M+1). Anal. Calcd for C₁₉H₁₉F₃N₄O₂S: C, 53.77; H, 4.51; N, 13.20. Found: C, 53.94; H, 4.43; N, 13.18.

N-(4-(1-(4-(Piperazin-1-ylmethyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)benzenesulfonamide (**12f**): The product was obtained as a yellow powder in 64% yield; mp 106–108 °C; IR (film) 3331 (broad NH), 2965 (C–H aromatic), 2935 (C–H aliphatic), 1329, 1161 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 2.51 (t, *J* = 5.0 Hz, 4H, piperazinyl H-3, H-5), 2.82 (t, *J* = 5.0 Hz, 4H, piperazinyl H-2, H-6), 2.99 (s, 2H, SO₂NH₂, D₂O exchangeable), 3.44 (s, 2H, CH₂), 6.69 (s, 1H, pyrazole H-4), 7.10 (d, *J* = 8.0 Hz, 2H,

piperazinylmethanephenyl H-3, H-5), 7.26 (d, *J* = 8.0 Hz, 2H, piperazinylmethanephenyl H-2, H-6), 7.39 (d, *J* = 8.7 Hz, 2H, sulfamoylphenyl H-2, H-6), 7.83 (d, *J* = 8.7 Hz, 2H, sulfamoylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 45.78, 54.15, 62.94, 106.48, 121.08 (q, *J* = 268 Hz, CF₃), 125.51, 127.34, 127.45, 128.75, 129.69, 139.82, 141.74, 142.37, 144.11 (q, *J* = 38 Hz, C–CF₃), 145.04; MS 466.1 (M+1). Anal. Calcd for C₂₁H₂₂F₃N₅O₂S: C, 54.18; H, 4.76; N, 15.05. Found: C, 54.09; H, 4.82; N, 15.27.

N-Ethyl-*N*-(4-(1-(4-(methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)nitrosamine (**13**): The ethanamine compound **12b** (724 mg, 1.7 mmol) was added to a solution of NaOMe (210 mg of 95% purity, 1.7 mmol) in THF (30 mL) with stirring at –65 °C. This mixture was purged with argon for 5 min, and the reaction was allowed to proceed under an atmosphere of nitric oxide gas (40 psi internal pressure) with stirring at –65 °C for 1 h. The reaction mixture was allowed to warm to 25 °C, and the solvent was removed from the filtrate to furnish product **13** as a pale yellow powder in 55% yield. In a second reaction, nitric oxide gas was bubbled into a solution of **12b** using a protic solvent system consisting of NaOMe in MeOH (25% w/v) and MeCN at 25 °C for 24 h which afforded **13** in 90% yield as a pale yellow powder; mp 117–119 °C; IR (film) 2977 (C–H aromatic), 2926 (C–H aliphatic), 1317, 1157 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) of the major rotamer δ 0.89 (t, *J* = 7.2 Hz, 3H, CH₂CH₃), 3.00 (s, 3H, SO₂CH₃), 3.44 (q, *J* = 7.2 Hz, 2H, CH₂CH₃), 5.22 (s, 2H, CH₂), 6.73 (s, 1H, pyrazole H-4), 7.20 (d, *J* = 8.1 Hz, 2H, substituted-aminomethanephenyl H-3, H-5), 7.25 (d, *J* = 8.1 Hz, 2H, substituted-aminomethanephenyl H-2, H-6), 7.44 (d, *J* = 8.5 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.84 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹H NMR (CDCl₃) of the minor rotamer δ 1.31 (t, *J* = 7.3 Hz, 3H, CH₂CH₃), 3.00 (s, 3H, SO₂CH₃), 4.11 (q, *J* = 7.3 Hz, 2H, CH₂CH₃), 4.74 (s, 2H, CH₂), 6.69 (s, 1H, pyrazole H-4), 7.05 (d, *J* = 8.1 Hz, 2H, substituted-aminomethanephenyl H-3, H-5), 7.11 (d, *J* = 8.1 Hz, 2H, substituted-aminomethanephenyl H-2, H-6), 7.43 (d, *J* = 8.5 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.85 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹³C NMR (CDCl₃) of the major rotamer δ 10.01, 37.60, 43.37, 54.10, 105.95, 119.90 (q, *J* = 268 Hz, CF₃), 124.71, 127.60, 127.69, 127.78, 128.43, 135.83, 139.10, 142.12, 143.20 (q, *J* = 38 Hz, C–CF₃), 143.44; ¹³C NMR (CDCl₃) of the minor rotamer δ 12.99, 43.37, 44.60, 46.16, 105.85, 119.90 (q, *J* = 268 Hz, CF₃), 126.89, 127.48, 127.58, 128.21, 128.47, 135.06, 139.02, 142.16, 143.18 (q, *J* = 38 Hz, C–CF₃), 143.57; MS 453.1 (M+1). Anal. Calcd for C₂₀H₁₉F₃N₄O₃S: C, 53.09; H, 4.23; N, 12.38. Found: C, 53.43; H, 4.48; N, 12.73.

N-(4-(1-(4-(Methylsulfonyl)phenyl)-3-(trifluoromethyl)-1H-pyrazol-5-yl)benzyl)-4-nitrosopiperazine (**14**): The title compound **14** was prepared, using a similar procedure to that described for the preparation of **13**, by using **12c** in place of **12b**, in 60% and 93% yields, respectively, using the two solvent systems described previously as a yellow powder; mp 130–132 °C; IR (film) 2968 (C–H aromatic), 2935 (C–H aliphatic), 1321, 1159 (SO₂) cm⁻¹; ¹H NMR (CDCl₃) δ 2.44 (t, *J* = 5.0 Hz, 2H, piperazinyl H-2, H-6), 2.68 (t, *J* = 5.0 Hz, 2H, piperazinyl H-2, H-6), 3.03 (s, 3H, SO₂CH₃), 3.59 (s, 2H, CH₂), 3.75 (t, *J* = 5.0 Hz, 2H, piperazinyl H-3, H-5), 4.21 (t, *J* = 5.0 Hz, 2H, piperazinyl H-3, H-5), 6.72 (s, 1H, pyrazole H-4), 7.16 (d, *J* = 8.0 Hz, 2H, piperazinylmethanephenyl H-3, H-5), 7.33 (d, *J* = 8.0 Hz, 2H, piperazinylmethanephenyl H-2, H-6), 7.47 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-2, H-6), 7.85 (d, *J* = 8.7 Hz, 2H, methanesulfonylphenyl H-3, H-5); ¹³C NMR (CDCl₃) δ 36.17, 44.40, 48.34, 51.57, 53.00, 61.6, 106.77, 120.98 (q, *J* = 268 Hz, CF₃), 125.73, 127.80, 128.57, 128.99, 129.62, 139.97, 140.42, 143.31, 144.23 (q, *J* = 38 Hz, C–CF₃), 144.88; MS 494.2 (M+1). Anal. Calcd for C₂₂H₂₂F₃N₅O₃S: C, 53.54; H, 4.49; N, 14.19. Found: C, 53.69; H, 4.46; N, 14.47.

22. *In vivo anti-inflammatory assay*: The test compounds **12a–f**, **13**, **14** and the reference drugs celecoxib were evaluated using the *in vivo* carrageenan-induced rat paw edema model reported previously (Winter, C. A.; Risley, E. A.; Nuss, G. W. *Proc. Soc. Exp. Biol. Med.* **1962**, *111*, 544).