

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/232530632>

ChemInform Abstract: Synthesis and Antitrypanosomal Activity of New 6,6,7-Trisubstituted Thiopyrano[2,3-d][1,3]thiazoles.

ARTICLE in BIOORGANIC & MEDICINAL CHEMISTRY LETTERS · OCTOBER 2012

Impact Factor: 2.42 · DOI: 10.1016/j.bmcl.2012.09.091 · Source: PubMed

CITATIONS

14

READS

29

5 AUTHORS, INCLUDING:



Nataliya Zelisko

Danylo Halytsky Lviv National Medical Univ...

4 PUBLICATIONS 22 CITATIONS

SEE PROFILE



Dmytro Atamanyuk

Mutabilis S.A.

22 PUBLICATIONS 275 CITATIONS

SEE PROFILE



Philippe Grellier

Muséum National d'Histoire Naturelle

197 PUBLICATIONS 3,614 CITATIONS

SEE PROFILE




Roman Lesyk

Danylo Halytsky Lviv National Medical Univ...

99 PUBLICATIONS 1,004 CITATIONS

SEE PROFILE



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl



Synthesis and antitrypanosomal activity of new 6,6,7-trisubstituted thiopyrano[2,3-d][1,3]thiazoles

Nataliya Zelisko^a, Dmytro Atamanyuk^{a,1}, Olexandr Vasylenko^b, Philippe Grellier^c, Roman Lesyk^{a,*}

^a Department of Pharmaceutical, Organic and Bioorganic Chemistry, Danylo Halytsky Lviv National Medical University, 69 Pekarska, Lviv 79010, Ukraine

^b Institute of Bioorganic Chemistry and Petrochemistry, National Academy of Science of Ukraine, 1 Murmanska, Kyiv 02094, Ukraine

^c National Museum of Natural History, UMR 7245 CNRS-MNHN, Team APE, CP 52, 57 Rue Cuvier, Paris 75005, France

ARTICLE INFO

Article history:

Received 11 July 2012

Revised 21 September 2012

Accepted 25 September 2012

Available online 2 October 2012

Keywords:

Human African trypanosomiasis

Thiopyrano[2,3-d][1,3]thiazoles

Trypanosoma brucei

Antitrypanosomal activity

ABSTRACT

A series of novel 6,6,7-trisubstituted thiopyrano[2,3-d][1,3]thiazoles-based molecules have been synthesized and evaluated as potential antitrypanosomal agents. The most active analogue **3b** inhibited *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* with an IC₅₀ of 0.26 and 0.42 μM, respectively. They could be considered as potent hits for further antitrypanosomal drug discovery efforts.

© 2012 Elsevier Ltd. All rights reserved.

Human African trypanosomiasis (HAT, African sleeping sickness) is a fatal infectious disease caused by the protozoa parasite *Trypanosoma brucei* and transmitted by the tsetse fly. Sixty millions people are at risk of this infection, which in West Africa is caused by *Trypanosoma brucei gambiense* while in East Africa by *Trypanosoma brucei rhodesiense*. Cattle act a reservoir for HAT and can be infected by the subspecies *Trypanosoma brucei brucei*, *Trypanosoma congolense* and *Trypanosoma evansi*, which have high economical impacts on local populations. Drugs therapy is a cornerstone of trypanosomicidal treatment. For the early stages of the disease suramin and the diamidine pentamidine are commonly used. Arsenical melarsoprol and eflornithine are reserved for the late stages of disease when the central nervous system is affected. Except for eflornithine, which has better safety profile, the other drugs cause many side effects and can induce severe toxicity reaction. Importantly, eflornithine is not effective in *T. b. rhodesiense* associated disease. In addition to toxicity issues, the problems of drug resistance have been reported for pentamidine, melarsoprol and eflornithine.¹ Thus, there is an urgent need for the development of the new drugs against this devastating illness.

It is well known that 4-thiazolidinones and related heterocyclic systems are important scaffolds in drug discovery. Their derivatives were demonstrated with antiinflammatory, antitumor, antimicrobial, antidiabetic and antibacterial actions.² Recently,

thiazolidinones derivatives have been identified as perspective agents in antitrypanosomal treatment, in particular among 2-hydrazono-4-thiazolidinones.^{3,4} Moreover 5-benzylidene-2-thioxo-4-thiazolidinone-3-acetic acids were reported as first small molecule inhibitors of *T. brucei* dolicholphosphate mannose synthase (DPMS), a validated drug target in African sleeping sickness.⁵ However, 5-arylidene-4-thiazolidinones can be considered as electrophilic and potentially reactive substances due to possible Michael addition of the nucleophilic protein residues to the exocyclic double bond. This property characterizes 5-arylidene-4-thiazolidinones as the frequent hitters or pan assay interference compounds (PAINS) that are useless in the modern drug discovery process because of their low selectivity.⁶ Our previous research demonstrated a number of arguments in favor of 7-arylthiopyrano[2,3-d][1,3]thiazoles design which could be one of approaches to address the selectivity issue.^{7,8} Thiopyrano[2,3-d][1,3]thiazoles can be considered as cyclic isosteric mimetics of their synthetic precursors 5-arylidene-4-thiazolidinones without Michael accepting functionalities (Fig. 1). The aim of present letter was to explore the antiparasitic activity of new thiopyrano[2,3-d][1,3]thiazoles against *Trypanosoma brucei brucei* (Tbb) and *Trypanosoma brucei gambiense* (Tbg).

The general synthetic pathways yielding target thiopyrano[2,3-d][1,3]thiazoles are shown in Schemes 1 and 2. Itaconic acid and its imides were studied as dienophiles in hetero-Diels–Alder reaction with 5-arylidene-4-thioxo-2-thiazolidinones **1a–g** (5-arylideneisorhodanines)⁹ as heterodienes. Diene synthesis reactions were performed in glacial acetic acid medium with catalytic amount of hydroquinone to prevent polymerization processes.^{10,11} As a

* Corresponding author. Tel.: +38 0322 75 59 66; fax: +38 0322 75 77 34.

E-mail address: dr_r_lesyk@org.lviv.net (R. Lesyk).

¹ Present address: Mutabilis, 102 Avenue Gaston Roussel, Romainville 93230, France.





Figure 1. Background for target compounds synthesis.



Scheme 1. Synthesis of *rel*-(6*R*,7*R*)-7-aryl-6-carboxymethylene-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acids (**2a-d**) and *rel*-(6'*R*,7'*R*)-3',7'-dihydro-2*H*,2'*H*,5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5'-triones (**3a-d**). Reagents and conditions: **1** (1.0 equiv), appropriate dienophile (1.2 equiv), hydroquinone, AcOH, reflux, 2 h, 44–73%.

result the novel *rel*-(6*R*,7*R*)-7-aryl-6-carboxymethyl-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acids (**2a-d**) and *rel*-(6'*R*,7'*R*)-3',7'-dihydro-2*H*,2'*H*,5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5'-triones (**3a-g**, **5a-c**) were synthesized. Variant of tandem acylation-*hetero*-Diels-Alder reaction was observed employing 5-(2-hydroxyphenylmethylidene)isothiazolidine **1g** as heterodienophile 'building block' with itaconic acid, yielding *rel*-2-[(5*aR*,11*bR*)-2,6-dioxo-3,5*a*,6,11*b*-tetrahydro-2*H*,5*H*-cromeno[4',3':4,5]thiopyrano[2,3-*d*]thiazol-5*a*-yl]acetic acid **4**.¹¹ Based on this heterocyclic acid we proposed a counter synthesis method of spiroderivatives **5a-c**.^{12,13} Thus, **4** undergoes recyclization in the presence of aromatic amines at 100 °C in acetic acid yielding abovementioned thiopyrano[2,3-*d*]thiazole derivatives **5a-c**. Study of ESI-MS, homonuclear ¹H and ¹³C NMR spectra, COSY and NOESY (1D with CH-Ar proton) experiments as well as heteronuclear HSQC and HMBC experiments confirmed the structures of synthesized compounds and the relative stereochemistry of [2 + 4]-cycloaddition and the consecutively stereoconfiguration of substitutions at the positions 6 and 7, in particular *trans* positioning of 6-carboxymethylene substituent relatively to 7-aryl moiety. For compound **2c** NOE coupling of CH proton at 3.71 ppm with the protons at 6.94 ppm (12%), 2.94 ppm (8%) and 2.84 ppm (4%) is in the accordance with shown structure, which is present as a racemic mixture. Moreover, the conclusion about stereochemistry of the mentioned itaconic acid based *hetero*-Diels-Alder reactions is consistent with our previous data of X-ray analysis of related 2-[7-(3,5-dibromo-2-

Scheme 2. Synthesis of *rel*-2-[(5*aR*,11*bR*)-2,6-dioxo-3,5*a*,6,11*b*-tetrahydro-2*H*,5*H*-cromeno[4',3':4,5]thiopyrano[2,3-*d*]thiazol-5*a*-yl]acetic acid (**4**) and *rel*-(6'*R*,7'*R*)-7'-(2-hydroxyphenyl)-1-aryl-3',7'-dihydro-2*H*,2'*H*,5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5'-triones (**5a-c**). Reagents and conditions: (a) **1g** (1.0 equiv), itaconic acid (1.2 equiv), hydroquinone, AcOH, reflux, 2 h, 45%; (b) **1g** (1.0 equiv), appropriate itaconimide (1.2 equiv), hydroquinone, AcOH, reflux, 2 h, 42–45%; (c) **4** (1 equiv), appropriate amine (1.3 equiv), AcOH, reflux, 10 h, 32–40%.

hydroxyphenyl)-6-ethoxycarbonyl-2-oxo-5*H*-2,3,6,7-tetrahydrothiopyrano[2,3-*d*][1,3]thiazol-6-yl]acetic acid.¹⁴

Compounds **2a-d**, **3a-d** and **5a-c** were studied for their antitrypanosomal activity against *T. brucei* and the results are summarized in **Tables 1** and **2**. Bloodstream forms of *Tbb* strain 90-13 and *Tbg* strain Feo were cultured in HMI9 medium supplemented with 10% FCS at 37 °C under an atmosphere of 5% CO₂.¹⁵ In all experiments, log-phase cell cultures were harvested by centrifugation at 3000×g and immediately used. Drug assays were based on the conversion of a redox-sensitive dye (resazurin) to a fluorescent product by viable cells as described previously.¹⁶ Compounds were first tested at concentrations of 10 and 1 µg/mL (**Table 1**) and IC₅₀s were determined¹⁷ for the most active ones (**Table 2**).

Some synthesized thiopyrano[2,3-*d*]thiazole spiroimides exhibited significant inhibitory activity against *T. brucei* as exemplified by **3b**, **3c** and **3d**. Among them the *rel*-(6'*R*,7'*R*)-7'-(3,4-dimethoxyphenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2*H*,2'*H*,5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5'-trione **3b** exhibited the most potent activity with an IC₅₀ of 0.26 and 0.42 µM against *Tbb* and *Tbg*, respectively. In general, these compounds were slightly more active on *Tbg* than on *Tbb*.

The maleimide fragment in spiroimide active hits (**3b-d**) is perpendicularly distorted due to spiro-attachment to the main

Table 1Thiopyrano[2,3-d]thiazole derivatives (**2a–d**, **3a–d**, **5a–c**), % inhibition of *Tbb* and *Tbg* bloodstream forms at 10 and 1 µg/mL concentrations. nd: not determined

Compound	R ¹	R ²	R ³	<i>Tbb</i> % inhibition		<i>Tbg</i> % inhibition	
				10 µg/mL	1 µg/mL	10 µg/mL	1 µg/mL
2a	BnO	MeO	—	<5	18,1	nd	nd
2b	F ₂ CHO	MeO	—	<5	11,3	nd	nd
2c	Me ₂ N	H	—	<5	5,3	nd	nd
2d	Et ₂ N	H	—	5,6	<5	nd	nd
3a	MeO	H	—	13,0	10,6	9,8	2,3
3b	MeO	MeO	—	93,6	82,4	nd	nd
3c	Et ₂ N	H	—	98,2	21,1	nd	nd
3d	Me ₂ N	H	—	98,2	36,8	nd	nd
5a	—	—	4-Cl	11,5	13,4	19,3	0,9
5b	—	—	4-OH	18,5	7,0	15,4	<5
5c	—	—	3-Me	24,5	5,5	15,3	<5

Table 2IC₅₀ of hit-compounds **3b–d** against *Tbb* and *Tbg*

Compound	<i>Tbb</i>		<i>Tbg</i>	
	IC ₅₀ (µM)	SD	IC ₅₀ (µM)	SD
3b	0,2624	0,0139	0,4234	0,1590
3c	2,3985	0,4123	3,6181	0,9259
3d	1,2751	0,3230	1,4711	0,0884
Pentamidine	0,0032	0,0004	0,0052	0,0009

scaffold. This may partially improve solubility and penetration of studied compounds with highly lipophylic motif, taking into account the entropic effect of desorganisation of crystalline lattice caused by deplanarization of the molecules.

One should note that the presence of two negative charges of dicarboxylic acids **2a–c** is not beneficial for the trypanosomal activity in the screened series of compounds which potentially can undergo deprotonation under physiologic conditions since penetration of the charged species through the lipophylic membrane of parasite is probably not favoured in this case. Spiroimide **3b** containing methoxy-substituted phenyl group possess prospective level of activity and can be used as a hit for further exploration of the SAR in these series.

For the first time 6,6-spiromaleimido-7-aryltrisubstituted thiopyrano[2,3-d][1,3]thiazoles were shown with activity against *Trypanosoma brucei brucei* and *Trypanosoma brucei gambiense* and, therefore, could be considered as potent hits for further antitrypanosomal drug discovery efforts.

Acknowledgment

Authors thank to Embassy of France in Ukraine and EGIDE for fellowship for Nataliya Zelisko.

References and notes

- Jacobs, R. T.; Nare, B.; Phillips, M. A. *Curr. Top. Med. Chem.* **2011**, *11*, 1255.
- Lesyk, R. B.; Zimenkovsky, B. S. *Curr. Org. Chem.* **2004**, *8*, 1547.
- Donnici, C. L.; Araújo, M. H.; Oliveira, H. S.; Moreira, D. R. M.; Pereira, V. R. A.; de Assis Souza, M.; de Castro, M. C. A. B.; Leite, A. C. L. *Bioorg. Med. Chem.* **2009**, *17*, 5038.
- Tenório, R. P.; Carvalho, C. S.; Pessanha, C. S.; de Lima, J. G.; de Faria, A. R.; Alves, A. J.; de Melo, E. J. T.; Góes, A. J. S. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2575.
- Smith, T. K.; Young, B. L.; Denton, H.; Hughes, D. L.; Wagner, G. K. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 1749.
- (a) Tomasic, T.; Masic, L. P. *Expert Opin. Drug Discov.* **2012**, *7*, 549; (b) Baell, J. B. *Future Med. Chem.* **2010**, *2*, 1529; (c) Baell, J. B.; Holloway, G. A. *J. Med. Chem.* **2010**, *53*, 2719; (d) Mendgen, T.; Steuer, C.; Klein, C. D. *J. Med. Chem.* **2012**, *55*, 743.
- Lesyk, R.; Zimenkovsky, B.; Atamanyuk, D.; Jensen, F.; Kiec-Kononowicz, K.; Gzella, A. *Bioorg. Med. Chem.* **2006**, *14*, 5230.
- Atamanyuk, D.; Zimenkovsky, B.; Lesyk, R. *J. Sulf. Chem.* **2008**, *29*, 151.

- (a) Komaritsa, I. D.; Baranov, S. N.; Grishchuk, A. P. *Khimiya Geterotsiklicheskikh Soedinenii* **1967**, *4*, 664; (b) *Chem. Abstr.* **1968**, *68*, 49495; (c) Grishchuk, A. P.; Komaritsa, I. D.; Baranov, C. N. U.S.S.R. Patent SU 197594, 1967 (in Russian); (d) *Chem. Abstr.* **1968**, *68*, 36112.
- Preparation of *rel*-(6*R*,7*R*)-7-aryl-6-carboxymethylene-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acids (**2a–d**), *rel*-(6*R*,7*R*)-3',7'-dihydro-2*H*,2',5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-triones (**3a–g**; **5a–c**, method A) and *rel*-2-[(5*aR*,11*bR*)-2,6-dioxo-3,5*a*,6,11*b*-tetrahydro-2*H*,5*H*-chromeno[4',3':4,5]thiopyrano[2,3-*d*]thiazol-5*a*-yl]acetic acid (**5**). A mixture of 10 mmol of appropriate 5-arylidene-4-thioxo-2-thiazolidinone and 11 mmol of itaconic acid or its appropriate imide was refluxed for 2 h with catalytic amount of hydroquinone (2–3 mg) for preventing of polymerization processes in 10 mL of glacial acetic acid, then left overnight at room temperature. The precipitated crystals were filtered off, washed with ethanol, and recrystallized from appropriate solvent.
- Spectral and analytical data for compounds (**2,3,4**) are as follows. *rel*-(6*R*,7*R*)-7-(4-Benzoyloxy-3-methoxyphenyl)-6-carboxymethylene-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acid (**2a**). Yield 69%, mp 186–189 °C (EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.82 (d, 1H, 5-H, *J* = 17.6 Hz), 2.96 (d, 1H, 5-H, *J* = 17.6 Hz), 3.19 (d, 1H, CH₂CO, *J* = 13.6 Hz), 3.35 (d, 1H, CH₂CO, *J* = 13.7 Hz), 3.72 (s, 3H, OCH₃), 3.81 (s, 1H, 7-H), 5.02 (s, 2H, OCH₂), 6.59 (d, 1H, arom., *J* = 8.0 Hz), 6.73 (s, 1H, arom.), 6.99 (d, 1H, arom., *J* = 8.4 Hz), 7.32–7.50 (m, 5H, arom.), 11.48 (s, 1H, NH), 12.62 (br.s, 2H, COOH). Anal. Calcd for C₂₃H₂₁NO₇S₂: C, 56.66; H, 4.34; N, 2.87. Found: C, 56.60; H, 4.20; N, 2.70. *rel*-(6*R*,7*R*)-7-(4-Difluoromethoxy-3-methoxyphenyl)-6-carboxymethyl-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acid (**2b**). Yield 50%, mp 210–212 °C (EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.84 (d, 1H, 5-H, *J* = 17.6 Hz), 2.97 (d, 1H, 5-H, *J* = 17.6 Hz), 3.21 (d, 1H, CH₂CO, *J* = 14.0 Hz), 3.34 (d, 1H, CH₂CO, *J* = 14.0 Hz), 3.79 (s, 3H, OCH₃), 3.79 (s, 1H, 7-H), 4.37 (s, 1H, OCHF₂), 6.68 (d, 1H, arom., *J* = 8.4 Hz), 6.88 (s, 1H, arom.), 7.13 (d, 1H, arom., *J* = 8.4 Hz), 11.52 (s, 1H, NH), 12.70 (br.s, 2H, COOH). ¹³C NMR (100 MHz, DMSO-*d*₆): 26.0, 26.5, 38.9, 46.8, 46.9, 56.2, 62.5, 104.3, 114.6, 117.0, 120.3, 121.1, 121.4, 138.5, 139.3, 150.4, 171.2, 172.5, 173.5. ESI-MS *m/z* 448 (M+H)⁺. Anal. Calcd for C₁₇H₁₅F₂NO₇S₂: C, 45.64; H, 3.38; N, 3.13. Found: C, 45.50; H, 3.20; N, 3.20. *rel*-(6*R*,7*R*)-7-(4-Dimethylaminophenyl)-6-carboxymethylene-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acid (**2c**). Yield 61%, mp 243–245 °C (MeOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.84 (d, 1H, 5-H, *J* = 17.7 Hz), 2.87 (s, 6H, 2*CH₃), 2.95 (d, 1H, 5-H, *J* = 17.5 Hz), 3.20 (d, 1H, CH₂CO, *J* = 13.7 Hz), 3.35 (d, 1H, CH₂CO, *J* = 13.7 Hz), 3.70 (s, 1H, 7-H), 6.64 (d, 2H, arom., *J* = 8.5 Hz), 6.92 (d, 2H, arom., *J* = 8.5 Hz), 11.44 (s, 1H, NH), 12.50 (br.s, 2H, COOH). ¹³C NMR (100 MHz, DMSO-*d*₆): 26.4, 39.0, 40.5, 46.6, 47.0, 105.6, 112.4, 119.4, 127.2, 129.9, 150.1, 171.3, 172.7, 173.6. Anal. Calcd for C₁₇H₁₈N₂O₅S₂: C, 51.76; H, 4.60; N, 7.10. Found: C, 51.60; H, 4.50; N, 7.20. *rel*-(6*R*,7*R*)-7-(4-Diethylaminophenyl)-6-carboxymethyl-2-oxo-3,5,6,7-tetrahydro-2*H*-thiopyrano[2,3-*d*][1,3]thiazole-6-carboxylic acid (**2d**). Yield 73%, mp 224–227 °C (DMF-EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 1.06 (t, 6H, 2*CH₃CH₂), 2.83 (d, 1H, 5-H, *J* = 17.4 Hz), 2.94 (d, 1H, 5-H, *J* = 17.5 Hz), 3.16 (d, 1H, CH₂CO, *J* = 13.7 Hz), 3.29 (q, 4H, 2*CH₃CH₂), 3.35 (d, 1H, CH₂CO, *J* = 13.7 Hz), 3.67 (s, 1H, 7-H), 6.56 (d, 2H, arom., *J* = 8.5 Hz), 6.89 (d, 2H, arom., *J* = 8.5 Hz), 11.42 (s, 1H, NH), 12.52 (br.s, 2H, COOH). ¹³C NMR (100 MHz, DMSO-*d*₆): 13.0, 26.5, 39.0, 44.0, 46.6, 47.0, 105.6, 111.4, 119.3, 126.0, 130.1, 147.3, 171.3, 172.7, 173.6. ESI-MS *m/z* 423 (M+H)⁺. Anal. Calcd for C₁₉H₂₂N₂O₅S₂: C, 54.01; H, 5.25; N, 6.63. Found: C, 54.20; H, 5.10; N, 6.70. *rel*-(6*R*,7*R*)-7'-(4-Methoxyphenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2*H*,2',5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**3a**). Yield 55%, mp 240–242 °C (DMF-EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 3.06 (d, 1H, 5'-H, *J* = 16.4 Hz), 3.10 (d, 1H, 5'-H, *J* = 16.4 Hz), 3.22 (d, 1H, 4-H, *J* = 13.7 Hz), 3.37 (d, 1H, 4-H, *J* = 13.7 Hz), 3.73 (s, 3H, OCH₃), 3.82 (s, 1H, 7'-H), 6.88 (d, 2H, arom., *J* = 8.6 Hz), 7.06 (d, 2H, arom., *J* = 8.6 Hz), 7.33 (d, 2H, arom., *J* = 8.9 Hz), 7.57 (d, 2H, arom., *J* = 8.9 Hz), 11.50 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 26.5, 41.1, 46.6, 47.1, 55.5, 105.0, 114.1, 120.0, 120.9, 127.1, 129.1, 130.5, 131.9, 138.4, 159.1, 169.3, 171.2, 173.7. ESI-MS *m/z* 472 and 474 (M+H)⁺. Anal. Calcd for C₂₂H₁₇ClN₂O₄S₂: C, 55.87; H, 3.62; N, 5.92. Found: C, 56.00; H, 3.50; N, 5.80. *rel*-(6*R*,7*R*)-7'-(3,4-Dimethoxyphenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2*H*,2',5*H*-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**3b**). Yield

- 44%, mp 258–260 °C (DMF-EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.96 (d, 1H, 5'-H, *J* = 18.1 Hz), 3.03 (d, 1H, 5'-H, *J* = 18.1 Hz), 3.50 (d, 1H, 4-H, *J* = 12.8 Hz), 3.63 (s, 3H, OCH₃), 3.70 (d, 1H, 4-H, *J* = 12.8 Hz), 3.74 (s, 3H, OCH₃), 4.35 (s, 1H, 7'-H), 6.74 (br.s, 1H, arom.), 6.80 (m, 3H, arom.), 6.93 (d, 1H, arom., *J* = 8.5 Hz), 7.52 (d, 2H, arom., *J* = 8.5 Hz), 11.56 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 33.8, 36.0, 47.0, 48.2, 56.1, 56.2, 106.4, 112.1, 120.9, 128.3, 128.7, 129.4, 130.9, 133.5, 149.0, 149.5, 170.9, 173.9, 178.4. ESI-MS *m/z* 503 and 505 (M+H)⁺. Anal. Calcd for C₂₃H₁₉ClN₂O₅S₂: C, 54.92; H, 3.81; N, 5.57. Found: C, 55.00; H, 3.70; N, 5.70. *rel*-(6'*R*,7'*R*)-7'-(4-Diethylaminophenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**3c**). Yield 52%, mp 242–244 °C (DMF-EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 1.04 (t, 6H, 2°CH₃), 2.91 (d, 1H, 5'-H, *J* = 18.4 Hz), 2.96 (d, 1H, 5'-H, *J* = 18.4 Hz), 3.30 (q, 4H, 2°CH₂), 3.44 (d, 1H, 4-H, *J* = 12.6 Hz), 3.66 (d, 1H, 4-H, *J* = 12.6 Hz), 4.22 (s, 1H, 7'-H), 6.55 (d, 2H, arom., *J* = 8.2 Hz), 6.82 (d, 2H, arom., *J* = 8.4 Hz), 6.96 (br.s, 2H, arom.), 7.44 (d, 2H, arom., *J* = 8.4 Hz), 11.49 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 12.8, 33.6, 35.8, 41.0, 44.1, 46.7, 48.2, 107.1, 111.7, 120.5, 121.4, 128.9, 129.2, 131.1, 133.4, 147.9, 170.9, 173.9, 178.5. ESI-MS *m/z* 514 and 516 (M+H)⁺. Anal. Calcd for C₂₅H₂₄ClN₃O₅S₂: C, 58.41; H, 4.71; N, 8.17. Found: C, 58.20; H, 4.70; N, 8.10. *rel*-(6'*R*,7'*R*)-7'-(4-Dimethylaminophenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**3d**). Yield 54%, mp 240–242 °C (DMF-EtOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.86 (s, 6H, 2°CH₃), 2.92 (d, 1H, 5'-H, *J* = 18.5 Hz), 2.97 (d, 1H, 5'-H, *J* = 18.5 Hz), 3.46 (d, 1H, 4-H, *J* = 12.8 Hz), 3.65 (d, 1H, 4-H, *J* = 12.8 Hz), 4.26 (s, 1H, 7'-H), 6.62 (d, 2H, arom., *J* = 8.0 Hz), 6.85 (d, 2H, arom., *J* = 8.4 Hz), 7.01 (br.s, 2H, arom.), 7.49 (d, 2H, arom., *J* = 8.4 Hz), 11.51 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 33.8, 35.9, 40.6, 46.7, 48.2, 107.1, 112.4, 120.7, 122.8, 128.9, 129.3, 131.0, 133.5, 150.8, 170.9, 173.8, 178.5. ESI-MS *m/z* 486 and 488 (M+H)⁺. Anal. Calcd for C₂₃H₂₀ClN₃O₅S₂: C, 56.84; H, 4.15; N, 8.65. Found: C, 56.70; H, 4.30; N, 8.40. *rel*-2-[(5*aR*,11*bR*)-2,6-Dioxo-3,5*a*,6,11*b*-tetrahydro-2H,5H-cromeno[4',3':4,5]thiopyrano[2,3-*d*]thiazol-5*a*-yl]acetic acid (**4**). Yield 45%, mp 240–242 °C (AcOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.60 (d, 1H, 5-H, *J* = 15.5 Hz), 2.72 (d, 1H, 5-H, *J* = 15.5 Hz), 3.40 (d, 1H, CH₂CO, *J* = 13.2 Hz), 3.46 (d, 1H, CH₂CO, *J* = 13.2 Hz), 4.41 (s, 1H, 11*b*-H), 7.16 (d, 1H, arom., *J* = 8.0 Hz), 7.26 (t, 1H, arom., *J* = 7.4 Hz), 7.36–7.44 (m, 2H, arom.), 11.53 (s, 1H, NH), 12.62 (br.s, 1H, COOH). ESI-MS *m/z* 350 (M+H)⁺. Anal. Calcd for C₁₅H₁₁NO₅S₂: C, 51.57; H, 3.17; N, 4.01. Found: C, 51.70; H, 3.30; N, 3.90.
12. Preparation of *rel*-(6'*R*,7'*R*)-7'-(2-hydroxyphenyl)-1-aryl-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-triones (**5a–c**, method **B**). A mixture of 3 mmol of **4** and 4 mmol of appropriate aromatic amine was refluxed for 10 h in 10 mL of glacial acetic acid, and then left overnight at room temperature. The precipitated crystals were filtered off, washed with ethanol, and recrystallized from appropriate solvent.
13. Spectral and analytical data for compounds (**5**) are as follows. *rel*-(6'*R*,7'*R*)-7'-(2-Hydroxyphenyl)-1-(4-chlorophenyl)-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**5a**). Yields 45% (method **A**), 32% (method **B**), mp 245–247 °C (MeCN). ¹H NMR (400 MHz, DMSO-*d*₆): 2.72 (d, 1H, 5'-H, *J* = 17.9 Hz), 3.14 (d, 1H, CH₂CO, *J* = 13.1 Hz), 3.24 (d, 1H, 5'-H, *J* = 17.9 Hz), 3.36 (d, 1H, CH₂CO, *J* = 13.1 Hz), 4.33 (s, 1H, 7'-H), 6.77 (m, 2H, arom.), 6.94 (d, 1H, arom., *J* = 7.3 Hz), 7.05 (t, 1H, arom., *J* = 7.7 Hz), 7.32 (d, 2H, arom., *J* = 8.7 Hz), 7.49 (d, 2H, arom., *J* = 8.7 Hz), 9.66 (s, 1H, OH), 11.44 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 28.3, 40.7, 40.9, 46.4, 104.8, 115.1, 119.5, 120.7, 125.4, 129.1, 129.2, 129.8, 131.7, 133.2, 155.7, 171.3, 173.8, 174.2, 176.3. ESI-MS *m/z* 459 and 461 (M+H)⁺. Anal. Calcd for C₂₁H₁₅ClN₂O₄S₂: C, 54.96; H, 3.29; N, 6.10. Found: C, 55.00; H, 3.40; N, 6.20. *rel*-(6'*R*,7'*R*)-7'-(2-Hydroxyphenyl)-1-(4-hydroxyphenyl)-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**5b**). Yields 42% (method **A**), 40% (method **B**), mp 247–250 °C (DMF-AcOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.68 (d, 1H, 5'-H, *J* = 17.9 Hz), 3.10 (d, 1H, 4-H, *J* = 13.1 Hz), 3.20 (d, 1H, 5'-H, *J* = 17.9 Hz), 3.37 (d, 1H, 4-H, *J* = 13.1 Hz), 4.30 (s, 1H, 7'-H), 6.76–6.81 (m, 4H, arom.), 6.94 (d, 1H, arom., *J* = 7.3 Hz), 7.02–7.07 (m, 4H, arom.), 9.54 (s, 1H, OH), 9.59 (s, 1H, OH), 11.40 (br.s, 1H, NH). Anal. Calcd for C₂₁H₁₆N₂O₅S₂: C, 57.26; H, 3.66; N, 6.36. Found: C, 57.40; H, 3.40; N, 6.30. *rel*-(6'*R*,7'*R*)-7'-(2-Hydroxyphenyl)-1-(3-methylphenyl)-3',7'-dihydro-2H,2'H,5H-spiro[pyrrolidin-3,6'-thiopyrano[2,3-*d*]thiazol]-2,2',5-trione (**5c**). Yields 45% (method **A**), 40% (method **B**), mp 247–250 °C (DMF-AcOH). ¹H NMR (400 MHz, DMSO-*d*₆): 2.33 (s, 3H, CH₃), 2.78 (d, 1H, 5'-H, *J* = 18.0 Hz), 3.21 (d, 1H, 5'-H, *J* = 18.0 Hz), 3.23 (d, 1H, 4-H, *J* = 13.0 Hz), 3.38 (d, 1H, 4-H, *J* = 13.0 Hz), 4.37 (s, 1H, 7'-H), 6.73–6.81 (m, 2H, arom.), 6.92 (d, 1H, arom., *J* = 7.5 Hz), 7.05–7.12 (m, 3H, arom.), 7.20 (d, 1H, arom., *J* = 7.2 Hz), 7.34 (t, 1H, arom., *J* = 7.8 Hz), 9.75 (s, 1H, OH), 11.59 (br.s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆): 21.3, 28.5, 40.7, 41.0, 46.3, 104.9, 115.1, 119.4, 120.6, 124.6, 125.5, 127.9, 129.0, 129.3, 129.8, 132.9, 138.7, 155.8, 171.2, 172.4, 174.4, 176.5. ESI-MS *m/z* 439 (M+H)⁺. Anal. Calcd for C₂₂H₁₈N₂O₄S₂: C, 60.26; H, 4.14; N, 6.39. Found: C, 60.00; H, 4.20; N, 6.20.
14. Kowiel, M.; Zelisko, N.; Atamanyuk, D.; Lesyk, R.; Gzella, A. K. *Acta Cryst.* **2012**, *E68*, o2721.
15. Bastos, I.; Motta, F. N.; Charneau, S.; Santana, J. M.; Dubost, L.; Augustyns, K.; Grellier, P. *Microb. Infect.* **2010**, *12*, 457.
16. Lethu, S.; Bosc, D.; Mouray E.; Grellier, P.; Dubois, J. J. *Enzyme Inhib. Med. Chem.* **2012**, in press, <http://dx.doi.org/10.3109/14756366.2011.643302>.
17. Drug stock solutions were prepared in pure DMSO. *T. brucei* bloodstream forms (10⁵ cells/mL) were cultured in 96-well plates either in the absence or in the presence of different concentrations of inhibitors in a final volume of 200 μL. After a 72-h incubation, resazurin solution was added in each well at the final concentration of 45 μM. Fluorescence was measured at 530 nm excitation and 590 nm emission wavelengths after a further 4-h incubation. The percentage of inhibition of parasite growth rate was calculated by comparing the fluorescence of parasites maintained in the presence of drug to that of in the absence of drug. DMSO was used as control. Drugs were first tested at the concentrations of 1 and 10 μg/mL in triplicate and the experiment repeated twice. For the most active ones, IC₅₀s were determined from the dose-response curves with drug concentrations ranging from 10 μg/mL to 5 ng/mL and presented in μM. IC₅₀ value is the mean ± the standard deviation of three independent experiments.