

Chapter 6

First article


Novel sulfanylphthalimide analogues as highly potent inhibitors of monoamine oxidase B

Mietha M. Van der Walt,^a Gisella Terre'Blanche,^a Anél Petzer,^b and Jacobus P. Petzer^a

a Pharmaceutical Chemistry, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa

b Unit for Drug Research and Development, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa

6.1 Graphical abstract



Published Manuscript: Bioorganic & Medicinal Chemistry Letters

Received Date: 12 July 2012

Revised Date: 16 August 2012

Accepted Date: 28 August 2012

Available online: 7 September 2012

DOI: <http://dx.doi.org/10.1016/j.bmcl.2012.08.113>

Reference: BMCL 19499

6.2 Author's contributions

Novel sulfanylpthalimide analogues as highly potent inhibitors of monoamine oxidase B

Mietha M. Van der Walt,^a Gisella Terre'Blanche,^a Anél Petzer,^b and Jacobus P. Petzer^a

a Pharmaceutical Chemistry, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa

b Unit for Drug Research and Development, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa

The above mentioned article is part of the doctoral thesis of M.M. van der Walt. As required, the table below indicates the contribution of each author to this research article. Also, the written declaration for consent by each author is provided. This article was submitted for publication in Bioorganic & Medicinal Chemistry Letters and the guidelines to authors are also given.

Author contributions:	Author name:	Description of contribution
research design:	J.P. Petzer	Concept of the study was provided.
performed research: a) Synthetic work b) Characterization via NMR and MS c) Compound purity determination by HPLC d) IC ₅₀ value determination e) Recovery of enzyme activity after dilution study	M.M. van der Walt SASOL centre for chemistry, NWU M.M. van der Walt M.M. van der Walt M.M. van der Walt A. Petzer	Performed all the syntheses. Carried out by the SASOL Centre for Chemistry, North-West University. NMR spectra was recorded by André Joubert and MS was recorded by Johan Jordaan. The degree of purity of each compound was determined by HPLC analysis. The IC ₅₀ values for the inhibition of MAO-A and MAO-B were measured. The recovery of enzyme activity after dilution was performed for compound 8a .
contributed new reagents/ analytic tools: a) Synthesis of compounds	G. Terre'Blanche	Providing the facilities, instrumentation and relevant reagents for performing the synthetic work.
b) IC ₅₀ value determinations and the recovery of enzyme activity	A. Petzer	Providing the facilities, instrumentation and relevant


after dilution study		reagents for performing the IC ₅₀ value determination and the recovery of enzyme activity after dilution study
analyzed data:		
a) Characterization by NMR and MS	M.M. van der Walt J.P. Petzer	The analyses of the NMR and MS data were performed by M.M. van der Walt, with critical feedback by J.P. Petzer.
b) Compound purity measurement determination	M.M. van der Walt J.P. Petzer	The analyses of the compound purity data were performed by M.M. van der Walt, with critical feedback by J.P. Petzer.
c) IC ₅₀ value measurements	M.M. van der Walt A. Petzer J.P. Petzer	Calculations of the IC ₅₀ values were performed by M.M. van der Walt, with critical feedback by J.P. Petzer and A. Petzer.
d) Recovery of enzyme activity after dilution study	M.M. van der Walt A. Petzer J.P. Petzer	The analyses and interpretation of the data was performed by M.M. van der Walt with critical feedback by J.P. Petzer and A. Petzer.
manuscript:		
a) Writing of manuscript	M.M. van der Walt J.P. Petzer	Contribution was equally for the mentioned authors.
b) Comments, suggestions and proof reading	G. Terre'Blanche A. Petzer	Contribution was equally for the mentioned authors


6.3 Declaration and consent of each co-author


Novel sulfanylpthalimide analogues as highly potent inhibitors of monoamine oxidase B

The following is a declaration by the co-authors that confirms their individual roles in the above mentioned article, as described in the table of author's contributions. The authors hereby give consent that this article may form part of this doctoral thesis.

I declare that I approve the inclusion of above mentioned article in this thesis, that my role in this study is as indicated in the author's contributions table and is representative of my actual contribution. Herewith, I grant consent that this article may be published as part of the doctoral thesis of Mietha Magdalena van der Walt.



Prof. J.P. Petzer

Dr. G. Terre'Blanche

Dr. A. Petzer

6.4 Journal publishing agreement

The following assignment of publishing rights was accepted on behalf of all the co-authors of the manuscript published (DOI: <http://dx.doi.org/10.1016/j.bmcl.2012.08.113>) in Bioorganic & Medicinal Chemistry Letters. Elsevier Ltd has granted the authors retention of these rights for scholarly purposes (section 6.4.2).

6.4.1 Assignment of publishing rights

I hereby assign to Elsevier Ltd the copyright in the manuscript identified above (government authors not electing to transfer agree to assign an exclusive publishing and distribution license) and any supplemental tables, illustrations or other information submitted therewith that are intended for publication as part of the manuscript (the "Article") in all forms and media (whether now known or hereafter developed), throughout the world, in all languages, for the full term of copyright, effective when and if the article is accepted for publication. This transfer includes the right to provide the Article in electronic and online forms and systems. With respect to supplemental data that I wish to make accessible through a link in the Article, I hereby grant a non-exclusive license for such linking. If I have agreed with Elsevier Ltd to make available such supplemental data on a site or through a service of Elsevier Ltd, I hereby grant a non-exclusive license for such publication, posting and making available, and further permit indexing and archiving.

6.4.2 Retention of rights for scholarly purposes

I understand that I retain or am hereby granted (without the need to obtain further permission) the Retained Rights (see description below), and that no rights in patents, trademarks or other intellectual property rights are transferred to Elsevier Ltd.

The Retained Rights include:

- the right to use the Preprint or Accepted Author Manuscript for Personal Use, Internal Institutional Use and for Permitted Scholarly Posting;
- the right to use the Published Journal Article for Personal Use and Internal Institutional Use.

6.5 Author's instructions – Bioorganic & Medicinal Chemistry Letters

In the following section the author's instructions for Bioorganic & Medicinal Chemistry Letters are provided as published in the "Guide for Authors" of this journal.

6.5.1 Guide for authors: Introduction

Bioorganic & Medicinal Chemistry Letters publishes research results of outstanding significance and timeliness in the fields of medicinal chemistry, chemical biology, bioorganic chemistry, bioinorganic chemistry, and related disciplines. Articles are in the form of communications reporting experimental or theoretical results of special interest. This journal has no page charges.

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/ethicalguidelines>.

Conflict of interest

All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work. See also <http://www.elsevier.com/conflictsofinterest>.

Submission declaration

Submission of an article implies that the work described has not been published previously (excepting the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly

by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Changes to authorship

This policy concerns the addition, deletion, or rearrangement of author names in the authorship of accepted manuscripts: *Before the accepted manuscript is published in an online issue:* Requests to add or remove an author, or to rearrange the author names, must be sent to the Journal Manager from the corresponding author of the accepted manuscript and must include: (a) the reason the name should be added or removed, or the author names rearranged and (b) written confirmation (e-mail, fax, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed. Requests that are not sent by the corresponding author will be forwarded by the Journal Manager to the corresponding author, who must follow the procedure as described above. Note that: (1) Journal Managers will inform the Journal Editors of any such requests and (2) publication of the accepted manuscript in an online issue is suspended until authorship has been agreed. *After the accepted manuscript is published in an online issue:* Any requests to add, delete, or rearrange author names in an article published in an online issue will follow the same policies as noted above and result in a corrigendum.

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright see <http://www.elsevier.com/copyright>). Acceptance of the agreement will ensure the widest possible dissemination of information. An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement. Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <http://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has pre-printed forms for use by authors in these cases: please consult <http://www.elsevier.com/permissions>.

Retained author rights

As an author you (or your employer or institution) retains certain rights; for details you are referred to: <http://www.elsevier.com/authorsrights>.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated. Please see <http://www.elsevier.com/funding>.

Funding body agreements and policies

Elsevier has established agreements and developed policies to allow authors whose articles appear in journals published by Elsevier, to comply with potential manuscript archiving requirements as specified as conditions of their grant awards. To learn more about existing agreements and policies please visit <http://www.elsevier.com/fundingbodies>.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's Web Shop <http://webshop.elsevier.com/languageediting/> or visit our customer support site <http://support.elsevier.com> for more information.

Submission

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail removing the need for a paper trail.

Manuscripts should be addressed to the appropriate regional editor:

- *submissions from Japan and other Asian countries*: Professor Masakatsu Shibasaki, Japanese Regional Editor, Graduate School of Pharmaceutical Sciences, The University of Tokyo, Japan
- *submissions from Europe*: Professor Stephen Neidle, Cancer Research UK Professorial Fellow and Professor of Chemical Biology, The School of Pharmacy, University of London, UK

- *submissions from USA, Canada, and all others*: Professor Dale L. Boger, Editor-in-Chief, The Scripps Research Institute, La Jolla, CA, USA.

All manuscripts will be centrally handled by the journal editorial office, which will forward manuscripts to the regional editors:

Editorial Office

Bioorganic & Medicinal Chemistry Letters

525 B Street, Suite 1900

San Diego, CA 92101-4495, USA

Facsimile: (1) 619-699-6801

E-mail: bmcl-eo@elsevier.com

Authors who are unable to submit their paper online should contact the editorial office prior to submission at the above address.

Submission address

Please submit your article via <http://ees.elsevier.com/bmcl>

6.5.2 Preparation

Use of word processing software

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier: <http://www.elsevier.com/guidepublication>). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Templates

Templates are provided to allow authors to view their paper in a style close to the final printed form. Their use is optional. All manuscripts will be fully typeset from the author's electronic files. It should be noted that due to defined typesetting standards and the complex requirements of electronic publishing, the publisher will not always be able to exactly match the layout the author

has submitted. In particular, in the finished journal article, figures and tables are usually placed at the top or bottom of pages. The template is only intended to be used in assisting with the preparation and submission of manuscripts.

It should be noted that use of the journal templates is not a requirement and their adoption will neither speed nor delay publication. Elsevier can handle most major word processing packages and in general most formatting applied by authors for style and layout is replaced when the article is being typeset.

These templates contain a large number of macros. To ensure successful PDF conversion during online submission, it is important that the author save a new document based on the template, rather than saving the template itself. To use the template, the author should save the final document as a Word file with a '.doc' extension (rather than the '.dot' extension).

Article structure

Subdivision

Text should either not be subdivided or done so in the simplest possible way consistent with clarity. Headings should be used sparingly. Common headings such as 'Introduction' and 'Results and discussion' should not be used, as the text within these sections should be self-explanatory. As such, the publisher recommends that only the 'Acknowledgments' and 'References and notes' headings be used.

Material and methods

Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

Theory/calculation

A Theory section should extend, not repeat, the background to the article already dealt with in the Introduction and lay the foundation for further work. In contrast, a Calculation section represents a practical development from a theoretical basis.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication.
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Graphical abstract

A Graphical abstract is mandatory for this journal. It should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership online.

Authors must provide images that clearly represent the work described in the article. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <http://www.elsevier.com/graphicalabstracts> for examples. Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images also in accordance with all technical requirements: Illustration Service.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article, using superscript Arabic numbers. Many word processors build footnotes into the text, and this feature maybe used. Should this not be the case, indicate the position of footnotes in the text and present the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list. Indicate each footnote in a table with a superscript lowercase letter.

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Save text in illustrations as 'graphics' or enclose the font.
- Only use the following fonts in your illustrations: Arial, Courier, Times, Symbol.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Produce images near to the desired size of the printed version.
- Submit each figure as a separate file.

A detailed guide on electronic artwork is available on our web site: <http://www.elsevier.com/artworkinstructions>. You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

Regardless of the application used, when your electronic artwork is finalised, please 'save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS: Vector drawings. Embed the font or save the text as 'graphics'.

TIFF: Colour or grayscale photographs (halftones): always use a minimum of 300 dpi.

TIFF: Bitmapped line drawings: use a minimum of 1000 dpi.

TIFF: Combinations bitmapped line/half-ton (colour or grayscale): a minimum of 500 dpi is required.

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is'.

Please do not:

- supply files that are optimised for screen use (e.g., GIF, BMP, PICT, WPG); the resolution is too low;
- supply files that are too low in resolution;
- submit graphics that are disproportionately large for the content.

Colour artwork

Please make sure that artwork files are in an acceptable format (TIFF, EPS or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable colour figures then Elsevier will ensure, at no additional charge that these figures will appear in colour on the Web (e.g. Science Direct and other sites) regardless of whether or not these illustrations are reproduced in colour in the printed version. For colour reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article. Please indicate your preference for colour: in print or on the Web only. For further information on the preparation of electronic artwork, please see <http://www.elsevier.com/artworkinstructions>.

Please note: Because of technical complications which can arise by converting colour figures to 'grayscale' (for the printed version should you not opt for colour in print) please submit in addition usable black and white versions of all the colour illustrations.

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Number tables consecutively in accordance with their appearance in the text. Place footnotes to tables below the table body and indicate them with superscript lowercase letters. Avoid vertical rules. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

References

In the text, references should be indicated by superscript Arabic numerals which run consecutively through the paper and appear after any punctuation. Please ensure that all references are cited in the text and vice versa. The reference list should preferably contain only literature references though other information (e.g., experimental details) can be placed in this section. Preferably, each reference should contain only one literature citation. Authors are expected to check the original source reference for accuracy. Journal titles should be

abbreviated according to American Chemical Society guidelines (The ACS Style Guide; Dodd, J. S., Ed.; American Chemical Society: Washington, DC, 1997). A list of currently accepted journal abbreviations may be found at BMCL Journal Abbreviations.

Journal abbreviations source

Journal names should be abbreviated according to Index Medicus journal abbreviations: <http://www.nlm.nih.gov/tsd/serials/lji.html>; List of title word abbreviations: <http://www.issn.org/2-22661-LTWA-online.php>; CAS (Chemical Abstracts Service): <http://www.cas.org/sent.html>.

Supplementary data

Elsevier accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high resolution images, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including Science Direct: <http://www.sciencedirect.com>. In order to ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at <http://www.elsevier.com/artworkinstructions>.

Mol files

Mol files (optional): Elsevier would like to enrich your online article by visualizing and providing details of chemical structures you define as the main chemical compounds described. For this purpose, corresponding mol files can be uploaded via the online submission system. Each compound needs to be submitted as a separate mol file. Please use your preferred drawing tool to export chemical structures as mol files and ensure that they are unique, complete and do not contain any R groups or other variables so that a correct In ChI key can be generated. For more information see <http://www.elsevier.com/mol>.

6.5.3 After acceptance

Use of the Digital Object Identifier

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alpha-numeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*): <http://dx.doi.org/10.1016/j.physletb.2010.09.059>

When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

Proofs

One set of page proofs (as PDF files) will be sent by e-mail to the corresponding author (if we do not have an e-mail address then paper proofs will be sent by post) or, a link will be provided in the e-mail so that authors can download the files themselves. Elsevier now provides authors with PDF proofs which can be annotated; for this you will need to download Adobe Reader version 7 (or higher) available free from <http://get.adobe.com/reader>. Instructions on how to annotate PDF files will accompany the proofs (also given online). The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/reader/tech-specs.html>.

If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return them to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and return by fax, or scan the pages and e-mail, or by post. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately – please let us have all your corrections within 48 hours. It is important to ensure that all corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility. Note that Elsevier may proceed with the publication of your article if no response is received.

Offprints

The corresponding author, at no cost, will be provided with a PDF file of the article via email (the PDF file is a watermarked version of the published article and includes a cover sheet with the journal cover image and a disclaimer outlining the terms and conditions of use). For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's Web Shop (<http://webshop.elsevier.com//myarticleservices/offprints>). Authors requiring printed copies of multiple articles may use Elsevier Web Shop's 'Create Your Own Book' service to collate multiple articles within a single cover (<http://webshop.elsevier.com//myarticleservices/offprints/myarticlesservices/booklets>).

6.6 Published article

Novel sulfanylpthalimide analogues as highly potent inhibitors of monoamine oxidase B

Mietha M. Van der Walt,^a Gisella Terre'Blanche,^a Anél Petzer,^b and Jacobus P. Petzer^a

^a *Pharmaceutical Chemistry, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa*

^b *Unit for Drug Research and Development, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa*

Abstract—Monoamine oxidase (MAO) plays an essential role in the catabolism of neurotransmitter amines. The two isoforms of this enzyme, MAO-A and –B, are considered to be drug targets for the therapy of depression and neurodegenerative diseases, respectively. Based on a recent report that the phthalimide moiety may be a useful scaffold for the design of potent MAO-B inhibitors, the present study examines a series of 5-sulfanylpthalimide analogues as potential inhibitors of both human MAO isoforms. The results document that 5-sulfanylpthalimides are highly potent and selective MAO-B inhibitors with all of the examined compounds possessing IC₅₀ values in the nanomolar range. The most potent inhibitor, 5-(benzylsulfanyl)phthalimide, exhibits an IC₅₀ value of 0.0045 µM for the inhibition of MAO-B with a 427-fold selectivity for MAO-B compared to MAO-A. We conclude that 5-sulfanylpthalimides represent an interesting class of MAO-B inhibitors and may serve as lead compounds for the design of antiparkinsonian therapy.

Keywords: isatin; phthalimide; sulfanylpthalimide; monoamine oxidase; MAO; inhibition.

Monoamine oxidase A and B (MAO-A and –B) are flavine adenine dinucleotide (FAD) containing enzymes, bound to the outer membrane of mitochondria.^{1,2} These enzymes catalyze the oxidative deamination of neurotransmitter and dietary amines thereby terminating their physiological actions.³ Although MAO-A and –B share 70% sequence identity, they exhibit different substrate and inhibitor specificities.⁴ MAO-A metabolizes serotonin, adrenaline and noradrenaline and inhibitors of this enzyme are in use for the treatment of clinical depression and anxiety.⁵ MAO-B preferentially metabolizes the dietary amine, 2-phenylethylamine, and may therefore act as a metabolic brain barrier, limiting the entry of this false neurotransmitter into the central nervous system.^{6,7} Since MAO-B also catabolizes dopamine in the brain, inhibitors of this enzyme are used in the treatment of Parkinson's disease.^{5,7,8} In Parkinson's disease, MAO-B inhibitors conserve the depleted supply of central dopamine and enhance dopamine levels following administration of levodopa, the metabolic precursor of dopamine.⁹ For these reasons, MAO-B inhibitors are frequently combined with levodopa in Parkinson's disease therapy. It should be noted that MAO-A also metabolizes dopamine in the primate brain, and MAO-A inhibitors may consequently also elevate dopamine levels in the central nervous system.⁹ In addition, MAO-A inhibitors may be employed to treat non-motor symptoms of Parkinson's disease such as depression and anxiety.^{3,10} MAO-A inhibitors in conjunction with levodopa should, however, be used with caution since this may lead to a severe hypertensive response.¹¹




Figure 1. The structures of isatin (1), phthalimide (2) and caffeine (3).

A wide variety of heterocyclic moieties have been employed in the design of MAO inhibitors. Among these are isatin (1), an endogenous small molecule inhibitor of MAO-A and –B, and caffeine (3) (Fig. 1).^{12–14} Phthalimide (2), an isomer of isatin, has also recently been reported to be a potentially useful scaffold for the design of MAO-B selective inhibitors.¹⁵ Although phthalimide is a weak MAO inhibitor, substitution on C5 yields structures endowed with highly potent and selective MAO-B inhibitory activities. In contrast, N-substitution yields structures that are essentially devoid of MAO inhibitory properties.¹⁵ The MAO-B inhibitory properties of isatin

and caffeine may similarly be enhanced by substitution on the C5 and C6 positions of isatin and the C8 position of caffeine. In this regard, the benzyloxy substituent has been shown to be particularly favorable, and benzyloxy substitution of isatin, caffeine and phthalimide yields compounds **4–6** which are several orders of magnitude more potent MAO-B inhibitors than the parent compounds (Fig. 2).^{13–15} In all instances, halogen substitution on the benzyloxy ring further improves inhibition potency. Modeling studies have shown that productive interactions of the benzyloxy side chain with the MAO-B entrance cavity may be responsible for this behavior. Interestingly, the benzylsulfanyl side chain appears to exhibit similar properties to that of the benzyloxy moiety, since a series of 8-(benzylsulfanyl)caffeine (**7**) analogues has recently been shown to exhibit similar MAO-B inhibition potencies to those of the 8-benzyloxycaffeine (**5**) analogues.¹⁶ Based on these analyses, the present study examines the possibility that benzylsulfanyl substitution on C5 of phthalimide (to yield **8a**) would also lead to highly potent MAO-B inhibition. For this purpose, the effect that substitution (Cl, Br, F and OCH₃) on the benzylsulfanyl ring has on MAO inhibition will be explored. In addition, this study also determines the effect on MAO inhibition by the phenylsulfanyl, (2-phenylethyl)sulfanyl, cyclohexylsulfanyl and (3-methylbutyl)sulfanyl substituents. This study therefore aims to discover new highly potent MAO-B inhibitors and to contribute to the structure-activity relationships (SARs) of MAO inhibition by phthalimide derived compounds.





Figure 2. The structures of 5-benzyloxyisatin (**4**), 8-benzyloxycaffeine (**5**), 5-benzyloxyphthalimide (**6**) and 8-(benzylsulfanyl)caffeine (**7**).


The 5-sulfanylphthalimides (**8a–k**) were conveniently synthesized according to a previously described protocol (Scheme 1).¹⁷ The appropriate thiol reagents were reacted with 5-nitrophthalimide in the presence of K₂CO₃ to yield the target compounds in low to good yields (4–76%). The 5-sulfanylphthalimides were purified via crystallization from an appropriate solvent. In each instance, the structures and purities of the target compounds were verified by ¹H NMR, ¹³C NMR, mass spectrometry and HPLC analysis as cited in the supplementary material. The presence of two ¹³C NMR signals at 167.6–168.8 ppm, which corresponds to the

carbonyl carbons at C1 and C3, and a ^1H NMR signal at 8.12–8.20 ppm (CDCl_3) or 11.28–11.36 ppm ($\text{DMSO}-d_6$), which corresponds to the phthalimide NH proton, confirmed the presence of the phthalimide ring (Table 1).



Scheme 1. Synthetic pathway to 5-sulfanylphthalimide analogues. Reagents and conditions: (a) K_2CO_3 , acetone, reflux, 24 h; (b) HCl (6 N).

Table 1. The ^1H NMR and ^{13}C NMR chemical shifts for the NH proton and carbonyl C1 and C3 of phthalimide analogues **8a–k**.




	R	NH	C1/C3	
8a	$-\text{S}-(\text{CH}_2)-\text{C}_6\text{H}_5$	11.28 ^a	168.8 ^a	168.8 ^a
8b	$-\text{S}-(\text{CH}_2)-(4\text{-Cl-C}_6\text{H}_4)$	11.29 ^a	168.8 ^a	168.8 ^a
8c	$-\text{S}-(\text{CH}_2)-(4\text{-Br-C}_6\text{H}_4)$	11.29 ^a	168.7 ^a	168.8 ^a
8d	$-\text{S}-(\text{CH}_2)-(4\text{-F-C}_6\text{H}_4)$	11.29 ^a	168.8 ^a	168.8 ^a
8e	$-\text{S}-(\text{CH}_2)-(4\text{-OCH}_3\text{-C}_6\text{H}_4)$	11.28 ^a	168.8 ^a	168.8 ^a
8f	$-\text{S-C}_6\text{H}_5$	8.20 ^b	167.7 ^b	167.8 ^b
8g	$-\text{S}-(4\text{-Cl-C}_6\text{H}_4)$	11.35 ^a	168.5 ^a	168.6 ^a
8h	$-\text{S}-(4\text{-Br-C}_6\text{H}_4)$	11.36 ^a	168.5 ^a	168.6 ^a
8i	$-\text{S}-(\text{CH}_2)_2-\text{C}_6\text{H}_5$	11.28 ^a	168.8 ^a	168.8 ^a
8j	$-\text{S-C}_6\text{H}_{11}$	11.30 ^a	167.6 ^b	167.8 ^b
8k	$-\text{S}-(\text{CH}_2)_2-\text{CH}(\text{CH}_3)_2$	8.12 ^b	167.9 ^b	168.0 ^b

^a NMR experiments conducted in $\text{DMSO}-d_6$.

^b NMR experiments conducted in CDCl_3 .

To examine the MAO inhibitory properties of the 5-sulfanylpthalimides, recombinant human MAO-A and -B were employed.¹⁸ The enzyme catalytic activities in the absence and presence of the test inhibitors were determined by fluorometrically measuring the MAO-catalyzed formation of 4-hydroxyquinoline from the mixed MAO-A/B substrate, kynuramine.^{14,18} This approach was suitable for evaluating the MAO inhibitory properties of all the 5-sulfanylpthalimides. The inhibition potencies of the test compounds were calculated from the sigmoidal dose-response curves and are expressed as the corresponding IC₅₀ values. These IC₅₀ values as well as the selectivity index (SI) values [SI = IC₅₀(MAO-A)/(IC₅₀(MAO-B))] are given in table 2. The results document that all of the 5-sulfanylpthalimides are potent inhibitors of human MAO-B with IC₅₀ values ranging from 0.0045–0.986 μM. In accordance with expectation (see introduction), benzylsulfanyl substitution of phthalimide to yield **8a**, resulted in highly potent MAO-B inhibition. An IC₅₀ value of 0.0045 μM was recorded for **8a**. In fact **8a** proved to be the most potent MAO-B inhibitor among the compounds of the present series. Compared to its C5 benzyloxy substituted phthalimide homologue, compound **6** (IC₅₀ = 0.043 μM), compound **8a** is approximately eightfold more potent as a MAO-B inhibitor.¹⁵ Compound **8a** also proved to be a more potent MAO-B inhibitor than 5-benzyloxyisatin (**4**) (IC₅₀ = 0.103 μM),¹³ 8-benzyloxycaffeine (**5**) (IC₅₀ = 1.77 μM)¹⁴ and 8-(benzylsulfanyl)caffeine (**7**) (IC₅₀ = 1.86 μM).¹⁶ Substitution on the benzylsulfanyl ring with Cl, Br, F and OCH₃, to yield **8b–e**, also resulted in highly potent inhibition, with these homologues exhibiting IC₅₀ values of 0.0056–0.020 μM. It is interesting to note that particularly the halogen substituted homologues **8b–d** are weaker MAO-B inhibitors than compound **8a**. This is in contrast to the results obtained with 8-benzyloxycaffeine (**5**), 5-benzyloxyphthalimide (**6**) and 8-(benzylsulfanyl)caffeine (**7**), where particularly bromine substitution on the phenyl ring leads to an enhancement in MAO-B inhibitory potency.^{14–16} From these results it may therefore be concluded that 5-(benzylsulfanyl)phthalimides are highly potent MAO-B inhibitors and superior to the lead structures, compounds **4–7**, of this study.

Table 2. The IC₅₀ values for the inhibition of recombinant human MAO-A and –B by 5-sulfanylpthalimides **8a–k**.



R	IC ₅₀ (μM) ^a			SI ^b
	MAO-A	MAO-B		
8a	–S–(CH ₂)–C ₆ H ₅	1.92 ± 0.172	0.0045 ± 0.0004	427
8b	–S–(CH ₂)–(4-Cl-C ₆ H ₄)	0.506 ± 0.032	0.0056 ± 0.0003	90
8c	–S–(CH ₂)–(4-Br-C ₆ H ₄)	0.273 ± 0.041	0.0074 ± 0.0029	37
8d	–S–(CH ₂)–(4-F-C ₆ H ₄)	0.958 ± 0.003	0.0068 ± 0.0009	141
8e	–S–(CH ₂)–(4-OCH ₃ -C ₆ H ₄)	1.63 ± 0.023	0.020 ± 0.0042	82
8f	–S–C ₆ H ₅	8.03 ± 0.622	0.986 ± 0.026	8.1
8g	–S–(4-Cl-C ₆ H ₄)	1.68 ± 0.343	0.457 ± 0.093	3.7
8h	–S–(4-Br-C ₆ H ₄)	1.01 ± 0.053	0.364 ± 0.050	2.8
8i	–S–(CH ₂) ₂ –C ₆ H ₅	2.27 ± 0.136	0.030 ± 0.0094	76
8j	–S–C ₆ H ₁₁	1.03 ± 0.062	0.179 ± 0.0082	5.8
8k	–S–(CH ₂) ₂ –CH(CH ₃) ₂	0.380 ± 0.042	0.015 ± 0.0033	26

^a All values are expressed as the mean ± SD of triplicate determinations.

^b The selectivity index is the selectivity for the MAO-B isoform and is given as the ratio of [IC₅₀(MAO-A)]/[IC₅₀(MAO-B)].

Although still considered as a potent MAO-B inhibitor, the phenylsulfanyl homologue **8f** (IC₅₀ = 0.986 μM) was the weakest inhibitor of the present series. In this case, however, halogen substitution in the phenyl ring to yield compounds **8g** (IC₅₀ = 0.457 μM) and **8h** (IC₅₀ = 0.364 μM), resulted in enhanced MAO-B inhibition. The (2-phenylethyl)sulfanyl substituted homologue **8i** was also found to be a potent MAO-B inhibitor with an IC₅₀ value of 0.030 μM, approximately sixfold weaker than **8a**. Since both **8f** and **8i** are weaker MAO-B inhibitor than **8a**, it may be concluded that the benzylsulfanyl side chain is particularly suitable for enhancing the MAO-B inhibitory potency of phthalimide, and that neither a reduction in side chain length (to yield **8f**), nor an increase in chain length (to yield **8i**) would further increase inhibitory activity. The general suitability of 5-sulfanylpthalimides for MAO-B inhibition was further demonstrated with the

finding that the cyclohexylsulfanyl (**8j**) and (3-methylbutyl)sulfanyl (**8k**) substituted homologues are also potent MAO-B inhibitors with IC_{50} values of 0.179 μ M and 0.015 μ M, respectively.

The 5-sulfanylpthalimides were found to also act as inhibitors of MAO-A. Four homologues, **8b–d** and **8k**, exhibited IC_{50} values in the nanomolar range, with **8c** being the most potent MAO-A inhibitor with an IC_{50} value of 0.273 μ M. As evident from the SI values, all of the 5-sulfanylpthalimides were, however, selective inhibitors of the MAO-B isoform. The most potent MAO-B inhibitor of the series, **8a**, is also the most selective inhibitor with an SI value of 427. This compound may therefore be considered the most suitable homologue of the series where MAO-B selectivity is desired. As already noted, MAO-A inhibition may lead to adverse effects when combined with certain antiparkinsonian therapies. In contrast, **8c** may be considered as a potent MAO inhibitor with comparatively low isoform selectivity.

Since it is reported that the benzyloxypthalimide class of MAO inhibitors interacts reversibly with MAO, the current study examined if this property is also shared by sulfanylpthalimides.¹⁵ For this purpose, the reversibility of MAO-B inhibition by the most potent sulfanylpthalimide MAO-B inhibitor, compound **8a**, was investigated by measuring the degree of enzyme recovery after dilution of the enzyme-inhibitor complex. MAO-B was preincubated with **8a** at concentrations of $10 \times IC_{50}$ and $100 \times IC_{50}$ for 30 min and then diluted to $0.1 \times IC_{50}$ and $1 \times IC_{50}$, respectively.¹⁹ The results show that after dilution of the enzyme-inhibitor complexes to concentrations of **8a** equal to $0.1 \times IC_{50}$ and $1 \times IC_{50}$, the MAO-B catalytic activities are recovered to levels of approximately 72% and 28%, respectively, of the control value (Fig. 3). This behavior is consistent with a reversible interaction of **8a** with MAO-B. In contrast, incubation of MAO-B with the irreversible inhibitor (R)-deprenyl ($10 \times IC_{50}$), the MAO-B activities were not recovered (2.1% of control). Interestingly, after dilution of the **8a**–MAO-B complex to $0.1 \times IC_{50}$ and $1 \times IC_{50}$, the enzyme activities are not recovered to 90% and 50%, respectively, as expected. This result suggests that, for the inhibition of MAO-B, **8a** may possess a quasi-reversible or tight-binding component.




Figure 3. The reversibility of the inhibition of MAO-B by **8a**. The enzyme was preincubated with **8a** at $10 \times \text{IC}_{50}$ and $100 \times \text{IC}_{50}$ for 30 min and then diluted to $0.1 \times \text{IC}_{50}$ and $1 \times \text{IC}_{50}$, respectively. For comparison, (R)-deprenyl, at $10 \times \text{IC}_{50}$ was similarly incubated with MAO-B and diluted to $0.1 \times \text{IC}_{50}$. The residual enzyme activities were subsequently measured.

In conclusion, the present study shows that 5-sulfanylpthalimides are potent and selective inhibitors of MAO-B. In this regard, the benzylsulfanyl side chain is particularly suitable for enhancing the MAO-B inhibitory potency of phthalimide. It is noteworthy that compound **8a** ($\text{IC}_{50} = 0.0045 \mu\text{M}$) is approximately 30,000-fold more potent than phthalimide (**2**) ($\text{IC}_{50} = 134 \mu\text{M}$).¹⁵ This illustrates the importance of the C5 side chain for MAO-B inhibitory activity. Based on their MAO-B inhibition potencies and appropriate selectivity profiles, this study concludes that 5-sulfanylpthalimides are suitable lead compounds for the development of antiparkinsonian drugs. From a design point of view it is noteworthy that a wide variety of C5 substituents yield 5-sulfanylpthalimides with potent MAO-B inhibitory actions. This suggests that structural modifications made to the C5 side chain in order to improve the properties of the compound are less likely to reduce MAO-B inhibition potency.

Acknowledgements

We are grateful to André Joubert of the SASOL Centre for Chemistry, North-West University for recording the NMR spectra, and the Mass Spectrometry Service, University of the Witwatersrand for the MS analyses. Financial support for this work was provided by the North-West University, the National Research Foundation and the Medical Research Council, South Africa.

References

1. Binda, C.; Newton-Vinson, P.; Hubálek, F.; Edmondson, D. E.; Mattevi, A. *Nat. Struct. Biol.* **2002**, *9*, 22.
2. Son, S.-Y.; Ma, J.; Kondou, Y.; Yoshimura, M.; Yamashita, E.; Tsukihara, T. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 5739.
3. Youdim, M. B. H.; Bakhle, Y. S. *Br. J. Pharmacol.* **2006**, *147*, S287.
4. Shih, J. C.; Chen, K.; Ridd, M. J. *Annu. Rev. Neurosci.* **1999**, *22*, 197.
5. Yamada, M.; Yasuhara, H. *Neurotoxicology*. **2004**, *25*, 215.
6. Lasbennes, F.; Sercombe, R.; Seylaz, J. J. *Cereb. Blood Flow Metab.* **1983**, *3*, 521.
7. Youdim, M. B. H.; Edmondson, D.; Tipton, K. F. *Nat. Rev. Neurosci.* **2006**, *7*, 295.
8. Nagatsu, T.; Sawada, M. *J. Neural Transm. Suppl.* **2006**, *71*, 53.
9. Finberg, J. P.; Wang, J.; Bankiewicz, K.; Harvey-White, J.; Kopin, I. J.; Goldstein, D. S. *J. Neural Transm. Suppl.* **1998**, *52*, 279.
10. Prediger, R. D. S.; Matheus, F. C.; Schwarzbald, M. L.; Lima, M. M. S.; Vital, M. A. B. F. *Neuropharmacology* **2012**, *62*, 115.
11. Zesiewicz, T. A.; Hauser, R. A. *In Parkinson's Disease: Diagnosis and Clinical Management*; Factor, S. A., Weiner, W. J., Eds.; Demos Medical Publishing: New York, 2002; 365.
12. Van der Walt, E. M.; Milczek, E. M.; Malan, S. F.; Edmondson, D. E.; Castagnoli, N., Jr.; Bergh, J. J.; Petzer, J. P. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 2509.
13. Manley-King, C. I.; Bergh, J. J.; Petzer, J. P. *Bioorg. Med. Chem.* **2011**, *19*, 261.
14. Strydom, B.; Malan, S. F.; Castagnoli, N., Jr.; Bergh, J. J.; Petzer, J. P. *Bioorg. Med. Chem.* **2010**, *18*, 1018.
15. Manley-King, C. I.; Bergh, J. J.; Petzer, J. P. *Bioorg. Med. Chem.* **2011**, *19*, 4829.
16. Booyesen, H. P.; Moraal, C.; Terre'Blanche, G.; Petzer, A.; Bergh, J. J.; Petzer, J. P. *Bioorg Med Chem.* **2011**, *19*, 7507.
17. Arend, M. P.; Flippin, L. A.; Guenzler-Pukall, V.; Ho, W. -B.; Turtle, E. D.; Du, X. U. S. Patent 20040254215, **2004**.
18. Novaroli, L.; Reist, M.; Favre, E.; Carotti, A.; Catto, M.; Carrupt, P. A. *Bioorg. Med. Chem.* **2005**, *13*, 6212.
19. Petzer, A.; Harvey, B. H.; Wegener, G.; Petzer, J. P. *Toxicol. Appl. Pharm.* **2012**, *258*, 403.

6.7 Supplementary material

Novel sulfanylpthalimide analogues as highly potent inhibitors of monoamine oxidase B

Mietha M. Van der Walt,^a Gisella Terre'Blanche,^a Anél Petzer,^b and Jacobus P. Petzer^{a*}

^a *Pharmaceutical Chemistry, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa*

^b *Unit for Drug Research and Development, School of Pharmacy, North-West University, Private Bag X6001, Potchefstroom, 2520, South Africa*

6.7.1 Experimental procedures

6.7.1.1 Chemicals and instrumentation

Unless otherwise noted, all the reagents were obtained from Sigma-Aldrich and were used without further purification. Proton (¹H) and carbon (¹³C) NMR spectra were recorded on a Bruker Avance III 600 spectrometer in CDCl₃ or DMSO-*d*₆. The chemical shifts are reported in parts per million (δ) relative to the signal of tetramethylsilane. Spin multiplicities are abbreviated as follows: s (singlet), d (doublet), dd (doublet of doublets), t (triplet), q (quartet) or m (multiplet). High resolution mass spectra (HRMS) were recorded with a Bruker micrOTOF-Q II mass spectrometer in atmospheric-pressure chemical ionization (APCI) mode. Melting points (mp) were measured with a Buchi M-545 melting point apparatus and are uncorrected. A Varian Cary Eclipse fluorescence spectrophotometer was used for fluorescence spectrophotometry. For the enzymology, kynuramine.2HBr and microsomes from insect cells containing recombinant human MAO-A and -B (5 mg protein/mL) were obtained from Sigma-Aldrich.

6.7.1.2 Synthesis of the 5-sulfanylpthalimide analogues (**8a–k**)

A mixture of the appropriate thiol (6 mmol), 4-nitrophthalimide (4 mmol) and K₂CO₃ (10 mmol) in 20 mL acetone was heated under reflux for 24 h. The reaction was cooled to room temperature and diluted with 150 mL water. The mixture was subsequently acidified to pH = 2 with 6 N HCl. The resulting precipitate was collected by filtration and dried at 50 °C.¹

5-(Benzylsulfanyl)phthalimide (**8a**)

The title compound (yellow crystals) was prepared from 4-nitrophthalimide and benzyl mercaptan in a yield of 65%: mp 177.5–177.6 °C (ethanol); ¹H NMR (Bruker Avance III 600, DMSO-*d*₆) δ 4.43 (s, 2H), 7.23–7.26 (m, 1H), 7.30–7.33 (m, 2H), 7.42 (d, 2H, J = 7.5 Hz), 7.67 (s, 3H), 11.28 (s, 1H); ¹³C NMR (Bruker Avance III 600, DMSO-*d*₆) δ 35.4, 120.2, 123.2, 127.4,

128.6, 128.9, 129.0, 131.5, 133.5, 136.3, 145.5, 168.8, 168.8; APCI-HRMS m/z : calcd for $C_{15}H_{12}NO_2S$ (MH^+), 270.0589, found 270.0568; Purity (HPLC): 100%.

5-[(4-Chlorobenzyl)sulfanyl]phthalimide (8b)

The title compound (white powder) was prepared from 4-nitrophthalimide and 4-chlorobenzyl mercaptan in a yield of 9%: mp 228.0–228.14 °C (ethanol); 1H NMR (Bruker Avance III 600, DMSO- d_6) δ 4.45 (s, 2H), 7.37 (d, 2H, J = 8.7 Hz), 7.44 (d, 2H, J = 8.7 Hz), 7.67 (m, 3H), 11.29 (s, 1H); ^{13}C NMR (Bruker Avance III 600, DMSO- d_6) δ 34.6, 120.4, 123.3, 128.5, 129.2, 130.7, 131.8, 131.9, 133.5, 135.7, 145.0, 168.8, 168.8; APCI-HRMS m/z : calcd for $C_{15}H_{11}ClNO_2S$ (MH^+), 304.0199, found 304.0174; Purity (HPLC): 99%.

5-[(4-Bromobenzyl)sulfanyl]phthalimide (8c)

The title compound (light yellow powder) was prepared from 4-nitrophthalimide and 4-bromobenzyl mercaptan in a yield of 7%: mp 255.7–255.9 °C (ethanol); 1H NMR (Bruker Avance III 600, DMSO- d_6) δ 4.43 (s, 2H), 7.38 (d, 2H, J = 8.3 Hz), 7.51 (d, 2H, J = 8.3 Hz), 7.68 (m, 3H), 11.29 (s, 1H); ^{13}C NMR (Bruker Avance III 600, DMSO- d_6) δ 34.6, 120.4, 120.5, 123.3, 129.2, 131.0, 131.4, 131.8, 133.5, 136.1, 145.0, 168.7, 168.8; APCI-HRMS m/z : calcd for $C_{15}H_{11}BrNO_2S$ (MH^+), 347.9694, found 347.9662; Purity (HPLC): 100%.

5-[(4-Fluorobenzyl)sulfanyl]phthalimide (8d)

The title compound (dark yellow powder) was prepared from 4-nitrophthalimide and 4-fluorobenzyl mercaptan in a yield of 70%: mp 179.8–180.2 °C (ethanol); 1H NMR (Bruker Avance III 600, DMSO- d_6) δ 4.43 (s, 2H), 7.13 (t, 2H, J = 8.7 Hz), 7.44–7.46 (m, 2H), 7.67 (s, 3H), 11.29 (s, 1H); ^{13}C NMR (Bruker Avance III 600, DMSO- d_6) δ 34.6, 115.3, 115.4, 120.4, 123.3, 129.1, 130.8, 130.9, 131.7, 132.6, 133.5, 145.2, 160.6, 162.2, 168.8, 168.8; APCI-HRMS m/z : calcd for $C_{15}H_{11}FNO_2S$ (MH^+), 288.0495, found 288.0484; Purity (HPLC): 96%.

5-[(4-Methoxybenzyl)sulfanyl]phthalimide (8e)

The title compound was prepared from 4-nitrophthalimide and 4-methoxybenzyl mercaptan in a yield of 63%: mp 172.7–172.8 °C (ethanol); 1H NMR (Bruker Avance III 600, DMSO- d_6) δ 3.70 (s, 3H), 4.37 (s, 2H), 6.87 (d, 2H, J = 8.7 Hz), 7.33 (d, 2H, J = 8.7 Hz), 7.66 (s, 3H), 11.28 (s, 1H); ^{13}C NMR (Bruker Avance III 600, DMSO- d_6) δ 35.0, 55.0, 114.0, 120.2, 123.2, 127.9, 128.9, 130.1, 131.5, 133.5, 145.8, 158.5, 168.8, 168.8; APCI-HRMS m/z : calcd for $C_{16}H_{14}NO_3S$ (MH^+), 300.0694, found 300.0675; Purity (HPLC): 100%.

5-(Phenylsulfanyl)phthalimide (8f)

The title compound (yellow crystals) was prepared from 4-nitrophthalimide and thiophenol in a yield of 4%: mp 166.6–168.0 °C (ethanol); ^1H NMR (Bruker Avance III 600, CDCl_3) δ 7.43–7.44 (m, 4H), 7.47 (m, 1H), 7.51–7.52 (m, 2H), 7.66 (d, 1H, $J = 7.9$ Hz), 8.20 (s, 1H); ^{13}C NMR (Bruker Avance III 600, CDCl_3) δ 121.5, 123.8, 129.1, 129.7, 130.1, 130.4, 131.9, 133.5, 134.6, 148.3, 167.7, 167.8; APCI-HRMS m/z : calcd for $\text{C}_{14}\text{H}_{10}\text{NO}_2\text{S}$ (MH^+), 256.0432, found 256.0404; Purity (HPLC): 100%.

5-[(4-Chlorophenyl)sulfanyl]phthalimide (8g)

The title compound (white powder) was prepared from 4-nitrophthalimide and 4-chlorothiophenol in a yield of 76%: mp 198.8–200.4 °C (ethanol); ^1H NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 7.40 (s, 1H), 7.54–7.56 (m, 5H), 7.73 (d, 1H, $J = 7.9$ Hz), 11.35 (s, 1H); ^{13}C NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 121.0, 123.9, 129.9, 130.1, 130.2, 132.6, 133.8, 134.5, 135.5, 144.8, 168.5, 168.6; APCI-HRMS m/z : calcd for $\text{C}_{14}\text{H}_9\text{ClNO}_2\text{S}$ (MH^+), 290.0043, found 290.0014; Purity (HPLC): 100%.

5-[(4-Bromophenyl)sulfanyl]phthalimide (8h)

The title compound (white powder) was prepared from 4-nitrophthalimide and 4-bromothiophenol in a yield of 46%: mp 203.2–203.3 °C (ethanol); ^1H NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 7.41 (d, 1H, $J = 1.5$ Hz), 7.48 (d, 2H, $J = 8.7$ Hz), 7.55 (dd, 1H, $J = 1.5, 7.9$ Hz), 7.69 (d, 2H, $J = 8.7$ Hz), 7.73 (d, 1H, $J = 7.9$ Hz), 11.36 (s, 1H); ^{13}C NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 121.2, 123.1, 123.9, 130.1, 130.5, 132.7, 133.1, 133.8, 135.6, 144.6, 168.5, 168.6; APCI-HRMS m/z : calcd for $\text{C}_{14}\text{H}_9\text{BrNO}_2\text{S}$ (MH^+), 333.9537, found 333.9538; Purity (HPLC): 99%.

5-[(2-Phenylethyl)sulfanyl]phthalimide (8i)

The title compound (dark yellow needles) was prepared from 4-nitrophthalimide and 2-phenylethyl mercaptan in a yield of 75%: mp 139.0–139.5 °C (ethanol); ^1H NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 2.92 (t, 2H, $J = 7.6$ Hz), 3.40 (t, 2H, $J = 7.2$ Hz), 7.19–7.21 (m, 1H), 7.26–7.30 (m, 4H), 7.63–7.69 (m, 3H), 11.28 (s, 1H); ^{13}C NMR (Bruker Avance III 600, $\text{DMSO}-d_6$) δ 32.3, 34.0, 120.0, 123.3, 126.4, 128.4, 128.6, 128.8, 131.4, 133.6, 139.6, 145.6, 168.8, 168.8; APCI-HRMS m/z : calcd for $\text{C}_{16}\text{H}_{14}\text{NO}_2\text{S}$ (MH^+), 284.0745, found 284.0720; Purity (HPLC): 100%.

5-(Cyclohexylsulfanyl)phthalimide (8j)

The title compound (light yellow powder) was prepared from 4-nitrophthalimide and cyclohexanethiol in a yield of 5%: mp 180.7–180.9 °C (ethanol); ^1H NMR (Bruker Avance III 600, DMSO- d_6) δ 1.22–1.43 (m, 6H), 1.58–1.60 (m, 1H), 1.69–1.72 (m, 2H), 1.94–1.97 (m, 2H), 3.59–3.63 (m, 1H), 7.67–7.71 (m, 2H), 11.30 (s, 1H); ^{13}C NMR (Bruker Avance III 600, CDCl_3) δ 25.6, 25.8, 32.9, 45.1, 122.4, 123.7, 128.8, 133.3, 133.5, 146.5, 167.6, 167.8; APCI-HRMS m/z : calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_2\text{S}$ (MH^+), 262.0902, found 262.0882; Purity (HPLC): 96%.

5-[(3-Methylbutyl)sulfanyl]phthalimide (8k)

The title compound (light yellow powder) was prepared from 4-nitrophthalimide and 3-methyl-1-butanethiol in a yield of 19%: mp 64–115 °C (ethanol); ^1H NMR (Bruker Avance III 600, CDCl_3) δ 0.94 (d, 6H, $J = 6.8$ Hz), 1.56–1.60 (q, 2H, $J = 7.2$ Hz), 1.72–1.76 (m, 1H), 3.01–3.03 (m, 2H), 7.51 (dd, 1H, $J = 1.5, 7.9$ Hz), 7.63 (d, 1H, $J = 1.5$ Hz), 7.69 (d, 1H, $J = 7.9$ Hz), 8.12 (s, 1H); ^{13}C NMR (Bruker Avance III 600, CDCl_3) δ 22.2, 27.5, 30.2, 37.2, 120.4, 123.6, 128.3, 131.5, 133.4, 147.9, 167.9, 168.0; APCI-HRMS m/z : calcd for $\text{C}_{13}\text{H}_{16}\text{NO}_2\text{S}$ (MH^+), 250.0902, found 250.0884; Purity (HPLC): 100%.

6.7.1.3 IC_{50} value determination

Recombinant human MAO-A and MAO-B containing microsomes (5 mg protein/mL) were pre-aliquoted and stored at -70 °C. All incubations were carried out in potassium phosphate buffer ($\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$ 100 mM, made isotonic with KCl, pH 7.4) to a final volume of 500 μL . The reactions contained various concentrations of the test inhibitor (0–100 μM) and kynuramine, a mixed MAO-A/B substrate. The final concentrations of kynuramine in the reactions were 45 μM and 30 μM for MAO-A and MAO-B, respectively. The stock solutions of all the test inhibitors were prepared in DMSO and added to the reactions to yield a final volume of 4% (v/v) DMSO. The reactions were initiated with the addition of MAO-A or MAO-B (0.0075 mg protein/mL) and allowed to incubate for 20 min at 37 °C. The reactions were subsequently terminated with the addition of 400 μL NaOH (2 N) and 1000 μL water, and the concentrations of the MAO generated 4-hydroxyquinoline were measured by fluorescence spectrophotometry ($\lambda_{\text{em}} = 310$; $\lambda_{\text{ex}} = 400$ nm).^{2,3} By employing a linear calibration curve (4-hydroxyquinoline: 0.047–1.56 μM), the enzyme catalytic rates were calculated and fitted to the one site competition model incorporated into the Prism software package (GraphPad). The IC_{50} values were determined in triplicate and are expressed as mean \pm standard deviation (SD).

6.7.1.4 Recovery of enzyme activity after dilution


Compound **8a** [$IC_{50}(\text{MAO-B}) = 0.0045 \mu\text{M}$], at concentrations equal to $0 \times IC_{50}$, $10 \times IC_{50}$ and $100 \times IC_{50}$, was preincubated with recombinant human MAO-B (0.75 mg/ml) for 30 min at 37 °C. The reaction solvent was K_2HPO_4/KH_2PO_4 (pH 7.4, 100 mM, made isotonic with KCl). DMSO at a final concentration of 4% was added as co-solvent to all preincubations. The reactions were subsequently diluted 100-fold with the addition of a solution of kynuramine to yield final concentrations of **8a** equal to $0 \times IC_{50}$, $0.1 \times IC_{50}$ and $1 \times IC_{50}$, and a final concentration of kynuramine equal to 30 μM . The final concentration of MAO-B was 0.0075 mg/mL. The reactions were incubated for a further 20 min at 37 °C, terminated and the residual rates of 4-hydroxyquinoline formation were measured as described above for the IC_{50} value determinations.⁴ As control experiments, (R)-deprenyl ($IC_{50} = 0.079 \mu\text{M}$) was similarly preincubated with recombinant human MAO-A and -B (0.75 mg/ml) at concentrations equal to $10 \times IC_{50}$ and diluted 100-fold with the addition of kynuramine to yield final concentrations of (R)-deprenyl equal to $0.1 \times IC_{50}$.⁴

6.7.2 References


1. Arend, M. P.; Flippin, L. A.; Guenzler-Pukall, V.; Ho, W. -B.; Turtle, E. D.; Du, X. U. S. Patent 20040254215, **2004**.
2. Strydom, B.; Malan, S. F.; Castagnoli, N., Jr.; Bergh, J. J.; Petzer, J. P. *Bioorg. Med. Chem.* **2010**, 18, 1018.
3. Novaroli, L.; Reist, M.; Favre, E.; Carotti, A.; Catto, M.; Carrupt, P. A. *Bioorg. Med. Chem.* **2005**, 13, 6212.
4. Petzer, A.; Harvey, B. H.; Wegener, G.; Petzer, J. P. *Toxicol. Appl. Pharm.* **2012**, 258, 403.

6.7.3. NMR spectra of the 5-sulfanylphthalimide analogues (**8a–k**)


^1H NMR (DMSO-*d*6): **5-(Benzylsulfanyl)phthalimide (8a)**




^{13}C NMR (DMSO-*d*6): **5-(Benzylsulfanyl)phthalimide (8a)**




¹H NMR (DMSO-*d*₆): **5-[(4-Chlorobenzyl)sulfanyl]phthalimide (8b)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Chlorobenzyl)sulfanyl]phthalimide (8b)**




¹H NMR (DMSO-*d*₆): **5-[(4-Bromobenzyl)sulfanyl]phthalimide (8c)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Bromobenzyl)sulfanyl]phthalimide (8c)**




¹H NMR (DMSO-*d*₆): **5-[(4-Fluorobenzyl)sulfanyl]phthalimide (8d)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Fluorobenzyl)sulfanyl]phthalimide (8d)**




¹H NMR (DMSO-*d*₆): **5-[(4-Methoxybenzyl)sulfanyl]phthalimide (8e)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Methoxybenzyl)sulfanyl]phthalimide (8e)**




^1H NMR (CDCl_3): **5-(Phenylsulfanyl)phthalimide (8f)**




^{13}C NMR (CDCl_3): **5-(Phenylsulfanyl)phthalimide (8f)**




¹H NMR (DMSO-*d*₆): **5-[(4-Chlorophenyl)sulfanyl]phthalimide (8g)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Chlorophenyl)sulfanyl]phthalimide (8g)**




¹H NMR (DMSO-*d*₆): **5-[(4-Bromophenyl)sulfanyl]phthalimide (8h)**




¹³C NMR (DMSO-*d*₆): **5-[(4-Bromophenyl)sulfanyl]phthalimide (8h)**




¹H NMR (DMSO-*d*₆): 5-[(2-Phenylethyl)sulfanyl]phthalimide (8i)




¹³C NMR (DMSO-*d*₆): 5-[(2-Phenylethyl)sulfanyl]phthalimide (8i)




¹H NMR (DMSO-*d*₆): **5-(Cyclohexylsulfanyl)phthalimide (8j)**




¹³C NMR (CDCl₃): **5-(Cyclohexylsulfanyl)phthalimide (8j)**



¹H NMR (CDCl₃): **5-[(3-Methylbutyl)sulfanyl]phthalimide (8k)**



¹³C NMR (CDCl₃): **5-[(3-Methylbutyl)sulfanyl]phthalimide (8k)**




6.7.4 HPLC traces of the 5-sulfanylpthalimide analogues (**8a–k**)

The degree of purity for each of the synthesized 5-sulfanylpthalimide analogues (**8a–k**) were estimated with HPLC analyses, which were carried out with an Agilent 1100 HPLC system equipped with a quaternary pump and an Agilent 1100 series diode array detector. Milli-Q water (Millipore) and HPLC grade acetonitrile (Merck) were used for the chromatography. A Venusil XBP C18 column (4.60 × 150 mm, 5 μm) was used and the mobile phase consisted initially of 30% acetonitrile and 70% MilliQ water at a flow rate of 1 mL/min. At the start of each HPLC run a solvent gradient program was initiated by linearly increasing the composition of the acetonitrile in the mobile phase to 85% acetonitrile over a period of 5 min. Each HPLC run lasted 15 min and a time period of 5 min was allowed for equilibration between runs. A volume of 20 μL of solutions of the test compounds in acetonitrile (1 mM) was injected into the HPLC system and the eluent was monitored at wavelengths of 254 nm.


5-(Benzylsulfanyl)phthalimide (**8a**)

Purity (HPLC): 100%.




5-[(4-Chlorobenzyl)sulfanyl]phthalimide (8b)

Purity (HPLC): 99%




5-[(4-Bromobenzyl)sulfanyl]phthalimide (8c)

Purity (HPLC): 100%




5-[(4-Fluorobenzyl)sulfanyl]phthalimide (8d)

Purity (HPLC): 96%




5-[(4-Methoxybenzyl)sulfanyl]phthalimide (8e)

Purity (HPLC): 100%




5-(Phenylsulfanyl)phthalimide (8f)

Purity (HPLC): 100%




5-[(4-Chlorophenyl)sulfanyl]phthalimide (8g)

Purity (HPLC): 100%




5-[(4-Bromophenyl)sulfanyl]phthalimide (8h)

Purity (HPLC): 99%




5-[(2-Phenylethyl)sulfanyl]phthalimide (8i)

Purity (HPLC): 100%



5-(Cyclohexylsulfanyl)phthalimide(8j)

Purity (HPLC): 96%



5-[(3-Methylbutyl)sulfanyl]phthalimide (8k)

Purity (HPLC): 100%

